

Oxytocin

Oxytocin — Neuropeptide Hormone

BONDING SEXUAL FUNCTION ORGASM SOCIAL ANTI-ANXIETY

WHAT IS IT?

Oxytocin is a 9-amino-acid neuropeptide produced in the hypothalamus and released by the posterior pituitary. Known as the 'bonding hormone' or 'love hormone,' it is released during physical touch, orgasm, childbirth, and breastfeeding. In performance and wellness contexts, exogenous oxytocin is used to enhance sexual experience, improve emotional connection, reduce social anxiety, accelerate orgasm, and support post-workout recovery via anti-inflammatory pathways.

DOSING AT A GLANCE

SC injection	Standard dose	Timing	Frequency
10–40 IU	20 IU	20–30 min prior	As needed

KEY BENEFITS

- Intensifies orgasm — facilitates and amplifies climax in men and women
- Increases emotional bonding and intimacy between partners
- Reduces social anxiety — improves confidence in social situations
- Enhances sexual arousal and physical sensitivity
- Anti-inflammatory — reduces cortisol and systemic inflammation post-exercise
- Accelerates wound healing (promotes myofibroblast activity)
- Improves trust, prosocial behavior, and relationship satisfaction
- Reduces compulsive eating and addictive behaviors (emerging data)
- Potential role in PTSD and social anxiety treatment

HOW IT WORKS

Oxytocin binds oxytocin receptors (OTR) throughout the brain and body. In the limbic system, it activates reward circuits (dopamine release) and reduces amygdala reactivity to threatening stimuli — explaining its anxiolytic and bonding effects. In sexual function, it amplifies genital sensation and facilitates spinal ejaculatory/orgasm reflexes. Peripherally, it reduces inflammation via NF-κB pathway inhibition.

PROTOCOLS

- **Sexual enhancement:** 20–40 IU SC injection 20–30 min before intimacy.
- **Social anxiety:** 20 IU SC 30–45 min before social event.
- **Post-workout recovery:** 20 IU SC post-training for anti-inflammatory effect.
- **Orgasm enhancement:** 20 IU for women 30 min pre-activity; pair with PT-141 for enhanced desire + intensity.
- **Note:** Take with a partner — pair-bonding effects work bidirectionally.

SIDE EFFECTS

- Mild injection site irritation (transient)
- Mild nausea at higher doses (>40 IU)
- Possible oxytocin 'crash' — emotional low after peak (dose dependent)
- In social contexts: can amplify negative emotions as well as positive (caution in contentious situations)
- Headache (rare)
- Uterine contractions at high doses — contraindicated in pregnancy

STORAGE

Reconstituted solution: refrigerate 2–8°C, use within 30 days. Lyophilized: -20°C. Protect from light.

■ **Contraindicated during pregnancy — causes uterine contractions. Effects are bidirectional on emotion — may amplify negative social experiences as well as positive ones.**

Educational use only. Not medical advice. Consult a qualified physician before using any compound. forgedbyfreedom.com