

CJC-1295 + Ipamorelin

CJC-1295 / Ipamorelin Stack

GH RELEASE

FAT LOSS

MUSCLE GAIN

SLEEP

ANTI-AGING

WHAT IS IT?

CJC-1295 (a GHRH analogue) + Ipamorelin (a GHRP/ghrelin mimetic) is the most popular GH secretagogue stack in use today. Together they produce synergistic GH pulse amplification that closely mimics the body's natural pulsatile GH release — without the cortisol and prolactin spikes of older GHRPs.

DOSING AT A GLANCE

CJC-1295 (no DAC)	Ipamorelin	Timing (optimal)	Cycle Length
100 mcg	100–200 mcg	Bedtime	3–6 months

KEY BENEFITS

- Significantly increased GH and IGF-1 levels
- Accelerated fat loss — especially visceral and subcutaneous
- Improved muscle recovery, body composition, and lean mass over time
- Dramatically improved deep sleep quality (GH-wave amplification during sleep)
- Anti-aging effects: improved skin elasticity, collagen production
- Increased bone density over long-term use
- Joint and connective tissue support via IGF-1 upregulation
- Improved metabolic rate and insulin sensitivity
- Very clean — Ipamorelin does not spike cortisol or prolactin like GHRP-6/Hexarelin

HOW IT WORKS

CJC-1295 (DAC version) has a half-life of ~8 days, providing sustained GHRH signal. Ipamorelin acts on ghrelin receptors to create GH pulses. Together, they simultaneously amplify GH pulse amplitude (CJC) and frequency (Ipamorelin), mimicking youth-level GH secretion without suppressing the natural HPG axis.

PROTOCOLS

- **Standard:** CJC-1295 100 mcg + Ipamorelin 100–200 mcg SC, before bed (fasted 2 hrs).
- **CJC-DAC version:** 1–2 mg CJC-DAC 1–2x/week + daily Ipamorelin 100–200 mcg.

- **2x Daily:** Add a morning dose pre-workout for accelerated body recomp.
- **Anti-Aging:** 5 days on / 2 days off cycling prevents receptor desensitization.

SIDE EFFECTS

- Water retention (mild) — especially first few weeks
- Tingling or numbness in hands (carpal tunnel-like, dose dependent)
- Temporary insulin sensitivity reduction (monitor glucose)
- Mild fatigue within 30–60 min post-injection (expected GH effect)
- Ipamorelin: slight hunger increase (much less than GHRP-6)

STORAGE

Lyophilized: room temp. Reconstituted in BW: refrigerate, use within 30 days.

■ ***Long-term use may cause pituitary desensitization. Cycle with breaks. Monitor IGF-1 levels.***

This document is for educational purposes only. Not medical advice. Consult a qualified physician before using any peptide or performance compound. forGEDbyfreedom.com