

IGF-1 LR3

Insulin-Like Growth Factor 1 Long Arg3

MUSCLE GROWTH

FAT LOSS

NUTRIENT
PARTITIONING

ADVANCED

HYPOGLYCEMIA
RISK

WHAT IS IT?

IGF-1 LR3 is a modified form of Insulin-Like Growth Factor 1 with a 13-amino-acid extension that reduces binding to IGF-BPs — increasing its half-life from minutes (native IGF-1) to approximately 20–30 hours. It promotes muscle cell hyperplasia (new cell formation), not just hypertrophy — making it uniquely anabolic.

DOSING AT A GLANCE

Daily Dose	Timing	Frequency	Cycle Max
20–100 mcg	Post-workout	Daily or ED	4–6 weeks

KEY BENEFITS

- Muscle hyperplasia — stimulates satellite cell proliferation and new muscle fiber formation
- Powerful nutrient partitioning — shuttles nutrients into muscle, away from fat
- Significant fat loss via IGF-1's lipolytic action
- Enhanced recovery and protein synthesis
- Synergistic with androgens and GH peptides
- Improved glucose uptake into muscle tissue
- Cognitive benefits — neuroprotective and neurogenic effects

HOW IT WORKS

IGF-1 LR3 binds IGF-1 receptors (IGF-1R) on muscle cells, activating PI3K/Akt/mTOR pathways for protein synthesis and satellite cell activation. The LR3 modification reduces IGF-BP binding by ~1000x, dramatically extending activity. It acts both systemically and locally at injection sites.

PROTOCOLS

- **Beginner:** 20–40 mcg IM post-workout, target worked muscles.
- **Intermediate:** 60–80 mcg post-workout, rotate injection sites.
- **Advanced:** 80–100 mcg/day split into 2 injections.

- **Cycle:** 4–6 weeks max — receptor downregulation occurs with extended use.
- **Stack:** IGF-1 LR3 + GH peptides + androgens for elite body recomp.

SIDE EFFECTS

- Hypoglycemia — most common risk; eat within 15–20 min of injection
- Jaw and organ growth concerns at sustained supraphysiologic doses
- Theoretical long-term cancer promotion risk (IGF-1 is mitogenic)
- Muscle cramps (electrolyte shift)
- Tingling, numbness (dose dependent)

STORAGE

Lyophilized: refrigerate. Reconstituted in AA: refrigerate, use within 21 days.

■ **Advanced compound. Hypoglycemia is a real risk — always have fast carbs on hand post-injection.**

This document is for educational purposes only. Not medical advice. Consult a qualified physician before using any peptide or performance compound. forgedbyfreedom.com