

PT-141

Bremelanotide (PT-141)

LIBIDO	SEXUAL FUNCTION	ERECTION	FEMALE DESIRE	FDA APPROVED (VYLEESI)
--------	-----------------	----------	---------------	------------------------

WHAT IS IT?

PT-141 (Bremelanotide) is a synthetic melanocortin peptide that activates MC3R and MC4R receptors in the brain to drive sexual desire and arousal at the neurological level — independent of the vascular mechanism targeted by ED drugs like Viagra. FDA approved (as Vyleesi) for hypoactive sexual desire disorder (HSDD) in women. Highly effective in men and women with low libido regardless of cause.

DOSING AT A GLANCE

Men (SC)	Women (FDA dose)	Timing	Frequency
0.5–2 mg	1.75 mg	1–2 hrs prior	As needed

KEY BENEFITS

- Dramatically increases sexual desire, motivation, and libido in men and women
- Promotes erections in men via central nervous system pathways
- Effective when PDE5 inhibitors fail or are insufficient alone
- Enhances arousal, sensitivity, and sexual satisfaction in women
- Works centrally — addresses psychological/neurological drivers of low libido
- Shown effective for HSDD, post-menopause low desire, hormone-related low libido
- Spontaneous — no need to time use around sexual activity (unlike PDE5i)

HOW IT WORKS

PT-141 acts on melanocortin receptors (MC3R, MC4R) in the hypothalamus, directly activating the brain's sexual arousal circuits. Unlike PDE5 inhibitors which work peripherally on blood vessels, PT-141 works at the CNS level — increasing desire and arousal rather than just facilitating erection mechanics.

PROTOCOLS

- **Men:** Start at 0.5–1 mg SC 1–2 hrs before activity. Increase to 2 mg if needed.
- **Women:** 1.75 mg SC 45 min before activity (FDA-approved Vyleesi dose).

- **Stack (Men):** PT-141 + low-dose Tadalafil (5 mg daily) for central + peripheral coverage.
- **Max use:** No more than 8 uses per month to avoid receptor desensitization.

SIDE EFFECTS

- Nausea (most common — take with anti-nausea OTC if needed)
- Flushing and transient blood pressure changes
- Spontaneous erections in men (dose dependent)
- Headache
- Hyperpigmentation with frequent long-term use (melanocortin effect)

STORAGE

Refrigerate reconstituted solution. Use within 30 days. Vyleesi autoinjector: room temp.

■ ***May cause transient blood pressure elevation. Use with caution if cardiovascular disease present.***

This document is for educational purposes only. Not medical advice. Consult a qualified physician before using any peptide or performance compound. forgedbyfreedom.com