

# Elys and Lamin B/C depletions in spermatogonia and spermatocytes

## 1. RNA-seq in larval testes. Differential expression and TPM comparison.

### 1.1. Comparison of all genes

We used a BDGP5.78 annotation for *Drosophila* genes:


chr	start	end	strand	id	gene_name	tss
chr2L	7529	9484	+	FBgn0031208	CG11023	7529
chr2L	9839	21376	-	FBgn0002121	l(2)gl	21376
chr2L	21823	25155	-	FBgn0031209	Ir21a	25155
chr2L	21952	24237	+	FBgn0263584	CR43609	21952
chr2L	25402	65404	-	FBgn0051973	Cda5	65404
chr2L	65999	66242	+	FBgn0266878	CR45339	65999

Using [salmon](#) we obtained read counts per gene and normalized TPM values:

chr	start	end	strand	gene_name	WT	ElysKD	LamKD
chr2R	18024473	18060339	+	a	6.7153633	5.6267533	5.8948233
chr3R	12632936	12655771	-	abd-A	7.6534000	9.6282000	9.5642133
chr3R	12752932	12797958	-	Abd-B	0.4616433	0.6147677	0.6040280
chr3L	16608966	16640982	-	Abl	9.4840567	12.5862000	10.4057300
chr2L	10973443	10975293	-	abo	27.6684000	41.1312667	36.6294333
chrX	264064	264980	+	ac	0.2590723	0.0034933	0.0034933


The main question that we investigate is whether the expression of genes on X-chromosome is higher than of those on autosomes in spermatocytes? And do Elys and/or Lamins knockdowns alter this discordance?

### All genes comparison between X and A




Median ratio of WT X to WT A is equal to 0.72

Let's see the ratios of TPMs in case of Elys and Lams knockdowns.  $p.v_1$  represents one-tailed comparison of each sample's medians with 1 via wilcoxon test,  $p.v_2$  represents two-tailed two-sample comparison using the same test





## 1.2. Comparisons of genes that have at least one TPM in all treatments

### Genes w/ TPM > 1 comparison between X and A




Median ratio of WT X to WT A for these genes is equal to 0.64






Here are similar boxplots for differentially expressed genes (via DESeq2 and TPM ratio > 2:




Dot plots for TPMs of differentially expressed genes (TPM ratio > 2):



## 1.4 Spermatocyte-specific genes

Boxplot with TPMs for SpC-specific genes (p-values are from two-sample two-sided wilcoxon test):


### SpC-specific genes comparison between X and A



Median ratios of WT X to WT A for SpC-specific genes is equal to 0.56

Ratios (p.v<sub>1</sub> represents testing if medians are lower than 1):

### Elys KD



### Lam KD

