

بسم الله الرحمن الرحيم

برنامه‌ریزی نیمه‌معین برای طراحی الگوریتم‌های تقریبی

جلسه ششم: دوگانی

قضیه نهایی دوگانی (برای برنامه نویسی نیمه معین)

$$\begin{array}{ll}\text{maximize} & C \bullet X \\ \text{subject to} & A_i \bullet X = b_i, \quad i = 1, 2, \dots, m \\ & X \succeq 0.\end{array}$$

4.1.1 Theorem. *If the semidefinite program (4.1) is feasible and has a finite value γ , and if there is a positive definite matrix \tilde{X} such that $A(\tilde{X}) = \mathbf{b}$, then the dual program*

$$\begin{array}{ll}\text{minimize} & \mathbf{b}^T \mathbf{y} \\ \text{subject to} & \sum_{i=1}^m y_i A_i - C \succeq 0\end{array} \tag{4.2}$$

is feasible and has finite value $\beta = \gamma$.

4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda\mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda\mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

مثال: مجموعه $K = \{0\}$ کنج محدب بسته است

4.2.2 Lemma. *The set $\text{PSD}_n \subseteq \text{SYM}_n$ of positive semidefinite matrices is a closed convex cone.*

4.2.2 Lemma. *The set $\text{PSD}_n \subseteq \text{SYM}_n$ of positive semidefinite matrices is a closed convex cone.*

• الف) بسته

• معادلا: مکملش باز است

• $\tilde{\mathbf{x}}^T M \tilde{\mathbf{x}} < 0$ هست که

• $M + \epsilon B$:

• $\max_B \tilde{\mathbf{x}}^T B \tilde{\mathbf{x}}$

4.2.2 Lemma. *The set $\text{PSD}_n \subseteq \text{SYM}_n$ of positive semidefinite matrices is a closed convex cone.*

• الف) بسته

• معادلا: مکملش باز است

• $\tilde{\mathbf{x}}^T M \tilde{\mathbf{x}} < 0$ هست که

• $M + \epsilon B$:

• $\max_B \tilde{\mathbf{x}}^T B \tilde{\mathbf{x}}$

• ب) λM :

• $\mathbf{x}^T \lambda M \mathbf{x} = \lambda \mathbf{x}^T M \mathbf{x} \geq 0$

4.2.2 Lemma. *The set $\text{PSD}_n \subseteq \text{SYM}_n$ of positive semidefinite matrices is a closed convex cone.*

• الف) بسته

• معادلا: مکملش باز است

• $\tilde{\mathbf{x}}^T M \tilde{\mathbf{x}} < 0$ هست که

• $M + \epsilon B$:

• $\max_B \tilde{\mathbf{x}}^T B \tilde{\mathbf{x}}$

• ب) λM :

• $\mathbf{x}^T \lambda M \mathbf{x} = \lambda \mathbf{x}^T M \mathbf{x} \geq 0$

• ج) $M + N$

• $\mathbf{x}^T (M + N) \mathbf{x} = \mathbf{x}^T M \mathbf{x} + \mathbf{x}^T N \mathbf{x} \geq 0$

4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda \mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

مثال: اگر K و L کنج محدب بسته باشند، آنگاه

$$K \oplus L := \{(\mathbf{x}, \mathbf{y}) \in V \oplus W : \mathbf{x} \in K, \mathbf{y} \in L\}$$

کنج محدب بسته است.

4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda\mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

مثال: اگر K و L کنج محدب بسته باشند، آنگاه

$$K \oplus L := \{(\mathbf{x}, \mathbf{y}) \in V \oplus W : \mathbf{x} \in K, \mathbf{y} \in L\}$$

کنج محدب بسته است.

• الف) بسته

• ب) λM :

• ج) $M + N$

4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda\mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

$$\nabla_n = \{(\mathbf{x}, r) \in \mathbb{R}^{n-1} \times \mathbb{R} : \|\mathbf{x}\| \leq r\}$$


(کنج بستنی) کنج محدب بسته است

4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda\mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

$$\nabla_n = \{(\mathbf{x}, r) \in \mathbb{R}^{n-1} \times \mathbb{R} : \|\mathbf{x}\| \leq r\}$$

(کنج بستنی) کنج محدب بسته است




4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda\mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

$$\nabla_n = \{(\mathbf{x}, r) \in \mathbb{R}^{n-1} \times \mathbb{R} : \|\mathbf{x}\| \leq r\}$$

(کنج بستنی) کنج محدب بسته است

• الف) بسته




4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda\mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

$$\nabla_n = \{(\mathbf{x}, r) \in \mathbb{R}^{n-1} \times \mathbb{R} : \|\mathbf{x}\| \leq r\}$$

(کنج بستنی) کنج محدب بسته است



• الف) بسته


• ب) ضرب

4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda \mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

$$\nabla_n = \{(\mathbf{x}, r) \in \mathbb{R}^{n-1} \times \mathbb{R} : \|\mathbf{x}\| \leq r\}$$

(کنج بستنی) کنج محدب بسته است




- الف) بسته
- ب) ضرب
- ج) جمع

4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda \mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

$$\nabla_n = \{(\mathbf{x}, r) \in \mathbb{R}^{n-1} \times \mathbb{R} : \|\mathbf{x}\| \leq r\}$$

(کنج بستنی) کنج محدب بسته است



- الف) بسته
- ب) ضرب
- ج) جمع

$$(x, r) + (y, s) = (x + y, r + s) \quad \bullet$$

$$\|x + y\| \leq \|x\| + \|y\|$$

4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda\mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

$$\triangleleft = \{(x, y, z) \in \mathbb{R}^3 : x \geq 0, y \geq 0, xy \geq z^2\}$$


(کنج بستنی وازگون) کنج محدب بسته است

4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda \mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

$$\triangleleft = \{(x, y, z) \in \mathbb{R}^3 : x \geq 0, y \geq 0, xy \geq z^2\}$$

(کنج بستنی واژگون) کنج محدب بسته است




4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda \mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

$$\triangleleft = \{(x, y, z) \in \mathbb{R}^3 : x \geq 0, y \geq 0, xy \geq z^2\}$$

(کنج بستنی وازگون) کنج محدب بسته است




• الف) بسته

4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda \mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

$$\triangleleft = \{(x, y, z) \in \mathbb{R}^3 : x \geq 0, y \geq 0, xy \geq z^2\}$$

(کنج بستنی وازگون) کنج محدب بسته است



• الف) بسته


• ب) ضرب

4.2.1 Definition. Let $K \subseteq V$ be a nonempty closed set.¹ K is called a *closed convex cone* if the following two conditions hold.

- (i) For all $\mathbf{x} \in K$ and all nonnegative real numbers λ , we have $\lambda \mathbf{x} \in K$.
- (ii) For all $\mathbf{x}, \mathbf{y} \in K$, we have $\mathbf{x} + \mathbf{y} \in K$.

$$\langle \rangle = \{(x, y, z) \in \mathbb{R}^3 : x \geq 0, y \geq 0, xy \geq z^2\}$$

(کنج بستنی وازگون) کنج محدب بسته است



- الف) بسته
- ب) ضرب
- ج) جمع

مثبت نیمه معین

$$\begin{pmatrix} x & z \\ z & y \end{pmatrix}$$

معادلا •