

بسم الله الرحمن الرحيم

نظريه علوم کامپیوتر

نظريه علوم کامپیوتر - بهار ۱۴۰۰-۱۴۰۱ - جلسه نهم: خودتولیدکنندگی

Theory of computation - 002 - S09 - self-reproducibility

Recall

Recall


8

- * Self-Reproducing TM

Recall

8


* Self-Reproducing TM



Recall


8

* Self-Reproducing TM




Recall

* Self-Reproducing TM




Recall

* Self-Reproducing TM



Recall


* Self-Reproducing TM



Recall

8


- * Self-Reproducing TM
- * The Recursion Theorem: TM T working on $\langle w, A \rangle \rightarrow$ TM R
 - * that has access to $\langle R \rangle$,
 - * behaves like T on $\langle w, R \rangle$



Recall

8

- * Self-Reproducing TM
- * The Recursion Theorem: TM T working on $\langle w, A \rangle \rightarrow$ TM R
 - * that has access to $\langle R \rangle$,
 - * behaves like T on $\langle w, R \rangle$




Theorem: For any TM T there is a TM R where for all w
 R on input w operates in the same way as T on input $\langle w, R \rangle$.


Recall

8

- * Self-Reproducing TM
- * The Recursion Theorem: TM T working on $\langle w, A \rangle \rightarrow$ TM R
 - * that has access to $\langle R \rangle$,
 - * behaves like T on $\langle w, R \rangle$




Theorem: For any TM T there is a TM R where for all w R on input w operates in the same way as T on input $\langle w, R \rangle$.




Recall

8

- * Self-Reproducing TM
- * The Recursion Theorem: TM T working on $\langle w, A \rangle \rightarrow$ TM R
 - * that has access to $\langle R \rangle$,
 - * behaves like T on $\langle w, R \rangle$




Theorem: For any TM T there is a TM R where for all w R on input w operates in the same way as T on input $\langle w, R \rangle$.




Recall

8

- * Self-Reproducing TM
- * The Recursion Theorem: TM T working on $\langle w, A \rangle \rightarrow$ TM R
 - * that has access to $\langle R \rangle$,
 - * behaves like T on $\langle w, R \rangle$




Theorem: For any TM T there is a TM R where for all w R on input w operates in the same way as T on input $\langle w, R \rangle$.




Recall

8

- * Self-Reproducing TM
- * The Recursion Theorem: TM T working on $\langle w, A \rangle \rightarrow$ TM R
 - * that has access to $\langle R \rangle$,
 - * behaves like T on $\langle w, R \rangle$




Theorem: For any TM T there is a TM R where for all w R on input w operates in the same way as T on input $\langle w, R \rangle$.




Recall

8

- * Self-Reproducing TM
- * The Recursion Theorem: TM T working on $\langle w, A \rangle \rightarrow$ TM R
 - * that has access to $\langle R \rangle$,
 - * behaves like T on $\langle w, R \rangle$




Theorem: For any TM T there is a TM R where for all w R on input w operates in the same way as T on input $\langle w, R \rangle$.




Recall

8

- * Self-Reproducing TM
- * The Recursion Theorem: TM T working on $\langle w, A \rangle \rightarrow$ TM R
 - * that has access to $\langle R \rangle$,
 - * behaves like T on $\langle w, R \rangle$



Theorem: For any TM T there is a TM R where for all w R on input w operates in the same way as T on input $\langle w, R \rangle$.



Ex 2: Fixed-point Theorem

Ex 2: Fixed-point Theorem

Theorem: For any computable function $f: \Sigma^* \rightarrow \Sigma^*$,
there is a TM R such that $L(R) = L(S)$ where $f(\langle R \rangle) = \langle S \rangle$.

Ex 2: Fixed-point Theorem

Theorem: For any computable function $f: \Sigma^* \rightarrow \Sigma^*$,
there is a TM R such that $L(R) = L(S)$ where $f(\langle R \rangle) = \langle S \rangle$.

In other words, consider f to be a program transformation function.
Then for some program R , its behavior is unchanged by f .

Ex 2: Fixed-point Theorem

Theorem: For any computable function $f: \Sigma^* \rightarrow \Sigma^*$,
there is a TM R such that $L(R) = L(S)$ where $f(\langle R \rangle) = \langle S \rangle$.

In other words, consider f to be a program transformation function.
Then for some program R , its behavior is unchanged by f .

Proof: Let R be the following TM.

Ex 2: Fixed-point Theorem

Theorem: For any computable function $f: \Sigma^* \rightarrow \Sigma^*$,
there is a TM R such that $L(R) = L(S)$ where $f(\langle R \rangle) = \langle S \rangle$.

In other words, consider f to be a program transformation function.
Then for some program R , its behavior is unchanged by f .

Proof: Let R be the following TM.

$R =$ "On input w

Ex 2: Fixed-point Theorem

Theorem: For any computable function $f: \Sigma^* \rightarrow \Sigma^*$,
there is a TM R such that $L(R) = L(S)$ where $f(\langle R \rangle) = \langle S \rangle$.

In other words, consider f to be a program transformation function.
Then for some program R , its behavior is unchanged by f .

Proof: Let R be the following TM.

R = "On input w

1. Get own description $\langle R \rangle$.

Ex 2: Fixed-point Theorem

Theorem: For any computable function $f: \Sigma^* \rightarrow \Sigma^*$,
there is a TM R such that $L(R) = L(S)$ where $f(\langle R \rangle) = \langle S \rangle$.

In other words, consider f to be a program transformation function.
Then for some program R , its behavior is unchanged by f .

Proof: Let R be the following TM.

$R =$ “On input w

1. Get own description $\langle R \rangle$.
2. Compute $f(\langle R \rangle)$ and call the result $\langle S \rangle$.

Ex 2: Fixed-point Theorem

Theorem: For any computable function $f: \Sigma^* \rightarrow \Sigma^*$,
there is a TM R such that $L(R) = L(S)$ where $f(\langle R \rangle) = \langle S \rangle$.

In other words, consider f to be a program transformation function.
Then for some program R , its behavior is unchanged by f .

Proof: Let R be the following TM.

$R =$ "On input w

1. Get own description $\langle R \rangle$.
2. Compute $f(\langle R \rangle)$ and call the result $\langle S \rangle$.
3. Simulate S on w ."

Ex 2: Fixed-point Theorem

Theorem: For any computable function $f: \Sigma^* \rightarrow \Sigma^*$,
there is a TM R such that $L(R) = L(S)$ where $f(\langle R \rangle) = \langle S \rangle$.

In other words, consider f to be a program transformation function.
Then for some program R , its behavior is unchanged by f .

Proof: Let R be the following TM.

R = "On input w

1. Get own description $\langle R \rangle$.
2. Compute $f(\langle R \rangle)$ and call the result $\langle S \rangle$.
3. Simulate S on w ."

Ex 3: *MIN*TM is T-unrecognizable

Ex 3: *MIN*TM is T-unrecognizable

10

Defn: M is a minimal TM if $|\langle M' \rangle| < |\langle M \rangle| \rightarrow L(M') \neq L(M)$.

Ex 3: *MIN*TM is T-unrecognizable

10

Defn: M is a minimal TM if $|\langle M' \rangle| < |\langle M \rangle| \rightarrow L(M') \neq L(M)$.

Thus, a minimal TM has the shortest description among all equivalent TMs.

Ex 3: *MIN*TM is T-unrecognizable

10

Defn: M is a minimal TM if $|\langle M' \rangle| < |\langle M \rangle| \rightarrow L(M') \neq L(M)$.

Thus, a minimal TM has the shortest description among all equivalent TMs.

Let $MIN_{TM} = \{ \langle M \rangle \mid M \text{ is a minimal TM} \}$.

Theorem: MIN_{TM} is T-unrecognizable.

Ex 3: MIN_{TM} is T-unrecognizable

10

Defn: M is a minimal TM if $|\langle M' \rangle| < |\langle M \rangle| \rightarrow L(M') \neq L(M)$.

Thus, a minimal TM has the shortest description among all equivalent TMs.

Let $MIN_{TM} = \{ \langle M \rangle \mid M \text{ is a minimal TM} \}$.

Theorem: MIN_{TM} is T-unrecognizable.

Proof by contradiction: Assume some TM E enumerates MIN_{TM} .

Ex 3: MIN_{TM} is T-unrecognizable

10

Defn: M is a minimal TM if $|\langle M' \rangle| < |\langle M \rangle| \rightarrow L(M') \neq L(M)$.

Thus, a minimal TM has the shortest description among all equivalent TMs.

Let $MIN_{TM} = \{ \langle M \rangle \mid M \text{ is a minimal TM} \}$.

Theorem: MIN_{TM} is T-unrecognizable.

Proof by contradiction: Assume some TM E enumerates MIN_{TM} .

Consider the following TM R :

Ex 3: MIN_{TM} is T-unrecognizable

10

Defn: M is a minimal TM if $|\langle M' \rangle| < |\langle M \rangle| \rightarrow L(M') \neq L(M)$.

Thus, a minimal TM has the shortest description among all equivalent TMs.

Let $MIN_{TM} = \{ \langle M \rangle \mid M \text{ is a minimal TM} \}$.

Theorem: MIN_{TM} is T-unrecognizable.

Proof by contradiction: Assume some TM E enumerates MIN_{TM} .

Consider the following TM R :

$R =$ "On input w

Ex 3: *MIN*TM is T-unrecognizable

10

Defn: M is a minimal TM if $|\langle M' \rangle| < |\langle M \rangle| \rightarrow L(M') \neq L(M)$.

Thus, a minimal TM has the shortest description among all equivalent TMs.

Let $MIN_{TM} = \{ \langle M \rangle \mid M \text{ is a minimal TM} \}$.

Theorem: MIN_{TM} is T-unrecognizable.

Proof by contradiction: Assume some TM E enumerates MIN_{TM} .

Consider the following TM R :

$R =$ "On input w

1. Get own description $\langle R \rangle$.

Ex 3: *MIN*TM is T-unrecognizable

10

Defn: M is a minimal TM if $|\langle M' \rangle| < |\langle M \rangle| \rightarrow L(M') \neq L(M)$.

Thus, a minimal TM has the shortest description among all equivalent TMs.

Let $MIN_{TM} = \{ \langle M \rangle \mid M \text{ is a minimal TM} \}$.

Theorem: MIN_{TM} is T-unrecognizable.

Proof by contradiction: Assume some TM E enumerates MIN_{TM} .

Consider the following TM R :

$R =$ "On input w

1. Get own description $\langle R \rangle$.
2. Run enumerator E until some TM B appears, where $|\langle R \rangle| < |\langle B \rangle|$.

Ex 3: *MIN*TM is T-unrecognizable

10

Defn: M is a minimal TM if $|\langle M' \rangle| < |\langle M \rangle| \rightarrow L(M') \neq L(M)$.

Thus, a minimal TM has the shortest description among all equivalent TMs.

Let $MIN_{TM} = \{ \langle M \rangle \mid M \text{ is a minimal TM} \}$.

Theorem: MIN_{TM} is T-unrecognizable.

Proof by contradiction: Assume some TM E enumerates MIN_{TM} .

Consider the following TM R :

$R =$ "On input w

1. Get own description $\langle R \rangle$.
2. Run enumerator E until some TM B appears, where $|\langle R \rangle| < |\langle B \rangle|$.
3. Simulate B on w ."

Ex 3: *MIN*TM is T-unrecognizable

10

Defn: M is a minimal TM if $|\langle M' \rangle| < |\langle M \rangle| \rightarrow L(M') \neq L(M)$.

Thus, a minimal TM has the shortest description among all equivalent TMs.

Let $MINTM = \{ \langle M \rangle \mid M \text{ is a minimal TM} \}$.

Theorem: *MIN*TM is T-unrecognizable.

Proof by contradiction: Assume some TM E enumerates *MIN*TM .

Consider the following TM R :

$R =$ “On input w

1. Get own description $\langle R \rangle$.
2. Run enumerator E until some TM B appears, where $|\langle R \rangle| < |\langle B \rangle|$.
3. Simulate B on w .”

Thus $L(R) = L(B)$ and $|\langle R \rangle| < |\langle B \rangle|$ so B isn't minimal, but $\langle B \rangle \in L(E)$, contradiction.

Ex 3: *MIN*TM is T-unrecognizable

10

Defn: M is a minimal TM if $|\langle M' \rangle| < |\langle M \rangle| \rightarrow L(M') \neq L(M)$.

Thus, a minimal TM has the shortest description among all equivalent TMs.

Let $MINTM = \{ \langle M \rangle \mid M \text{ is a minimal TM} \}$.

Theorem: *MIN*TM is T-unrecognizable.

Proof by contradiction: Assume some TM E enumerates *MIN*TM .

Consider the following TM R :

R = “On input w

1. Get own description $\langle R \rangle$.
2. Run enumerator E until some TM B appears, where $|\langle R \rangle| < |\langle B \rangle|$.
3. Simulate B on w .”

Thus $L(R) = L(B)$ and $|\langle R \rangle| < |\langle B \rangle|$ so B isn't minimal, but $\langle B \rangle \in L(E)$, contradiction.

Ex 3: *MIN*TM is T-unrecognizable

10

Defn: M is a minimal TM if $|\langle M' \rangle| < |\langle M \rangle| \rightarrow L(M') \neq L(M)$.

Thus, a minimal TM has the shortest description among all equivalent TMs.

Let $MIN_{TM} = \{ \langle M \rangle \mid M \text{ is a minimal TM} \}$.

Theorem: MIN_{TM} is T-unrecognizable.

Proof by contradiction: Assume some TM E enumerates

Consider the following TM R :

$R =$ "On input w

1. Get own description $\langle R \rangle$.
2. Run enumerator E until some TM B appears, where $|\langle R \rangle| < |\langle B \rangle|$.
3. Simulate B on w ."

Thus $L(R) = L(B)$ and $|\langle R \rangle| < |\langle B \rangle|$ so B isn't minimal, but $\langle B \rangle \in L(E)$, contradiction.

Check-in 11.3

Let A be an infinite subset of MIN_{TM} .

Is it possible that A is T-recognizable?

- (a) Yes.
- (b) No.

Check-in 11.3

Other applications

Other applications

11

1. Computer viruses.

Other applications

11

1. Computer viruses.
2. A true but unprovable mathematical statement due to Kurt Gödel:
“This statement is unprovable.”

Other applications

11

1. Computer viruses.
2. A true but unprovable mathematical statement due to Kurt Gödel:
“This statement is unprovable.”

Other applications

11

1. Computer viruses.
2. A true but unprovable mathematical statement due to Kurt Gödel:
“This statement is unprovable.”

Intro to Mathematical Logic

12

Goal: A mathematical study of mathematical reasoning itself.

Formally defines the language of mathematics, mathematical truth, and provability.

Intro to Mathematical Logic

12

Goal: A mathematical study of mathematical reasoning itself.

Formally defines the language of mathematics, mathematical truth, and provability.

Gödel's First Incompleteness Theorem:

In any reasonable formal system, some true statements are not provable.

Intro to Mathematical Logic

12

Goal: A mathematical study of mathematical reasoning itself.

Formally defines the language of mathematics, mathematical truth, and provability.

Gödel's First Incompleteness Theorem:

In any reasonable formal system, some true statements are not provable.

Proof: We use two properties of formal proofs:

Intro to Mathematical Logic

12

Goal: A mathematical study of mathematical reasoning itself.

Formally defines the language of mathematics, mathematical truth, and provability.

Gödel's First Incompleteness Theorem:

In any reasonable formal system, some true statements are not provable.

Proof: We use two properties of formal proofs:

1) Soundness: If ϕ has a proof π then ϕ is true.

Intro to Mathematical Logic

12

Goal: A mathematical study of mathematical reasoning itself.

Formally defines the language of mathematics, mathematical truth, and provability.

Gödel's First Incompleteness Theorem:

In any reasonable formal system, some true statements are not provable.

Proof: We use two properties of formal proofs:

- 1) Soundness: If ϕ has a proof π then ϕ is true.
- 2) Checkability: The language $\{ \langle \pi, \phi \rangle \mid \pi \text{ is a proof of statement } \phi \}$ is decidable.

Intro to Mathematical Logic

12

Goal: A mathematical study of mathematical reasoning itself.

Formally defines the language of mathematics, mathematical truth, and provability.

Gödel's First Incompleteness Theorem:

In any reasonable formal system, some true statements are not provable.

Proof: We use two properties of formal proofs:

- 1) Soundness: If ϕ has a proof π then ϕ is true.
- 2) Checkability: The language $\{ \langle \pi, \phi \rangle \mid \pi \text{ is a proof of statement } \phi \}$ is decidable.

Checkability implies the set of provable statements $\{ \langle \phi \rangle \mid \phi \text{ has a proof} \}$ is T-recognizable.

Intro to Mathematical Logic

12

Goal: A mathematical study of mathematical reasoning itself.

Formally defines the language of mathematics, mathematical truth, and provability.

Gödel's First Incompleteness Theorem:

In any reasonable formal system, some true statements are not provable.

Proof: We use two properties of formal proofs:

- 1) Soundness: If ϕ has a proof π then ϕ is true.
- 2) Checkability: The language $\{ \langle \pi, \phi \rangle \mid \pi \text{ is a proof of statement } \phi \}$ is decidable.

Checkability implies the set of provable statements $\{ \langle \phi \rangle \mid \phi \text{ has a proof} \}$ is T-recognizable.

Similarly, if we can always prove $\langle M, w \rangle \in \overline{ATM}$ when it is true, then \overline{ATM} is T-recognizable (false!).

Intro to Mathematical Logic

12

Goal: A mathematical study of mathematical reasoning itself.

Formally defines the language of mathematics, mathematical truth, and provability.

Gödel's First Incompleteness Theorem:

In any reasonable formal system, some true statements are not provable.

Proof: We use two properties of formal proofs:

- 1) Soundness: If ϕ has a proof π then ϕ is true.
- 2) Checkability: The language $\{ \langle \pi, \phi \rangle \mid \pi \text{ is a proof of statement } \phi \}$ is decidable.

Checkability implies the set of provable statements $\{ \langle \phi \rangle \mid \phi \text{ has a proof} \}$ is T-recognizable.

Similarly, if we can always prove $\langle M, w \rangle \in \overline{ATM}$ when it is true, then \overline{ATM} is T-recognizable (false!).

Therefore, some true statements of the form $\langle M, w \rangle \in \overline{ATM}$ are unprovable.

Intro to Mathematical Logic

12

Goal: A mathematical study of mathematical reasoning itself.

Formally defines the language of mathematics, mathematical truth, and provability.

Gödel's First Incompleteness Theorem:

In any reasonable formal system, some true statements are not provable.

Proof: We use two properties of formal proofs:

- 1) Soundness: If ϕ has a proof π then ϕ is true.
- 2) Checkability: The language $\{ \langle \pi, \phi \rangle \mid \pi \text{ is a proof of statement } \phi \}$ is decidable.

Checkability implies the set of provable statements $\{ \langle \phi \rangle \mid \phi \text{ has a proof} \}$ is T-recognizable.

Similarly, if we can always prove $\langle M, w \rangle \in \overline{ATM}$ when it is true, then \overline{ATM} is T-recognizable (false!).

Therefore, some true statements of the form $\langle M, w \rangle \in \overline{ATM}$ are unprovable.

Next, we use the Recursion Theorem to give a specific example of a true but unprovable statement.

Intro to Mathematical Logic

12

Goal: A mathematical study of mathematical reasoning itself.

Formally defines the language of mathematics, mathematical truth, and provability.

Gödel's First Incompleteness Theorem:

In any reasonable formal system, some true statements are not provable.

Proof: We use two properties of formal proofs:

- 1) Soundness: If ϕ has a proof π then ϕ is true.
- 2) Checkability: The language $\{ \langle \pi, \phi \rangle \mid \pi \text{ is a proof of statement } \phi \}$ is decidable.

Checkability implies the set of provable statements $\{ \langle \phi \rangle \mid \phi \text{ has a proof} \}$ is T-recognizable.

Similarly, if we can always prove $\langle M, w \rangle \in \overline{ATM}$ when it is true, then \overline{ATM} is T-recognizable (false!).

Therefore, some true statements of the form $\langle M, w \rangle \in \overline{ATM}$ are unprovable.

Next, we use the Recursion Theorem to give a specific example of a true but unprovable statement.

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

Theorem: (1) ϕ_U has no proof

(2) ϕ_U is true

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

Theorem: (1) ϕ_U has no proof

(2) ϕ_U is true

Proof:

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

Theorem: (1) ϕ_U has no proof

(2) ϕ_U is true

Proof:

(1) If ϕ_U has a proof .

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

Theorem: (1) ϕ_U has no proof

(2) ϕ_U is true

Proof:

(1) If ϕ_U has a proof \rightarrow TM R accepts 0

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

Theorem: (1) ϕ_U has no proof

(2) ϕ_U is true

Proof:

(1) If ϕ_U has a proof \rightarrow TM R accepts 0 $\rightarrow \langle R, 0 \rangle \in \overline{ATM}$ is false

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

Theorem: (1) ϕ_U has no proof

(2) ϕ_U is true

Proof:

(1) If ϕ_U has a proof \rightarrow TM R accepts 0 $\rightarrow \overbrace{\langle R, 0 \rangle}^{\phi_U} \in \overline{ATM}$ is false

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

Theorem: (1) ϕ_U has no proof

(2) ϕ_U is true

Proof:

(1) If ϕ_U has a proof \rightarrow TM R accepts 0 $\rightarrow \overbrace{\langle R, 0 \rangle}^{\phi_U} \in \overline{ATM}$ is false $\rightarrow \phi_U$ cannot have a proof.

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

Theorem: (1) ϕ_U has no proof

(2) ϕ_U is true

Proof:

(1) If ϕ_U has a proof \rightarrow TM R accepts 0 $\rightarrow \overbrace{\langle R, 0 \rangle \in \overline{ATM}}^{\phi_U}$ is false $\rightarrow \phi_U$ cannot have a proof.

(2) If ϕ_U is false \rightarrow

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

Theorem: (1) ϕ_U has no proof

(2) ϕ_U is true

Proof:

- (1) If ϕ_U has a proof \rightarrow TM R accepts 0 $\rightarrow \overbrace{\langle R, 0 \rangle \in \overline{ATM}}^{\phi_U}$ is false $\rightarrow \phi_U$ cannot have a proof.
- (2) If ϕ_U is false $\rightarrow \langle R, 0 \rangle \notin \overline{ATM}$

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

Theorem: (1) ϕ_U has no proof

(2) ϕ_U is true

Proof:

- (1) If ϕ_U has a proof \rightarrow TM R accepts 0 $\rightarrow \overbrace{\langle R, 0 \rangle \in ATM}^{\phi_U}$ is false $\rightarrow \phi_U$ cannot have a proof.
- (2) If ϕ_U is false $\rightarrow \langle R, 0 \rangle \notin ATM \rightarrow R$ accepts 0.

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

Theorem: (1) ϕ_U has no proof

(2) ϕ_U is true

Proof:

- (1) If ϕ_U has a proof \rightarrow TM R accepts 0 $\rightarrow \overbrace{\langle R, 0 \rangle \in \overline{ATM}}^{\phi_U}$ is false $\rightarrow \phi_U$ cannot have a proof.
- (2) If ϕ_U is false $\rightarrow \langle R, 0 \rangle \notin \overline{ATM} \rightarrow R$ accepts 0 $\rightarrow R$ found a proof that ϕ_U is true

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

Theorem: (1) ϕ_U has no proof

(2) ϕ_U is true

Proof:

(1) If ϕ_U has a proof \rightarrow TM R accepts 0 $\rightarrow \overbrace{\langle R, 0 \rangle \in ATM}^{\phi_U}$ is false $\rightarrow \phi_U$ cannot have a proof.

(2) If ϕ_U is false $\rightarrow \langle R, 0 \rangle \notin \overline{ATM} \rightarrow R$ accepts 0 $\rightarrow R$ found a proof that ϕ_U is true $\rightarrow \phi_U$ is true.

A True but Unprovable Statement

13

Implement Gödel statement “This statement is unprovable.”

Let ϕ_U be the statement $\langle R, 0 \rangle \in \overline{ATM}$ where R is the following TM:

$R =$ “On any input

1. Obtain $\langle R \rangle$ and use it to obtain ϕ_U .
2. For each possible proof $\pi = \pi_1, \pi_2, \dots$

Test if π is a proof that ϕ_U is true.

If yes, then *accept*. Otherwise, continue.”

Theorem: (1) ϕ_U has no proof

(2) ϕ_U is true

Proof:

(1) If ϕ_U has a proof \rightarrow TM R accepts 0 $\rightarrow \overbrace{\langle R, 0 \rangle \in ATM}^{\phi_U}$ is false $\rightarrow \phi_U$ cannot have a proof.

(2) If ϕ_U is false $\rightarrow \langle R, 0 \rangle \notin \overline{ATM} \rightarrow R$ accepts 0 $\rightarrow R$ found a proof that ϕ_U is true $\rightarrow \phi_U$ is true.

Quick review of this topic

14

1. Self-reference and The Recursion Theorem
2. Various applications.
3. Sketch of Gödel's First Incompleteness Theorem in mathematical logic.

