

بسم الله الرحمن الرحيم

نظريه علوم کامپیوتر

نظريه علوم کامپیوتر - بهار ۱۴۰۰-۱۴۰۱ - جلسه ششم: محاسبه پذیری و محاسبه ناپذیری (۲)

Theory of computation - 002 - S06 - non-computability (2)

Recall

- Languages on machines which are decidable
 - $ADFA = \{ \langle B, w \rangle \mid B \text{ is a DFA and } B \text{ accepts } w \}$
 - $ANFA = \{ \langle B, w \rangle \mid B \text{ is a NFA and } B \text{ accepts } w \}$
 - $EDFA = \{ \langle B \rangle \mid B \text{ is a DFA and } L(B) = \emptyset \}$
 - $EQDFA = \{ \langle A, B \rangle \mid A \text{ and } B \text{ are DFAs and } L(A) = L(B) \}$
 - $ACFG = \{ \langle G, w \rangle \mid G \text{ is a CFG and } w \in L(G) \}$
 - $ECFG = \{ \langle G \rangle \mid G \text{ is a CFG and } L(G) = \emptyset \}$
 - $EQCFG = \{ \langle G, H \rangle \mid G, H \text{ are CFGs and } L(G) = L(H) \}$
 - $AMBIGCFG = \{ \langle G \rangle \mid G \text{ is an ambiguous CFG} \}$

Acceptance Problem for TMs

12

Let $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Acceptance Problem for TMs

12

Let $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Acceptance Problem for TMs

12

Let $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof: Later

Acceptance Problem for TMs

12

Let $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof: Later

Theorem: ATM is T-recognizable

Acceptance Problem for TMs

12

Let $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof: Later

Theorem: ATM is T-recognizable

Proof: The following TM U recognizes ATM

Acceptance Problem for TMs

12

Let $ATM = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w\}$

Theorem: ATM is not decidable

Proof: Later

Theorem: ATM is T-recognizable

Proof: The following TM U recognizes ATM

$U =$ “On input $\langle M, w \rangle$

Acceptance Problem for TMs

12

Let $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof: Later

Theorem: ATM is T-recognizable

Proof: The following TM U recognizes ATM

$U =$ “On input $\langle M, w \rangle$

1. Simulate M on input w .

Acceptance Problem for TMs

12

Let $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof: Later

Theorem: ATM is T-recognizable

Proof: The following TM U recognizes ATM

$U =$ “On input $\langle M, w \rangle$

1. Simulate M on input w .
2. *Accept* if M halts and accepts.

Acceptance Problem for TMs

12

Let $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof: Later

Theorem: ATM is T-recognizable

Proof: The following TM U recognizes ATM

$U =$ “On input $\langle M, w \rangle$

1. Simulate M on input w .
2. *Accept* if M halts and accepts.
3. *Reject* if M halts and rejects.

Acceptance Problem for TMs

12

Let $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof: Later

Theorem: ATM is T-recognizable

Proof: The following TM U recognizes ATM

$U =$ “On input $\langle M, w \rangle$

1. Simulate M on input w .
2. *Accept* if M halts and accepts.
3. *Reject* if M halts and rejects.
4. *Reject* if M never halts.”

Acceptance Problem for TMs

12

Let $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof: Later

Theorem: ATM is T-recognizable

Proof: The following TM U recognizes ATM

$U =$ “On input $\langle M, w \rangle$

1. Simulate M on input w .
2. *Accept* if M halts and accepts.
3. *Reject* if M halts and rejects.
4. ~~*Reject* if M never halts.~~ Not a legal TM action.

Acceptance Problem for TMs

12

Let $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof: Later

Theorem: ATM is T-recognizable

Proof: The following TM U recognizes ATM

$U =$ “On input $\langle M, w \rangle$

1. Simulate M on input w .
2. *Accept* if M halts and accepts.
3. *Reject* if M halts and rejects.
4. ~~Reject if M never halts.~~ Not a legal TM action.

Turing’s original “Universal Computing Machine”



Acceptance Problem for TMs

12

Let $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof: Later

Theorem: ATM is T-recognizable

Proof: The following TM U recognizes ATM

$U =$ “On input $\langle M, w \rangle$

1. Simulate M on input w .
2. *Accept* if M halts and accepts.
3. *Reject* if M halts and rejects.
4. ~~*Reject* if M never halts.~~ Not a legal TM action.

Turing’s original “Universal Computing Machine”



Von Neumann said U inspired the concept of a stored program computer.

Acceptance Problem for TMs

12

Let $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof: Later

Theorem: ATM is T-recognizable

Proof: The following TM U recognizes ATM

$U =$ “On input $\langle M, w \rangle$

1. Simulate M on input w .
2. *Accept* if M halts and accepts.
3. *Reject* if M halts and rejects.
4. ~~*Reject* if M never halts.~~ Not a legal TM action.

Turing’s original “Universal Computing Machine”



Von Neumann said U inspired the concept of a stored program computer.

The Size of Infinity

The Size of Infinity

13

How to compare the relative sizes of infinite sets?

The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

Defn: Say that set A and B have the same size if there is a one-to-one and onto function $f: A \rightarrow B$

The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

Defn: Say that set A and B have the same size if there is a one-to-one and onto function $f: A \rightarrow B$

$x \neq y \rightarrow$
 $f(x) \neq f(y)$
“injective”

The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

Defn: Say that set A and B have the same size if there is a one-to-one and onto function $f: A \rightarrow B$

$x \neq y \rightarrow f(x) \neq f(y)$
“injective”

$\text{Range}(f) = B$
“surjective”

The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

Defn: Say that set A and B have the same size if there is

a one-to-one and onto function $f: A \rightarrow B$

$x \neq y \rightarrow$
 $f(x) \neq f(y)$
“injective”

$\text{Range}(f) = B$
“surjective”

We call such an f a 1-1 correspondence

The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

Defn: Say that set A and B have the same size if there is

a one-to-one and onto function $f: A \rightarrow B$

$x \neq y \rightarrow$
 $f(x) \neq f(y)$
“injective”

$\text{Range}(f) = B$
“surjective”

We call such an f a 1-1 correspondence

Informally, two sets have the same size if we can pair up their members.

The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

Defn: Say that set A and B have the same size if there is

a one-to-one and onto function $f: A \rightarrow B$

$x \neq y \rightarrow$
 $f(x) \neq f(y)$
"injective"

$\text{Range}(f) = B$
"surjective"

We call such an f a 1-1 correspondence

Informally, two sets have the same size if we can pair up their members.

This definition works for finite sets.

The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

Defn: Say that set A and B have the same size if there is a one-to-one and onto function $f: A \rightarrow B$

$x \neq y \rightarrow$
 $f(x) \neq f(y)$
“injective”

$\text{Range}(f) = B$
“surjective”

We call such an f a 1-1 correspondence

Informally, two sets have the same size if we can pair up their members.

This definition works for finite sets.



The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

Defn: Say that set A and B have the same size if there is

a one-to-one and onto function $f: A \rightarrow B$

$x \neq y \rightarrow$
 $f(x) \neq f(y)$
“injective”

$\text{Range}(f) = B$
“surjective”

We call such an f a 1-1 correspondence

Informally, two sets have the same size if we can pair up their members.

This definition works for finite sets.



The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

Defn: Say that set A and B have the same size if there is a one-to-one and onto function $f: A \rightarrow B$


$x \neq y \rightarrow f(x) \neq f(y)$
"injective"

$\text{Range}(f) = B$
"surjective"

We call such an f a 1-1 correspondence

Informally, two sets have the same size if we can pair up their members.

This definition works for finite sets.



The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

Defn: Say that set A and B have the same size if there is a one-to-one and onto function $f: A \rightarrow B$


$x \neq y \rightarrow f(x) \neq f(y)$ "injective"
 $\text{Range}(f) = B$ "surjective"

We call such an f a 1-1 correspondence

Informally, two sets have the same size if we can pair up their members.

This definition works for finite sets.

Apply it to infinite sets too.



The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

Defn: Say that set A and B have the same size if there is

a one-to-one and onto function $f: A \rightarrow B$

$x \neq y \rightarrow$
 $f(x) \neq f(y)$
“injective”


$\text{Range}(f) = B$
“surjective”

We call such an f a 1-1 correspondence

Informally, two sets have the same size if we can pair up their members.

This definition works for finite sets.

Apply it to infinite sets too.



The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

Defn: Say that set A and B have the same size if there is

a one-to-one and onto function $f: A \rightarrow B$

$x \neq y \rightarrow$
 $f(x) \neq f(y)$
“injective”


$\text{Range}(f) = B$
“surjective”

We call such an f a 1-1 correspondence

Informally, two sets have the same size if we can pair up their members.

This definition works for finite sets.

Apply it to infinite sets too.



© Source unknown. All rights reserved. This content is excluded from our Creative Commons license. For more information, see <https://ocw.mit.edu/fairuse>.

The Size of Infinity

13

How to compare the relative sizes of infinite sets?

Cantor (~1890s) had the following idea.

Defn: Say that set A and B have the same size if there is

a one-to-one and onto function $f: A \rightarrow B$

$x \neq y \rightarrow$
 $f(x) \neq f(y)$
“injective”


$\text{Range}(f) = B$
“surjective”

We call such an f a 1-1 correspondence

Informally, two sets have the same size if we can pair up their members.

This definition works for finite sets.

Apply it to infinite sets too.



© Source unknown. All rights reserved. This content is excluded from our Creative Commons license. For more information, see <https://ocw.mit.edu/fairuse>.

Countable Sets

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let

$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let

$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let

$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

$\mathbb{N} \quad \mathbb{Z}$


Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let

$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size



Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

\mathbb{N}	\mathbb{Z}
1	
2	
3	
4	
5	
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	
3	
4	
5	
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	
4	
5	
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	
5	
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

$$\text{Let } \mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$$

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

$\mathbb{N} \quad \mathbb{Q}^+$

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

$$\text{Let } \mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathcal{N}	\mathcal{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

N	Q^+

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

\mathbb{N}	\mathbb{Q}^+

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

\mathbb{N}	\mathbb{Q}^+
1	
2	
3	
4	
5	
6	
7	

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	
2	
3	
4	
5	
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	
3	
4	
5	
6	
7	

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	
3	
4	
5	
6	
7	

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	
4	
5	
6	
7	

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	
4	
5	
6	
7	

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	
4	
5	
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	
4	
5	
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	
5	
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	
5	
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	2/3
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	2/3
7	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	2/3
7	1/3

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	2/3
7	1/3

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	2/3
7	1/3

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	2/3
7	1/3

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	2/3
7	1/3

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	2/3
7	1/3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	2/3
7	1/3

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	2/3
7	1/3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	2/3
7	1/3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

Defn: A set is countable if it is finite or it has the same size as \mathbb{N} .

Both \mathbb{Z} and \mathbb{Q}^+ are countable.

Countable Sets

14

Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and let
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Show \mathbb{N} and \mathbb{Z} have the same size

Let $\mathbb{Q}^+ = \left\{ \frac{m}{n} : m, n \in \mathbb{N} \right\}$

Show \mathbb{N} and \mathbb{Q}^+ have the same size

\mathbb{N}	\mathbb{Z}
1	0
2	-1
3	1
4	-2
5	2
6	-3
7	3

\mathbb{N}	\mathbb{Q}^+
1	1/1
2	2/1
3	1/2
4	3/1
5	3/2
6	2/3
7	1/3

	1	2	3	4	...
1	1/1	1/2	1/3	1/4	
2	2/1	2/2	2/3	2/4	...
3	3/1	3/2	3/3	3/4	
4	4/1	4/2	4/3	4/4	

Defn: A set is countable if it is finite or it has the same size as \mathbb{N} .

Both \mathbb{Z} and \mathbb{Q}^+ are countable.

\mathbb{R} is Uncountable – Diagonalization

15

Let $\mathbb{R} =$ all real numbers (expressible by infinite decimal expansion)

\mathbb{R} is Uncountable – Diagonalization

15

Let $\mathbb{R} =$ all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

\mathbb{R} is Uncountable – Diagonalization

15

Let $\mathbb{R} =$ all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	
2	
3	
4	
5	
6	
7	

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	
3	
4	
5	
6	
7	

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	
4	
5	
6	
7	

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	
5	
6	
7	

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	
6	
7	

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	
7	

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2. 7 18281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2. 7 18281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\text{not in list}}{8}$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.8$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2. 7 18281828...
2	3.1 4 1592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#7}{8}\overset{\#4}{5}$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#7}{8}\overset{\#4}{5}$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#7}{8}\overset{\#4}{5}\overset{\#0}{1}$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#7}{8}\overset{\#4}{5}\overset{\#0}{1}$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#1}{8}\overset{\#4}{5}\overset{\#0}{1}\overset{\#2}{6}$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#1}{8}\overset{\#4}{5}\overset{\#0}{1}\overset{\#2}{6}$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#1}{8}\overset{\#4}{5}\overset{\#0}{1}\overset{\#2}{6}\overset{\#5}{1}$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#1}{8}\overset{\#4}{5}\overset{\#0}{1}\overset{\#2}{6}\overset{\#5}{1}$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2. 7 18281828...
2	3.1 4 1592653...
3	0.00 0 000000...
4	1.414 2 13562...
5	0.1428 5 7242...
6	0.20787 9 576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#1}{8}\overset{\#4}{5}\overset{\#0}{1}\overset{\#2}{6}\overset{\#5}{1}\overset{\#9}{8}$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#1}{8}\overset{\#4}{5}\overset{\#0}{1}\overset{\#2}{6}\overset{\#5}{1}\overset{\#9}{8}$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#1}{8}\overset{\#4}{5}\overset{\#0}{1}\overset{\#2}{6}\overset{\#5}{1}\overset{\#9}{8}\overset{\#8}{2}...$$

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#1}{8}\overset{\#4}{5}\overset{\#0}{1}\overset{\#2}{6}\overset{\#5}{1}\overset{\#9}{8}\overset{\#8}{2}...$$

differs from the n^{th} number in the n^{th} digit
so cannot be the n^{th} number for any n .

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2. 7 18281828...
2	3.1 4 1592653...
3	0.00 0 000000...
4	1.414 2 13562...
5	0.1428 5 7242...
6	0.20787 9 576...
7	1.234567 8 90...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#1}{8}\overset{\#4}{5}\overset{\#0}{1}\overset{\#2}{6}\overset{\#5}{1}\overset{\#9}{8}\overset{\#8}{2}...$$

differs from the n^{th} number in the n^{th} digit
so cannot be the n^{th} number for any n .

Hence x is not paired with any n . It is missing from the list.

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#1}{8}\overset{\#4}{5}\overset{\#0}{1}\overset{\#2}{6}\overset{\#5}{1}\overset{\#9}{8}\overset{\#8}{2}...$$

differs from the n^{th} number in the n^{th} digit
so cannot be the n^{th} number for any n .

Hence x is not paired with any n . It is missing from the list.

Therefore f is not a 1-1 correspondence.

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

Diagonalization

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#1}{8}\overset{\#4}{5}\overset{\#0}{1}\overset{\#2}{6}\overset{\#5}{1}\overset{\#9}{8}\overset{\#8}{2}...$$

differs from the n^{th} number in the n^{th} digit
so cannot be the n^{th} number for any n .

Hence x is not paired with any n . It is missing from the list.

Therefore f is not a 1-1 correspondence.

\mathbb{R} is Uncountable – Diagonalization

15

Let \mathbb{R} = all real numbers (expressible by infinite decimal expansion)

Theorem: \mathbb{R} is uncountable

Proof by contradiction via diagonalization: Assume \mathbb{R} is countable

Diagonalization

So there is a 1-1 correspondence $f: \mathbb{N} \rightarrow \mathbb{R}$

1	2.718281828...
2	3.141592653...
3	0.000000000...
4	1.414213562...
5	0.142857242...
6	0.207879576...
7	1.234567890...

Demonstrate a number $x \in \mathbb{R}$ that is missing from the list.

$$x = 0.\overset{\#1}{8}\overset{\#4}{5}\overset{\#0}{1}\overset{\#2}{6}\overset{\#5}{1}\overset{\#9}{8}\overset{\#8}{2}...$$

differs from the n^{th} number in the n^{th} digit
so cannot be the n^{th} number for any n .

Hence x is not paired with any n . It is missing from the list.

Therefore f is not a 1-1 correspondence.

\mathbb{R} is Uncountable – Corollaries

16

Let \mathcal{L} = all languages

\mathbb{R} is Uncountable – Corollaries

16

Let \mathcal{L} = all languages

Corollary 1: \mathcal{L} is uncountable

\mathbb{R} is Uncountable – Corollaries

16

Let \mathcal{L} = all languages

Corollary 1: \mathcal{L} is uncountable

Proof: There's a 1-1 correspondence from \mathcal{L} to \mathbb{R} so they are the same size.

	{,	0,	1,	00,	01,	10,	11,	000,	...
	{	0,		00,	01,				...
	.0	1	0	1	1	0	0	0	...

\mathbb{R} is Uncountable – Corollaries

16

Let \mathcal{L} = all languages

Corollary 1: \mathcal{L} is uncountable

Proof: There's a 1-1 correspondence from \mathcal{L} to \mathbb{R} so they are the same size.

Observation: $\Sigma^* = \{\varepsilon, 0, 1, 00, 01, 10, 11, 000, \dots\}$ is countable.

	{,	0,	1,	00,	01,	10,	11,	000,	...
	{	0,		00,	01,				...
	.0	1	0	1	1	0	0	0	...

\mathbb{R} is Uncountable – Corollaries

16

Let \mathcal{L} = all languages

Corollary 1: \mathcal{L} is uncountable

Proof: There's a 1-1 correspondence from \mathcal{L} to \mathbb{R} so they are the same size.

Observation: $\Sigma^* = \{\varepsilon, 0, 1, 00, 01, 10, 11, 000, \dots\}$ is countable.

Let \mathcal{M} = all Turing machines

	{,	0,	1,	00,	01,	10,	11,	000,	...
	{	0,		00,	01,				...
	.0	1	0	1	1	0	0	0	...

\mathbb{R} is Uncountable – Corollaries

16

Let \mathcal{L} = all languages

Corollary 1: \mathcal{L} is uncountable

Proof: There's a 1-1 correspondence from \mathcal{L} to \mathbb{R} so they are the same size.

Observation: $\Sigma^* = \{\varepsilon, 0, 1, 00, 01, 10, 11, 000, \dots\}$ is countable.

Let \mathcal{M} = all Turing machines

Observation: \mathcal{M} is countable.

	{,	0,	1,	00,	01,	10,	11,	000,	...
	{	0,		00,	01,				...
	.0	1	0	1	1	0	0	0	...

\mathbb{R} is Uncountable – Corollaries

16

Let \mathcal{L} = all languages

Corollary 1: \mathcal{L} is uncountable

Proof: There's a 1-1 correspondence from \mathcal{L} to \mathbb{R} so they are the same size.

Observation: $\Sigma^* = \{\varepsilon, 0, 1, 00, 01, 10, 11, 000, \dots\}$ is countable.

Let \mathcal{M} = all Turing machines

Observation: \mathcal{M} is countable.

Because $\{\langle M \rangle \mid M \text{ is a TM}\} \subseteq \Sigma^*$.

	{,	0,	1,	00,	01,	10,	11,	000,	...
	{	0,		00,	01,				...
	.0	1	0	1	1	0	0	0	...

\mathbb{R} is Uncountable – Corollaries

16

Let \mathcal{L} = all languages

Corollary 1: \mathcal{L} is uncountable

Proof: There's a 1-1 correspondence from \mathcal{L} to \mathbb{R} so they are the same size.

Observation: $\Sigma^* = \{\varepsilon, 0, 1, 00, 01, 10, 11, 000, \dots\}$ is countable.

Let \mathcal{M} = all Turing machines

Observation: \mathcal{M} is countable.

Because $\{\langle M \rangle \mid M \text{ is a TM}\} \subseteq \Sigma^*$.

	{,	0,	1,	00,	01,	10,	11,	000,	...
	{	0,		00,	01,				...
	.0	1	0	1	1	0	0	0	...

Corollary 2: Some language is not decidable.

\mathbb{R} is Uncountable – Corollaries

16

Let \mathcal{L} = all languages

Corollary 1: \mathcal{L} is uncountable

Proof: There's a 1-1 correspondence from \mathcal{L} to \mathbb{R} so they are the same size.

Observation: $\Sigma^* = \{\varepsilon, 0, 1, 00, 01, 10, 11, 000, \dots\}$ is countable.

Let \mathcal{M} = all Turing machines

Observation: \mathcal{M} is countable.

Because $\{\langle M \rangle \mid M \text{ is a TM}\} \subseteq \Sigma^*$.

	{,	0,	1,	00,	01,	10,	11,	000,	...
	{	0,		00,	01,				...
	.0	1	0	1	1	0	0	0	...

Corollary 2: Some language is not decidable.

Because there are more languages than TMs.

\mathbb{R} is Uncountable – Corollaries

16

Let \mathcal{L} = all languages

Corollary 1: \mathcal{L} is uncountable

Proof: There's a 1-1 correspondence from \mathcal{L} to \mathbb{R} so they are the same size.

Observation: $\Sigma^* = \{\varepsilon, 0, 1, 00, 01, 10, 11, 000, \dots\}$ is countable.

Let \mathcal{M} = all Turing machines

Observation: \mathcal{M} is countable.

Because $\{\langle M \rangle \mid M \text{ is a TM}\} \subseteq \Sigma^*$.

	{	0,	1,	00,	01,	10,	11,	000,	...
	{	0,		00,	01,				...
	.0	1	0	1	1	0	0	0	...

Corollary 2: Some language is not decidable.

Because there are more languages than TMs.

We will show some specific language A_{TM} is not decidable.

\mathbb{R} is Uncountable – Corollaries

16

Let \mathcal{L} = all languages

Corollary 1: \mathcal{L} is uncountable

Proof: There's a 1-1 correspondence from \mathcal{L} to \mathbb{R} so they are the same size.

Observation: $\Sigma^* = \{\varepsilon, 0, 1, 00, 01, 10, 11, 000, \dots\}$ is countable.

Let \mathcal{M} = all Turing machines

Observation: \mathcal{M} is countable.

Because $\{\langle M \rangle \mid M \text{ is a TM}\} \subseteq \Sigma^*$.

	{,	0,	1,	00,	01,	10,	11,	000,	...
	{	0,		00,	01,				...
	.0	1	0	1	1	0	0	0	...

Corollary 2: Some language is not decidable.

Because there are more languages than TMs.

We will show some specific language A_{TM} is not decidable.

\mathbb{R} is Uncountable – Corollaries

16

Let \mathcal{L} = all languages

Corollary 1: \mathcal{L} is uncountable

Proof: There's a 1-1 correspondence from \mathcal{L} to \mathbb{R} so they are the same size.

Observation: $\Sigma^* = \{\epsilon, 0, 1, 00, 01, 10, 11, 000, \dots\}$ is countable

Let \mathcal{M} = all Turing machines

Observation: \mathcal{M} is countable.

Because $\{\langle M \rangle \mid M \text{ is a TM}\} \subseteq \Sigma^*$.

Corollary 2: Some language is not decidable.

Because there are more languages than TMs.

We will show some specific language A_{TM} is not decidable.

Check-in 8.1

Hilbert's 1st question asked if there is a set of intermediate size between \mathbb{N} and \mathbb{R} . Gödel and Cohen showed that we cannot answer this question by using the standard axioms of mathematics.

How can we interpret their conclusion?

- (a) We need better axioms to describe reality.
- (b) Infinite sets have no mathematical reality so Hilbert's 1st question has no answer.

ATM is undecidable

17

Recall $ATM = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w\}$

ATM is undecidable

17

Recall $ATM = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w\}$

Theorem: ATM is not decidable

ATM is undecidable

17

Recall $ATM = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w\}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

ATM is undecidable

17

Recall $ATM = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w\}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

ATM is undecidable

17

Recall $ATM = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w\}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Why is this proof a diagonalization?

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions: ...

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs		All TM descriptions:			
↓					...
		acc	rej	acc	acc ...

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				
					...
	acc	rej	acc	acc	...
	rej	rej	rej	rej	

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				
					...
	acc	rej	acc	acc	...
	rej	rej	rej	rej	
	acc	acc	acc	acc	...

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				
					...
	acc	rej	acc	acc	...
	rej	rej	rej	rej	
	acc	acc	acc	acc	...
	rej	rej	acc	acc	

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				

	acc	rej	acc	acc	...
	rej	rej	rej	rej	
	acc	acc	acc	acc	...
	rej	rej	acc	acc	

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				

	acc	rej	acc	acc	...
	rej	rej	rej	rej	
	acc	acc	acc	acc	...
	rej	rej	acc	acc	
	rej				

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				
					...
	acc	rej	acc	acc	...
	rej	rej	rej	rej	
	acc	acc	acc	acc	...
	rej	rej	acc	acc	
	rej				

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				
					...
	acc	rej	acc	acc	...
	rej	rej	rej	rej	
	acc	acc	acc	acc	...
	rej	rej	acc	acc	
	rej	acc			

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				

	acc	rej	acc	acc	...
	rej	rej	rej	rej	
	acc	acc	acc	acc	...
	rej	rej	acc	acc	
	rej	acc			

ATM is undecidable

17

Recall $ATM = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w\}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				
					...
	acc	rej	acc	acc	...
	rej	rej	rej	rej	
	acc	acc	acc	acc	...
	rej	rej	acc	acc	
	rej	acc	rej		

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				

	acc	rej	acc	acc	...
	rej	rej	rej	rej	
	acc	acc	acc	acc	...
	rej	rej	acc	acc	
	rej	acc	rej		

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				
					...
	acc	rej	acc	acc	...
	rej	rej	rej	rej	
	acc	acc	acc	acc	...
	rej	rej	acc	acc	
	rej	acc	rej	rej	

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				
					...
	acc	rej	acc	acc	...
	rej	rej	rej	rej	
	acc	acc	acc	acc	...
	rej	rej	acc	acc	
	rej	acc	rej	rej	?

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ "On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts."

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				...
	acc	rej	acc	acc	...
	rej	rej	rej	rej	
	acc	acc	acc	acc	...
	rej	rej	acc	acc	
	rej	acc	rej	rej	?

ATM is undecidable

17

Recall $ATM = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ accepts } w \}$

Theorem: ATM is not decidable

Proof by contradiction: Assume some TM H decides ATM .

So H on $\langle M, w \rangle = \begin{cases} \text{Accept} & \text{if } M \text{ accepts } w \\ \text{Reject} & \text{if not} \end{cases}$

Use H to construct TM D

$D =$ “On input $\langle M \rangle$

1. Simulate H on input $\langle M, \langle M \rangle \rangle$
2. *Accept* if H rejects. *Reject* if H accepts.”

D accepts $\langle M \rangle$ iff M doesn't accept $\langle M \rangle$.

D accepts $\langle D \rangle$ iff D doesn't accept $\langle D \rangle$.

Contradiction.

Why is this proof a diagonalization?

All TMs ↓	All TM descriptions:				...
	acc	rej	acc	acc	...
	rej	rej	rej	rej	
	acc	acc	acc	acc	...
	rej	rej	acc	acc	
	rej	acc	rej	rej	?

\overline{ATM} is T-unrecognizable

18

Theorem: If A and \overline{A} are T-recognizable then A is decidable

\overline{ATM} is T-unrecognizable

18

Theorem: If A and \overline{A} are T-recognizable then A is decidable

Proof: Let TM M_1 and M_2 recognize A and \overline{A} .

\overline{ATM} is T-unrecognizable

18

Theorem: If A and \overline{A} are T-recognizable then A is decidable

Proof: Let TM M_1 and M_2 recognize A and \overline{A} .

Construct TM T deciding A .

\overline{ATM} is T-unrecognizable

18

Theorem: If A and \overline{A} are T-recognizable then A is decidable

Proof: Let TM M_1 and M_2 recognize A and \overline{A} .

Construct TM T deciding A .

$T =$ “On input w

\overline{ATM} is T-unrecognizable

18

Theorem: If A and \overline{A} are T-recognizable then A is decidable

Proof: Let TM M_1 and M_2 recognize A and \overline{A} .

Construct TM T deciding A .

$T =$ “On input w

1. Run M_1 and M_2 on w in parallel until one accepts.

\overline{ATM} is T-unrecognizable

18

Theorem: If A and \overline{A} are T-recognizable then A is decidable

Proof: Let TM M_1 and M_2 recognize A and \overline{A} .

Construct TM T deciding A .

$T =$ “On input w

1. Run M_1 and M_2 on w in parallel until one accepts.
2. If M_1 accepts then *accept*.
If M_2 accepts then *reject*.”

\overline{ATM} is T-unrecognizable

18

Theorem: If A and \overline{A} are T-recognizable then A is decidable

Proof: Let TM M_1 and M_2 recognize A and \overline{A} .

Construct TM T deciding A .

$T =$ “On input w

1. Run M_1 and M_2 on w in parallel until one accepts.
2. If M_1 accepts then *accept*.
If M_2 accepts then *reject*.”

Corollary: \overline{ATM} is T-unrecognizable

\overline{ATM} is T-unrecognizable

18

Theorem: If A and \overline{A} are T-recognizable then A is decidable

Proof: Let TM M_1 and M_2 recognize A and \overline{A} .

Construct TM T deciding A .

$T =$ “On input w

1. Run M_1 and M_2 on w in parallel until one accepts.
2. If M_1 accepts then *accept*.
If M_2 accepts then *reject*.”

Corollary: \overline{ATM} is T-unrecognizable

Proof: ATM is T-recognizable but also undecidable

\overline{ATM} is T-unrecognizable

18

Theorem: If A and \overline{A} are T-recognizable then A is decidable

Proof: Let TM M_1 and M_2 recognize A and \overline{A} .


Construct TM T deciding A .

$T =$ "On input w

1. Run M_1 and M_2 on w in parallel until one accepts.
2. If M_1 accepts then *accept*.
If M_2 accepts then *reject*."

Corollary: \overline{ATM} is T-unrecognizable

Proof: ATM is T-recognizable but also undecidable



\overline{ATM} is T-unrecognizable

18

Theorem: If A and \overline{A} are T-recognizable then A is decidable

Proof: Let TM M_1 and M_2 recognize A and \overline{A} .


Construct TM T deciding A .

$T =$ “On input w

1. Run M_1 and M_2 on w in parallel until one accepts.
2. If M_1 accepts then *accept*.
If M_2 accepts then *reject*.”

Corollary: \overline{ATM} is T-unrecognizable

Proof: ATM is T-recognizable but also undecidable



\overline{ATM} is T-unrecognizable

18

Theorem: If A and \overline{A} are T-recognizable then A is decidable

Proof: Let TM M_1 and M_2 recognize A and \overline{A} .

Construct TM T deciding A .

$T =$ “On input w

1. Run M_1 and M_2 on w in parallel until one accepts
2. If M_1 accepts then *accept*.
If M_2 accepts then *reject*.”

Corollary: \overline{ATM} is T-unrecognizable

Proof: ATM is T-recognizable but also undecidable

Check-in 8.3

From what we’ve learned, which closure properties can we prove for the class of T-recognizable languages? Choose all that apply.

- (a) Closed under union.
- (b) Closed under intersection.
- (c) Closed under complement.
- (d) Closed under concatenation.
- (e) Closed under star.

The Reducibility Method

The Reducibility Method

19

Use our knowledge that ATM is undecidable to show other problems are undecidable.

The Reducibility Method

19

Use our knowledge that ATM is undecidable to show other problems are undecidable.

Defn: $HALT_{TM} = \{ \langle M, w \rangle \mid M \text{ halts on input } w \}$

The Reducibility Method

19

Use our knowledge that ATM is undecidable to show other problems are undecidable.

Defn: $HALT_{TM} = \{ \langle M, w \rangle \mid M \text{ halts on input } w \}$

Theorem: $HALT_{TM}$ is undecidable

The Reducibility Method

19

Use our knowledge that ATM is undecidable to show other problems are undecidable.

Defn: $HALT_{TM} = \{ \langle M, w \rangle \mid M \text{ halts on input } w \}$

Theorem: $HALT_{TM}$ is undecidable

Proof by contradiction, showing that ATM is reducible to $HALT_{TM}$:

The Reducibility Method

19

Use our knowledge that ATM is undecidable to show other problems are undecidable.

Defn: $HALT_{TM} = \{ \langle M, w \rangle \mid M \text{ halts on input } w \}$

Theorem: $HALT_{TM}$ is undecidable

Proof by contradiction, showing that ATM is reducible to $HALT_{TM}$:

Assume that $HALT_{TM}$ is decidable and show that ATM is decidable (false!).

The Reducibility Method

19

Use our knowledge that ATM is undecidable to show other problems are undecidable.

Defn: $HALT_{TM} = \{ \langle M, w \rangle \mid M \text{ halts on input } w \}$

Theorem: $HALT_{TM}$ is undecidable

Proof by contradiction, showing that ATM is reducible to $HALT_{TM}$:

Assume that $HALT_{TM}$ is decidable and show that ATM is decidable (false!).

Let TM R decide $HALT_{TM}$.

The Reducibility Method

19

Use our knowledge that ATM is undecidable to show other problems are undecidable.

Defn: $HALT_{TM} = \{ \langle M, w \rangle \mid M \text{ halts on input } w \}$

Theorem: $HALT_{TM}$ is undecidable

Proof by contradiction, showing that ATM is reducible to $HALT_{TM}$:

Assume that $HALT_{TM}$ is decidable and show that ATM is decidable (false!).

Let TM R decide $HALT_{TM}$.

Construct TM S deciding ATM .

The Reducibility Method

19

Use our knowledge that ATM is undecidable to show other problems are undecidable.

Defn: $HALT_{TM} = \{ \langle M, w \rangle \mid M \text{ halts on input } w \}$

Theorem: $HALT_{TM}$ is undecidable

Proof by contradiction, showing that ATM is reducible to $HALT_{TM}$:

Assume that $HALT_{TM}$ is decidable and show that ATM is decidable (false!).

Let TM R decide $HALT_{TM}$.

Construct TM S deciding ATM .

$S =$ “On input $\langle M, w \rangle$

The Reducibility Method

19

Use our knowledge that ATM is undecidable to show other problems are undecidable.

Defn: $HALT_{TM} = \{ \langle M, w \rangle \mid M \text{ halts on input } w \}$

Theorem: $HALT_{TM}$ is undecidable

Proof by contradiction, showing that ATM is reducible to $HALT_{TM}$:

Assume that $HALT_{TM}$ is decidable and show that ATM is decidable (false!).

Let TM R decide $HALT_{TM}$.

Construct TM S deciding ATM .

$S =$ “On input $\langle M, w \rangle$

1. Use R to test if M on w halts. If not, reject.

The Reducibility Method

19

Use our knowledge that ATM is undecidable to show other problems are undecidable.

Defn: $HALT_{TM} = \{ \langle M, w \rangle \mid M \text{ halts on input } w \}$

Theorem: $HALT_{TM}$ is undecidable

Proof by contradiction, showing that ATM is reducible to $HALT_{TM}$:

Assume that $HALT_{TM}$ is decidable and show that ATM is decidable (false!).

Let TM R decide $HALT_{TM}$.

Construct TM S deciding ATM .

$S =$ “On input $\langle M, w \rangle$

1. Use R to test if M on w halts. If not, reject.
2. Simulate M on w until it halts (as guaranteed by R).

The Reducibility Method

19

Use our knowledge that ATM is undecidable to show other problems are undecidable.

Defn: $HALT_{TM} = \{ \langle M, w \rangle \mid M \text{ halts on input } w \}$

Theorem: $HALT_{TM}$ is undecidable

Proof by contradiction, showing that ATM is reducible to $HALT_{TM}$:

Assume that $HALT_{TM}$ is decidable and show that ATM is decidable (false!).

Let TM R decide $HALT_{TM}$.

Construct TM S deciding ATM .

$S =$ “On input $\langle M, w \rangle$

1. Use R to test if M on w halts. If not, reject.
2. Simulate M on w until it halts (as guaranteed by R).
3. If M has accepted then *accept*.
If M has rejected then *reject*.

The Reducibility Method

19

Use our knowledge that ATM is undecidable to show other problems are undecidable.

Defn: $HALT_{TM} = \{ \langle M, w \rangle \mid M \text{ halts on input } w \}$

Theorem: $HALT_{TM}$ is undecidable

Proof by contradiction, showing that ATM is reducible to $HALT_{TM}$:

Assume that $HALT_{TM}$ is decidable and show that ATM is decidable (false!).

Let TM R decide $HALT_{TM}$.

Construct TM S deciding ATM .

$S =$ “On input $\langle M, w \rangle$

1. Use R to test if M on w halts. If not, reject.
2. Simulate M on w until it halts (as guaranteed by R).
3. If M has accepted then *accept*.
If M has rejected then *reject*.

TM S decides ATM , a contradiction. Therefore $HALT_{TM}$ is undecidable.

The Reducibility Method

19

Use our knowledge that ATM is undecidable to show other problems are undecidable.

Defn: $HALT_{TM} = \{ \langle M, w \rangle \mid M \text{ halts on input } w \}$

Theorem: $HALT_{TM}$ is undecidable

Proof by contradiction, showing that ATM is reducible to $HALT_{TM}$:

Assume that $HALT_{TM}$ is decidable and show that ATM is decidable (false!).

Let TM R decide $HALT_{TM}$.

Construct TM S deciding ATM .

$S =$ “On input $\langle M, w \rangle$

1. Use R to test if M on w halts. If not, reject.
2. Simulate M on w until it halts (as guaranteed by R).
3. If M has accepted then *accept*.
If M has rejected then *reject*.

TM S decides ATM , a contradiction. Therefore $HALT_{TM}$ is undecidable.

Reducibility – Concept

Reducibility – Concept

20

If we have two languages (or problems) A and B , then A is reducible to B means that we can use B to solve A .

Reducibility – Concept

20

If we have two languages (or problems) A and B , then A is reducible to B means that we can use B to solve A .

Example 1: Measuring the area of a rectangle is reducible to measuring the lengths of its sides.

Reducibility – Concept

20

If we have two languages (or problems) A and B , then A is reducible to B means that we can use B to solve A .

Example 1: Measuring the area of a rectangle is reducible to measuring the lengths of its sides.

Example 2: We showed that $ANFA$ is reducible to $ADFA$.

Reducibility – Concept

20

If we have two languages (or problems) A and B , then A is reducible to B means that we can use B to solve A .

Example 1: Measuring the area of a rectangle is reducible to measuring the lengths of its sides.

Example 2: We showed that $ANFA$ is reducible to $ADFA$.

Example 3: From Pset 2, $PUSHER$ is reducible to $ECFG$.
(Idea- Convert push states to accept states.)

Reducibility – Concept

20

If we have two languages (or problems) A and B , then A is reducible to B means that we can use B to solve A .

Example 1: Measuring the area of a rectangle is reducible to measuring the lengths of its sides.

Example 2: We showed that $ANFA$ is reducible to $ADFA$.

Example 3: From Pset 2, $PUSHER$ is reducible to $ECFG$.
(Idea- Convert push states to accept states.)

If A is reducible to B then solving B gives a solution to A .

Reducibility – Concept

20

If we have two languages (or problems) A and B , then A is reducible to B means that we can use B to solve A .

Example 1: Measuring the area of a rectangle is reducible to measuring the lengths of its sides.

Example 2: We showed that $ANFA$ is reducible to $ADFA$.

Example 3: From Pset 2, $PUSHER$ is reducible to $ECFG$.
(Idea- Convert push states to accept states.)

If A is reducible to B then solving B gives a solution to A .
- then B is easy $\rightarrow A$ is easy.

Reducibility – Concept

20

If we have two languages (or problems) A and B , then A is reducible to B means that we can use B to solve A .

Example 1: Measuring the area of a rectangle is reducible to measuring the lengths of its sides.

Example 2: We showed that ANFA is reducible to ADFA .

Example 3: From Pset 2, *PUSHER* is reducible to *ECFG* .
(Idea- Convert push states to accept states.)

If A is reducible to B then solving B gives a solution to A .

- then B is easy $\rightarrow A$ is easy.
- then A is hard $\rightarrow B$ is hard.

Reducibility – Concept

20

If we have two languages (or problems) A and B , then A is reducible to B means that we can use B to solve A .

Example 1: Measuring the area of a rectangle is reducible to measuring the lengths of its sides.

Example 2: We showed that ANFA is reducible to ADFA .

Example 3: From Pset 2, *PUSHER* is reducible to *ECFG* .
(Idea- Convert push states to accept states.)

If A is reducible to B then solving B gives a solution to A .

- then B is easy $\rightarrow A$ is easy.

- then A is hard $\rightarrow B$ is hard.

Reducibility – Concept

20

If we have two languages (or problems) A and B , then A is reducible to B means that we can use B to solve A .

Example 1: Measuring the area of a rectangle is reducible to measuring the lengths of its sides.

Example 2: We showed that ANFA is reducible to ADFA .

Example 3: From Pset 2, *PUSHER* is reducible to *ECFG* .
(Idea- Convert push states to accept states.)

If A is reducible to B then solving B gives a solution to A .

- then B is easy $\rightarrow A$ is easy.

- then A is hard $\rightarrow B$ is hard.

this is the form we will use

Reducibility – Concept

20

If we have two languages (or problems) A and B , then A is reducible to B means that we can use B to solve A .

Example 1: Measuring the area of a rectangle is reducible to measuring the lengths of its sides.

Example 2: We showed that ANFA is reducible to ADFA .

Example 3: From Pset 2, *PUSHER* is reducible to *ECFG* .
(Idea- Convert push states to accept states.)

If A is reducible to B then solving B gives a solution to A .

- then B is easy $\rightarrow A$ is easy.

- then A is hard $\rightarrow B$ is hard.

this is the form we will use

Reducibility – Concept

20

If we have two languages (or problems) A and B , then A is reducible to B means that we can use B to solve A .

Example 1: Measuring the area of a rectangle is reducible to measuring the lengths of its sides.

Example 2: We showed that ANFA is reducible to A DFA

Example 3: From Pset 2, *PUSHER* is reducible to *ECFG*.
(Idea- Convert push states to accept states.)

If A is reducible to B then solving B gives a solution to A
- then B is easy $\rightarrow A$ is easy.

- then A is hard $\rightarrow B$ is hard.

this is the form we will use

Check-in 9.1

Is Biology reducible to Physics?

- (a) Yes, all aspects of the physical world may be explained in terms of Physics, at least in principle.
- (b) No, some things in the world, maybe life, the brain, or consciousness, are beyond the realm of Physics.
- (c) I'm on the fence on this question!

ETM is undecidable

21

Let $ETM = \{ \langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset \}$

Theorem: ETM is undecidable

ETM is undecidable

21

Let $ETM = \{ \langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset \}$

Theorem: ETM is undecidable

Proof by contradiction. Show that ATM is reducible to ETM .

ETM is undecidable

21

Let $ETM = \{ \langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset \}$

Theorem: ETM is undecidable

Proof by contradiction. Show that ATM is reducible to ETM .

Assume that ETM is decidable and show that ATM is decidable (false!).

ETM is undecidable

21

Let $ETM = \{ \langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset \}$

Theorem: ETM is undecidable

Proof by contradiction. Show that ATM is reducible to ETM .

Assume that ETM is decidable and show that ATM is decidable (false!).

Let TM R decide ETM .

ETM is undecidable

21

Let $ETM = \{ \langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset \}$

Theorem: ETM is undecidable

Proof by contradiction. Show that ATM is reducible to ETM .

Assume that ETM is decidable and show that ATM is decidable (false!).

Let TM R decide ETM .

Construct TM S deciding ATM .

ETM is undecidable

21

Let $ETM = \{ \langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset \}$

Theorem: ETM is undecidable

Proof by contradiction. Show that ATM is reducible to ETM .

Assume that ETM is decidable and show that ATM is decidable (false!).

Let TM R decide ETM .

Construct TM S deciding ATM .

$S =$ "On input $\langle M, w \rangle$

ETM is undecidable

21

Let $ETM = \{ \langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset \}$

Theorem: ETM is undecidable

Proof by contradiction. Show that ATM is reducible to ETM .

Assume that ETM is decidable and show that ATM is decidable (false!).

Let TM R decide ETM .

Construct TM S deciding ATM .

$S =$ “On input $\langle M, w \rangle$

1. Transform M to new TM $M_w =$ “On input x
 1. If $x \neq w$, *reject*.
 2. else run M on w
 3. *Accept* if M accepts.”

ETM is undecidable

21

Let $ETM = \{ \langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset \}$

Theorem: ETM is undecidable

Proof by contradiction. Show that ATM is reducible to ETM .

Assume that ETM is decidable and show that ATM is decidable (false!).

Let TM R decide ETM .

Construct TM S deciding ATM .

$S =$ “On input $\langle M, w \rangle$

1. Transform M to new TM $M_w =$ “On input x
 1. If $x \neq w$, *reject*.
 2. else run M on w
 3. *Accept* if M accepts.”

M_w works like M except that it always rejects strings x where $x \neq w$.

ETM is undecidable

21

Let $ETM = \{ \langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset \}$

Theorem: ETM is undecidable

Proof by contradiction. Show that ATM is reducible to ETM .

Assume that ETM is decidable and show that ATM is decidable (false!).

Let TM R decide ETM .

Construct TM S deciding ATM .

$S =$ “On input $\langle M, w \rangle$

1. Transform M to new TM $M_w =$ “On input x
 1. If $x \neq w$, *reject*.
 2. else run M on w
 3. *Accept* if M accepts.”

M_w works like M except that it always rejects strings x where $x \neq w$.

So

$$L(M_w) = \begin{cases} \{w\} & \text{if } M \text{ accepts } w \\ \emptyset & \text{if } M \text{ rejects } w \end{cases}$$

ETM is undecidable

21

Let $ETM = \{ \langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset \}$

Theorem: ETM is undecidable

Proof by contradiction. Show that ATM is reducible to ETM .

Assume that ETM is decidable and show that ATM is decidable (false!).

Let TM R decide ETM .

Construct TM S deciding ATM .

$S =$ “On input $\langle M, w \rangle$

1. Transform M to new TM $M_w =$ “On input x
 1. If $x \neq w$, *reject*.
 2. else run M on w
 3. *Accept* if M accepts.”
2. Use R to test whether $L(M_w) = \emptyset$

M_w works like M except that it always rejects strings x where $x \neq w$.

So

$$L(M_w) = \begin{cases} \{w\} & \text{if } M \text{ accepts } w \\ \emptyset & \text{if } M \text{ rejects } w \end{cases}$$

ETM is undecidable

21

Let $ETM = \{ \langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset \}$

Theorem: ETM is undecidable

Proof by contradiction. Show that ATM is reducible to ETM .

Assume that ETM is decidable and show that ATM is decidable (false!).

Let TM R decide ETM .

Construct TM S deciding ATM .

$S =$ “On input $\langle M, w \rangle$

1. Transform M to new TM $M_w =$ “On input x
 1. If $x \neq w$, *reject*.
 2. else run M on w
 3. *Accept* if M accepts.”
2. Use R to test whether $L(M_w) = \emptyset$
3. If YES [so M rejects w] then *reject*.
If NO [so M accepts w] then *accept*.

M_w works like M except that it always rejects strings x where $x \neq w$.

So

$$L(M_w) = \begin{cases} \{w\} & \text{if } M \text{ accepts } w \\ \emptyset & \text{if } M \text{ rejects } w \end{cases}$$

ETM is undecidable

21

Let $ETM = \{ \langle M \rangle \mid M \text{ is a TM and } L(M) = \emptyset \}$

Theorem: ETM is undecidable

Proof by contradiction. Show that ATM is reducible to ETM .

Assume that ETM is decidable and show that ATM is decidable (false!).

Let TM R decide ETM .

Construct TM S deciding ATM .

$S =$ “On input $\langle M, w \rangle$

1. Transform M to new TM $M_w =$ “On input x
 1. If $x \neq w$, *reject*.
 2. else run M on w
 3. *Accept* if M accepts.”
2. Use R to test whether $L(M_w) = \emptyset$
3. If YES [so M rejects w] then *reject*.
If NO [so M accepts w] then *accept*.

M_w works like M except that it always rejects strings x where $x \neq w$.

So

$$L(M_w) = \begin{cases} \{w\} & \text{if } M \text{ accepts } w \\ \emptyset & \text{if } M \text{ rejects } w \end{cases}$$