

بسم الله الرحمن الرحيم


جلسه هجدهم

خلاصه سازی برای مدداده

احساس فُشَرْدَگَى تىكى



مرور




مرور

$$C = (c+2)k$$

(ϵ, Ck) -RIP

نک_ k

$$x \rightarrow y = \Pi x$$




الگوریتم بازیابی (ورودی: y و Π)


$$\|\hat{x} - x\|_1 \leq c(k) \quad \min_{y \text{ is } k\text{-sparse}} \|y - x\|_1$$

$$\begin{aligned} & \min ||z||_1 \\ \text{s.t. } & \quad \Pi z = y \end{aligned}$$


مرور



مرور



مرور




Lemma 2 Assume A satisfies the nullspace property of order $2k$ with constant $C < 2$. Then for x^* that minimizes $\|x^*\|_1$ subject to $Ax^* = Ax$ we have

$$\|x - x^*\|_1 \leq \frac{2C}{2-C} Err_1^k(x)$$

: $A\eta = 0$ که η

$$|\mathcal{T}| = 2k \text{ برای هر } \|\eta\|_1 \leq C \|\eta_{-\mathcal{T}}\|_1$$



Lemma 1 Suppose that A satisfies RIP of order $(c + 2)k$ with constant δ , $c > 1$. Then A satisfies the nullspace property of order $2k$ with constant $C = 1 + \sqrt{2/c}(1 + \delta)(1 - \delta)$.

Lemma 2 Assume A satisfies the nullspace property of order $2k$ with constant $C < 2$. Then for x^* that minimizes $\|x^*\|_1$ subject to $Ax^* = Ax$ we have

$$\|x - x^*\|_1 \leq \frac{2C}{2-C} Err_1^k(x)$$

: $A\eta = 0$ که η

$|T| = 2k$ برای هر $\|\eta\|_1 \leq C \|\eta_{-T}\|_1$

جایگزین

RIP₁



Lemma 1 Suppose that A satisfies RIP of order $(c + 2)k$ with constant δ , $c > 1$. Then A satisfies the nullspace property of order $2k$ with constant $C = 1 + \sqrt{2/c}(1 + \delta)(1 - \delta)$.



Definition 28. We say a matrix $\Pi \in \mathbb{R}^{m \times n}$ satisfies the (ε, k) -restricted isometry property (or RIP for short) if for all k -sparse vectors x of unit Euclidean norm,

$$1 - \varepsilon \leq \|\Pi x\|_2^2 \leq 1 + \varepsilon.$$

Definition 63 (RIP_1). A matrix A is (ϵ, k) - RIP_1 if for all k sparse vector v ,

$$(1 - \epsilon) \|v\|_1 \leq \|Av\|_1 \leq (1 + \epsilon) \|v\|_1$$







دوست داریم:

Definition 63 (RIP_1). A matrix A is (ϵ, k) - RIP_1 if for all k sparse vector v ,

$$(1 - \epsilon) \|v\|_1 \leq \|Av\|_1 \leq (1 + \epsilon) \|v\|_1$$



دوست داریم:


Definition 63 (RIP_1). A matrix A is (ϵ, k) - RIP_1 if for all k sparse vector v ,

$$(1 - \epsilon) \|v\|_1 \leq \|Av\|_1 \leq (1 + \epsilon) \|v\|_1$$

$$d(1 - \epsilon) \|v\|_1 \leq \|Av\|_1 \leq d \|v\|_1$$


Definition 64 (Expander). A (l, ϵ) -unbalanced expander is a bipartite simple graph $G = (U, V, E)$, $|U| = n, |V| = m$, with left degree d such that for any $X \subset U$ with $|X| \leq l$, the set of neighbors $N(X)$ of X has size $|N(X)| \geq (1 - \epsilon)d|X|$.

$$\deg() = d$$



Definition 64 (Expander). A (l, ϵ) -unbalanced expander is a bipartite simple graph $G = (U, V, E)$, $|U| = n$, $|V| = m$, with left degree d such that for any $X \subset U$ with $|X| \leq l$, the set of neighbors $N(X)$ of X has size $|N(X)| \geq (1 - \epsilon)d|X|$.

$$\deg() = d$$




Claim 69. For any $n/2 \geq l \geq 1$, $\epsilon > 0$, there exists a (l, ϵ) -unbalanced expander with left degree $d = O(\log(n/l)/\epsilon)$ and right set size $O(ld/\epsilon) = O(l \log(n/l)/\epsilon^2)$.


Definition 64 (Expander). A (l, ϵ) -unbalanced expander is a bipartite simple graph $G = (U, V, E)$, $|U| = n, |V| = m$, with left degree d such that for any $X \subset U$ with $|X| \leq l$, the set of neighbors $N(X)$ of X has size $|N(X)| \geq (1 - \epsilon)d|X|$.

Theorem 65 ([BGIKS08]). For $A \in \{0, 1\}^{m \times n}$, if the underlying bipartite graph is a $(k, d(1 - \frac{\epsilon}{2}))$ expander, then for all k sparse vector v

$$d(1 - \epsilon) \|v\|_1 \leq \|Av\|_1 \leq d \|v\|_1$$




Proof. $\|Av\|_1 \leq d \|v\|_1$



$$d(1 - \epsilon) \|v\|_1 \leq \|Av\|_1:$$


Proof. $\|Av\|_1 \leq d \|v\|_1$



$$d(1 - \epsilon) \|v\|_1 \leq \|Av\|_1:$$

Claim 66. $\|Av\|_1 \geq \sum_{(i,j)=e \in E} r(e)|v_i|$

Proof. $\|Av\|_1 \leq d \|v\|_1$




$$d(1 - \epsilon) \|v\|_1 \leq \|Av\|_1:$$

Claim 66. $\|Av\|_1 \geq \sum_{(i,j)=e \in E} r(e)|v_i|$

Claim 67. $\sum_{(i,j)=e \in E} r(e)|v_i| \geq (1 - \epsilon)d \|v\|_1$

Proof. $\|Av\|_1 \leq d \|v\|_1$



$$d(1 - \epsilon) \|v\|_1 \leq \|Av\|_1:$$

Claim 66. $\|Av\|_1 \geq \sum_{(i,j)=e \in E} r(e)|v_i|$

Claim 67. $\sum_{(i,j)=e \in E} r(e)|v_i| \geq (1 - \epsilon)d \|v\|_1$

$$\sum_{(i,j)=e \in E} r(e)|v_i| \geq \sum_{(i,j)=e \in E} r'(e)|v_i| \geq d \|v\|_1 - \epsilon d \|v\|_1$$

Definition 64 (Expander). A (l, ϵ) -unbalanced expander is a bipartite simple graph $G = (U, V, E)$, $|U| = n$, $|V| = m$, with left degree d such that for any $X \subset U$ with $|X| \leq l$, the set of neighbors $N(X)$ of X has size $|N(X)| \geq (1 - \epsilon)d|X|$.


Theorem 65 ([BGIKS08]). For $A \in \{0, 1\}^{m \times n}$, if the underlying bipartite graph is a $(k, d(1 - \frac{\epsilon}{2}))$ expander, then for all k sparse vector v

$$d(1 - \epsilon) \|v\|_1 \leq \|Av\|_1 \leq d \|v\|_1$$

Definition 64 (Expander). A (l, ϵ) -unbalanced expander is a bipartite simple graph $G = (U, V, E)$, $|U| = n$, $|V| = m$, with left degree d such that for any $X \subset U$ with $|X| \leq l$, the set of neighbors $N(X)$ of X has size $|N(X)| \geq (1 - \epsilon)d|X|$.

Theorem 65 ([BGIKS08]). For $A \in \{0, 1\}^{m \times n}$, if the underlying bipartite graph is a $(k, d(1 - \frac{\epsilon}{2}))$ expander, then for all k sparse vector v

$$d(1 - \epsilon) \|v\|_1 \leq \|Av\|_1 \leq d \|v\|_1$$




Lemma 70. Consider any $\eta \in \mathbb{R}^n$ such that $A\eta = 0$, and let S be any set of k coordinates of η . Then we have

$$\|\eta_S\|_1 \leq \alpha(\epsilon)\|\eta\|_1$$

Definition 64 (Expander). A (l, ϵ) -unbalanced expander is a bipartite simple graph $G = (U, V, E)$, $|U| = n$, $|V| = m$, with left degree d such that for any $X \subset U$ with $|X| \leq l$, the set of neighbors $N(X)$ of X has size $|N(X)| \geq (1 - \epsilon)d|X|$.


Theorem 65 ([BGIKS08]). For $A \in \{0, 1\}^{m \times n}$, if the underlying bipartite graph is a $(k, d(1 - \frac{\epsilon}{2}))$ expander, then for all k sparse vector v

$$d(1 - \epsilon) \|v\|_1 \leq \|Av\|_1 \leq d \|v\|_1$$



Lemma 70. Consider any $\eta \in \mathbb{R}^n$ such that $A\eta = 0$, and let S be any set of k coordinates of η . Then we have

$$\|\eta_S\|_1 \leq \alpha(\epsilon)\|\eta\|_1$$




Lemma 2 Assume A satisfies the nullspace property of order $2k$ with constant $C < 2$. Then for x^* that minimizes $\|x^*\|_1$ subject to $Ax^* = Ax$ we have

$$\|x - x^*\|_1 \leq \frac{2C}{2 - C} Err_1^k(x)$$

Definition 64 (Expander). A (l, ϵ) -unbalanced expander is a bipartite simple graph $G = (U, V, E)$, $|U| = n$, $|V| = m$, with left degree d such that for any $X \subset U$ with $|X| \leq l$, the set of neighbors $N(X)$ of X has size $|N(X)| \geq (1 - \epsilon)d|X|$.

Theorem 65 ([BGIKS08]). For $A \in \{0, 1\}^{m \times n}$, if the underlying bipartite graph is a $(k, d(1 - \frac{\epsilon}{2}))$ expander, then for all k sparse vector v

$$d(1 - \epsilon) \|v\|_1 \leq \|Av\|_1 \leq d \|v\|_1$$




Lemma 70. Consider any $\eta \in \mathbb{R}^n$ such that $A\eta = 0$, and let S be any set of k coordinates of η . Then we have

$$\|\eta_S\|_1 \leq \alpha(\epsilon)\|\eta\|_1$$

Definition 64 (Expander). A (l, ϵ) -unbalanced expander is a bipartite simple graph $G = (U, V, E)$, $|U| = n$, $|V| = m$, with left degree d such that for any $X \subset U$ with $|X| \leq l$, the set of neighbors $N(X)$ of X has size $|N(X)| \geq (1 - \epsilon)d|X|$.


Theorem 65 ([BGIKS08]). For $A \in \{0, 1\}^{m \times n}$, if the underlying bipartite graph is a $(k, d(1 - \frac{\epsilon}{2}))$ expander, then for all k sparse vector v


$$d(1 - \epsilon) \|v\|_1 \leq \|Av\|_1 \leq d \|v\|_1$$




Lemma 70. Consider any $\eta \in \mathbb{R}^n$ such that $A\eta = 0$, and let S be any set of k coordinates of η . Then we have


$$\|\eta_S\|_1 \leq \alpha(\epsilon)\|\eta\|_1$$









$$\|A'\eta_S\|_1$$



$$\|A'\eta_S\|_1 = \|A\eta_S\|_1$$




$$\|A'\eta_S\|_1 = \|A\eta_S\|_1 \geq d(1-2\epsilon)\|\eta_S\|_1$$



$$\|A'\eta_S\|_1 = \|A\eta_S\|_1 \geq d(1-2\epsilon)\|\eta_S\|_1$$


$$0 = \|A'\eta\|_1 = \|A'\eta_S + A'\eta_{-S}\|_1$$



$$\|A'\eta_S\|_1 = \|A\eta_S\|_1 \geq d(1-2\epsilon)\|\eta_S\|_1$$




$$0 = \|A'\eta\|_1 = \|A'\eta_S + A'\eta_{-S}\|_1$$



 
 $\|A'\eta_S\|_1 = \|A\eta_S\|_1 \geq d(1-2\epsilon)\|\eta_S\|_1$

$$0 = \|A'\eta\|_1 = \|A'\eta_S + A'\eta_{-S}\|_1$$



$$\|A'\eta_S\|_1 = \|A\eta_S\|_1 \geq d(1-2\epsilon)\|\eta_S\|_1$$

$$0 = \|A'\eta\|_1 = \|A'\eta_S + A'\eta_{-S}\|_1$$


$$\begin{aligned}
0 = \|A'\eta\|_1 &\geq \|A'\eta_S\|_1 - \sum_{l \geq 1} \sum_{(i,j) \in E, i \in S_l, j \in N(S)} |\eta_i| \\
&\geq d(1 - 2\epsilon) \|\eta_S\|_1 - \sum_{l \geq 1} |E(S_l : N(S))| \min_{i \in S_{l-1}} |\eta_i| \\
&\geq d(1 - 2\epsilon) \|\eta_S\|_1 - \sum_{l \geq 1} |E(S_l : N(S))| \cdot \|\eta_{S_{l-1}}\|_1 / k \\
&\leq \\
&|N(S \cup S_l)| \geq d(1 - \epsilon) |S \cup S_l| \quad \text{dashed box} \rightarrow d\epsilon 2k \\
&\geq d(1 - 2\epsilon) \|\eta_S\|_1 - d\epsilon 2k \sum_{l \geq 1} \|\eta_{S_{l-1}}\|_1 / k \\
&\geq d(1 - 2\epsilon) \|\eta_S\|_1 - 2d\epsilon \|\eta\|_1
\end{aligned}$$

$$d(1 - 2\epsilon) \|\eta_S\|_1 \leq 2d\epsilon \|\eta\|_1$$

Definition 64 (Expander). A (l, ϵ) -unbalanced expander is a bipartite simple graph $G = (U, V, E)$, $|U| = n$, $|V| = m$, with left degree d such that for any $X \subset U$ with $|X| \leq l$, the set of neighbors $N(X)$ of X has size $|N(X)| \geq (1 - \epsilon)d|X|$.


Theorem 65 ([BGIKS08]). For $A \in \{0, 1\}^{m \times n}$, if the underlying bipartite graph is a $(k, d(1 - \frac{\epsilon}{2}))$ expander, then for all k sparse vector v

$$d(1 - \epsilon) \|v\|_1 \leq \|Av\|_1 \leq d \|v\|_1$$



Lemma 70. Consider any $\eta \in \mathbb{R}^n$ such that $A\eta = 0$, and let S be any set of k coordinates of η . Then we have

$$\|\eta_S\|_1 \leq \alpha(\epsilon)\|\eta\|_1$$



Lemma 2 Assume A satisfies the nullspace property of order $2k$ with constant $C < 2$. Then for x^* that minimizes $\|x^*\|_1$ subject to $Ax^* = Ax$ we have

$$\|x - x^*\|_1 \leq \frac{2C}{2 - C} Err_1^k(x)$$

