

## Review

# Magnesium Ion: A New Switch in Tumor Treatment

Leyi Huang <sup>1,2,†</sup> , Renxi Lin <sup>2,3,†</sup>, Jiaxi Chen <sup>2</sup>, Yuanlin Qi <sup>2,\*</sup>  and Ling Lin <sup>2,4,\*</sup>

<sup>1</sup> Laboratory of Gynecologic Oncology, Fujian Maternity and Child Health Hospital, College of Clinical Medicine for Obstetrics & Gynecology and Pediatrics, Fujian Medical University, Fuzhou 350001, China; lyhuang23@163.com

<sup>2</sup> Department of Biochemistry and Molecular Biology, Fujian Medical University, Fuzhou 350122, China; linrenxi89@fjmu.edu.cn (R.L.); 19959179207@163.com (J.C.)

<sup>3</sup> Experimental Teaching Center of Basic Medicine, Fujian Medical University, Fuzhou 350122, China

<sup>4</sup> Key Laboratory of Brain Aging and Neurodegenerative Disease, Fujian Medical University, Fuzhou 350122, China

\* Correspondence: ylqi@fjmu.edu.cn (Y.Q.); llinzheng2022@gmail.com (L.L.)

† These authors contributed equally to this work.

**Abstract:** The magnesium ion is an essential cation in the human body and participates in numerous physiological activities. A deficiency in magnesium ions is closely associated with tumor development, and supplementation with magnesium ions has been shown to partially inhibit tumor growth. However, the specific mechanisms by which magnesium ions suppress tumor proliferation remain unclear. Currently, studies have revealed that mitochondria may serve as a crucial intermediate link in the regulation of tumors by magnesium ions. Mitochondria might intervene in the proliferation and invasion of tumor cells by modulating energy metabolism and oxidative stress levels. Regrettably, there has been no comprehensive review of the role of magnesium in cancer therapy to date. Therefore, this article provides a comprehensive scrutiny of the relationship between magnesium ions and tumors, aiming to offer insights for clinical tumor treatment strategies involving magnesium ion intervention.

**Keywords:** magnesium ion; mitochondria; mitochondrial function; energy metabolism; cancer therapy



**Citation:** Huang, L.; Lin, R.; Chen, J.; Qi, Y.; Lin, L. Magnesium Ion: A New Switch in Tumor Treatment. *Biomedicines* **2024**, *12*, 1717. <https://doi.org/10.3390/biomedicines12081717>

Academic Editors: Jan Ježek and Veronique Baud

Received: 30 April 2024

Revised: 12 July 2024

Accepted: 26 July 2024

Published: 1 August 2024



**Copyright:** © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Magnesium ions ( $Mg^{2+}$ ), the fourth most abundant element in the human body and the second most prevalent cation in cellular environments after potassium ions, actively participate in vital physiological processes. They regulate enzymatic reactions involved in substance synthesis, influencing cell growth, differentiation, energy metabolism, excitability transmission in neurons, and cardiac excitability [1]. This is facilitated by magnesium ion transport proteins, with normal plasma magnesium concentrations ranging between 0.75 and 0.85 mmol/L [2]. Factors such as dietary intake, renal excretion, and skeletal muscle storage can impact magnesium homeostasis [3]. As a calcium ion antagonist, magnesium ion deficiency in neurons leads to increased activity of N-methyl-D-aspartic acid receptor (NMDA), causing excessive influx of calcium ions, reactive oxygen species (ROS) production, and mitochondrial damage. This heightened neuronal excitability is implicated in various diseases [4], including Parkinson's and Alzheimer's diseases, migraine, anxiety, and depression [3].

Due to the critical role of magnesium in physiological functions, some researchers have begun to consider the correlation between magnesium and cancer. Interestingly, besides magnesium ions, certain metal ions are also believed to be closely associated with cancer therapy. A recent study from China found that iron promotes DNA damage repair, thereby promoting platinum resistance in ovarian cancer; thus, iron depletion may emerge as a new direction for ovarian cancer treatment [5]. Similarly, copper can induce oxidative stress, inhibit the ubiquitin–proteasome system, and promote tumor cell death

through mechanisms like “cuproptosis” [6]. Furthermore, copper can serve as a nanocarrier to deliver therapeutic drugs, exerting anti-tumor efficacy. Therefore, it is reasonable to speculate that magnesium may also hold significant potential in cancer therapy.

In 1969, Bois et al. [7] successfully induced thymic lymphoma in rats through a magnesium-deficient diet. In 1989, Kurata et al. [8] confirmed in mice that magnesium supplementation in the diet effectively reduced the incidence of liver cancer, while a magnesium-deficient diet could induce tumor formation. A higher magnesium intake in the diet (400 mg/day) has been shown to reduce the incidence of colorectal cancer in postmenopausal women, and elevated blood magnesium levels are negatively correlated with the risk of colorectal cancer in females [9]. Moreover, recent research by Lötscher et al. [10], published in *Cell*, indicated that mice fed a magnesium-deficient diet exhibited significantly reduced efficacy of chimeric antigen receptor T-cell immunotherapy (CAR-T) against tumors, and B-cell lymphoma patients with lower serum magnesium levels had a shorter survival period.

As a cofactor for DNA repair enzymes, magnesium plays a crucial role in maintaining genomic stability and regulating cell proliferation, differentiation, and apoptosis. Magnesium deficiency may compromise DNA stability, contributing to tumor initiation and poor prognosis [11,12]. Yan’s team reported that magnesium glycinate inhibits liver cancer cell proliferation by blocking the mitogen-activated protein kinase (MAPK) signaling pathway in vitro [13]. Additionally, magnesium supplementation can inhibit the transcription of genes related to liver cancer cell proliferation by dephosphorylating p-Smad2/3 and blocking the transforming growth factor  $\beta$  (TGF- $\beta$ ) signaling pathway [14].

Meanwhile, magnesium ions play a crucial regulatory role in mitochondrial function. According to the research conducted by Koning et al. [15], magnesium sulfate can safeguard the mitochondrial respiratory chain, concurrently reducing the generation of ROS and inflammatory factors. Additionally, magnesium glycyrrhizinate is effective in preserving mitochondrial membrane potential and ultrastructure, while mitigating mitochondrial DNA (mtDNA) damage [16]. Given the pivotal role of mitochondrial oxidative stress and mtDNA damage in tumor progression [17], magnesium ions hold promise as a novel target for cancer therapy. Further investigation into their potential mechanisms in anticancer treatment is anticipated to offer new therapeutic avenues and hope for cancer patients.

## 2. Methods

To identify eligible studies for this review, we performed a computerized search of the PubMed and Web of Science databases, covering all publications available up to December 2023. Our search employed keywords including “magnesium”, “cancer”, “mitochondrial”, and their synonyms. Moreover, we scrutinized the reference lists of pertinent articles and reviews to identify studies not indexed in these databases. We meticulously evaluated all abstracts and full-text articles, selecting those deemed relevant for screening and inclusion in this review.

## 3. Magnesium Ion Transport Proteins and Their Implications in Tumor Development

Mg<sup>2+</sup> is predominantly distributed within cellular compartments, and its intracellular distribution and regulation rely heavily on magnesium ion transport proteins located on cellular and organelle membranes. Currently, major magnesium ion transport proteins include members of the transient receptor potential melastatin (TRPM) channel protein family, the human solute carrier (SLC) superfamily, magnesium transporter (MagT) proteins, cyclin M (CNNM) family proteins, and mitochondrial RNA splicing 2 (Mrs2) family genes.

### 3.1. TRPM Family

The TRPM family is widely expressed in numerous mammalian cells and is implicated in various biological functions, including redox reactions, inflammation, and insulin secretion [18]. This channel protein family comprises numerous members with varying permeabilities to calcium ions ( $\text{Ca}^{2+}$ ) and magnesium ions ( $\text{Mg}^{2+}$ ). TRPM7 and TRPM9 play crucial roles in intestinal magnesium ion absorption [19]. Moreover, the TRPM family participates in tumorigenesis by regulating the calcium ion balance [20], modulating cellular oxidative phosphorylation levels [21], and influencing ROS generation [22] within mitochondria.

In prostate cancer, the expression of TRPM8 and TRPM2 is significantly greater in tumor cells than in normal cells, whereas in lung cancer, TRPM5 upregulation is predominant [23]. Additionally, the TRPM family plays a significant role in various cancers, such as breast cancer, pancreatic cancer, and melanoma, and it is a potential target for antitumor therapies [18,23]. The TRPM family may provide insights into its potential as a therapeutic target for combating tumorigenesis.

### 3.2. SLC Family

The SLC family comprises crucial membrane transport proteins on the cell membrane and consists of 52 subfamilies with more than 400 members. These proteins are widely distributed on cell membranes and various organelle membranes and function as typical transmembrane proteins [24]. The SLC41 protein family facilitates the transmembrane transport of various metal ions, glucose, lipids, neurotransmitters, and drugs [25], thus playing a crucial role in the development of various diseases, including metabolic disorders, neurological diseases, and tumors [26].

The SLC41 family was discovered in 2003 and comprises SLC41A1, SLC41A2, and SLC41A3 [26]. The SLC41 family is capable of transmembrane transport of  $\text{Mg}^{2+}$  and, along with other magnesium ion transporters, such as TRPM7, participates in the regulation of  $\text{Mg}^{2+}$  homeostasis within the organism [27]. Variations in the SLC41A1 gene within the PARK16 locus in Parkinson's disease patients have a significant impact on the pathogenesis of Parkinson's disease as well as on ion homeostasis and the function of dopamine neurons [24,26]. Additionally, the overexpression or downregulation of SLC41A1 is closely associated with the occurrence of preeclampsia and renal diseases [24]. A study by Njiaju et al. [28] reported the association between SLC41A2 and chemotherapeutic drug sensitivity, in which knocking down SLC41A2 significantly increased the sensitivity of tumor cells to paclitaxel. Conversely, high SLC41A3 expression is closely correlated with poor prognosis in hepatocellular carcinoma patients, leading to a significantly shorter survival period in patients with elevated SLC41A3 expression than in those with normal SLC41A3 expression [29]. In summary, the SLC41 family participates in the onset of various diseases through the regulation of  $\text{Mg}^{2+}$  homeostasis and has emerged as a potential target for drug therapy.

### 3.3. MagT Family

The MagT protein family primarily includes MagT1, which is widely expressed in various human organs and tissues. It exhibits high similarity with human tumor suppressor candidate 3 (TUSC3). Both MagT1 and TUSC3 function in magnesium ion transport and protein N-terminal glycosylation. MagT1 was identified as a crucial target gene for X-linked immunodeficiency with magnesium defect (XMEN) disease [30]. The absence of the MagT1 protein leads to impaired N-glycosylation of platelet glycoproteins, resulting in platelet dysfunction and accelerated arterial thrombosis formation [31]. Simultaneously, the MagT1 protein regulates the intracellular magnesium ion concentration, influencing macrophage polarization and immune cell function [32].

MagT1 also plays a significant role in regulating the growth of tumor cells. Zheng et al. [33] reported a close correlation between MagT1 expression and poor prognosis in patients with colorectal cancer, with patients expressing lower levels of MagT1 experiencing longer survival. In breast cancer research, Li et al. [34] discovered that downregulation of MagT1 protein expression significantly inhibited the progression of breast cancer by suppressing Ki67. Additionally, MagT1 has been identified as an essential gene for cervical cancer cell proliferation [35], and low MagT1 expression inhibits the growth of cervical cancer cells and promotes cell apoptosis [36].

### 3.4. CNNM Family


The CNNM protein family is primarily distributed in brain tissues, kidneys, and intestines. As cyclin M is a conserved  $Mg^{2+}$  transport protein, the main function of this protein family is to transport magnesium ions out of cells. Additionally, the CNNM family proteins promote the absorption of  $Mg^{2+}$  in the intestines and kidneys [37]. The mechanism of magnesium transport by the CNNM family is complex. According to Bai et al. [38], the CNNM family proteins were regulated by the phosphatase of regenerating liver (PRLS), which is closely associated with tumor metastasis. CNNM selectively binds to the TRPM7 protein, thereby facilitating  $Mg^{2+}$  uptake. The intricate role of the CNNM family in  $Mg^{2+}$  transport is crucial for maintaining the intracellular  $Mg^{2+}$  balance [37]. Dysfunction of the CNNM family proteins often leads to hereditary hypomagnesemia, resulting in conditions such as hypertension, pulmonary arterial hypertension, epilepsy, Parkinson's disease, and other disorders [39]. In colon cancer tissues, the expression of the CNNM4 protein is significantly decreased compared to that in normal tissues. Critically, a lower level of CNNM4 implies a greater degree of malignancy of the tumor [40].

### 3.5. MRS Family

The Mrs2 protein is a magnesium ion transport protein located on the mitochondrial membrane. When stimulated by the glycolysis end-product lactate, Mrs2 transports magnesium ions stored in the endoplasmic reticulum to the mitochondria [41]. The spatiotemporal changes in magnesium ions within cells induced by Mrs2 result in reduced mitochondrial oxygen consumption and increased ROS production [42]. Using electron microscopy, He et al. [43] discovered that in addition to  $Mg^{2+}$ , Mrs2 also serves as a nonselective channel for calcium ions and can permeate sodium and potassium ions. The well-known Warburg effect demonstrates that most tumor cells exist in a microenvironment characterized by lactate accumulation. Therefore, subcellular magnesium ion transport mediated by Mrs2 is likely to become a crucial “switch” in regulating tumor cell growth [44].

Furthermore, Mrs2 plays a significant role in regulating mitochondrial function and cellular metabolism. According to the latest research by Madaris et al. [45], in mice induced with a classic Western diet (a high-sugar, high-fat, high-cholesterol diet), knocking out the mitochondrial Mrs2 channel effectively inhibited Western diet-induced obesity, metabolic syndrome, and spontaneous tumor incidence. This difference may be due to the loss of Mrs2, leading to the reprogramming of systemic energy metabolism regulated by hypoxia-inducible factor-1 $\alpha$  (HIF-1 $\alpha$ ) in the liver and adipose tissue, enhancing mitochondrial activity, reducing hepatic fat accumulation, and inducing browning of white adipose tissue.

Due to the unique structural location of Mrs2 and its crucial role in subcellular magnesium ion transport and mitochondrial metabolic functions, Mrs2 has a great potential for elucidating the mechanism of action of magnesium ions against cancer. Mrs2 may serve as a new target for energy metabolism-related tumor therapy (Figure 1).



**Figure 1.** Magnesium ion transporters regulate intracellular magnesium ion levels. On the cell membrane, MagT1 and TRPM7 primarily mediate the transport of magnesium ions into the cell, while CNNM2 is a bidirectional magnesium ion transporter. SLC41A1 functions as a sodium/magnesium exchanger. Meanwhile, on the mitochondrial membrane, there are two magnesium ion transporters. Mrs2 is mainly responsible for transporting magnesium ions from the endoplasmic reticulum into the mitochondria in response to lactate stimulation, whereas SLC41A3 is responsible for transporting magnesium ions out of the mitochondria. **Abbreviations:** TRPM, transient receptor potential melastatin channel protein family; SLC, human solute carrier superfamily; MagT, magnesium transporter proteins; CNNM, cyclin M family proteins; Mrs2, mitochondrial RNA splicing 2 family genes. By Figdraw 2.0.

## 4. The Impact of Magnesium Ions on Tumor Development

### 4.1. Breast Cancer

Breast cancer is a significant gynecological tumor. Interestingly, high magnesium intake improves the prognosis of breast cancer. A study involving 1170 breast cancer patients from New York State found that dietary magnesium intake is negatively correlated with overall mortality [46]. This correlation may be due to the protective effects of magnesium ions on the cardiovascular system, as well as their roles in maintaining genomic stability [47], regulating cell differentiation, proliferation, and apoptosis, as well as preventing angiogenesis [48].

The inhibitory effect of magnesium on breast cancer has also been confirmed in basic experiments. A study by Yang et al. [49] found that increasing the intracellular  $Mg^{2+}$  concentration in breast cancer cells, induced by methyl jasmonate, led to a decrease in the expression of the magnesium ion transporter TRPM7, an increase in ROS levels, and the promotion of apoptosis in breast cancer cells. In animal models, knocking out the



magnesium ion transporter MagT1 significantly downregulated Ki67, thereby markedly inhibiting the progression of breast cancer [34]. Additionally, a clinical controlled study found that plasma magnesium levels in female breast cancer patients were significantly lower than those in the normal population [50]. Therefore, we have reason to believe that magnesium ions play an important role in the occurrence and development of breast cancer, and that appropriate dietary supplementation of magnesium could effectively inhibit the progression of breast cancer.

#### 4.2. Colorectal Cancer

In colorectal cancer research, many investigators have found that magnesium intake greatly influences the risk and prognosis of the disease. According to Gorczyca et al. [51], daily supplementation of 400 mg of magnesium can reduce the incidence of colorectal cancer in postmenopausal women. Additionally, Polter et al. [52] identified through a prospective study that serum magnesium levels were negatively correlated with colon cancer risk among female participants. Moreover, higher magnesium intake effectively improves chemotherapy-induced peripheral neuropathy in colorectal cancer patients [53]. Furthermore, Wesselink et al. [54] discovered through a large multicenter prospective cohort study that magnesium intake lowers overall mortality in colorectal cancer patients. This effect may be attributed to magnesium's crucial role in regulating the synthesis metabolism of 25-hydroxyvitamin D [55]. High levels of magnesium and 25-hydroxyvitamin D inhibit inflammation and enhance patient survival rates [56,57].

In basic research, Li et al. [58] proposed that magnesium supplementation can induce apoptosis in colorectal cancer cells by activating the caspase-3 pathway. This finding was supported by animal experiments, where continuous magnesium injection for three weeks inhibited subcutaneous tumor growth in nude mice, and hematoxylin–eosin staining (HE) confirmed significant apoptotic features in the tumors of magnesium-supplemented mice [58]. These results provide a theoretical basis for the clinical use of magnesium-related drugs in the treatment of colorectal cancer, demonstrating the potential clinical value of magnesium ion therapies.

#### 4.3. Other Cancers

In addition to breast cancer and colorectal cancer, magnesium ions have also been shown to be associated with many other common cancers. A prospective cohort study by the Shrubsole team from the United States found that higher dietary magnesium intake is associated with a lower risk of liver cancer [59]. Additionally, an increase in magnesium concentration in lung cancer tissues can enhance the anti-tumor immunity of T cells [10]. In cell experiments, magnesium can effectively inhibit ovarian cancer and osteosarcoma cells [60,61]. However, these basic research findings still require further clinical trials to confirm their therapeutic efficacy.

### 5. Mechanisms of Magnesium Ion Therapy in Tumor Treatment

#### 5.1. The Impact of Magnesium Ions on Mitochondrial Function

The intracellular magnesium ion distribution is regulated primarily by various magnesium ion transporter families. Mrs2 and solute carrier family 41 member A3 (SLC41A3) are the main magnesium transporters on the mitochondrial membrane. In 2016, Mastroiuto et al. [62] first observed that the overexpression of SLC41A3 in human embryonic kidney 293 (HEK293) cells led to a 60% increase in mitochondrial magnesium ion efflux compared to that in normal cells. SLC41A3 functions as a  $\text{Na}^+/\text{Mg}^{2+}$  exchanger, and the efflux of magnesium ions depends not only on the sodium ion concentration but also on temperature. At a temperature of 16 °C, the ion flow nearly disappears [62]. Moreover, overexpression of SLC41A3 may deplete mitochondrial magnesium ions, thereby impairing cellular respiration and reducing mitochondrial ATP production [63]. The endoplasmic reticulum and mitochondria are important storage sites for magnesium ions within cells, and another mitochondrial magnesium ion transporter, Mrs2, acts as a messenger facili-

tating magnesium ion transfer between these two organelles. Under lactate stimulation, Mrs2 can transport magnesium ions from the endoplasmic reticulum to the mitochondria, further stimulating mitochondrial ROS production [41].

The mitochondrial magnesium ion regulation system composed of SLC41A3 and Mrs2 allows cells to flexibly adjust magnesium ion levels within mitochondria under different physiological and environmental conditions to meet the metabolic needs of the cell. The functions and regulatory mechanisms of Mrs2 and SLC41A3 provide essential safeguards for maintaining cellular energy balance and normal metabolism, underscoring their irreplaceable roles in cell survival and functional maintenance.

#### 5.1.1. The Impact of Magnesium Ions on Mitochondrial Energy Metabolism

In 2016, Ha et al. discovered [64] that magnesium-rich seawater effectively increases the activity of mitochondrial enzymes and the expression of mtDNA in adipocytes, suggesting a potential regulatory and antiobesity role of magnesium ions. As a crucial co-factor for many key enzymes in glucose metabolism pathways, magnesium ions play a significant regulatory role in the energy metabolism of mitochondria. Free magnesium ions can indirectly stimulate the dephosphorylation of pyruvate dehydrogenase (PDH) by pyruvate dehydrogenase phosphatase, thereby upregulating the activity of the pyruvate decarboxylase component within the pyruvate dehydrogenase complex (PDHC). This, in turn, affects ATP synthesis by enhancing the activity of the rate-limiting enzyme in the tricarboxylic acid cycle, the  $\alpha$ -ketoglutarate dehydrogenase complex (OGDC), ultimately regulating cellular energy metabolism [65].


#### 5.1.2. The Impact of Magnesium Ions on Oxidative Stress

In both hypertonic and hypotonic environments, magnesium ions and potassium ions play a protective role in the permeability of the mitochondrial outer membrane, effectively safeguarding mitochondrial activity [66]. Conversely, magnesium deficiency inhibits the magnesium transporter protein on the mitochondrial membrane (Mrs2) [67]. As a result, the magnesium ion content within the mitochondria decreases, leading to reduced electron transport chain activity, increased mitochondrial ROS, inhibition of key antioxidant enzymes, and subsequent initiation of oxidative stress [19]. Simultaneously, magnesium ion deficiency, which antagonize calcium ions, increases intracellular calcium overload, activating numerous calcium-dependent kinases and proteins, such as nitric oxide synthase and calcium-dependent calcium-binding proteins, further augmenting ROS production [68]. Additionally, magnesium-deficiency-induced ROS overload activates transcription factors such as nuclear factor kappa-B (NF- $\kappa$ B), which induces lipid peroxidation and stimulates the secretion of proinflammatory cytokines, including interleukin (IL)-1, IL-6, and tumor necrosis factor (TNF)- $\alpha$ , triggering inflammatory responses and exacerbating ROS production [69]. Supplementation with magnesium sulfate has been shown to effectively protect newborn neurons by preserving mitochondrial respiration and reducing ROS production and inflammation [15].

#### 5.1.3. Relationships between Mitochondrial Function and Cancer

The well-known Warburg effect suggests that tumor cells preferentially utilize glycolysis for glucose metabolism, indicating a potential association between mitochondrial dysfunction and the pathogenesis of cancer. Due to the limited energy provided by glycolysis, tumor cells require increased mitochondrial activity to meet their energy demands [70]. Hasumi [71] and Lang [72] confirmed a substantial increase in mitochondrial quantity in malignant pheochromocytoma, accompanied by a significant increase in respiratory frequency compared to that in normal cells. Cannino et al. [73] observed an increased number and enlarged volume of mitochondria in tumor cells, with notable structural differences from those of normal mitochondria, including swelling, shrinkage, and alterations in the outer membrane. These abnormalities further impact mitochondrial respiratory function and stimulate the accumulation of lactate.

Therefore, mitochondria have emerged as novel targets for cancer therapy. Magnesium can inhibit cancer cell energy supply, promote apoptosis, and impede tumor proliferation and invasion [74] by suppressing mitochondrial respiratory chain activity [75], inducing excessive ROS production [76], damaging mitochondrial DNA [77], and interfering with tumor cell metabolic pathways [78]. Additionally, magnesium supplementation can protect the mitochondrial membrane integrity of normal cells and stabilize mitochondrial DNA, reducing the production of ROS, thereby preventing oxidative stress and inflammatory responses, as well as ultimately contributing to tumor prevention (Figure 2).



**Figure 2.** The potential application of magnesium ions in cancer therapy. This figure illustrates the potential application of magnesium ions in cancer therapy. By blocking the Mrs2 protein in tumor cells, the magnesium content in mitochondria will decrease. This deficiency will further reduce electron transport chain activity, increase ROS production, and activate inflammatory responses. For tumor cells, excessive oxidative stress and inflammation will induce cell death. Additionally, mitochondrial magnesium deficiency will lead to reduced ATP production, causing energy deprivation and subsequent cell death in tumor cells. Conversely, supplementing with adequate magnesium can increase intracellular magnesium ion content, enhance mitochondrial magnesium levels, reduce ROS production, and protect mitochondrial membranes and mitochondrial DNA. This helps maintain mitochondrial function stability and prevent tumorigenesis. Green arrows indicate a decrease in the substance content, while red arrows indicate an increase in the substance content. **Abbreviations:** TRPM, transient receptor potential melastatin channel protein family; MagT, magnesium transporter proteins; Mrs2, mitochondrial RNA splicing 2 family genes; PDHC, pyruvate dehydrogenase complex; OGDC, oxoglutarate dehydrogenase complex; ROS, reactive oxygen species; NF-κB, nuclear factor kappa-B; IL, interleukin; TNF-α, tumor necrosis factor. By Figdraw 2.0.



### 5.2. Magnesium Ions and Inflammation

Magnesium ions participate in immune responses through various mechanisms. Magnesium maintains the normal function of cell surface proteins on CD8<sup>+</sup> T cells and regulates T-cell-mediated immune responses, which are crucial for cancer immunotherapy [10]. Additionally, magnesium is an essential cofactor for the synthesis of immunoglobulins, C3 convertase, and the adhesion of immune cells [79]. Magnesium deficiency can increase chronic inflammation; enhance immune stress; and lead to the activation of TNF, IL-1, and IL-6, thereby stimulating tumor proliferation [68]. Therefore, maintaining adequate magnesium levels is crucial for cancer patients.

### 5.3. Magnesium Regulates Apoptosis in Tumor Cells

According to research by Wei et al. [80], magnesium can activate the adenosine 5'-monophosphate (AMP)-activated protein kinase (AMPK)/mammalian target of rapamycin (mTOR)/ULK1 pathway, thereby participating in the activation of the autophagy pathway and inducing apoptosis in osteosarcoma cells. Furthermore, since magnesium can degrade into magnesium ions, hydroxide ions, and hydrogen gas, it can partially block the cell cycle, inhibit the proliferation of gallbladder cancer cells, and induce apoptosis [81]. The research of Yuan et al. [82] also confirmed that magnesium isoglycyrrhizinate can inhibit the apoptosis of bladder cancer cells HTB9 and BIU87 and suppress bladder cancer progression by upregulating miR-26b expression and inactivating the NADPH oxidase 4 (NOX4)/NF- $\kappa$ B/HIF-1 $\alpha$  signaling pathway. In breast cancer cells, an increase in intracellular magnesium similarly induces apoptosis in MCF-7 cells [49]. Therefore, magnesium may exert antitumor effects by inducing apoptosis.

## 6. Prospects of Magnesium Ions in Tumor Therapy

### 6.1. Clinical Applications of Magnesium in Cancer Therapy

Currently, magnesium ions have been employed as anti-tumor implant materials for interventional therapy. During degradation, magnesium generates magnesium hydroxide, hydrogen gas, and magnesium ions. Magnesium hydroxide alkalinizes the acidic tumor microenvironment, while hydrogen gas and magnesium ions exhibit significant anti-inflammatory and antioxidant effects [83].


An increase in the blood magnesium concentration can inhibit mitochondrial oxidative stress, restraining tumor cell proliferation and invasion [84]. Additionally, magnesium ions can modulate the function of T cells, promoting the activation of immune responses and inhibiting tumor growth [10]. Regrettably, the clinical application of magnesium in cancer therapy remains rather limited, despite some clinical trials having confirmed its inhibitory effects on tumor progression. As a key “switch” in regulating tumor growth, magnesium ions are poised to become a new target for cancer therapy and hold potential for widespread clinical applications in the future.

### 6.2. Other Roles of Magnesium Ions in Cancer Therapy

In addition to directly inhibiting tumor growth, magnesium can be used in the treatment of cancer patients for pain relief, antispasmodic effects, and thrombosis prevention. According to the study by Na et al. [85], intraoperative intravenous injection of magnesium sulfate can effectively reduce the hypercoagulable state in patients undergoing laparoscopic colorectal cancer surgery. Additionally, Wu et al. [86] found that oral administration of magnesium-L-threonate can enhance the analgesic effect of opioids, reduce the required dosage of opioid drugs, and alleviate opioid-induced constipation, thereby improving the quality of life for patients with advanced cancer. For patients undergoing mastectomy, the administration of 250 mL of magnesium sulfate at 50 mg/kg prior to anesthesia induction can reduce postoperative analgesic requirements and alleviate postoperative pain to some extent [87]. Furthermore, because magnesium ions relieve smooth muscle spasms, postoperative administration of magnesium sulfate can help alleviate catheter-related bladder discomfort in surgical patients [88,89].

## 7. Conclusions

In general, the dysfunction of mitochondria is closely associated with the occurrence and progression of tumors. Given the pivotal role of mitochondria as the “engine” in cellular energy metabolism, targeted regulation of mitochondrial function is poised to emerge as a novel direction in cancer therapy. Magnesium ions can play a role in the prevention and treatment of tumors by influencing ROS production, oxidative stress levels, mtDNA stability, and the activity of key mitochondrial metabolic enzymes. Magnesium ion therapy holds promise as a novel therapeutic modality for metabolic tumors, offering advantages such as low toxicity and anti-inflammatory properties, thus exhibiting significant clinical application potential (Figure 3).



**Figure 3.** The relationship between low-magnesium ions and cancers. Magnesium deficiency will affect the tricarboxylic acid (TCA) cycle, promoting lactate accumulation. It also leads to elevated calcium ions, which enhances oxidative stress. The production of ROS will further upregulate NF- $\kappa$ B, stimulating the production of inflammatory factors and lipid peroxidation. These effects may potentially lead to tumorigenesis. Green arrows indicate an increase in the substance content, while red arrows indicate a decrease in the substance content.

**Author Contributions:** Conceptualization, L.H., Y.Q. and L.L.; writing—review and editing, L.H., R.L. and J.C.; supervision, Y.Q. and L.L. All authors have read and agreed to the published version of the manuscript.

**Funding:** This study was funded by grants from the Natural Science Foundation of Fujian, China (grant no. 2018J0838, Ling Lin); the Natural Science Foundation of Fujian, China (grant no. 2022J011036,

Leyi Huang); and the Startup Fund for scientific research, Fujian Medical University (grant no. 2021QH1172, Leyi Huang).

**Institutional Review Board Statement:** This manuscript is a review article and does not involve a research protocol requiring approval from the relevant institutional review board or ethics committee.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** We express our gratitude to Home for Researchers for providing the platform for creating figures. We also thank Yuxuan Huang, Weibin Lin, Yihan Shen, Shaoqing Zheng, Zeyu Wu, Yiyang Wang, and Sihao Chen from Fujian Medical University for their assistance in literature collection and figure creation. Above all, the authors are grateful to all the patients and investigators who contribute to the progress of medicine.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## Abbreviations

ROS	Reactive oxygen species
TRPM	Transient receptor potential melastatin channel protein family
SLC	Human solute carrier superfamily
MagT	Magnesium transporter proteins
CNNM	Cyclin M family proteins
Mrs2	Mitochondrial RNA splicing 2 family genes
PDHC	Pyruvate dehydrogenase complex
OGDC	Oxoglutarate dehydrogenase complex
NOS	Nitric oxide synthase
LPO	Lipid hydroperoxide
NF- $\kappa$ B	Nuclear factor kappa-B
IL	Interleukin
TNF	Tumor necrosis factor
HE	Hematoxylin–eosin staining

## References

- de Baaij, J.H.; Hoenderop, J.G.; Bindels, R.J. Magnesium in man: Implications for health and disease. *Physiol. Rev.* **2015**, *95*, 1–46. [[CrossRef](#)] [[PubMed](#)]
- Al Alawi, A.M.; Al Badi, A.; Al Huraizi, A.; Falhammar, H. Magnesium: The recent research and developments. *Adv. Food Nutr. Res.* **2021**, *96*, 193–218. [[PubMed](#)]
- Fiorentini, D.; Cappadone, C.; Farruggia, G.; Prata, C. Magnesium: Biochemistry, Nutrition, Detection, and Social Impact of Diseases Linked to Its Deficiency. *Nutrients* **2021**, *13*, 1136. [[CrossRef](#)] [[PubMed](#)]
- Huang, L. *Study on the Relationship between the Changes of Retinal Dopamine Neurons and Magnesium Ions in 6-OHDA-Induced Parkinsonian Rats*; Fujian Medical University: Fuzhou, China, 2021. [[CrossRef](#)]
- Zhang, Q.; Chen, C.; Zou, X.; Wu, W.; Di, Y.; Li, N.; Fu, A. Iron promotes ovarian cancer malignancy and advances platinum resistance by enhancing DNA repair via FTH1/FTL/POLQ/RAD51 axis. *Cell Death Dis.* **2024**, *15*, 329. [[CrossRef](#)] [[PubMed](#)]
- Lin, Q.; Fan, H.; Zou, Y.; Chen, S. Research progress of copper and its compounds in anti-tumor therapy. *Chin. J. Oncol. Prev. Treat.* **2023**, *15*, 328–334.
- Bois, P.; Sandborn, E.; Messier, P. A study of thymic lymphosarcoma developing in magnesium-deficient rats. *Cancer Res.* **1969**, *29*, 763–775. [[PubMed](#)]
- Auwerckx, J.; Rybarczyk, P.; Kischel, P.; Dhennin-Duthille, I.; Chatelain, D.; Sevestre, H.; Van Seuning, I.; Ouadid-Ahidouch, H.; Jonckheere, N.; Gautier, M. Mg Transporters in Digestive Cancers. *Nutrients* **2021**, *13*, 210. [[CrossRef](#)]
- Folsom, A.R.; Hong, C.P. Magnesium Intake and Reduced Risk of Colon Cancer in a Prospective Study of Women. *Am. J. Epidemiol.* **2006**, *163*, 232–235. [[CrossRef](#)] [[PubMed](#)]
- Lötscher, J.; Martí I Lindez, A.; Kirchhammer, N.; Cribioli, E.; Giordano Attianese, G.; Trefny, M.; Lenz, M.; Rothschild, S.; Strati, P.; Künzli, M.; et al. Magnesium sensing via LFA-1 regulates CD8 T cell effector function. *Cell* **2022**, *185*, 585–602.e529. [[CrossRef](#)]
- Castiglioni, S.; Maier, J. Magnesium and cancer: A dangerous liason. *Magnes. Res.* **2011**, *24*, S92–S100. [[CrossRef](#)]
- Blaszczak, U.; Duda-Chodak, A. Magnesium: Its role in nutrition and carcinogenesis. *Rocz. Panstw. Zakl. Hig.* **2013**, *64*, 165–171. [[PubMed](#)]

13. Liu, Y.; Li, X.; Zou, Q.; Liu, L.; Zhu, X.; Jia, Q.; Wang, L.; Yan, R. Inhibitory effect of magnesium cantharidate on human hepatoma SMMC-7721 cell proliferation by blocking MAPK signaling pathway. *Xi Bao Yu Fen. Zi Mian Yi Xue Za Zhi=Chin. J. Cell. Mol. Immunol.* **2017**, *33*, 347–351.
14. Liu, Y.; Xu, Y.; Ma, H.; Wang, B.; Xu, L.; Zhang, H.; Song, X.; Gao, L.; Liang, X.; Ma, C. Hepatitis B virus X protein amplifies TGF- $\beta$  promotion on HCC motility through down-regulating PPM1a. *Oncotarget* **2016**, *7*, 33125–33135. [[CrossRef](#)] [[PubMed](#)]
15. Koning, G.; Leverin, A.; Nair, S.; Schwendimann, L.; Ek, J.; Carlsson, Y.; Gressens, P.; Thornton, C.; Wang, X.; Mallard, C.; et al. Magnesium induces preconditioning of the neonatal brain via profound mitochondrial protection. *J. Cereb. Blood Flow. Metab. Off. J. Int. Soc. Cereb. Blood Flow. Metab.* **2019**, *39*, 1038–1055. [[CrossRef](#)] [[PubMed](#)]
16. Wang, X.; Zhu, H.; Hu, J.; Li, H.; Guo, S.; Chen, B.; Liu, C.; Wang, G.; Zhou, F. Magnesium Isoglycyrrhizinate Reduces the Target-Binding Amount of Cisplatin to Mitochondrial DNA and Renal Injury through SIRT3. *Int. J. Mol. Sci.* **2022**, *23*, 13093. [[CrossRef](#)] [[PubMed](#)]
17. Reuter, S.; Gupta, S.; Chaturvedi, M.; Aggarwal, B. Oxidative stress, inflammation, and cancer: How are they linked? *Free Radic. Biol. Med.* **2010**, *49*, 1603–1616. [[CrossRef](#)] [[PubMed](#)]
18. Ciaglia, T.; Vestuto, V.; Bertamino, A.; González-Muñiz, R.; Gómez-Monterrey, I. On the modulation of TRPM channels: Current perspectives and anticancer therapeutic implications. *Front. Oncol.* **2023**, *12*, 1065935. [[CrossRef](#)]
19. Arancibia-Hernández, Y.; Hernández-Cruz, E.; Pedraza-Chaverri, J. Magnesium (Mg) Deficiency, Not Well-Recognized Non-Infectious Pandemic: Origin and Consequence of Chronic Inflammatory and Oxidative Stress-Associated Diseases. *Cell. Physiol. Biochem. Int. J. Exp. Cell. Physiol. Biochem. Pharmacol.* **2023**, *57*, 1–23. [[CrossRef](#)]
20. Gao, Y.; Liao, P. TRPM4 channel and cancer. *Cancer Lett.* **2019**, *454*, 66–69. [[CrossRef](#)]
21. Chen, Y.; Liu, L.; Xia, L.; Wu, N.; Wang, Y.; Li, H.; Chen, X.; Zhang, X.; Liu, Z.; Zhu, M.; et al. TRPM7 silencing modulates glucose metabolic reprogramming to inhibit the growth of ovarian cancer by enhancing AMPK activation to promote HIF-1 $\alpha$  degradation. *J. Exp. Clin. Cancer Res. CR* **2022**, *41*, 44. [[CrossRef](#)]
22. Miller, B. TRPM2 in Cancer. *Cell Calcium* **2019**, *80*, 8–17. [[CrossRef](#)] [[PubMed](#)]
23. Hantute-Ghesquier, A.; Haustrate, A.; Prevorskaya, N.; Lehen'kyi, V.Y. TRPM Family Channels in Cancer. *Pharmaceuticals* **2018**, *11*, 58. [[CrossRef](#)]
24. Schweigel-Röntgen, M.; Kolisek, M. SLC41 transporters—Molecular identification and functional role. *Curr. Top. Membr.* **2014**, *73*, 383–410. [[CrossRef](#)] [[PubMed](#)]
25. Nguyen, Y.T.K.; Ha, H.T.T.; Nguyen, T.H.; Nguyen, L.N. The role of SLC transporters for brain health and disease. *Cell Mol. Life Sci.* **2021**, *79*, 20. [[CrossRef](#)]
26. Nemoto, T.; Tagashira, H.; Kita, T.; Kita, S.; Iwamoto, T. Functional characteristics and therapeutic potential of SLC41 transporters. *J. Pharmacol. Sci.* **2023**, *151*, 88–92. [[CrossRef](#)] [[PubMed](#)]
27. Sahni, J.; Scharenberg, A. The SLC41 family of MgtE-like magnesium transporters. *Mol. Asp. Med.* **2013**, *34*, 620–628. [[CrossRef](#)]
28. Njiaju, U.; Gamazon, E.; Gorsic, L.; Delaney, S.; Wheeler, H.; Im, H.; Dolan, M. Whole-genome studies identify solute carrier transporters in cellular susceptibility to paclitaxel. *Pharmacogenetics Genom.* **2012**, *22*, 498–507. [[CrossRef](#)]
29. Li, Q.; Xiong, D.; Wang, H.; Jin, W.; Ma, Y.; Fan, X. High Expression of SLC41A3 Correlates with Poor Prognosis in Hepatocellular Carcinoma. *Oncotargets Ther.* **2021**, *14*, 2975–2988. [[CrossRef](#)]
30. Ding, H.; Li, Y.; Fang, M.; Chen, J.; Liu, L.; Lu, Z.; Hou, J.; Luo, M. Epigenetic activation of the TUSC3 gene as a potential therapy for XMEN disease. *J. Allergy Clin. Immunol.* **2023**, *151*, 1622–1633. [[CrossRef](#)]
31. Gotru, S.; Mammadova-Bach, E.; Sogkas, G.; Schuhmann, M.; Schmitt, K.; Kraft, P.; Herterich, S.; Mantimin, M.; Pinarci, A.; Beck, S.; et al. MAGT1 Deficiency Dysregulates Platelet Cation Homeostasis and Accelerates Arterial Thrombosis and Ischemic Stroke in Mice. *Arterioscler. Thromb. Vasc. Biol.* **2023**, *43*, 1494–1509. [[CrossRef](#)]
32. Oh, M.; Jang, J.; Lee, J. Polarization of THP-1-Derived Macrophage by Magnesium and MAGT1 Inhibition in Wound Healing. *Arch. Plast. Surg.* **2023**, *50*, 432–442. [[CrossRef](#)] [[PubMed](#)]
33. Zheng, K.; Yang, Q.; Xie, L.; Qiu, Z.; Huang, Y.; Lin, Y.; Tu, L.; Cui, C. Overexpression of MAGT1 is associated with aggressiveness and poor prognosis of colorectal cancer. *Oncol. Lett.* **2019**, *18*, 3857–3862. [[CrossRef](#)] [[PubMed](#)]
34. Li, L.; Zhang, X.; Li, Y.; Xiao, B.; Pei, S.; Jiang, H.; Zhang, X. Transcription factor KLF16 activates MAGT1 to regulate the tumorigenesis and progression of breast cancer. *Int. J. Mol. Med.* **2022**, *50*, 115. [[CrossRef](#)] [[PubMed](#)]
35. Zhang, J.; Yang, X.; Ni, H. Long non-coding RNA FLVCR1-AS1 functions as a ceRNA to aggravate cervical cancer cell growth by the miR-381-3p/MAGT1 axis. *Arch. Gynecol. Obstet.* **2022**, *306*, 2093–2103. [[CrossRef](#)] [[PubMed](#)]
36. Bi, C.; Zhang, X.; Chen, Y.; Dong, Y.; Shi, Y.; Lei, Y.; Lv, D.; Cao, X.; Li, W.; Shi, H. MAGT1 is required for HeLa cell proliferation through regulating p21 expression, S-phase progress, and ERK/p38 MAPK MYC axis. *Cell Cycle (Georget. Tex.)* **2021**, *20*, 2233–2247. [[CrossRef](#)] [[PubMed](#)]
37. Funato, Y.; Miki, H. Molecular function and biological importance of CNNM family Mg<sup>2+</sup> transporters. *J. Biochem.* **2019**, *165*, 219–225. [[CrossRef](#)]
38. Bai, Z.; Feng, J.; Franken, G.A.C.; Al'Saadi, N.; Cai, N.; Yu, A.S.; Lou, L.; Komiya, Y.; Hoenderop, J.G.J.; de Baaij, J.H.F.; et al. CNNM proteins selectively bind to the TRPM7 channel to stimulate divalent cation entry into cells. *PLoS Biol.* **2021**, *19*, e3001496. [[CrossRef](#)] [[PubMed](#)]



39. Hirata, Y.; Funato, Y.; Takano, Y.; Miki, H.  $Mg^{2+}$ -dependent interactions of ATP with the cystathionine- $\beta$ -synthase (CBS) domains of a magnesium transporter. *J. Biol. Chem.* **2014**, *289*, 14731–14739. [[CrossRef](#)]
40. Funato, Y.; Yamazaki, D.; Mizukami, S.; Du, L.; Kikuchi, K.; Miki, H. Membrane protein CNNM4-dependent  $Mg^{2+}$  efflux suppresses tumor progression. *J. Clin. Investig.* **2014**, *124*, 5398–5410. [[CrossRef](#)]
41. Daw, C.; Ramachandran, K.; Enslow, B.; Maity, S.; Bursic, B.; Novello, M.; Rubannelsonkumar, C.; Mashal, A.; Ravichandran, J.; Bakewell, T.; et al. Lactate Elicits ER-Mitochondrial Mg Dynamics to Integrate Cellular Metabolism. *Cell* **2020**, *183*, 474–489.e417. [[CrossRef](#)]
42. Li, M.; Li, Y.; Lu, Y.; Li, J.; Lu, X.; Ren, Y.; Wen, T.; Wang, Y.; Chang, S.; Zhang, X.; et al. Molecular basis of Mg permeation through the human mitochondrial Mrs2 channel. *Nat. Commun.* **2023**, *14*, 4713. [[CrossRef](#)] [[PubMed](#)]
43. He, Z.; Chen, Q.; He, W.; Cao, J.; Yao, S.; Huang, Q.; Zheng, Y. Hepatocellular carcinoma subtypes based on metabolic pathways reveals potential therapeutic targets. *Front. Oncol.* **2023**, *13*, 1086604. [[CrossRef](#)] [[PubMed](#)]
44. Liberti, M.; Locasale, J. The Warburg Effect: How Does it Benefit Cancer Cells? *Trends Biochem. Sci.* **2016**, *41*, 211–218. [[CrossRef](#)] [[PubMed](#)]
45. Madaris, T.; Venkatesan, M.; Maity, S.; Stein, M.; Vishnu, N.; Venkateswaran, M.; Davis, J.; Ramachandran, K.; Uthayabalan, S.; Allen, C.; et al. Limiting Mrs2-dependent mitochondrial Mg uptake induces metabolic programming in prolonged dietary stress. *Cell Rep.* **2023**, *42*, 112155. [[CrossRef](#)] [[PubMed](#)]
46. Tao, M.H.; Dai, Q.; Millen, A.E.; Nie, J.; Edge, S.B.; Trevisan, M.; Shields, P.G.; Freudenheim, J.L. Associations of intakes of magnesium and calcium and survival among women with breast cancer: Results from Western New York Exposures and Breast Cancer (WEB) Study. *Am. J. Cancer Res.* **2016**, *6*, 105–113. [[CrossRef](#)] [[PubMed](#)]
47. Hartwig, A. Role of magnesium in genomic stability. *Mutat. Res.* **2001**, *475*, 113–121. [[CrossRef](#)] [[PubMed](#)]
48. Wolf, F.I.; Maier, J.A.; Nasulewicz, A.; Feillet-Coudray, C.; Simonacci, M.; Mazur, A.; Cittadini, A. Magnesium and neoplasia: From carcinogenesis to tumor growth and progression or treatment. *Arch. Biochem. Biophys.* **2007**, *458*, 24–32. [[CrossRef](#)]
49. Yang, D.K.; Tungalag, T.; Lee, S.J.; Kim, S.J. Methyl Jasmonate-induced Increase in Intracellular Magnesium Promotes Apoptosis in Breast Cancer Cells. *Anticancer. Res.* **2024**, *44*, 1087–1095. [[CrossRef](#)] [[PubMed](#)]
50. Bezerra, D.L.C.; Mendes, P.M.V.; Melo, S.R.S.; Dos Santos, L.R.; Santos, R.O.; Vieira, S.C.; Henriques, G.S.; Freitas, B.; Marreiro, D.D.N. Hypomagnesemia and Its Relationship with Oxidative Stress Markers in Women with Breast Cancer. *Biol. Trace Elem. Res.* **2021**, *199*, 4466–4474. [[CrossRef](#)]
51. Gorczyca, A.M.; He, K.; Xun, P.; Margolis, K.L.; Wallace, J.P.; Lane, D.; Thomson, C.; Ho, G.Y.; Shikany, J.M.; Luo, J. Association between magnesium intake and risk of colorectal cancer among postmenopausal women. *Cancer Causes Control* **2015**, *26*, 1761–1769. [[CrossRef](#)]
52. Polter, E.J.; Onyeaghala, G.; Lutsey, P.L.; Folsom, A.R.; Joshi, C.E.; Platz, E.A.; Prizment, A.E. Prospective Association of Serum and Dietary Magnesium with Colorectal Cancer Incidence. *Cancer Epidemiol. Biomark. Prev.* **2019**, *28*, 1292–1299. [[CrossRef](#)] [[PubMed](#)]
53. Wesselink, E.; Winkels, R.M.; van Baar, H.; Geijsen, A.; van Zutphen, M.; van Halteren, H.K.; Hansson, B.M.E.; Radema, S.A.; de Wilt, J.H.W.; Kampman, E.; et al. Dietary Intake of Magnesium or Calcium and Chemotherapy-Induced Peripheral Neuropathy in Colorectal Cancer Patients. *Nutrients* **2018**, *10*, 398. [[CrossRef](#)] [[PubMed](#)]
54. Wesselink, E.; Kok, D.E.; Bours, M.J.L.; de Wilt, J.H.W.; van Baar, H.; van Zutphen, M.; Geijsen, A.; Keulen, E.T.P.; Hansson, B.M.E.; van den Ouweland, J.; et al. Vitamin D, magnesium, calcium, and their interaction in relation to colorectal cancer recurrence and all-cause mortality. *Am. J. Clin. Nutr.* **2020**, *111*, 1007–1017. [[CrossRef](#)] [[PubMed](#)]
55. Uwitonze, A.M.; Razzaque, M.S. Role of Magnesium in Vitamin D Activation and Function. *J. Am. Osteopath. Assoc.* **2018**, *118*, 181–189. [[CrossRef](#)] [[PubMed](#)]
56. Krishnan, A.V.; Feldman, D. Mechanisms of the anti-cancer and anti-inflammatory actions of vitamin D. *Annu. Rev. Pharmacol. Toxicol.* **2011**, *51*, 311–336. [[CrossRef](#)] [[PubMed](#)]
57. Nielsen, F.H. Magnesium deficiency and increased inflammation: Current perspectives. *J. Inflamm. Res.* **2018**, *11*, 25–34. [[CrossRef](#)] [[PubMed](#)]
58. Li, H.; Feng, X.; Li, H.; Ma, S.; Song, W.; Yang, B.; Jiang, T.; Yang, C. The Supplement of Magnesium Element to Inhibit Colorectal Tumor Cells. *Biol. Trace Elem. Res.* **2023**, *201*, 2895–2903. [[CrossRef](#)] [[PubMed](#)]
59. Shah, S.C.; Zhu, X.; Dai, Q.; Peek, R.M.; Shrubsole, M.J. Magnesium intake is associated with a reduced risk of incident liver cancer, based on an analysis of the NIH-American Association of Retired Persons (NIH-AARP) Diet and Health Study prospective cohort. *Am. J. Clin. Nutrition.* **2021**, *113*, 630–638. [[CrossRef](#)]
60. Zan, R.; Ji, W.; Qiao, S.; Wu, H.; Wang, W.; Ji, T.; Yang, B.; Zhang, S.; Luo, C.; Song, Y.; et al. Biodegradable magnesium implants: A potential scaffold for bone tumor patients. *Sci. China Mater.* **2020**, *64*, 1007–1020. [[CrossRef](#)]
61. Qiao, S.; Wang, Y.; Zan, R.; Wu, H.; Zhang, X. iodegradable Mg Implants Suppress the Growth of Ovarian Tumor. *ACS Biomater. Sci. Eng.* **2020**, *6*, 1755–1763. [[CrossRef](#)]
62. Mastrototaro, L.; Smorodchenko, A.; Aschenbach, J.; Kolisek, M.; Sponder, G. Solute carrier 41A3 encodes for a mitochondrial Mg(2+) efflux system. *Sci. Rep.* **2016**, *6*, 27999. [[CrossRef](#)]



63. Pilchova, I.; Klacanova, K.; Tatarkova, Z.; Kaplan, P.; Racay, P. The Involvement of Mg in Regulation of Cellular and Mitochondrial Functions. *Oxidative Med. Cell. Longev.* **2017**, *2017*, 6797460. [[CrossRef](#)] [[PubMed](#)]
64. Ha, B.; Moon, D.; Kim, H.; Shon, Y. Magnesium and calcium-enriched deep-sea water promotes mitochondrial biogenesis by AMPK-activated signals pathway in 3T3-L1 preadipocytes. *Biomed. Pharmacother.=Biomed. Pharmacother.* **2016**, *83*, 477–484. [[CrossRef](#)]
65. Panov, A.; Scarpa, A. Independent modulation of the activity of alpha-ketoglutarate dehydrogenase complex by  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ . *Biochemistry* **1996**, *35*, 427–432. [[CrossRef](#)]
66. Gorgoglione, V.; Laraspata, D.; La Piana, G.; Marzulli, D.; Lofrumento, N. Protective effect of magnesium and potassium ions on the permeability of the external mitochondrial membrane. *Arch. Biochem. Biophys.* **2007**, *461*, 13–23. [[CrossRef](#)]
67. Liu, M.; Dudley, S. Magnesium, Oxidative Stress, Inflammation, and Cardiovascular Disease. *Antioxidants* **2020**, *9*, 907. [[CrossRef](#)] [[PubMed](#)]
68. Mazur, A.; Maier, J.; Rock, E.; Gueux, E.; Nowacki, W.; Rayssiguier, Y. Magnesium and the inflammatory response: Potential physiopathological implications. *Arch. Biochem. Biophys.* **2007**, *458*, 48–56. [[CrossRef](#)] [[PubMed](#)]
69. Xie, C.; Li, X.; Zhu, J.; Wu, J.; Geng, S.; Zhong, C. Magnesium isoglycyrrhizinate suppresses LPS-induced inflammation and oxidative stress through inhibiting NF- $\kappa$ B and MAPK pathways in RAW264.7 cells. *Bioorganic Med. Chem.* **2019**, *27*, 516–524. [[CrossRef](#)]
70. Wang, Y.; Patti, G. The Warburg effect: A signature of mitochondrial overload. *Trends Cell Biol.* **2023**, *33*, 1014–1020. [[CrossRef](#)]
71. Hasumi, H.; Baba, M.; Hasumi, Y.; Huang, Y.; Oh, H.; Hughes, R.; Klein, M.; Takikita, S.; Nagashima, K.; Schmidt, L.; et al. Regulation of mitochondrial oxidative metabolism by tumor suppressor FLCN. *J. Natl. Cancer Inst.* **2012**, *104*, 1750–1764. [[CrossRef](#)]
72. Lang, M.; Vocke, C.; Merino, M.; Schmidt, L.; Linehan, W. Mitochondrial DNA mutations distinguish bilateral multifocal renal oncocytomas from familial Birt-Hogg-Dubé tumors. *Mod. Pathol. Off. J. United States Can. Acad. Pathol. Inc.* **2015**, *28*, 1458–1469. [[CrossRef](#)] [[PubMed](#)]
73. Cannino, G.; Ciscato, F.; Masgras, I.; Sánchez-Martín, C.; Rasola, A. Metabolic Plasticity of Tumor Cell Mitochondria. *Front. Oncol.* **2018**, *8*, 333. [[CrossRef](#)] [[PubMed](#)]
74. Winter, J.; Yadav, T.; Rutter, J. Stressed to death: Mitochondrial stress responses connect respiration and apoptosis in cancer. *Mol. Cell* **2022**, *82*, 3321–3332. [[CrossRef](#)] [[PubMed](#)]
75. Jain, I.; Zazzeron, L.; Goli, R.; Alexa, K.; Schatzman-Bone, S.; Dhillon, H.; Goldberger, O.; Peng, J.; Shalem, O.; Sanjana, N.; et al. Hypoxia as a therapy for mitochondrial disease. *Science* **2016**, *352*, 54–61. [[CrossRef](#)] [[PubMed](#)]
76. Perillo, B.; Di Donato, M.; Pezone, A.; Di Zazzo, E.; Giovannelli, P.; Galasso, G.; Castoria, G.; Migliaccio, A. ROS in cancer therapy: The bright side of the moon. *Exp. Mol. Med.* **2020**, *52*, 192–203. [[CrossRef](#)] [[PubMed](#)]
77. Li, C.; Zhang, Y.; Liu, J.; Kang, R.; Klionsky, D.; Tang, D. Mitochondrial DNA stress triggers autophagy-dependent ferroptotic death. *Autophagy* **2021**, *17*, 948–960. [[CrossRef](#)] [[PubMed](#)]
78. Lee, J.; Yesilkanal, A.; Wynne, J.; Frankenberger, C.; Liu, J.; Yan, J.; Elbaz, M.; Rabe, D.; Rustandy, F.; Tiwari, P.; et al. Effective breast cancer combination therapy targeting BACH1 and mitochondrial metabolism. *Nature* **2019**, *568*, 254–258. [[CrossRef](#)] [[PubMed](#)]
79. Caruso, A.; Vollmer, J.; Machacek, M.; Kortvely, E. Modeling the activation of the alternative complement pathway and its effects on hemolysis in health and disease. *PLoS Comput. Biol.* **2020**, *16*, e1008139. [[CrossRef](#)] [[PubMed](#)]
80. Wei, X.; Tang, Z.; Wu, H.; Zuo, X.; Dong, H.; Tan, L.; Wang, W.; Liu, Y.; Wu, Z.; Shi, L.; et al. Biofunctional magnesium-coated Ti6Al4V scaffolds promote autophagy-dependent apoptosis in osteosarcoma by activating the AMPK/mTOR/ULK1 signaling pathway. *Mater. Today Bio* **2021**, *12*, 100147. [[CrossRef](#)]
81. Peng, H.; Fan, K.; Zan, R.; Gong, Z.J.; Sun, W.; Sun, Y.; Wang, W.; Jiang, H.; Lou, J.; Ni, J.; et al. Degradable magnesium implants inhibit gallbladder cancer. *Acta Biomater.* **2021**, *128*, 514–522. [[CrossRef](#)]
82. Yuan, Z.; Guo, G.; Sun, G.; Li, Q.; Wang, L.; Qiao, B. Magnesium isoglycyrrhizinate suppresses bladder cancer progression by modulating the miR-26b/Nox4 axis. *Bioengineered* **2022**, *13*, 7986–7999. [[CrossRef](#)] [[PubMed](#)]
83. Zan, R.; Wang, H.; Cai, W.; Ni, J.; Luthringer-Feyerabend, B.; Wang, W.; Peng, H.; Ji, W.; Yan, J.; Xia, J.; et al. Controlled release of hydrogen by implantation of magnesium induces P53-mediated tumor cells apoptosis. *Bioact. Mater.* **2022**, *9*, 385–396. [[CrossRef](#)]
84. Xu, B.; Song, Y.; Yang, K.; Li, Y.; Chen, B.; Liao, X.; Jia, Q. Magnesium metal and its corrosion products: Promising materials for tumor interventional therapy. *J. Magnes. Alloys* **2023**, *11*, 763–775. [[CrossRef](#)]
85. Na, H.S.; Shin, H.J.; Kang, S.B.; Hwang, J.W.; Do, S.H. Effects of magnesium sulphate on coagulation after laparoscopic colorectal cancer surgery, measured by rotational thromboelastometry (ROTEM(R)). *Anaesthesia* **2014**, *69*, 1314–1321. [[CrossRef](#)] [[PubMed](#)]
86. Wu, S.; Jin, T.; Ma, B.; Ji, Y.; Huang, X.; Wang, P.; Liu, X.; Krylov, B.V.; Liu, X.; Ma, K. Oral application of magnesium-L-threonate enhances analgesia and reduces the dosage of opioids needed in advanced cancer patients-A randomized, double-blind, placebo-controlled trial. *Cancer Med.* **2023**, *12*, 4343–4351. [[CrossRef](#)] [[PubMed](#)]
87. Hatice Akbudak, I.; Yilmaz, S.; Ilhan, S.; Yuksel Tanriverdi, S.; Erdem, E. The effect of preemptive magnesium sulfate on postoperative pain in patients undergoing mastectomy: A clinical trial. *Eur. Rev. Med. Pharmacol. Sci.* **2023**, *27*, 7907–7913. [[CrossRef](#)] [[PubMed](#)]

88. Park, J.Y.; Hong, J.H.; Kim, D.H.; Yu, J.; Hwang, J.H.; Kim, Y.K. Magnesium and Bladder Discomfort after Transurethral Resection of Bladder Tumor: A Randomized, Double-blind, Placebo-controlled Study. *Anesthesiology* **2020**, *133*, 64–77. [[CrossRef](#)]
89. Jiang, W.; Zeng, X.; Zhou, X.; Liao, O.; Ju, F.; Zhao, Z.; Zhang, X. Effect of magnesium sulfate perioperative infusion on postoperative catheter-related bladder discomfort in male patients undergoing laparoscopic radical resection of gastrointestinal cancer: A prospective, randomized and controlled study. *BMC Anesth.* **2023**, *23*, 396. [[CrossRef](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.