

A guide to the arithmetic certificate for $K = \mathbf{Q}(\sqrt{-126740891})$, $p = 5$

Generated by `tools/certificate_guide.gp`

What this document is

This guide restates the content of `certificate.gp` in the notation of Section 2.5 of the paper. Each of the 18 entries certifies one value $D_x(e_j)$ of a secondary norm operator: it stores the unramified cyclic quintic L_x/K , the normalized generator σ_x , the pair (a', J) representing the torsion class e_j , the auxiliary ideal I' , the compactly represented element t , a residue prime for the sign check, and the expected norm class $[N_x(I')]$. The two defining equations are

$$I'^{(1-\sigma_x)^2}(t) J \mathcal{O}_{L_x} = \mathcal{O}_{L_x}, \quad N_x(t) = a'.$$

Facts marked **Checked** are recomputed exactly by the generating script; the full verification is performed by the shared verifier `certificate/verify_certificate.c` as described in the README files of the `certificate` directory.

Notation and conventions

Ideals as matrices. An ideal of the ring of integers is stored by its *Hermite normal form* (HNF): an upper-triangular integer matrix whose columns are the coordinates, with respect to the fixed integral basis, of a $\mathbf{Z}$ -basis of the ideal. For example, the ideal $J = \begin{pmatrix} 415 & 176 \\ 0 & 1 \end{pmatrix}$ of Entry 1 below, over K with integral basis $(1, s)$, denotes the ideal $\mathbf{Z}415 + \mathbf{Z}(176 + s)$. The determinant of the matrix is the norm of the ideal. Elements are likewise given by their coordinate vectors in the integral basis.

Which integral basis. The one PARI returns for the field in question – and that basis is LLL-reduced, so it is not canonical: another machine may reduce differently, and the same coordinate vector would then denote a different algebraic number. The certificate therefore records the basis it was written against, as its last field per entry and in the header for K . The verifier expresses the stored basis in its own, checks that the resulting matrix is integral with determinant ± 1 – so that both bases span the same ring of integers – and converts the coordinates before checking anything. Where the two bases agree, that matrix is the identity and nothing changes.

Characters. The header fixes three generators $\kappa_1, \kappa_2, \kappa_3$ of $\text{Cl}(K)$ (as HNF ideals). A character x on $\text{Cl}(K)/5$ is recorded by its value vector $(x(\bar{\kappa}_1), x(\bar{\kappa}_2), x(\bar{\kappa}_3))$; thus $(1, 0, 0)$ is the character dual to $\bar{\kappa}_1$. The column number j records which basis element e_j of $\text{Cl}(K)[5]$ the stored pair (a', J) represents.

Automorphisms. σ_x is a field automorphism of L_x . Writing θ for the chosen root of the stored absolute polynomial, so that $L_x = \mathbf{Q}(\theta)$, the automorphism is determined by the single value $\sigma_x(\theta)$, and the certificate stores the coordinate vector of $\sigma_x(\theta)$ in the integral basis of L_x .

Base field data

The base field is $K = \mathbf{Q}(s)$ with $s^2 - s + 31685223 = 0$, of discriminant -126740891 . Class group invariants $(195 \ 5 \ 5)$, class number 4875; the three stored generator ideals (HNF) are $\begin{pmatrix} 3 & 2 \\ 0 & 1 \end{pmatrix}$, $\begin{pmatrix} 5041 & 4727 \\ 0 & 1 \end{pmatrix}$, $\begin{pmatrix} 4631 & 1002 \\ 0 & 1 \end{pmatrix}$, and the torsion units are generated by -1 of order 2.

Entry 1: character a , column e_1

Prescribed character vector $(1 \ 0 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 4y^4 + (-s - 51)y^3 + (22s - 130606)y^2 + (-73s + 1282752)y + (944s - 8886784)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 8y^9 - 87y^8 - 260778y^7 + 35298066y^6 - 1408732864y^5 + 36955715157y^4 - 495711807334y^3 + 5451302849399y^2 - 27164204420768y + 107202383621888$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 0 \ 0 \ 0 \ -1 \ 0 \ 1 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{3102493}{12309502009375} + \left(\frac{291}{12309502009375}\right)s, \quad J = \begin{pmatrix} 415 & 176 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 415$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $2517218714932 = 2^2 \cdot 629304678733$:

$$\begin{pmatrix} 1258609357466 & 782598244 & 295823383280 & 1040807781051 & 374714318371 & 881523773442 & 446703420967 & 28559662847 & 374797064752 & 1061214248295 \\ 0 & 2 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 7 factors with signed exponents of absolute value up to 2; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 3$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (0 \ 4 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 2: character a , column e_2

Prescribed character vector $(1 \ 0 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 4y^4 + (-s - 51)y^3 + (22s - 130606)y^2 + (-73s + 1282752)y + (944s - 8886784)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$\begin{aligned} & y^{10} - 8y^9 - 87y^8 - 260778y^7 + 35298066y^6 - 1408732864y^5 + \\ & 36955715157y^4 - 495711807334y^3 + 5451302849399y^2 - 2716420442 \\ & 0768y + 107202383621888 \end{aligned}$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 0 \ 0 \ 0 \ -1 \ 0 \ 1 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{219809146}{3255243551009881201} + \left(\frac{318135}{3255243551009881201} \right) s, \quad J = \begin{pmatrix} 5041 & 4727 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 5041$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $1770080412983 = 79 \cdot 22406081177$:

$$\begin{pmatrix} 1770080412983 & 1510714998347 & 1120869054289 & 112190521040 & 339361891663 & 1204542237616 & 782737303535 & 236549689332 & 1220294746920 & 361780794306 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 429 factors with signed exponents of absolute value up to 47181946632638440857675912; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 5$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (0 \ 0 \ 2)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 3: character a , column e_3

Prescribed character vector $(1 \ 0 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 4y^4 + (-s - 51)y^3 + (22s - 130606)y^2 + (-73s + 1282752)y + (944s - 8886784)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$\begin{aligned} & y^{10} - 8y^9 - 87y^8 - 260778y^7 + 35298066y^6 - 1408732864y^5 + \\ & 36955715157y^4 - 495711807334y^3 + 5451302849399y^2 - 2716420442 \\ & 0768y + 107202383621888 \end{aligned}$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 0 \ 0 \ 0 \ -1 \ 0 \ 1 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{993784579}{2129972052005212151} + \left(-\frac{189862}{2129972052005212151}\right)s, \quad J = \begin{pmatrix} 4631 & 1002 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 4631$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $68983994758064 = 2^4 \cdot 17 \cdot 19273 \cdot 13159219$:

$$\begin{pmatrix} 8622999344758 & 6781122146730 & 4516605923366 & 5130270560533 & 2352677385784 & 5332181867908 & 2595910204892 & 3898891182244 & 3260057641759 & 4861137617409 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 1 & 0 & 1 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 2 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 434 factors with signed exponents of absolute value up to 29768691156393232794117728; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 5$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (0 \ 4 \ 2)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 4: character b , column e_1

Prescribed character vector $(0 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 3343y^3 + (-12s + 6)y^2 + 4877203y + (-7872s + 3936)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 6686y^8 + 20930055y^6 + 37171651334y^4 + 29773334866921y^2 + 1963482050497536$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 0 \ -1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{3102493}{12309502009375} + \left(\frac{291}{12309502009375}\right)s, \quad J = \begin{pmatrix} 415 & 176 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 415$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $2310192872676 = 2^2 \cdot 3^3 \cdot 13 \cdot 1645436519$:

$$\begin{pmatrix} 385032145446 & 72463898364 & 185662190274 & 247564315267 & 309220926069 & 305524751703 & 246419462003 & 379415088796 & 330071315739 & 26879822676 \\ 0 & 6 & 2 & 0 & 1 & 0 & 0 & 4 & 0 & 3 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 168 factors with signed exponents of absolute value up to 916676568633958379; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 5$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (2 \ 0 \ 2)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 5: character b , column e_2

Prescribed character vector $(0 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 3343y^3 + (-12s + 6)y^2 + 4877203y + (-7872s + 3936)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 6686y^8 + 20930055y^6 + 37171651334y^4 + 29773334866921y^2 + 1963482050497536$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 0 \ -1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{219809146}{3255243551009881201} + \left(\frac{318135}{3255243551009881201} \right) s, \quad J = \begin{pmatrix} 5041 & 4727 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 5041$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $52984325659248 = 2^4 \cdot 3^3 \cdot 4639 \cdot 26438651$:

$$\begin{pmatrix} 1471786823868 & 139866380796 & 547896065202 & 242670393129 & 1144414346761 & 731424363391 & 40963888507 & 980917637800 & 385201276751 & 99080988602 \\ 0 & 12 & 0 & 4 & 9 & 2 & 4 & 4 & 4 & 7 \\ 0 & 0 & 3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 173 factors with signed exponents of absolute value up to 586528490631369531; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 5$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (4 \ 0 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 6: character b , column e_3

Prescribed character vector $(0 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 3343y^3 + (-12s + 6)y^2 + 4877203y + (-7872s + 3936)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 6686y^8 + 20930055y^6 + 37171651334y^4 + 29773334866921y^2 + 1963482050497536$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 0 \ -1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{993784579}{2129972052005212151} + \left(-\frac{189862}{2129972052005212151} \right) s, \quad J = \begin{pmatrix} 4631 & 1002 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 4631$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $791337809793024 = 2^{12} \cdot 3^3 \cdot 13^2 \cdot 181 \cdot 233923$:

$$\begin{pmatrix} 118890896904 & 106903926584 & 78981328684 & 16017718450 & 16397287810 & 110003149150 & 63538484455 & 19080543696 & 22351785982 & 36465493692 \\ 0 & 104 & 8 & 96 & 26 & 60 & 16 & 96 & 32 & 80 \\ 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 168 factors with signed exponents of absolute value up to 1526123369717270750; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 5$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (2 \ 0 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 7: character c , column e_1

Prescribed character vector $(0 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 2y^4 + (-s + 1257)y^3 + (10s + 156160)y^2 + (77s - 563510)y + (-186s - 499926)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$\begin{aligned} & y^{10} - 4y^9 + 2517y^8 + 307304y^7 + 31512412y^6 - 240007962y^5 \\ & + 21262501393y^4 - 116663455444y^3 + 231321854097y^2 - 3440986085 \\ & 34y + 1346200966620 \end{aligned}$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(1 \ -1 \ -1 \ -1 \ 1 \ 0 \ 0 \ 0 \ -1 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{3102493}{12309502009375} + \left(\frac{291}{12309502009375} \right) s, \quad J = \begin{pmatrix} 415 & 176 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 415$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $43691997566160 = 2^4 \cdot 3^6 \cdot 5 \cdot 773 \cdot 969181$:

$$\begin{pmatrix} 67425922170 & 18635752350 & 65079686430 & 23704313484 & 57490566268 & 44112431072 & 60551202973 & 19707348334 & 55412101289 & 29530809219 \\ 0 & 18 & 0 & 6 & 8 & 8 & 6 & 5 & 4 & 6 \\ 0 & 0 & 6 & 0 & 5 & 0 & 0 & 4 & 0 & 1 \\ 0 & 0 & 0 & 6 & 5 & 4 & 0 & 2 & 0 & 3 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 175 factors with signed exponents of absolute value up to 68363451248169; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 11$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (0 \ 2 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 8: character c , column e_2

Prescribed character vector $(0 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 2y^4 + (-s + 1257)y^3 + (10s + 156160)y^2 + (77s - 563510)y + (-186s - 499926)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$\begin{aligned} & y^{10} - 4y^9 + 2517y^8 + 307304y^7 + 31512412y^6 - 240007962y^5 \\ & + 21262501393y^4 - 116663455444y^3 + 231321854097y^2 - 3440986085 \\ & 34y + 1346200966620 \end{aligned}$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(1 \ -1 \ -1 \ -1 \ 1 \ 0 \ 0 \ 0 \ -1 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{219809146}{3255243551009881201} + \left(\frac{318135}{3255243551009881201} \right) s, \quad J = \begin{pmatrix} 5041 & 4727 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 5041$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $4738365657936 = 2^4 \cdot 3^5 \cdot 41 \cdot 3529 \cdot 8423$:

$$\begin{pmatrix} 21936878046 & 14301118134 & 4266067230 & 18788553654 & 11090729859 & 21500582275 & 19103428410 & 21895977265 & 13777821995 & 5232960196 \\ 0 & 6 & 0 & 0 & 0 & 0 & 2 & 2 & 4 & 5 \\ 0 & 0 & 6 & 0 & 0 & 0 & 2 & 2 & 0 & 3 \\ 0 & 0 & 0 & 2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 3 & 0 & 2 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 184 factors with signed exponents of absolute value up to 121393102922285; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 11$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (4 \ 0 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 9: character c , column e_3

Prescribed character vector $(0 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 2y^4 + (-s + 1257)y^3 + (10s + 156160)y^2 + (77s - 563510)y + (-186s - 499926)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$\begin{aligned} & y^{10} - 4y^9 + 2517y^8 + 307304y^7 + 31512412y^6 - 240007962y^5 \\ & + 21262501393y^4 - 116663455444y^3 + 231321854097y^2 - 3440986085 \\ & 34y + 1346200966620 \end{aligned}$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(1 \ -1 \ -1 \ -1 \ 1 \ 0 \ 0 \ 0 \ -1 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{993784579}{2129972052005212151} + \left(-\frac{189862}{2129972052005212151}\right)s, \quad J = \begin{pmatrix} 4631 & 1002 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 4631$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $4738365657936 = 2^4 \cdot 3^5 \cdot 41 \cdot 3529 \cdot 8423$:

$$\begin{pmatrix} 21936878046 & 14301118134 & 4266067230 & 18788553654 & 11090729859 & 21500582275 & 19103428410 & 21895977265 & 13777821995 & 5232960196 \\ 0 & 6 & 0 & 0 & 0 & 0 & 2 & 2 & 4 & 5 \\ 0 & 0 & 6 & 0 & 0 & 0 & 2 & 2 & 0 & 3 \\ 0 & 0 & 0 & 2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 3 & 0 & 2 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 185 factors with signed exponents of absolute value up to 66477891663197; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 11$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of +1. Certified value: $D_x(e_3) = (4 \ 0 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 10: character $a + b$, column e_1

Prescribed character vector $(1 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 1448y^3 + (-14s + 7)y^2 - 721792y + (13184s - 6592)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 2896y^8 + 653120y^6 + 8300613291y^4 - 11175679657344y^2 + 5507457485287424$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 1 \ 0 \ -1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{3102493}{12309502009375} + \left(\frac{291}{12309502009375}\right)s, \quad J = \begin{pmatrix} 415 & 176 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 415$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $41352840998453 = 19391 \cdot 31123 \cdot 68521$:

$$\begin{pmatrix} 41352840998453 & 40977124414409 & 2954119126460 & 36216802919142 & 24950752149833 & 8245624489039 & 31609257452534 & 31425286576557 & 26519198189036 & 9481127578612 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 738 factors with signed exponents of absolute value up to 438725811777816756244842555562531376; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 11$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of +1. Certified value: $D_x(e_1) = (3 \ 2 \ 1)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 11: character $a + b$, column e_2

Prescribed character vector $(1 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 1448y^3 + (-14s + 7)y^2 - 721792y + (13184s - 6592)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 2896y^8 + 653120y^6 + 8300613291y^4 - 11175679657344y^2 + 5507457485287424$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 1 \ 0 \ -1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{219809146}{3255243551009881201} + \left(\frac{318135}{3255243551009881201} \right) s, \quad J = \begin{pmatrix} 5041 & 4727 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 5041$. **Checked:** $(a')^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $6861571930844 = 2^2 \cdot 191161 \cdot 8973551$:

$$\begin{pmatrix} 3430785965422 & 2806524789482 & 1362905018369 & 386679145455 & 1398429903459 & 440085774721 & 878042214922 & 2606445249317 & 2443686250258 & 673814074885 \\ 0 & 2 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 738 factors with signed exponents of absolute value up to 250641831182934151437263084473360029; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 11$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (4 \ 1 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 12: character $a + b$, column e_3

Prescribed character vector $(1 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 1448y^3 + (-14s + 7)y^2 - 721792y + (13184s - 6592)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 2896y^8 + 653120y^6 + 8300613291y^4 - 11175679657344y^2 + 5507457485287424$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 1 \ 0 \ -1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{993784579}{2129972052005212151} + \left(-\frac{189862}{2129972052005212151}\right)s, \quad J = \begin{pmatrix} 4631 & 1002 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 4631$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $81484156074064 = 2^4 \cdot 519091 \cdot 9810919$:

$$\begin{pmatrix} 10185519509258 & 4610258658268 & 5354354227806 & 6991330801755 & 5591520358224 & 8111059034923 & 5031267787071 & 2395650226613 & 7541295183605 & 5317919191739 \\ 0 & 2 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 2 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 738 factors with signed exponents of absolute value up to 226667133424010189826279948868396661; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 11$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (3 \ 2 \ 3)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 13: character $a + c$, column e_1

Prescribed character vector $(1 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 2y^4 - 1584y^3 + 184401y^2 + (-332s - 7592554)y + (17296s + 100545784)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 4y^9 - 3164y^8 + 375138y^7 - 13413988y^6 - 352702624y^5 \\ + 57655248033y^4 - 3118766762016y^3 + 98226544621260y^2 - 1890853 \\ 628692256y + 19589879383825088$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 0 \ -1 \ -1 \ -1 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{3102493}{12309502009375} + \left(\frac{291}{12309502009375}\right)s, \quad J = \begin{pmatrix} 415 & 176 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 415$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $925818819357461 = 71 \cdot 263 \cdot 269 \cdot 184314553$:

$$\begin{pmatrix} 925818819357461 & 357809260895362 & 841987457645357 & 747458974860657 & 202630608866020 & 49494708977954 & 393208806204374 & 33768649003996 & 790702762627633 & 745356448182395 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 376 factors with signed exponents of absolute value up to 2200257927144; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 3$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (1 \ 2 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 14: character $a + c$, column e_2

Prescribed character vector $(1 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 2y^4 - 1584y^3 + 184401y^2 + (-332s - 7592554)y + (17296s + 100545784)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 4y^9 - 3164y^8 + 375138y^7 - 13413988y^6 - 352702624y^5 \\ + 57655248033y^4 - 3118766762016y^3 + 98226544621260y^2 - 1890853 \\ 628692256y + 19589879383825088$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 0 \ -1 \ -1 \ -1 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{219809146}{3255243551009881201} + \left(\frac{318135}{3255243551009881201} \right) s, \quad J = \begin{pmatrix} 5041 & 4727 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 5041$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $67523144841488 = 2^4 \cdot 7^2 \cdot 181 \cdot 647 \cdot 735451$:

$$\begin{pmatrix} 1205770443598 & 675329168994 & 54975896324 & 750477486612 & 329087020088 & 55919660406 & 622745718921 & 1038544992322 & 1041552168861 & 869585195634 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 14 & 0 & 9 & 3 & 11 & 0 & 9 \\ 0 & 0 & 0 & 0 & 2 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 376 factors with signed exponents of absolute value up to 2475892142333; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 3$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (2 \ 1 \ 3)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 15: character $a + c$, column e_3

Prescribed character vector $(1 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 2y^4 - 1584y^3 + 184401y^2 + (-332s - 7592554)y + (17296s + 100545784)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$\begin{aligned} & y^{10} - 4y^9 - 3164y^8 + 375138y^7 - 13413988y^6 - 352702624y^5 \\ & + 57655248033y^4 - 3118766762016y^3 + 98226544621260y^2 - 1890853 \\ & 628692256y + 19589879383825088 \end{aligned}$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 0 \ -1 \ -1 \ -1 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{993784579}{2129972052005212151} + \left(-\frac{189862}{2129972052005212151} \right) s, \quad J = \begin{pmatrix} 4631 & 1002 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 4631$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $2173202744215724 = 2^2 \cdot 13^2 \cdot 683 \cdot 4706876953$:

$$\begin{pmatrix} 1086601372107862 & 889783596872784 & 178905456239736 & 947155631605467 & 703368614732151 & 741830416383116 & 287360132477867 & 969881359777380 & 1039919908294475 & 493121059999661 \\ 0 & 2 & 0 & 1 & 1 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 376 factors with signed exponents of absolute value up to 2826787881291; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 3$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (2 \ 0 \ 3)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 16: character $b + c$, column e_1

Prescribed character vector $(0 \ 1 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 3371y^3 + (12s - 6)y^2 + 1686955y + (-1956s + 978)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 6742y^8 + 14737551y^6 - 6810778534y^4 + 1358386075249y^2 + 121225634387244$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{3102493}{12309502009375} + \left(\frac{291}{12309502009375}\right)s, \quad J = \begin{pmatrix} 415 & 176 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 415$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $19095996670704 = 2^4 \cdot 3^3 \cdot 163 \cdot 853 \cdot 317923$:

$$\begin{pmatrix} 795666527946 & 50581444248 & 618726207240 & 413828566666 & 254329067945 & 714243757729 & 565712375551 & 397215881105 & 72194133087 & 549256689123 \\ 0 & 6 & 0 & 4 & 5 & 0 & 2 & 2 & 5 & 5 \\ 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 322 factors with signed exponents of absolute value up to 73996438; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 5$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (1 \ 0 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 17: character $b + c$, column e_2

Prescribed character vector $(0 \ 1 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 3371y^3 + (12s - 6)y^2 + 1686955y + (-1956s + 978)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 6742y^8 + 14737551y^6 - 6810778534y^4 + 1358386075249y^2 + 121225634387244$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{219809146}{3255243551009881201} + \left(\frac{318135}{3255243551009881201} \right) s, \quad J = \begin{pmatrix} 5041 & 4727 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 5041$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $101681239707324 = 2^2 \cdot 3^6 \cdot 41 \cdot 163 \cdot 5217733$:

$$\begin{pmatrix} 627661973502 & 513565071063 & 32688214848 & 252257413392 & 266483833300 & 403564942337 & 515256888573 & 125387165426 & 574009040640 & 551650777958 \\ 0 & 3 & 0 & 2 & 0 & 2 & 0 & 2 & 2 & 0 \\ 0 & 0 & 9 & 3 & 8 & 7 & 6 & 1 & 1 & 8 \\ 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 3 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 308 factors with signed exponents of absolute value up to 66703973; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 5$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (4 \ 1 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 18: character $b + c$, column e_3

Prescribed character vector $(0 \ 1 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 3371y^3 + (12s - 6)y^2 + 1686955y + (-1956s + 978)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 6742y^8 + 14737551y^6 - 6810778534y^4 + 1358386075249y^2 + 121225634387244$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{993784579}{2129972052005212151} + \left(-\frac{189862}{2129972052005212151}\right)s, \quad J = \begin{pmatrix} 4631 & 1002 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 4631$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $320462439669756 = 2^2 \cdot 3^8 \cdot 41 \cdot 297826439$:

$$\begin{pmatrix} 659387735946 & 381308619558 & 314026880400 & 515409886578 & 517692919915 & 539071366721 & 239337232470 & 163353702779 & 516330574221 & 248760653069 \\ 0 & 6 & 0 & 0 & 1 & 4 & 0 & 0 & 4 & 1 \\ 0 & 0 & 9 & 0 & 5 & 1 & 6 & 7 & 4 & 5 \\ 0 & 0 & 0 & 3 & 1 & 2 & 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 3 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 333 factors with signed exponents of absolute value up to 1419641881; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 5$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (3 \ 1 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.