

A guide to the arithmetic certificate for $K = \mathbf{Q}(\sqrt{-207666763})$, $p = 5$

Generated by `tools/certificate_guide.gp`

What this document is

This guide restates the content of `certificate.gp` in the notation of Section 2.5 of the paper. Each of the 18 entries certifies one value $D_x(e_j)$ of a secondary norm operator: it stores the unramified cyclic quintic L_x/K , the normalized generator σ_x , the pair (a', J) representing the torsion class e_j , the auxiliary ideal I' , the compactly represented element t , a residue prime for the sign check, and the expected norm class $[N_x(I')]$. The two defining equations are

$$I'^{(1-\sigma_x)^2}(t) J \mathcal{O}_{L_x} = \mathcal{O}_{L_x}, \quad N_x(t) = a'.$$

Facts marked **Checked** are recomputed exactly by the generating script; the full verification is performed by the shared verifier `certificate/verify_certificate.c` as described in the README files of the `certificate` directory.

Notation and conventions

Ideals as matrices. An ideal of the ring of integers is stored by its *Hermite normal form* (HNF): an upper-triangular integer matrix whose columns are the coordinates, with respect to the fixed integral basis, of a $\mathbf{Z}$ -basis of the ideal. For example, the ideal $J = \begin{pmatrix} 2629 & 2019 \\ 0 & 1 \end{pmatrix}$ of Entry 1 below, over K with integral basis $(1, s)$, denotes the ideal $\mathbf{Z} 2629 + \mathbf{Z} (2019 + s)$. The determinant of the matrix is the norm of the ideal. Elements are likewise given by their coordinate vectors in the integral basis.

Which integral basis. The one PARI returns for the field in question – and that basis is LLL-reduced, so it is not canonical: another machine may reduce differently, and the same coordinate vector would then denote a different algebraic number. The certificate therefore records the basis it was written against, as its last field per entry and in the header for K . The verifier expresses the stored basis in its own, checks that the resulting matrix is integral with determinant ± 1 – so that both bases span the same ring of integers – and converts the coordinates before checking anything. Where the two bases agree, that matrix is the identity and nothing changes.

Characters. The header fixes three generators $\kappa_1, \kappa_2, \kappa_3$ of $\text{Cl}(K)$ (as HNF ideals). A character x on $\text{Cl}(K)/5$ is recorded by its value vector $(x(\bar{\kappa}_1), x(\bar{\kappa}_2), x(\bar{\kappa}_3))$; thus $(1, 0, 0)$ is the character dual to $\bar{\kappa}_1$. The column number j records which basis element e_j of $\text{Cl}(K)[5]$ the stored pair (a', J) represents.

Automorphisms. σ_x is a field automorphism of L_x . Writing θ for the chosen root of the stored absolute polynomial, so that $L_x = \mathbf{Q}(\theta)$, the automorphism is determined by the single value $\sigma_x(\theta)$, and the certificate stores the coordinate vector of $\sigma_x(\theta)$ in the integral basis of L_x .

Base field data

The base field is $K = \mathbf{Q}(s)$ with $s^2 - s + 51916691 = 0$, of discriminant -207666763 . Class group invariants $(50 \ 10 \ 5)$, class number 2500; the three stored generator ideals (HNF) are $\begin{pmatrix} 1169 & 1073 \\ 0 & 1 \end{pmatrix}$, $\begin{pmatrix} 3541 & 3207 \\ 0 & 1 \end{pmatrix}$, $\begin{pmatrix} 2347 & 1592 \\ 0 & 1 \end{pmatrix}$, and the torsion units are generated by -1 of order 2.

Entry 1: character a , column e_1

Prescribed character vector $(1 \ 0 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 2917y^3 + (-28s + 14)y^2 - 4675205y + (7300s - 3650)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 5834y^8 - 841521y^6 + 13427539578y^4 + 633998613425y^2 + 2766640450067500$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ -1 \ 0 \ -1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{7819837}{125589384071564149} + \left(\frac{49172}{125589384071564149}\right)s, \quad J = \begin{pmatrix} 2629 & 2019 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 2629$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $5236775471326400 = 2^6 \cdot 5^2 \cdot 24917 \cdot 131355487$:

$$\begin{pmatrix} 32729846695790 & 19034682206160 & 22079543210832 & 495030723360 & 10871148595935 & 21514060838939 & 27314933676474 & 25387588672776 & 7312846335509 & 29597474726 \\ 0 & 10 & 2 & 2 & 5 & 3 & 6 & 2 & 7 & 5 \\ 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 2 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 558 factors with signed exponents of absolute value up to 801381994235608608567; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (0 \ 4 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 2: character a , column e_2

Prescribed character vector $(1 \ 0 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 2917y^3 + (-28s + 14)y^2 - 4675205y + (7300s - 3650)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 5834y^8 - 841521y^6 + 13427539578y^4 + 633998613425y^2 + 2766640450067500$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ -1 \ 0 \ -1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{4846798}{52598468275676629} + \left(-\frac{130739}{894173960686502693}\right)s, \quad J = \begin{pmatrix} 3893 & 3297 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 3893$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $172833814467504 = 2^4 \cdot 3^2 \cdot 29 \cdot 43^2 \cdot 877 \cdot 25523$:

$$\begin{pmatrix} 167474626422 & 136747804292 & 96070959763 & 159118133215 & 127971040846 & 167155402907 & 31191146131 & 122903460055 & 63929919471 & 125894012955 \\ 0 & 2 & 1 & 1 & 0 & 1 & 1 & 0 & 1 & 1 \\ 0 & 0 & 129 & 69 & 84 & 75 & 60 & 85 & 61 & 52 \\ 0 & 0 & 0 & 2 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 558 factors with signed exponents of absolute value up to 290868339864622474700; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (0 \ 3 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 3: character a , column e_3

Prescribed character vector $(1 \ 0 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 2917y^3 + (-28s + 14)y^2 - 4675205y + (7300s - 3650)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 5834y^8 - 841521y^6 + 13427539578y^4 + 633998613425y^2 + 2766640450067500$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ -1 \ 0 \ -1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{263382107}{71214011112376507} + \left(-\frac{5957}{71214011112376507}\right)s, \quad J = \begin{pmatrix} 2347 & 1592 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 2347$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $157259949880716 = 2^2 \cdot 3^2 \cdot 41 \cdot 106544681491$:

$$\begin{pmatrix} 26209991646786 & 14772190165388 & 11175848723407 & 6113726627562 & 16339450043246 & 15877291207427 & 19323808575215 & 1360019728689 & 1923107208954 & 9051844451582 \\ 0 & 2 & 1 & 0 & 0 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 3 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 559 factors with signed exponents of absolute value up to 461134681829006158597; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (0 \ 0 \ 2)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 4: character b , column e_1

Prescribed character vector $(0 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 2y^4 + (-s - 9971)y^3 + (122s + 382006)y^2 + (-4553s + 2863618)y + (53932s - 179108972)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$\begin{aligned}
& y^{10} - 4y^9 - 19939y^8 + 804020y^7 + 155541918y^6 - 20656844102* \\
& y^5 + 1335109198071y^4 - 57517974247918y^3 + 1630760624355461y^2 \\
& - 26521365163449172y + 183078398985894064
\end{aligned}$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ -1 \ -2 \ 0 \ 1 \ 0 \ -1 \ 1 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{7819837}{125589384071564149} + \left(\frac{49172}{125589384071564149}\right)s, \quad J = \begin{pmatrix} 2629 & 2019 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 2629$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $13487528469564 = 2^2 \cdot 3^2 \cdot 374653568599$:

$$\begin{pmatrix}
2247921411594 & 1320318670650 & 724268856741 & 1276803325930 & 768135962880 & 1842146593225 & 56656261100 & 703367373339 & 1809821823879 & 367551907464 \\
0 & 2 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\
0 & 0 & 3 & 1 & 0 & 1 & 0 & 2 & 0 & 1 \\
0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1
\end{pmatrix}$$

The element t is stored in compact form as a product of 412 factors with signed exponents of absolute value up to 190979094329227970; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (1 \ 0 \ 1)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 5: character b , column e_2

Prescribed character vector $(0 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 2y^4 + (-s - 9971)y^3 + (122s + 382006)y^2 + (-4553s + 2863618)y + (53932s - 179108972)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$\begin{aligned}
& y^{10} - 4y^9 - 19939y^8 + 804020y^7 + 155541918y^6 - 20656844102* \\
& y^5 + 1335109198071y^4 - 57517974247918y^3 + 1630760624355461y^2 \\
& - 26521365163449172y + 183078398985894064
\end{aligned}$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ -1 \ -2 \ 0 \ 1 \ 0 \ -1 \ 1 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{4846798}{52598468275676629} + \left(-\frac{130739}{894173960686502693}\right) s, \quad J = \begin{pmatrix} 3893 & 3297 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 3893$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $480825388634112 = 2^{12} \cdot 3^2 \cdot 23^2 \cdot 31^2 \cdot 25657$:

$$\begin{pmatrix} 219521292 & 43900836 & 30824292 & 51188568 & 215878162 & 163335276 & 81011170 & 145532660 & 13839618 & 84714980 \\ 0 & 2852 & 1200 & 428 & 1870 & 669 & 1792 & 1014 & 202 & 926 \\ 0 & 0 & 4 & 0 & 2 & 1 & 2 & 2 & 0 & 2 \\ 0 & 0 & 0 & 12 & 8 & 3 & 8 & 0 & 0 & 10 \\ 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 412 factors with signed exponents of absolute value up to 230929667948059606; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (0 \ 0 \ 2)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 6: character b , column e_3

Prescribed character vector $(0 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 2y^4 + (-s - 9971)y^3 + (122s + 382006)y^2 + (-4553s + 2863618)y + (53932s - 179108972)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$\begin{aligned} & y^{10} - 4y^9 - 19939y^8 + 804020y^7 + 155541918y^6 - 20656844102* \\ & y^5 + 1335109198071y^4 - 57517974247918y^3 + 1630760624355461y^2 \\ & - 26521365163449172y + 183078398985894064 \end{aligned}$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ -1 \ -2 \ 0 \ 1 \ 0 \ -1 \ 1 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{263382107}{71214011112376507} + \left(-\frac{5957}{71214011112376507}\right) s, \quad J = \begin{pmatrix} 2347 & 1592 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 2347$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $447235425161188 = 2^2 \cdot 31 \cdot 71 \cdot 50799116897$:

$$\begin{pmatrix} 223617712580594 & 78913820643228 & 13320601971187 & 95923605060799 & 174246576228594 & 7650752745118 & 40917203278250 & 59123789450697 & 129381710437483 & 164787727277979 \\ 0 & 2 & 1 & 1 & 0 & 1 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 406 factors with signed exponents of absolute value up to 15092670489291852; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (2 \ 0 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 7: character c , column e_1

Prescribed character vector $(0 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 2685y^3 + (8s - 4)y^2 - 115192y + (-14512s + 7256)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 5370y^8 + 6978841y^6 + 3941249248y^4 - 12041371061760y^2 + 10933558714571968$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-2 \ -2 \ -1 \ 1 \ -1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{7819837}{125589384071564149} + \left(\frac{49172}{125589384071564149}\right)s, \quad J = \begin{pmatrix} 2629 & 2019 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 2629$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $472418802044352 = 2^6 \cdot 3^2 \cdot 820171531327$:

$$\begin{pmatrix} 9842058375924 & 4795925116977 & 5748750334125 & 9385006924065 & 6432328895314 & 3365793032988 & 6484189016580 & 9055923204835 & 3399077634763 & 6890598207123 \\ 0 & 3 & 2 & 1 & 0 & 1 & 1 & 0 & 2 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 555 factors with signed exponents of absolute value up to 17983191749811102177046877848; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (1 \ 2 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 8: character c , column e_2

Prescribed character vector $(0 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 2685y^3 + (8s - 4)y^2 - 115192y + (-14512s + 7256)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 5370y^8 + 6978841y^6 + 3941249248y^4 - 12041371061760y^2 + 10933558714571968$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-2 \ -2 \ -1 \ 1 \ -1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{4846798}{52598468275676629} + \left(-\frac{130739}{894173960686502693}\right)s, \quad J = \begin{pmatrix} 3893 & 3297 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 3893$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $1025500962545392 = 2^4 \cdot 3779863 \cdot 16956649$:

$$\begin{pmatrix} 128187620318174 & 76351201928371 & 29871509207124 & 46568070590057 & 77447081442819 & 52075310822958 & 101185230121110 & 118540138883141 & 111958359526236 & 104785119408516 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 1 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 555 factors with signed exponents of absolute value up to 2950280129673359394696484510; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (0 \ 0 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 9: character c , column e_3

Prescribed character vector $(0 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 2685y^3 + (8s - 4)y^2 - 115192y + (-14512s + 7256)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 5370y^8 + 6978841y^6 + 3941249248y^4 - 12041371061760y^2 + 10933558714571968$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-2 \ -2 \ -1 \ 1 \ -1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{263382107}{71214011112376507} + \left(-\frac{5957}{71214011112376507}\right)s, \quad J = \begin{pmatrix} 2347 & 1592 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 2347$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $195196712041728 = 2^8 \cdot 3^2 \cdot 53 \cdot 4027 \cdot 396947$:

$$\begin{pmatrix} 2033299083768 & 647053578768 & 1708728845772 & 2018826042163 & 511467635472 & 779856678401 & 1645845131375 & 1556699459521 & 851644016300 & 199496301142 \\ 0 & 6 & 2 & 4 & 3 & 4 & 1 & 5 & 0 & 4 \\ 0 & 0 & 4 & 2 & 1 & 2 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 555 factors with signed exponents of absolute value up to 3706076345513484036059920557; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (3 \ 0 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 10: character $a + b$, column e_1

Prescribed character vector $(1 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 1773y^3 + (-6s + 3)y^2 + 471172y + (246s - 123)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 3546y^8 + 4085873y^6 + 198224955y^4 + 68744982490y^2 + 3141790457427$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-1 \ -2 \ 0 \ 1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{7819837}{125589384071564149} + \left(\frac{49172}{125589384071564149}\right)s, \quad J = \begin{pmatrix} 2629 & 2019 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 2629$. **Checked:** $(a')^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $2567277714495600 = 2^4 \cdot 3^2 \cdot 5^2 \cdot 17 \cdot 41948982263$:

$$\begin{pmatrix} 42787961908260 & 32330408476937 & 21665077039160 & 42183805838486 & 12499427747577 & 17674029298520 & 32922766497790 & 33258541220661 & 17001352519264 & 4601842637137 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 20 & 8 & 19 & 12 & 0 & 6 & 8 & 4 \\ 0 & 0 & 0 & 3 & 0 & 0 & 0 & 2 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 569 factors with signed exponents of absolute value up to 83762484408491022597482433672526721909892402029549269; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (2 \ 3 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 11: character $a + b$, column e_2

Prescribed character vector $(1 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 1773y^3 + (-6s + 3)y^2 + 471172y + (246s - 123)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 3546y^8 + 4085873y^6 + 198224955y^4 + 68744982490y^2 + 3141790457427$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-1 \ -2 \ 0 \ 1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{4846798}{52598468275676629} + \left(-\frac{130739}{894173960686502693}\right)s, \quad J = \begin{pmatrix} 3893 & 3297 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 3893$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $172901988132112 = 2^4 \cdot 10806374258257$:

$$\begin{pmatrix} 21612748516514 & 8440681956086 & 6568077776434 & 9687159368273 & 11557475186659 & 5132517287924 & 7671656471784 & 9552393025204 & 10723021894315 & 6032691845309 \\ 0 & 2 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 1 & 1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 569 factors with signed exponents of absolute value up to 39698891915110589788079390078814316827447423421199605; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (2 \ 3 \ 1)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 12: character $a + b$, column e_3

Prescribed character vector $(1 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 1773y^3 + (-6s + 3)y^2 + 471172y + (246s - 123)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 3546y^8 + 4085873y^6 + 198224955y^4 + 68744982490y^2 + 3141790457427$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-1 \ -2 \ 0 \ 1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{263382107}{71214011112376507} + \left(-\frac{5957}{71214011112376507}\right) s, \quad J = \begin{pmatrix} 2347 & 1592 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 2347$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $316873415640924 = 2^2 \cdot 3^2 \cdot 8802039323359$:

$$\begin{pmatrix} 52812235940154 & 767918682478 & 33135257267796 & 4758394963721 & 8361057858601 & 7981999651364 & 48834669333325 & 12418268700210 & 15317595250013 & 27059393822802 \\ 0 & 2 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 3 & 0 & 0 & 0 & 2 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 571 factors with signed exponents of absolute value up to 86566199040747872800119432108328734223231632965839732; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (2 \ 3 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 13: character $a + c$, column e_1

Prescribed character vector $(1 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 4133y^3 + 3874204y + (-15840s + 7920)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 8266y^8 + 24830097y^6 - 32024170264y^4 + 15009456633616y^2 + 13026188442643200$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-2 \ -1 \ -1 \ 1 \ -2 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{7819837}{125589384071564149} + \left(\frac{49172}{125589384071564149}\right) s, \quad J = \begin{pmatrix} 2629 & 2019 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 2629$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $242637573445388 = 2^2 \cdot 11^2 \cdot 17 \cdot 34679 \cdot 850349$:

$$\begin{pmatrix} 11028980611154 & 7659469918848 & 10984123010233 & 2267018097449 & 2310058771341 & 5333625976444 & 4660914356093 & 6108953828904 & 6943534721901 & 5239591664535 \\ 0 & 11 & 0 & 1 & 1 & 1 & 8 & 6 & 4 & 2 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 260 factors with signed exponents of absolute value up to 342696798556; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (1 \ 3 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 14: character $a + c$, column e_2

Prescribed character vector $(1 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 4133y^3 + 3874204y + (-15840s + 7920)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 8266y^8 + 24830097y^6 - 32024170264y^4 + 15009456633616y^2 + 13026188442643200$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-2 \ -1 \ -1 \ 1 \ -2 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{4846798}{52598468275676629} + \left(-\frac{130739}{894173960686502693}\right)s, \quad J = \begin{pmatrix} 3893 & 3297 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 3893$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $11348065753200 = 2^4 \cdot 3^4 \cdot 5^2 \cdot 11 \cdot 17 \cdot 31^2 \cdot 1949$:

$$\begin{pmatrix} 338950590 & 140457714 & 195579299 & 322460941 & 61668943 & 271880988 & 49538952 & 291454771 & 210272570 & 108091801 \\ 0 & 186 & 165 & 104 & 182 & 150 & 96 & 8 & 90 & 24 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 5 & 4 & 0 & 0 & 0 & 4 & 3 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 6 & 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 & 6 & 1 & 5 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 267 factors with signed exponents of absolute value up to 628617816494; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of +1. Certified value: $D_x(e_2) = (1 \ 2 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 15: character $a + c$, column e_3

Prescribed character vector $(1 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 4133y^3 + 3874204y + (-15840s + 7920)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 8266y^8 + 24830097y^6 - 32024170264y^4 + 15009456633616y^2 + 13026188442643200$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-2 \ -1 \ -1 \ 1 \ -2 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{263382107}{71214011112376507} + \left(-\frac{5957}{71214011112376507}\right)s, \quad J = \begin{pmatrix} 2347 & 1592 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 2347$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $242637573445388 = 2^2 \cdot 11^2 \cdot 17 \cdot 34679 \cdot 850349$:

$$\begin{pmatrix} 11028980611154 & 7659469918848 & 10984123010233 & 2267018097449 & 2310058771341 & 5333625976444 & 4660914356093 & 6108953828904 & 6943534721901 & 5239591664535 \\ 0 & 11 & 0 & 1 & 1 & 1 & 8 & 6 & 4 & 2 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 260 factors with signed exponents of absolute value up to 645690957532; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (1 \ 3 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 16: character $b + c$, column e_1

Prescribed character vector $(0 \ 1 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 680y^3 + (10s - 5)y^2 + 1860234y + (488s - 244)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 1360y^8 + 4182868y^6 + 7721587315y^4 + 3967177436476y^2 + 12363648401968$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-2 \ 2 \ -1 \ 1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{7819837}{125589384071564149} + \left(\frac{49172}{125589384071564149}\right)s, \quad J = \begin{pmatrix} 2629 & 2019 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 2629$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $7498623075836593 = 881 \cdot 5647 \cdot 1507258799$:

$$\begin{pmatrix} 7498623075836593 & 2226408355231278 & 3054029924954316 & 3276148222234553 & 4521754286467904 & 1193483831953610 & 4021169656363274 & 6469538317214402 & 1057819705331739 & 1984151582128351 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 2 factors with signed exponents of absolute value up to 2; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 3$ with residue degree 2; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (2 \ 1 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 17: character $b + c$, column e_2

Prescribed character vector $(0 \ 1 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 680y^3 + (10s - 5)y^2 + 1860234y + (488s - 244)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 1360y^8 + 4182868y^6 + 7721587315y^4 + 3967177436476y^2 + 12363648401968$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-2 \ 2 \ -1 \ 1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = \frac{4846798}{52598468275676629} + \left(-\frac{130739}{894173960686502693}\right)s, \quad J = \begin{pmatrix} 3893 & 3297 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 3893$. **Checked:** $(a')^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $2295123594311664 = 2^4 \cdot 3^2 \cdot 41 \cdot 151 \cdot 367 \cdot 727 \cdot 9649$:

$$\begin{pmatrix} 191260299525972 & 40895211500676 & 129747611993031 & 114203607435321 & 35830775754078 & 175604131326864 & 112896247386547 & 160591600909124 & 120141892312509 & 171567491119770 \\ 0 & 4 & 2 & 2 & 2 & 0 & 2 & 2 & 2 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 3 & 1 & 0 & 1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 7 factors with signed exponents of absolute value up to 1; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 5$ with residue degree 2; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (1 \ 3 \ 2)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 18: character $b + c$, column e_3

Prescribed character vector $(0 \ 1 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 680y^3 + (10s - 5)y^2 + 1860234y + (488s - 244)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 1360y^8 + 4182868y^6 + 7721587315y^4 + 3967177436476y^2 + 12363648401968$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-2 \ 2 \ -1 \ 1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{263382107}{71214011112376507} + \left(-\frac{5957}{71214011112376507}\right)s, \quad J = \begin{pmatrix} 2347 & 1592 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 2347$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $5105013679476 = 2^2 \cdot 3^2 \cdot 773 \cdot 183448817$:

$$\begin{pmatrix} 850835613246 & 684662461341 & 378402333096 & 266130923787 & 543142920168 & 550332345141 & 167406616789 & 76440803006 & 298864856100 & 578069679333 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 3 & 1 & 0 & 1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 508 factors with signed exponents of absolute value up to 4094218485288585229915261665829; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (3 \ 1 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.