

A guide to the arithmetic certificate for $K = \mathbf{Q}(\sqrt{-65818135})$, $p = 5$

Generated by `tools/certificate_guide.gp`

What this document is

This guide restates the content of `certificate.gp` in the notation of Section 2.5 of the paper. Each of the 18 entries certifies one value $D_x(e_j)$ of a secondary norm operator: it stores the unramified cyclic quintic L_x/K , the normalized generator σ_x , the pair (a', J) representing the torsion class e_j , the auxiliary ideal I' , the compactly represented element t , a residue prime for the sign check, and the expected norm class $[N_x(I')]$. The two defining equations are

$$I'^{(1-\sigma_x)^2}(t) J \mathcal{O}_{L_x} = \mathcal{O}_{L_x}, \quad N_x(t) = a'.$$

Facts marked **Checked** are recomputed exactly by the generating script; the full verification is performed by the shared verifier `certificate/verify_certificate.c` as described in the README files of the `certificate` directory.

Notation and conventions

Ideals as matrices. An ideal of the ring of integers is stored by its *Hermite normal form* (HNF): an upper-triangular integer matrix whose columns are the coordinates, with respect to the fixed integral basis, of a $\mathbf{Z}$ -basis of the ideal. For example, the ideal $J = \begin{pmatrix} 376 & 102 \\ 0 & 1 \end{pmatrix}$ of Entry 1 below, over K with integral basis $(1, s)$, denotes the ideal $\mathbf{Z}376 + \mathbf{Z}(102 + s)$. The determinant of the matrix is the norm of the ideal. Elements are likewise given by their coordinate vectors in the integral basis.

Which integral basis. The one PARI returns for the field in question – and that basis is LLL-reduced, so it is not canonical: another machine may reduce differently, and the same coordinate vector would then denote a different algebraic number. The certificate therefore records the basis it was written against, as its last field per entry and in the header for K . The verifier expresses the stored basis in its own, checks that the resulting matrix is integral with determinant ± 1 – so that both bases span the same ring of integers – and converts the coordinates before checking anything. Where the two bases agree, that matrix is the identity and nothing changes.

Characters. The header fixes three generators $\kappa_1, \kappa_2, \kappa_3$ of $\text{Cl}(K)$ (as HNF ideals). A character x on $\text{Cl}(K)/5$ is recorded by its value vector $(x(\bar{\kappa}_1), x(\bar{\kappa}_2), x(\bar{\kappa}_3))$; thus $(1, 0, 0)$ is the character dual to $\bar{\kappa}_1$. The column number j records which basis element e_j of $\text{Cl}(K)[5]$ the stored pair (a', J) represents.

Automorphisms. σ_x is a field automorphism of L_x . Writing θ for the chosen root of the stored absolute polynomial, so that $L_x = \mathbf{Q}(\theta)$, the automorphism is determined by the single value $\sigma_x(\theta)$, and the certificate stores the coordinate vector of $\sigma_x(\theta)$ in the integral basis of L_x .

Base field data

The base field is $K = \mathbf{Q}(s)$ with $s^2 - s + 16454534 = 0$, of discriminant -65818135 . Class group invariants $(30 \ 10 \ 10)$, class number 3000; the three stored generator ideals (HNF) are $\begin{pmatrix} 167 & 122 \\ 0 & 1 \end{pmatrix}$, $\begin{pmatrix} 3236 & 2849 \\ 0 & 1 \end{pmatrix}$, $\begin{pmatrix} 1532 & 1449 \\ 0 & 1 \end{pmatrix}$, and the torsion units are generated by -1 of order 2.

Entry 1: character a , column e_1

Prescribed character vector $(1 \ 0 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 1835y^3 + (-4s + 2)y^2 - 755006y + (4162s - 2081)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 3670y^8 + 1857213y^6 + 3034144560y^4 + 22163904296y^2 + 285029448523735$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 1 \ 0 \ -1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{2727091}{7515177189376} + \left(\frac{69}{7515177189376}\right)s, \quad J = \begin{pmatrix} 376 & 102 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 376$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $75609622983685 = 5 \cdot 7 \cdot 2160274942391$:

$$\begin{pmatrix} 75609622983685 & 47283923323083 & 27054639213321 & 34692623375662 & 20080767651362 & 49907583744599 & 45626435912409 & 52729161695551 & 43425964277059 & 44132736917913 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 270 factors with signed exponents of absolute value up to 902920854; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 29$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (0 \ 3 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 2: character a , column e_2

Prescribed character vector $(1 \ 0 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 1835y^3 + (-4s + 2)y^2 - 755006y + (4162s - 2081)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 3670y^8 + 1857213y^6 + 3034144560y^4 + 22163904296y^2 + 285029448523735$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 1 \ 0 \ -1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{423739183}{478923089887538176} + \left(-\frac{134871}{478923089887538176}\right)s, \quad J = \begin{pmatrix} 3436 & 2514 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 3436$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $103402703869507 = 7^3 \cdot 41117 \cdot 7331897$:

$$\begin{pmatrix} 14771814838501 & 8966377870163 & 8606216883170 & 8130438787871 & 13343964720712 & 869263280160 & 11400917715298 & 5969108397036 & 6826106309166 & 3300076709857 \\ 0 & 7 & 4 & 0 & 2 & 0 & 4 & 3 & 4 & 5 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 276 factors with signed exponents of absolute value up to 3169729789; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 29$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (0 \ 2 \ 1)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 3: character a , column e_3

Prescribed character vector $(1 \ 0 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 1835y^3 + (-4s + 2)y^2 - 755006y + (4162s - 2081)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 3670y^8 + 1857213y^6 + 3034144560y^4 + 22163904296y^2 + 285029448523735$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 1 \ 0 \ -1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{9195119}{5023060313370901} + \left(-\frac{17324}{5023060313370901}\right)s, \quad J = \begin{pmatrix} 1381 & 823 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 1381$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $3628600103807 = 7^4 \cdot 1511287007$:

$$\begin{pmatrix} 74053063343 & 7809615821 & 24962335090 & 38114576417 & 68393645561 & 31076094436 & 27497178538 & 64080793374 & 56748121474 & 59310507186 \\ 0 & 7 & 0 & 0 & 1 & 0 & 5 & 2 & 5 & 5 \\ 0 & 0 & 7 & 0 & 1 & 0 & 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 279 factors with signed exponents of absolute value up to 2498889388; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 29$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (0 \ 4 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 4: character b , column e_1

Prescribed character vector $(0 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 4428y^3 + (52s - 26)y^2 - 4028269y + (-7412s + 3706)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 8856y^8 + 11550646y^6 + 8818708996y^4 + 3543006704241y^2 + 903974962796860$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ -1 \ -1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{2727091}{7515177189376} + \left(\frac{69}{7515177189376}\right)s, \quad J = \begin{pmatrix} 376 & 102 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 376$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $2684026368000 = 2^{12} \cdot 3^8 \cdot 5^3 \cdot 17 \cdot 47$:

$$\begin{pmatrix} 287640 & 284220 & 117360 & 3120 & 101202 & 164874 & 204713 & 234078 & 192594 & 187204 \\ 0 & 180 & 84 & 120 & 144 & 48 & 4 & 100 & 97 & 84 \\ 0 & 0 & 12 & 0 & 6 & 6 & 3 & 9 & 9 & 0 \\ 0 & 0 & 0 & 30 & 12 & 6 & 28 & 20 & 17 & 26 \\ 0 & 0 & 0 & 0 & 6 & 0 & 3 & 3 & 5 & 5 \\ 0 & 0 & 0 & 0 & 0 & 6 & 2 & 4 & 0 & 5 \\ 0 & 0 & 0 & 0 & 0 & 0 & 4 & 3 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 207 factors with signed exponents of absolute value up to 472655332384026; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of +1. Certified value: $D_x(e_1) = (4 \ 0 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 5: character b , column e_2

Prescribed character vector $(0 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 4428y^3 + (52s - 26)y^2 - 4028269y + (-7412s + 3706)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 8856y^8 + 11550646y^6 + 8818708996y^4 + 3543006704241y^2 + 903974962796860$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ -1 \ -1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{423739183}{478923089887538176} + \left(-\frac{134871}{478923089887538176}\right) s, \quad J = \begin{pmatrix} 3436 & 2514 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 3436$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $374974030373760 = 2^7 \cdot 3^4 \cdot 5 \cdot 38167 \cdot 189517$:

$$\begin{pmatrix} 1735990881360 & 181384163088 & 1366340857329 & 393394629930 & 1735646724250 & 942187251161 & 1175044230886 & 1588495672502 & 1501797634731 & 866510699935 \\ 0 & 12 & 0 & 6 & 11 & 6 & 8 & 8 & 0 & 2 \\ 0 & 0 & 3 & 0 & 1 & 1 & 1 & 0 & 0 & 2 \\ 0 & 0 & 0 & 6 & 5 & 0 & 1 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 208 factors with signed exponents of absolute value up to 70760531059040; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of +1. Certified value: $D_x(e_2) = (0 \ 0 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 6: character b , column e_3

Prescribed character vector $(0 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 4428y^3 + (52s - 26)y^2 - 4028269y + (-7412s + 3706)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 8856y^8 + 11550646y^6 + 8818708996y^4 + 3543006704241y^2 + 903974962796860$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ -1 \ -1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{9195119}{5023060313370901} + \left(-\frac{17324}{5023060313370901}\right) s, \quad J = \begin{pmatrix} 1381 & 823 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 1381$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $67209581260800 = 2^{12} \cdot 3^4 \cdot 5^2 \cdot 11^2 \cdot 167 \cdot 401$:

$$\begin{pmatrix} 530378640 & 17582880 & 51540192 & 488681036 & 318011218 & 144283882 & 462488274 & 432881326 & 361755195 & 431406302 \\ 0 & 40 & 0 & 0 & 24 & 0 & 19 & 11 & 38 & 27 \\ 0 & 0 & 396 & 2 & 394 & 152 & 116 & 175 & 60 & 68 \\ 0 & 0 & 0 & 2 & 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 205 factors with signed exponents of absolute value up to 1189098449411209; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (3 \ 0 \ 3)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 7: character c , column e_1

Prescribed character vector $(0 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 499y^3 + (-20s + 10)y^2 + 518780y + (9630s - 4815)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 998y^8 + 1286561y^6 + 7099555940y^4 - 6069153712100y^2 + 1525942450920375$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{2727091}{7515177189376} + \left(\frac{69}{7515177189376}\right)s, \quad J = \begin{pmatrix} 376 & 102 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 376$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $62815232425905 = 3^2 \cdot 5 \cdot 29 \cdot 283 \cdot 170085787$:

$$\begin{pmatrix} 20938410808635 & 11959530549276 & 1777670601763 & 16445998551337 & 12824007308869 & 16347671763141 & 4643088919581 & 10145134163698 & 1355123928888 & 7583120609209 \\ 0 & 3 & 0 & 0 & 1 & 1 & 0 & 1 & 2 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 423 factors with signed exponents of absolute value up to 47343022488470165008569184540; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (4 \ 2 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 8: character c , column e_2

Prescribed character vector $(0 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 499y^3 + (-20s + 10)y^2 + 518780y + (9630s - 4815)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 998y^8 + 1286561y^6 + 7099555940y^4 - 6069153712100y^2 + 1525942450920375$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{423739183}{478923089887538176} + \left(-\frac{134871}{478923089887538176}\right)s, \quad J = \begin{pmatrix} 3436 & 2514 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 3436$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $62815232425905 = 3^2 \cdot 5 \cdot 29 \cdot 283 \cdot 170085787$:

$$\begin{pmatrix} 20938410808635 & 11959530549276 & 1777670601763 & 16445998551337 & 12824007308869 & 16347671763141 & 4643088919581 & 10145134163698 & 1355123928888 & 7583120609209 \\ 0 & 3 & 0 & 0 & 1 & 1 & 0 & 1 & 2 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 418 factors with signed exponents of absolute value up to 51636876843506689906245173985; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of +1. Certified value: $D_x(e_2) = (4 \ 2 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 9: character c , column e_3

Prescribed character vector $(0 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 499y^3 + (-20s + 10)y^2 + 518780y + (9630s - 4815)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 998y^8 + 1286561y^6 + 7099555940y^4 - 6069153712100y^2 + 1525942450920375$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{9195119}{5023060313370901} + \left(-\frac{17324}{5023060313370901}\right)s, \quad J = \begin{pmatrix} 1381 & 823 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 1381$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $25097312214875 = 5^3 \cdot 16943 \cdot 11850233$:

$$\begin{pmatrix} 1003892488595 & 977215999005 & 476422651313 & 818852914870 & 310751809317 & 717630864284 & 617464530756 & 25036455751 & 271350166264 & 328971762283 \\ 0 & 5 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 4 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 5 & 4 & 0 & 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 419 factors with signed exponents of absolute value up to 56927146183146050017979406335; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of +1. Certified value: $D_x(e_3) = (0 \ 1 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 10: character $a + b$, column e_1

Prescribed character vector $(1 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 1408y^3 + (2s - 1)y^2 + 700226y + (2118s - 1059)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 2816y^8 + 3382916y^6 + 2037654551y^4 + 629719261006y^2 + 73813787857935$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{2727091}{7515177189376} + \left(\frac{69}{7515177189376}\right)s, \quad J = \begin{pmatrix} 376 & 102 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 376$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $414256155720235 = 5 \cdot 47 \cdot 4523 \cdot 389739587$:

$$\begin{pmatrix} 414256155720235 & 216419549108434 & 43138126896018 & 231518822387282 & 32584464217621 & 186269003539725 & 344622710039400 & 44631480178161 & 284266807696088 & 48357685988998 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 617 factors with signed exponents of absolute value up to 14337465750007525015251837496378193492201913120583600376342498; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (1 \ 4 \ 3)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 11: character $a + b$, column e_2

Prescribed character vector $(1 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 1408y^3 + (2s - 1)y^2 + 700226y + (2118s - 1059)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 2816y^8 + 3382916y^6 + 2037654551y^4 + 629719261006y^2 + 73813787857935$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{423739183}{478923089887538176} + \left(-\frac{134871}{478923089887538176}\right)s, \quad J = \begin{pmatrix} 3436 & 2514 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 3436$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $69306691386400 = 2^5 \cdot 5^2 \cdot 13^2 \cdot 17 \cdot 30154321$:

$$\begin{pmatrix} 66641049410 & 56978271200 & 49062074452 & 59342101014 & 16581128900 & 14481436396 & 33317123900 & 44019062526 & 7830719769 & 20504256314 \\ 0 & 10 & 6 & 2 & 4 & 4 & 2 & 7 & 8 & 0 \\ 0 & 0 & 26 & 24 & 14 & 21 & 23 & 22 & 14 & 25 \\ 0 & 0 & 0 & 2 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 625 factors with signed exponents of absolute value up to 7792410538702590794282180684740183341891447753266426908361176; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (3 \ 2 \ 1)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 12: character $a + b$, column e_3

Prescribed character vector $(1 \ 1 \ 0)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 + 1408y^3 + (2s - 1)y^2 + 700226y + (2118s - 1059)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} + 2816y^8 + 3382916y^6 + 2037654551y^4 + 629719261006y^2 + 73813787857935$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 1 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{9195119}{5023060313370901} + \left(-\frac{17324}{5023060313370901}\right)s, \quad J = \begin{pmatrix} 1381 & 823 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 1381$. **Checked:** $(a')^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $9408299058511 = 17 \cdot 47^2 \cdot 250533887$:

$$\begin{pmatrix} 200176575713 & 83728177730 & 82961865025 & 166916732692 & 54008305564 & 153337427120 & 10631634609 & 164332355001 & 128498306374 & 178559875355 \\ 0 & 47 & 8 & 46 & 0 & 20 & 0 & 30 & 0 & 4 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 617 factors with signed exponents of absolute value up to 14128213608025858302661376278261485375132975994065079540896671; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of +1. Certified value: $D_x(e_3) = (4 \ 1 \ 4)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 13: character $a + c$, column e_1

Prescribed character vector $(1 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - y^4 + (-s - 459)y^3 + (-32s + 94026)y^2 + (20s + 2384885)y + (2920s + 4198015)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 2y^9 - 918y^8 + 188939y^7 + 21247444y^6 + 970324130y^5 + 22829023855y^4 + 327392360075y^3 + 3408867996825y^2 + 21952503402250y + 157933526841625$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 0 \ 1 \ 0 \ 0 \ 1 \ 0 \ -1 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{2727091}{7515177189376} + \left(\frac{69}{7515177189376}\right) s, \quad J = \begin{pmatrix} 376 & 102 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 376$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $34337724625 = 5^3 \cdot 4513 \cdot 60869$:

$$\begin{pmatrix} 1373508985 & 1201641340 & 607693915 & 670993078 & 407497101 & 201241587 & 521146528 & 940738925 & 998955740 & 1007758079 \\ 0 & 5 & 0 & 0 & 1 & 1 & 0 & 0 & 3 & 0 \\ 0 & 0 & 5 & 0 & 0 & 4 & 4 & 1 & 1 & 3 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 566 factors with signed exponents of absolute value up to 10315579000624460349595; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of +1. Certified value: $D_x(e_1) = (0 \ 2 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 14: character $a + c$, column e_2

Prescribed character vector $(1 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - y^4 + (-s - 459) y^3 + (-32s + 94026) y^2 + (20s + 2384885) y + (2920s + 4198015)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$\begin{aligned} & y^{10} - 2y^9 - 918y^8 + 188939y^7 + 21247444y^6 + 970324130y^5 + \\ & 22829023855y^4 + 327392360075y^3 + 3408867996825y^2 + 2195250340 \\ & 2250y + 157933526841625 \end{aligned}$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 0 \ 1 \ 0 \ 0 \ 1 \ 0 \ -1 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{423739183}{478923089887538176} + \left(-\frac{134871}{478923089887538176}\right) s, \quad J = \begin{pmatrix} 3436 & 2514 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 3436$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $44777455405 = 5 \cdot 17 \cdot 526793593$:

$$\begin{pmatrix} 44777455405 & 27194151013 & 5550849355 & 34580100438 & 15093253168 & 21803145289 & 32529052248 & 24294814835 & 31310347811 & 15977002534 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 596 factors with signed exponents of absolute value up to 20518856656736022742717; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of +1. Certified value: $D_x(e_2) = (3 \ 0 \ 2)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 15: character $a + c$, column e_3

Prescribed character vector $(1 \ 0 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - y^4 + (-s - 459)y^3 + (-32s + 94026)y^2 + (20s + 2384885)y + (2920s + 4198015)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 2y^9 - 918y^8 + 188939y^7 + 21247444y^6 + 970324130y^5 + 22829023855y^4 + 327392360075y^3 + 3408867996825y^2 + 21952503402250y + 157933526841625$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(0 \ 0 \ 0 \ 1 \ 0 \ 0 \ 1 \ 0 \ -1 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{9195119}{5023060313370901} + \left(-\frac{17324}{5023060313370901}\right)s, \quad J = \begin{pmatrix} 1381 & 823 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 1381$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $159260256386111 = 159260256386111$:

$$\begin{pmatrix} 159260256386111 & 60742635824748 & 2085117131728 & 40112618317426 & 118164677014251 & 124257340947545 & 99753669023153 & 70306056152814 & 41757426177213 & 76508174887371 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 595 factors with signed exponents of absolute value up to 92858689301476038427018; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (0 \ 0 \ 0)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 16: character $b + c$, column e_1

Prescribed character vector $(0 \ 1 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 3691y^3 + (18s - 9)y^2 + 183575y + (-550s + 275)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 7382y^8 + 13990631y^6 + 3976118285y^4 - 292099987625y^2 + 4977496459375$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-1 \ 2 \ 0 \ 1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{2727091}{7515177189376} + \left(\frac{69}{7515177189376}\right)s, \quad J = \begin{pmatrix} 376 & 102 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 376$. **Checked:** $(a') J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $41287842623375 = 5^3 \cdot 1307 \cdot 2179 \cdot 115979$:

$$\begin{pmatrix} 1651513704935 & 197082804290 & 681018395530 & 575917572615 & 522103902825 & 860373300322 & 1146319549058 & 323950215745 & 1334181509219 & 655367054469 \\ 0 & 5 & 0 & 1 & 0 & 0 & 2 & 2 & 0 & 1 \\ 0 & 0 & 5 & 1 & 4 & 4 & 3 & 1 & 0 & 3 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 471 factors with signed exponents of absolute value up to 265325962754132; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_1) = (2 \ 2 \ 3)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 17: character $b + c$, column e_2

Prescribed character vector $(0 \ 1 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 3691y^3 + (18s - 9)y^2 + 183575y + (-550s + 275)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 7382y^8 + 13990631y^6 + 3976118285y^4 - 292099987625y^2 + 4977496459375$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-1 \ 2 \ 0 \ 1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{423739183}{478923089887538176} + \left(-\frac{134871}{478923089887538176}\right)s, \quad J = \begin{pmatrix} 3436 & 2514 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 3436$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $175224734029175 = 5^2 \cdot 271 \cdot 44563 \cdot 580379$:

$$\begin{pmatrix} 35044946805835 & 34948528340994 & 31952374147505 & 20022178114138 & 31720664882343 & 5856137943217 & 28804890454724 & 3076278950829 & 24578025479534 & 1991843914734 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 5 & 0 & 0 & 4 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 469 factors with signed exponents of absolute value up to 21007029185640; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_2) = (2 \ 3 \ 2)$ in the coordinates of $\text{Cl}(K)/5$.

Entry 18: character $b + c$, column e_3

Prescribed character vector $(0 \ 1 \ 1)$, i.e. the values of x on the classes of the three stored generators of $\text{Cl}(K)$.

The extension and its normalized generator

Relative polynomial of L_x over K :

$$y^5 - 3691y^3 + (18s - 9)y^2 + 183575y + (-550s + 275)$$

Absolute polynomial of L_x over $\mathbf{Q}$ (degree 10):

$$y^{10} - 7382y^8 + 13990631y^6 + 3976118285y^4 - 292099987625y^2 + 4977496459375$$

The automorphism σ_x is determined by the image of the primitive element θ : the coordinates of $\sigma_x(\theta)$ in the integral basis of L_x are $(-1 \ 2 \ 0 \ 1 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0)$.

The pair (a', J) representing the torsion class

$$a' = -\frac{9195119}{5023060313370901} + \left(-\frac{17324}{5023060313370901}\right)s, \quad J = \begin{pmatrix} 1381 & 823 \\ 0 & 1 \end{pmatrix}.$$

$N(J) = 1381$. **Checked:** $(a')J^5 = \mathcal{O}_K$: pass.

The auxiliary pair (t, I')

I' is the ideal of L_x with the following HNF matrix in the integral basis of the absolute model; its absolute norm is $175224734029175 = 5^2 \cdot 271 \cdot 44563 \cdot 580379$:

$$\begin{pmatrix} 35044946805835 & 34948528340994 & 31952374147505 & 20022178114138 & 31720664882343 & 5856137943217 & 28804890454724 & 3076278950829 & 24578025479534 & 1991843914734 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 5 & 0 & 0 & 4 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The element t is stored in compact form as a product of 466 factors with signed exponents of absolute value up to 194953005306918; it is never expanded (its principal ideal and residues are evaluated factor by factor).

Sign check and certified value

Residue prime: the stored prime ideal above $\ell = 7$ with residue degree 1; there the quotient $N_x(t)/a'$ reduces to the residue of $+1$. Certified value: $D_x(e_3) = (2 \ 3 \ 2)$ in the coordinates of $\text{Cl}(K)/5$.