

Genetic Connectivity in a Montane Salamander (*Pseudoeurycea leprosa*)

PURDUE UNIVERSITY
FORT WAYNE

Department of Biology

Emily Brink, Galen Burrell, Nicki Felger, Emmanuel Green

Molecular Ecology and Evolution (FNR 505000) Data Analysis Project

Introduction

- Salamander migration is restricted in montane habitats.
- Populations become isolated spatially and genetically.
- Velo-Antón et al. (2013) used microsatellites to investigate genetic structure between populations
- Using STRUCTURE analysis, they found 7 distinct genetic clusters in 23 sampled populations
- Considerable population admixture in two regions of the study area:
 - Central populations
 - Eastern populations

Objective

- Compare two methods of assessing genetic variation
 - STRUCTURE
 - Discriminant Analysis of Principle Components (DAPC)
- Does our DAPC analysis support/refute author conclusions?

Methods

- 351 individuals sampled from 23 localities – 9 microsatellites amplified and analyzed
- DAPC generated using *Adegenet* package in Rstudio
- Calculated DAPC for 7 clusters (same as original paper)

Synthesis of Results from Research Studies





Fig 1: Two maps describing salamander populations sampled in mountains of central Mexico. Results of Bayesian phylogeny (left) and STRUCTURE analysis (right) by Velo-Antón et al. (2013).

Fig 2: Results of DAPC (right) using K=7 clusters provides evidence for four distinct genetic groups. Distance between clusters indicates genetic differentiation.



Clusters Assigned by DAPC Analysis




Fig 3: Table displaying assignment of individuals from sampled populations to DAPC clusters (left). Original populations are color coded relative to their assignment in STRUCTURE. Red line indicates central/eastern division.

Conclusions

- Our analysis agrees with findings in original paper
- Based on DAPC, 4 genetically differentiated groups (Fig. 2)
 - Central and eastern groups in different clusters
- Significant admixture within central and eastern groups also predicted by DAPC (Fig. 3)
- DAPC suggested 9 clusters as the optimum number for the data – reveals more diversity in eastern populations
- STRUCTURE discerned greater diversity within central populations
- More sampling in these populations would give greater detail on genetic structure
- Future spatial PCA analysis – need individual GPS data

Acknowledgements

Our deepest gratitude extends to our professor Dr. Mark Jordan and our fellow classmates.

Literature Cited

Velo-Antón, G., Parra, J. L., Parra-Olea, G., & Zamudio, K. R. (2013). Tracking climate change in a dispersal-limited species: reduced spatial and genetic connectivity in a montane salamander. *Molecular Ecology*, 22(12), 3261–3278.

Jombart, T., & Collins, C. (2015). A tutorial for discriminant analysis of principal components (DAPC) using adegenet 2.0.0. London: Imperial College London, MRC Centre for Outbreak Analysis and Modelling.

Jombart, T., Devillard, S., & Balloux, F. (2010). Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. *BMC Genetics*, 11(1), 94.

R Core Team (2021). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.