

Rigidification and the Coherent Nerve for enriched Quasicategories

Harry Gindi

Abstract

We introduce, for \mathcal{C} a regular Cartesian Reedy category a model category whose fibrant objects are an analogue of quasicategories enriched in simplicial presheaves on \mathcal{C} . We then develop a coherent realization and nerve for this model structure and demonstrate using an enriched version of the necklaces of Dugger and Spivak that our model category is Quillen-equivalent to the category of categories enriched in simplicial presheaves on \mathcal{C} . We then show that for any Cartesian-closed left-Bousfield localization of the category of simplicial presheaves on \mathcal{C} , the coherent nerve and realization descend to a Quillen equivalence on the localizations of these model categories. As an application, we demonstrate a version of Yoneda's lemma for these enriched quasicategories.

Contents

| | | |
|----------|---|-----------|
| 1 | Formal \mathcal{C}-quasicategories | 4 |
| 1.1 | The wreath product with Δ | 4 |
| 1.2 | The generalized intertwiner and $\widehat{\Delta} \int \widehat{\mathcal{C}}$ | 6 |
| 1.3 | The horizontal Joyal model structure | 8 |
| 1.4 | The corner tensor construction | 9 |
| 1.5 | The regular Reedy structure of $\Theta[\mathcal{C}]$ | 16 |
| 1.6 | The anodyne theorem for horizontal inner anodynes | 17 |
| 1.7 | Comparison with Rezk's complete $\Theta[\mathcal{C}]$ -spaces | 24 |
| 1.8 | Recognition of horizontal Joyal fibrations | 29 |
| 2 | The Coherent Nerve, Horizontal case | 33 |
| 2.1 | The coherent realization for $\Theta[\mathcal{C}]$ | 33 |
| 2.2 | Necklaces and the coherent realization | 37 |
| 2.3 | Gadgets | 39 |
| 2.4 | Quillen functoriality | 40 |
| 2.5 | The Hom by cosimplicial resolutions for \mathcal{C} -precategories | 43 |
| 2.6 | The Hom by cosimplicial resolution | 45 |
| 2.7 | The horizontal comparison theorem | 50 |

| | | |
|----------|--|-----------|
| 3 | The Coherent Nerve, Local case | 53 |
| 3.1 | The $(\mathcal{C}, \mathcal{P})$ -enriched model structure | 53 |
| 3.2 | The Yoneda embedding and Yoneda's lemma | 56 |
| 3.3 | Weighted limits and colimits | 58 |
| 3.4 | Examples | 59 |

Introduction

In his thesis, David Oury introduced machinery to give a novel proof that his constructed model structure on Θ_2 -sets is Cartesian-monoidal closed. Charles Rezk constructed a model structure on Θ_2 -spaces that is known to be Quillen biequivalent to Oury's model structure. However, Rezk's construction allows for enrichment in a larger class of model categories, namely Cartesian-closed model categories whose underlying categories are simplicial presheaves on a small category \mathcal{C} satisfying some tame restrictions.

Bergner and Rezk also showed by means of a zig-zag of Quillen equivalences that Θ_n -spaces model the same homotopy theory as $\text{Psh}_\Delta(\Theta_{n-1})$ -enriched categories. Because the equivalence is indirect, however, many of the ideas from Lurie's work on $(\infty, 1)$ -categories cannot be adapted in a straightforward manner. In order to rectify this, we construct a generalized version of the coherent realization and nerve using Θ_n -sets (or more generally $\Theta[\mathcal{C}]$ -sets), and we demonstrate that this adjunction is a Quillen equivalence using an enriched version of Dugger and Spivak's calculus of necklaces.

In fact, our result is strictly stronger than the result of Bergner and Rezk because it allows us to account at the very least for the new case $\Theta = \Theta_\omega$, which is regular Cartesian Reedy by [Ber07], which the Bergner-Rezk approach could not handle, since one of the categories appearing in the zig-zag (the height- n analogue of Segal categories) does not make sense for $\mathcal{C} \neq \Theta_n$ for n finite. Their approach goes through rigidification results for simplicial algebras of algebraic theories due to Badzioch.

Our results work in more generality, although unfortunately not in complete generality. We are forced to impose stringent conditions on \mathcal{C} , but we stop well-short of requiring it to be a model of an algebraic theory.

Formal \mathcal{C} -quasicategories

In the first chapter, we apply a general construction to define what we call *labeled simplicial sets* with respect to a monoidal category \mathcal{V} . We then specialize to the case where \mathcal{V} is the Cartesian-monoidal category of presheaves of sets on a small category \mathcal{C} , which we additionally require to be a special kind of Reedy category that axiomatizes a form of the Eilenberg-Zilber shuffle decomposition for products of simplicial sets. We then define $\Theta[\mathcal{C}]$ to be the full subcategory of the labeled simplicial sets whose underlying simplicial sets are simplices and whose edges are all labeled by representable presheaves on \mathcal{C} .

We define the category of \mathcal{C} -cellular sets to be the category of presheaves of sets on this category. We then apply machinery of Cisinski and Oury to construct the *horizontal Joyal model structure* on the category of \mathcal{C} -cellular sets that has many of the familiar nice properties of the Joyal model structure. We call the fibrant objects of this category the *formal \mathcal{C} -quasicategories*.

Among the important technical results proven in this chapter, we direct the attention of the reader to Theorem 1.6, proving that the model structure is Cartesian-monoidal as well as 1.7, proving a

useful equivalence with an analogous model structure constructed by Rezk.

The chapter culminates with a key technical result that gives a characterization of the fibrant objects by a simple lifting property and the fibrations between them as the *isofibrations*, namely the horizontal inner fibrations that have the right lifting property with respect to the inclusion of a vertex into a freestanding isomorphism, extending an important theorem of Joyal to this setting.

The Coherent Nerve, Horizontal case

In the second chapter, we define an extension of Lurie’s coherent realization functor \mathfrak{C}_Δ to our setting. We leverage the equivalence between \mathcal{C} -indexed simplicially-enriched categories with a constant set of objects and $\text{Psh}_\Delta(\mathcal{C})$ -enriched categories to induce this functor pointwise from \mathfrak{C}_Δ . We then work to give an explicit calculation of this functor on representables and more generally on \mathcal{C} -labeled simplices.

We use the pointwise characterization of this realization to straightforwardly extend the results of Dugger and Spivak [DS11a] on alternative realizations to our setting, while on the other hand, we make use of the explicit characterization to demonstrate directly that the coherent realization and its right adjoint, the coherent nerve, form a Quillen pair.

$$\mathfrak{C} : \widehat{\Theta[\mathcal{C}]} \rightleftarrows \mathbf{Cat}_{\text{Psh}_\Delta(\mathcal{C})_{\text{inj}}} : \mathfrak{N}$$

For the next step in this chapter, we introduce cosimplicial resolutions in order to compute mapping objects for formal \mathcal{C} -quasicategories. We extend ideas from [DS11b] to demonstrate that the coherent nerve and realization actually specify a Quillen equivalence.

The Coherent Nerve, Local case

In this chapter, we give a way to perform a left-Bousfield localization of the horizontal Joyal model structure with respect to Cartesian presentations of the form $(\mathcal{C}, \mathcal{S})$ (though still under the hypothesis that \mathcal{C} is regular Cartesian Reedy). The local objects are exactly the formal \mathcal{C} -quasicategories whose mapping objects are \mathcal{S} -local. Using our comparison theorem with Rezk’s model structure, we can apply his result to show that this model structure is again Cartesian monoidal.

To prove the main result of the paper, we use the compatibility of the coherent realization and nerve with the formation of mapping objects to demonstrate that they remain Quillen equivalences after simultaneous localization.

As a corollary of the main result, we apply a theorem of Lurie to construct a Yoneda embedding. We then demonstrate that it is fully faithful and also prove Yoneda’s lemma, which we then leverage to define representability. From representability, we propose a definition of weighted limits and colimits.

Looking forward

A major challenge in the theory of higher categories is the problem of coherence, that is to say, defining functors and appropriately-natural transformations valued in a higher category of higher categories. It was observed early as the 1970 that a powerful way to deal with coherence problems even for functors from a 1-category to the 2-category of categories was to perform a rectification of that theory to the theory of Cartesian fibrations.

Lurie extended this point of view to the theory of $(\infty, 1)$ -categories for two reasons: Less crucially, one can use the theory of Cartesian fibrations to work with $(\infty, 2)$ -categorical notions without ever

actually giving a definition of $(\infty, 2)$ -category. Much more important than this shortcut, however, is the fact that Cartesian fibrations greatly simplify coherence problems.

Unfortunately, our paper does not even begin to scratch the surface of the fibrational point of view, and as a consequence, it is much more difficult to work in our setting in light of the consequent coherence problems. We expect that to understand the fibrational point of view, attempts will have to be made to understand higher-categorical lax structure. Lax structure is better-understood in the strict setting due to recent work of Ara, Maltsiniotis, and Steiner, but all attempts thusfar to extend these highly combinatorial results to the theory of weak higher categories have produced no tangible results. We suspect that this might change in the future when an equivalence theorem between the Complicial model of Verity and the Θ -style model studied here is proven.

We hope also that new approaches to dealing with coherence problems might be discovered, and if they can be made to work, we expect that the results of this paper will be even more useful.

Acknowledgements

First and foremost, we would like to thank George Raptis for advising us in the writing of the version of this paper to be submitted as a Master's thesis at Universität Regensburg. His suggestions have been invaluable, and we are extremely grateful for the time he has spent reading this paper closely.

We would also like to thank Denis-Charles Cisinski for his help with axiomatizing the Eilenberg-Zilber shuffle decomposition 1.1.3, which is an important technical condition without which many of the constructions in this paper would not function.

We would like to give a special thanks Alexander Campbell for his invaluable help in correcting the proof of Theorem 1.8.6, which previously cited an incorrect result.

Finally, we would like to acknowledge Andrea Gagna for spending his time discussing these ideas with us over several long conversations as well as Eric Peterson for his sage advice and friendship over the years.

1 Formal \mathcal{C} -quasicategories

1.1 The wreath product with Δ

In this section, we will consider a slightly more general definition of the wreath product with Δ , as in Oury's thesis. Segal observed long ago that a monoidal category is classified precisely by a pseudofunctor $M_\bullet : \Delta^{\text{op}} \rightarrow \mathbf{Cat}$ such that $M_0 = *$ is the terminal category and the maps $M_n \rightarrow (M_1)^n$ induced by the inclusion of the spine $Sp[n] \hookrightarrow \Delta^n$ are all isomorphisms. This brings us to our first definition:

Definition 1.1.1. Suppose \mathcal{V} is a monoidal category. Then we construct a Grothendieck fibration

$$\Delta \int \mathcal{V} \rightarrow \Delta$$

by applying the Grothendieck construction to the functor

$$\mathcal{V}_\bullet : \Delta^{\text{op}} \rightarrow \mathbf{Cat}$$

classifying \mathcal{V} . We call the total space of this fibration the *wreath product* of Δ with \mathcal{V} .

Now, notice that since \mathbf{Cat} is complete, the functor \mathcal{V}_\bullet extends uniquely along the co-Yoneda embedding to a continuous functor

$$\widehat{\mathcal{V}_\bullet^{\text{op}}}^{\text{op}} : \widehat{\Delta}^{\text{op}} \rightarrow \mathbf{Cat},$$

which is exactly the right Kan extension of \mathcal{V}_\bullet along the co-Yoneda embedding $\Delta^{\text{op}} \hookrightarrow \widehat{\Delta}^{\text{op}}$. Applying the Grothendieck construction to the functor $\widehat{\mathcal{V}_\bullet^{\text{op}}}^{\text{op}}$, we define the Grothendieck fibration

$$\widehat{\Delta} \int \mathcal{V} \rightarrow \widehat{\Delta}.$$

The total space of this fibration is called the category of \mathcal{V} -labeled simplicial sets.

Proposition 1.1.2. *The pullback of the fibration*

$$\widehat{\Delta} \int \mathcal{V} \rightarrow \widehat{\Delta}$$

along the Yoneda embedding $\Delta \hookrightarrow \widehat{\Delta}$ is exactly the fibration

$$\Delta \int \mathcal{V} \rightarrow \Delta,$$

and therefore, the induced map

$$\Delta \int \mathcal{V} \hookrightarrow \widehat{\Delta} \int \mathcal{V}$$

is a fully faithful embedding.

Proof. As the functor \mathcal{V}_\bullet factors as the composite

$$\Delta^{\text{op}} \hookrightarrow \widehat{\Delta}^{\text{op}} \xrightarrow{\widehat{\mathcal{V}_\bullet^{\text{op}}}^{\text{op}}} \mathbf{Cat},$$

where the first functor is fully faithful, it follows that $\Delta \int \mathcal{V} \rightarrow \Delta$ is the pullback of the fibration $\widehat{\Delta} \int \mathcal{V} \rightarrow \widehat{\Delta}$ along the fully faithful Yoneda embedding. Ergo, the map in question is fully faithful. \square

For the purposes of this paper, we do not need this level of generality. We specialize as follows:

Definition 1.1.3. A small regular skeletal Reedy category \mathcal{C} (see [Cis06, 8.2.3]) is called a *regular Cartesian Reedy category* if it satisfies two conditions:

(CR1) The class of regular presheaves on \mathcal{C} (see [Cis06, 8.2]) is closed under finite products.

(CR2) If I is a finite set and $c \rightarrow \prod_{i \in I} c_i$ is a nondegenerate section, then $\dim c \leq \sum_{i \in I} \dim c_i$.

Note 1.1.4. The axioms for regular Cartesian Reedy categories imply that \mathcal{C} is a Reedy multicategory in the sense of [BR11]. It also asserts a weak form of the Eilenberg-Zilber shuffle decomposition. There may be a way to prove (CR2) from (CR1), but we were unable to do so.

In the sequel, we assume that $\mathcal{V} = \widehat{\mathcal{C}}$ is the category of presheaves of sets on a small regular Cartesian Reedy category \mathcal{C} admitting a terminal object. Then we give the following definition:

Definition 1.1.5. For any regular Cartesian Reedy category \mathcal{C} admitting a terminal object, we define the category of \mathcal{C} -cells

$$\Theta[\mathcal{C}] \subseteq \Delta \int \widehat{\mathcal{C}}$$

to be the full subcategory spanned by the objects of the form $[n](h_{c_1}, \dots, h_{c_n})$ for $c_1, \dots, c_n \in \mathcal{C}$, and where h_\bullet denotes the Yoneda embedding.

Remark 1.1.6. The requirement that \mathcal{C} have a terminal object is a technical condition that ensures that Δ embeds fully and faithfully in $\Theta[\mathcal{C}]$. The condition that \mathcal{C} is regular Cartesian Reedy is probably not necessary, but it will ensure later on that the generating cofibrations of the injective model structure on simplicial presheaves $\text{Psh}_\Delta(\mathcal{C})$ admit a very simple description.

Remark 1.1.7. When \mathcal{C} has a terminal object, the inclusion $* \rightarrow \mathcal{C}$ is a fully faithful right-adjoint. The construction Θ preserves fully faithful right-adjoints, so we have a right-adjoint functor

$$\Delta = \Theta[*] \hookrightarrow \Theta[\mathcal{C}].$$

Passing to presheaf categories, this functor also extends to a colimit-preserving functor

$$\mathcal{H} : \widehat{\Delta} \hookrightarrow \widehat{\Theta[\mathcal{C}]}$$

In particular, on representables we have that

$$\mathcal{H}(\Delta^n) = [n](*, \dots, *).$$

Therefore, we have an adjoint triple,

$$\begin{array}{ccc} & \xrightarrow{\pi} & \\ \widehat{\Theta[\mathcal{C}]} & \xleftarrow[\mathcal{H}]{\perp} & \widehat{\Delta} \\ & \xrightarrow[\mathcal{N}]{\perp} & \end{array}$$

Definition 1.1.8. We define a special cosimplicial object in $\widehat{\Theta[\mathcal{C}]}$ by the formula

$$E^\bullet = \mathcal{H}(\text{cosk}_0 \Delta^\bullet).$$

This cosimplicial object will be a cosimplicial resolution of a point, once we define our model structures.

1.2 The generalized intertwiner and $\widehat{\Delta} \int \widehat{\mathcal{C}}$

Rezk introduced an intertwining functor by means of an explicit construction, but Oury gave an even more powerful version, which we recall here:

Definition 1.2.1. Recall that we have a fully-faithful embedding

$$L : \Theta[\mathcal{C}] \hookrightarrow \Delta \int \widehat{\mathcal{C}} \hookrightarrow \widehat{\Delta} \int \widehat{\mathcal{C}}.$$

We define the *intertwiner* to be the functor

$$\square : \widehat{\Delta} \int \widehat{\mathcal{C}} \rightarrow \widehat{\Theta[\mathcal{C}]}$$

by the formula

$$(S, \Omega) \mapsto S \square \Omega = \text{Hom}_{\widehat{\Delta} \int \widehat{\mathcal{C}}}(L(\cdot), (S, \Omega)).$$

Note 1.2.2. The restriction of the intertwiner to $\Delta \int \widehat{\mathcal{C}}$ is exactly the intertwiner of Rezk. When we apply the intertwiner to an object belonging to the full subcategory $\Delta \int \widehat{\mathcal{C}}$, that is, $(S, \Omega) = [n](A_1, \dots, A_n)$, we will switch to Rezk's notation, namely

$$V[n](A_1, \dots, A_n) \stackrel{\text{def}}{=} S \square \Omega$$

Definition 1.2.3. An object (S, Ω) of $\widehat{\Delta} \int \widehat{\mathcal{C}}$ is called *normalized* if the image of Ω_1 does not contain the empty presheaf on \mathcal{C} .

Proposition 1.2.4. *The restriction of the intertwiner to the full subcategory of normalized objects in $\widehat{\Delta} \int \widehat{\mathcal{C}}$ is fully faithful.*

Proof. Recall before we begin that a map $(S, \Omega) \rightarrow (S', \Omega')$ is given by a morphism of simplicial sets $f : S \rightarrow S'$ and a natural modification $\zeta : \Omega \rightarrow \Omega' \circ f$.

In order to prove fullness, let $\gamma : S \square \Omega \rightarrow S' \square \Omega'$ be a map in $\widehat{\Theta}[\widehat{\mathcal{C}}]$. We notice that $\text{Hom}([n](\emptyset, \dots, \emptyset), (S, \Omega))$ is naturally isomorphic to S_n , and proceed by diagram chase. Since by assumption (S, Ω) is normalized, every map $[n](\emptyset, \dots, \emptyset) \rightarrow (S, \Omega)$ factors uniquely through at least one map $[n](c_1, \dots, c_n) \rightarrow (S, \Omega)$.

Choosing such a factorization, the natural transformation γ sends this to a map

$$[n](c_1, \dots, c_n) \rightarrow (S', \Omega'),$$

and finally, precomposing this map with the unique map $[n](\emptyset, \dots, \emptyset) \rightarrow [n](c_1, \dots, c_n)$, we obtain a map $[n](\emptyset, \dots, \emptyset) \rightarrow (S', \Omega')$. Taking these together gives a map $S_n \rightarrow S'_n$, naturally in n .

Now assume that $S' = S$ and that the map induced by γ is the identity. Then notice that a map

$$[n](c_1, \dots, c_n) \rightarrow (S, \Omega)$$

is completely determined by its action on the degree n part, but this amounts to picking an n -simplex of S together with its labeling (A_1, \dots, A_n) , and a map $(c_1, \dots, c_n) \rightarrow (A_1, \dots, A_n)$. Then the natural transformation gives a natural map $((A_1)_{c_1}, \dots, (A_n)_{c_n}) \rightarrow ((A'_1)_{c_1}, \dots, (A'_n)_{c_n})$, taking the naturality in n and the c_i , these together determine a natural modification $\Omega \rightarrow \Omega'$.

To see faithfulness, notice that the construction in the proof of fullness defines a left-inverse to the definition of the map on morphisms defined by the intertwiner. \square

Definition 1.2.5. We call a presheaf of sets on $\Theta[\mathcal{C}]$ a \mathcal{C} -cellular set.

Note 1.2.6. Although the case when $\mathcal{C} = \Theta_{n-1}$ (respectively $\mathcal{C} = \Theta = \Theta_\omega$) are not strictly the focus of this paper, note that $\Theta[\Theta_{n-1}] = \Theta_n$ (respectively $\Theta[\Theta] = \Theta$). In these cases, we call presheaves of sets on $\Theta[\mathcal{C}]$ *n-cellular sets* (respectively, *cellular sets*).

Definition 1.2.7. We say that a \mathcal{C} -cellular set X is *sober* if it is the image of a normalized object of $\widehat{\Delta} \int \widehat{\mathcal{C}}$. If $f : X \rightarrow Y$ is the image under the intertwiner of a Cartesian map of normalized labeled simplicial sets, we call f *Cartesian*.

Proposition 1.2.8. *All representable \mathcal{C} -cellular sets are sober.*

Proof. By construction. \square

Proposition 1.2.9. *The class of sober \mathcal{C} -cellular sets is closed under Cartesian product.*

Proof. From the construction of the intertwiner, we see that if $\widehat{\Delta} \int \widehat{\mathcal{C}}$ has all Cartesian products, then the intertwiner preserves them, since

$$\mathrm{Hom}_{\widehat{\Delta} \int \widehat{\mathcal{C}}}(L(\cdot), (S, \Omega) \times (S', \Omega')) = \mathrm{Hom}_{\widehat{\Delta} \int \widehat{\mathcal{C}}}(L(\cdot), (S, \Omega)) \times \mathrm{Hom}_{\widehat{\Delta} \int \widehat{\mathcal{C}}}(L(\cdot), (S', \Omega')).$$

Then we define the Cartesian product of (S, Ω) and (S', Ω') by the formula

$$S \times S' \xrightarrow{\Omega \times \Omega'} \widehat{\mathcal{C}}_{\bullet} \times \widehat{\mathcal{C}}_{\bullet} \xrightarrow{\times} \widehat{\mathcal{C}}_{\bullet}.$$

It is clear that this satisfies the universal property of the product. \square

Proposition 1.2.10. *The functor associated with the fibration $\Theta[\mathcal{C}] \rightarrow \Delta$ induces an adjunction*

$$\widehat{\Theta[\mathcal{C}]} \xrightleftharpoons[\mathcal{H}]{\pi} \widehat{\Delta},$$

as we saw earlier. If $X = S \square \Omega$ is sober, and $f : S' \rightarrow S$ is a map of simplicial sets, then the image of the Cartesian lift $\tilde{f} : (S', f^*(\Omega)) \rightarrow (S, \Omega)$ under the intertwiner is exactly the pullback of $\mathcal{H}(f)$ along the component at X of the unit of the adjunction $\mu_X : X \rightarrow \mathcal{H}\pi X = \mathcal{H}S$.

Proof. By inspection of the definition of $S \square \Omega$, we can see that $\pi(S \square \Omega) = S$.

We can see that \mathcal{H} factors as $\square \circ \mathfrak{t}$, where \mathfrak{t} is the right-adjoint to the projection $\widehat{\Delta} \int \widehat{\mathcal{C}} \rightarrow \widehat{\Delta}$, which exists by explicit computation as the functor sending the simplicial set S to the object $(S, \Omega_{\mathfrak{t}})$ where $\Omega_{\mathfrak{t}}$ is the labeling sending all simplices of S to the terminal presheaf on \mathcal{C} . We can see that the pullback of (S, Ω) along $f : S' \rightarrow S$ satisfies the universal property of the fibre product of the unit map $(S, \Omega) \rightarrow \mathfrak{t}(S)$ with the map $\mathfrak{t}(f)$, so such pullbacks exist in $\widehat{\Delta} \int \widehat{\mathcal{C}}$ and are obviously preserved by \square , which by construction preserves whatever limits exist. The proposition follows immediately from these two observations. \square

1.3 The horizontal Joyal model structure

We define a Cisinski model structure on $\widehat{\Theta[\mathcal{C}]}$ and state several results that will be proven over the next few sections.

Definition 1.3.1. There is a Cisinski model structure called the *horizontal Joyal model structure* on $\widehat{\Theta[\mathcal{C}]}$ where the separating interval is given by

$$E^1 = \mathcal{H}(\mathrm{cosk}_0 \Delta^1),$$

and the set of generating anodynes is given by

$$\mathcal{J} = \{\square_n^{\perp}(\lambda_k^n, \delta^{c_1}, \dots, \delta^{c_n}) : 0 < k < n \text{ and } c_1, \dots, c_n \in \mathrm{Ob} \mathcal{C}\},$$

where $\lambda_k^n : \Lambda_k^n \hookrightarrow \Delta^n$ is the simplicial horn inclusion, and where $\delta^c : \partial c \hookrightarrow c$ is the inclusion of the boundary of c (recall that \mathcal{C} was taken to be a regular Cartesian Reedy category, so this makes sense).

We call $\mathrm{rlp}(\mathcal{J})$ the class of *horizontal inner fibrations*, and we call $\mathrm{llp}(\mathrm{rlp}(\mathcal{J}))$ the class of *horizontal inner anodynes*.

Remark 1.3.2. The precise definition and construction of the corner-intertwiner \square_n^\perp is deferred to Section 1.4, but in this particular case, we can compute it by hand in terms of the intertwiner to be

$$V_{\Lambda_k^n}(c_1, \dots, c_n) \cup \left(\bigcup_{i=1}^n V[n](c_1, \dots, \partial c_i \dots, c_n) \right) \hookrightarrow [n](c_1, \dots, c_n),$$

where $V_{\Lambda_k^n}(c_1, \dots, c_n)$ is the pullback of $[n](c_1, \dots, c_n)$ by the inclusion $\Lambda_k^n \hookrightarrow \Delta^n$ (whenever $K \subseteq \Delta^n$, we can apply this formula to compute the corner tensor).

Definition 1.3.3. We call an object with the right lifting property with respect to \mathcal{J} a *formal \mathcal{C} -quasicategory*.

Note 1.3.4. In the case where \mathcal{C} is the terminal category, these are precisely the quasicategories, since the horns in the definition above become exactly the simplicial inner horn inclusions.

The following results are stated here without proof. All proofs are heavily inspired by [Our10] and provided in full in the sections 1.5, 1.6, and 1.8.

Proposition 1.3.5. *The class of all monomorphisms of $\widehat{\Theta[\mathcal{C}]}$ is exactly $\text{Cell}(\mathcal{M})$, where*

$$\mathcal{M} = \{\square_n^\perp(\delta^n, \delta^{c_1}, \dots, \delta^{c_n}) : n \geq 0 \text{ and } c_1, \dots, c_n \in \text{Ob } \mathcal{C}\}.$$

Proposition 1.3.6. *For any inner anodyne inclusion $\iota : K \hookrightarrow \Delta^n$ and any family f_1, \dots, f_n of monomorphisms of $\widehat{\mathcal{C}}$, the map*

$$\square_n^\perp(\iota, f_1, \dots, f_n)$$

is horizontal inner anodyne.

Theorem 1.3.7. *The horizontal Joyal model structure is Cartesian-closed, and in particular,*

$$\text{Cell}(\mathcal{M}) \times^\perp \text{Cell}(\mathcal{J}) \subseteq \text{Cell}(\mathcal{J}).$$

Theorem 1.3.8. *A horizontal inner fibration between formal \mathcal{C} -quasicategories is a fibration for the horizontal Joyal model structure if and only if it has the right lifting property with respect to the map $\Delta^0 \hookrightarrow E^1$. In particular, the formal \mathcal{C} -quasicategories are exactly the fibrant objects for the horizontal Joyal model structure.*

Proposition 1.3.9. *Given a necklace T , there is a unique embedding $\iota_T : T \hookrightarrow \Delta[T]$, where $\Delta[T]$ is the unique representable whose spine is exactly the spine of T . This map sends the spine of T isomorphically onto the spine of $\Delta[T]$ and is a horizontal inner anodyne.*

1.4 The corner tensor construction

The overwhelming majority of this section is due to Oury, although we had to redo some of the proofs, since they contained mistakes. Following [Our10, 3.1] we define the corner tensor, a vast generalization of the corner product.

Definition 1.4.1. Let \mathcal{V} be a cocomplete symmetric monoidal closed category, and let all categories, functors, and natural transformations in what follows to be \mathcal{V} -enriched. Suppose we have a category \mathcal{T} and an n -ary functor

$$\wedge : \mathcal{T}^{\otimes n} \rightarrow \mathcal{T}$$

be a functor. Let $(\mathcal{A}_i)_{i=1}^n$ and \mathcal{D} be categories admitting enough colimits such that all tensors with $\text{Hom}_{\mathcal{T}}$ exist and coends over \mathcal{T} exist. Let

$$\nabla_i : \mathcal{A}_i \otimes \mathcal{A}_i \rightarrow \mathcal{A}, \quad \nabla : \mathcal{D} \otimes \mathcal{D} \rightarrow \mathcal{D}, \quad \square : \mathcal{A}_1 \otimes \cdots \otimes \mathcal{A}_n \rightarrow \mathcal{D}$$

be functors. Then we define the following functors:

$$\nabla_i^\perp : \mathcal{A}_i^\mathcal{T} \otimes \mathcal{A}_i^\mathcal{T} \rightarrow \mathcal{A}^\mathcal{T}, \quad \nabla^\perp : \mathcal{D}^\mathcal{T} \otimes \mathcal{D}^\mathcal{T} \rightarrow \mathcal{D}^\mathcal{T}, \quad \square^\perp : \mathcal{A}_1^\mathcal{T} \otimes \cdots \otimes \mathcal{A}_n^\mathcal{T} \rightarrow \mathcal{D}^\mathcal{T}$$

by the Day convolution, for example,

$$\square^\perp(M_1, \dots, M_n)t = \int^{u_1, \dots, u_n} \mathcal{T}(\wedge(u_1, \dots, u_n), t) \otimes \square(M_1(u_1), \dots, M_n(u_n)).$$

Lemma 1.4.2. *The functor \square^\perp preserves all colimits preserved by \square in each variable and by \otimes in its second variable.*

Proof. By coend manipulation. □

We specialize now to the case where \mathcal{V} is just the category of sets and where $\mathcal{T} = [1]$ is the categorical 1-simplex. The functor $\wedge : [1]^n \rightarrow [1]$ is given by taking the infimum.

Note 1.4.3. We will often consider arrows in a category \mathcal{D} as functors $[1] \rightarrow \mathcal{D}$. By abuse of notation, we will denote the functor $[1] \rightarrow \mathcal{D}$ classifying an arrow $f : A \rightarrow B$ simply by f . We will denote the evaluation of this functor on the objects of $x \in [1]$ by $f(x)$, such that in the case of a map $f : A \rightarrow B$,

$$(1.4.4) \quad f(0) = A \quad \text{and} \quad f(1) = B$$

Definition 1.4.5. Given $\square : \mathcal{A}_1 \times \cdots \times \mathcal{A}_n \rightarrow \mathcal{D}$, where each category appearing is cocomplete, we define the *corner tensor* $\square^\perp : \mathcal{A}_1^{[1]} \times \cdots \times \mathcal{A}_n^{[1]} \rightarrow \mathcal{D}^{[1]}$ by the formula

$$\square^\perp(f_1, \dots, f_n)(t) = \int^{u_1, \dots, u_n} [1](u_1 \wedge \cdots \wedge u_n, t) \cdot \square(f_1(u_1), \dots, f_n(u_n)).$$

If $(g, h) : f_i \rightarrow f'_i$ is a commutative square, let

$$(g^\perp, h^\perp) : \square^\perp(f_1, \dots, f_i, \dots, f_n) \rightarrow \square^\perp(f_1, \dots, f'_i, \dots, f_n)$$

be the induced commutative square.

Example 1.4.6. If we take \square to be a bifunctor $\mathcal{D} \times \mathcal{D} \rightarrow \mathcal{D}$ in an appropriately cocomplete category, then given $f_1 : A \rightarrow B$ and $f_2 : C \rightarrow D$, their corner tensor is the familiar corner product:

$$f_1 \square^\perp f_2 = A \square D \coprod_{A \square C} B \square C \rightarrow B \square D.$$

Remark 1.4.7. The previous example is in some sense universal, and it will allow us to reduce certain questions about big corner tensors to binary ones. In particular, we will frequently use the observation that, in the situation of the example the map $f_1(0) \square f_2(1) \rightarrow (f_1 \square^\perp f_2)(0)$ is a pushout of the map $f_1(0) \square f_2(0) \rightarrow f_1(1) \square f_2(0)$.

Lemma 1.4.8. *If for any $i \in \{0, \dots, n\}$, the map $f_i : A_i \rightarrow B_i$ is an identity map, then the corner tensor $\square^\perp(f_1, \dots, f_n)$ is an identity map.*

Proof. Assume for simplicity that $i = 1$, which is without loss of generality by reindexing. Then the result follows by setting

$$U(s, t) = \int^{u_1, \dots, u_n} ([1](u_1, s) \times [1](u_2 \wedge \dots \wedge u_n, t)) \cdot \square(f_1(u_1), f_2(u_2), \dots, f_n(u_n)).$$

Notice that by Yoneda reduction,

$$\int^{s, t} [1](s \wedge t, x) \times [1](u_1, s) \times [1](u_2 \wedge \dots \wedge u_n, t) = [1](u_1 \wedge u_2 \wedge \dots \wedge u_n, x),$$

so we have that

$$\square^\perp(f_1, \dots, f_n)(x) = \int^{s, t} [1](s \wedge t, x) U(s, t),$$

which exhibits

$$\square^\perp(f_1, \dots, f_n)(0) = U(0, 1) \coprod_{U(0, 0)} U(1, 0),$$

but $U(0, 1) = \square(B_1, \dots, B_n)$, which demonstrates that the map $U(0, 0) \rightarrow U(1, 0)$ is the identity by cofinality. The map $U(0, 1) \rightarrow \square^\perp(f_1, \dots, f_n)(0)$ is the pushout of an identity map, and the map $U(0, 1) \rightarrow \square^\perp(f_1, \dots, f_n)(1)$ is also the identity, so it follows that the map $\square^\perp(f_1, \dots, f_n)$ must also be the identity. \square

Lemma 1.4.9. *Suppose \square preserves pushouts in the first argument and*

$$\begin{array}{ccc} A_1 & \xrightarrow{g} & A'_1 \\ f_1 \downarrow & & \downarrow f'_1 \\ B_1 & \xrightarrow{h} & B'_1 \end{array},$$

is a pushout square in \mathcal{A}_1 , then the induced square

$$(g^\perp, h^\perp) : \square^\perp(f_1, \dots, f_n) \rightarrow \square^\perp(f'_1, \dots, f'_n)$$

is a pushout square.

Proof. Let Q denote $\square^\perp(f_1, \dots, f_n)(0)$ and $Q' = \square^\perp(f'_1, \dots, f'_n)(0)$, with $g^\perp : Q \rightarrow Q'$ being the induced map on the domains of corner tensors. Let $(\alpha, \alpha') : \square(g, B_2, \dots, B_n) \rightarrow g^\perp$ be the induced commutative square

$$\begin{array}{ccc} \square(A_1, B_2, \dots, B_n) & \xrightarrow{\square(g, B_2, \dots, B_n)} & \square(A'_1, B_2, \dots, B_n) \\ \alpha \downarrow & & \downarrow \alpha' \\ Q & \xrightarrow{g^\perp} & Q' \end{array}.$$

Then we can factor the square

$$\begin{array}{ccc}
\Box(A_1, B_2, \dots, B_n) & \xrightarrow{\Box(g, B_2, \dots, B_n)} & \Box(A'_1, B_2, \dots, B_n) \\
\downarrow \Box(f_1, B_2, \dots, B_n) & & \downarrow \Box(f'_1, B_2, \dots, B_n) \\
\Box(B_1, B_2, \dots, B_n) & \xrightarrow{h^-} & \Box(B'_1, B_2, \dots, B_n)
\end{array}$$

as the composite square

$$\begin{array}{ccc}
\Box(A_1, B_2, \dots, B_n) & \xrightarrow{\Box(g, B_2, \dots, B_n)} & \Box(A'_1, B_2, \dots, B_n) \\
\downarrow \alpha & & \downarrow \alpha' \\
Q & \xrightarrow{g^\perp} & Q' \\
\downarrow \Box^\perp(f_1, \dots, f_n) & & \downarrow \Box^\perp(f'_1, \dots, f_n) \\
\Box(B_1, B_2, \dots, B_n) & \xrightarrow{h^-} & \Box(B'_1, B_2, \dots, B_n)
\end{array} .$$

where the outer square is a pushout square by the fact that \Box preserves colimits in the first argument. Then to prove the lemma, it suffices to prove the claim that (α, α') is a pushout square. Then we define

$$U(s, t) = \int^{e_1, \dots, e_n} ([1](e_1, s) \times [1](e_2 \wedge \dots \wedge e_n, t)) \cdot \Box(f_1(e_1), f_2(e_2), \dots, f_n(e_n)).$$

and

$$U'(s, t) = \int^{e_1, \dots, e_n} ([1](e_1, s) \times [1](e_2 \wedge \dots \wedge e_n, t)) \cdot \Box(f'_1(e_1), f_2(e_2), \dots, f_n(e_n)).$$

Then notice that

$$\alpha : \Box(A_1, B_2, \dots, B_n) \rightarrow Q$$

is the pushout of $u : U(0, 0) \rightarrow U(1, 0)$ and

$$\alpha' : \Box(A'_1, B_2, \dots, B_n) \rightarrow Q$$

is the pushout of $u' : U'(0, 0) \rightarrow U'(1, 0)$. So we have a commutative square (in which the vertices are arrows and the arrows are commutative squares)

$$\begin{array}{ccc}
u & \xrightarrow{\quad} & \alpha \\
\downarrow & & \downarrow \\
u' & \xrightarrow{\quad} & \alpha'
\end{array} .$$

where $u \rightarrow \alpha$ and $u' \rightarrow \alpha'$ are coCartesian. Then by the pasting law for pushouts, it suffices to show that the map $u \rightarrow u'$ is also coCartesian. By cofinality, we can see that the map $u : U(0, 0) \rightarrow U(1, 0)$ is induced by the natural family of maps

$$\square(f_1, f_2(e_2), \dots, f_n(e_n)) : \square(A_1, f_2(e_2), \dots, f_n(e_n)) \rightarrow \square(B_1, f_2(e_2), \dots, f_n(e_n)),$$

that $u' : U'(0, 0) \rightarrow U'(1, 0)$ is induced by the natural family of maps

$$\square(f'_1, f_2(e_2), \dots, f_n(e_n)) : \square(A'_1, f_2(e_2), \dots, f_n(e_n)) \rightarrow \square(B'_1, f_2(e_2), \dots, f_n(e_n)),$$

that $U(0, 0) \rightarrow U'(0, 0)$ is induced by the natural family of maps

$$\square(g, f_2(e_2), \dots, f_n(e_n)) : \square(A_1, f_2(e_2), \dots, f_n(e_n)) \rightarrow \square(A'_1, f_2(e_2), \dots, f_n(e_n)),$$

and that $U(1, 0) \rightarrow U'(1, 0)$ is induced by the natural family of maps

$$\square(h, f_2(e_2), \dots, f_n(e_n)) : \square(B_1, f_2(e_2), \dots, f_n(e_n)) \rightarrow \square(B'_1, f_2(e_2), \dots, f_n(e_n)).$$

But these form a natural pushout square, which implies that $u \rightarrow u'$ is a pushout because \square preserves colimits in the first argument and the pushout moves outside of the coend. This proves the claim that $\alpha \rightarrow \alpha'$ is a pushout, which proves the lemma. \square

Note 1.4.10. The above holds for any $i \in \{1, \dots, n\}$ as well, but for simplicity of notation, we proved it only in the case $i = 1$.

Corollary 1.4.11. *If \square preserves pushouts in the i^{th} argument, then if f_i is a pushout of f'_i , it follows that $\square^\perp(f_1, \dots, f_i, \dots, f_n)$ is a pushout of $\square^\perp(f_1, \dots, f'_i, \dots, f_n)$.*

Note 1.4.12. The next few lemmata involve very complicated diagrams. As such, we will leave the objects appearing in these diagrams as implicit.

Lemma 1.4.13. *Suppose again that \square preserves pushouts in its first argument, and suppose we have a coCartesian square in \mathcal{A}*


$$\begin{array}{ccc} \cdot & \xrightarrow{h} & \cdot \\ \downarrow u & & \downarrow v \\ \cdot & \xrightarrow{g} & \cdot \end{array},$$

and suppose that we have a commutative square

$$\begin{array}{ccc} \cdot & \xrightarrow{g} & \cdot \\ \downarrow p & & \downarrow q \\ \cdot & \xrightarrow{\text{id}} & \cdot \end{array}.$$

Given family of maps $\mathbf{f} = (f_i \in \mathcal{A}_i^{[1]})_{i=2}^n$, let $Q_{\bullet, \mathbf{f}}$ denote the evaluation of $\square^\perp(\bullet, \mathbf{f})$ at 0. Then $g^\perp : Q_{p, \mathbf{f}} \rightarrow Q_{q, \mathbf{f}}$ is a pushout of $\square^\perp(h, \mathbf{f})$.


Proof. The data allow us to construct a commutative cube



Since the front and back faces of this cube are coCartesian, it gives a coCartesian square in $\mathcal{A}^{[1]}$,

$$\begin{array}{ccc} h & \xrightarrow{(h, \text{id})} & \text{id} \\ \downarrow (u, qv) & & \downarrow (v, qv) \\ p & \xrightarrow{(g, \text{id})} & q \end{array}$$

Then applying $\square^\perp(\bullet, \mathbf{f})$, we have a commutative cube



its front and back faces remain pushouts, since \square^\perp preserves all colimits in each argument preserved by \square . Then the map $g^\perp : Q_{p, \mathbf{f}} \rightarrow Q_{q, \mathbf{f}}$ is a pushout of the map $h^\perp : Q_{h, \mathbf{f}} \rightarrow Q_{\text{id}, \mathbf{f}}$, but by Lemma 1.4.8, we see that $\square^\perp(\text{id}, \mathbf{f}) = \text{id}$, so by commutativity, it follows that $h^\perp = \square^\perp(h, \mathbf{f})$, and therefore, g^\perp is a pushout of $\square^\perp(h, \mathbf{f})$. \square

Assume in the sequel that \square preserves connected colimits in each argument.

Lemma 1.4.14 ([Our10, Lemma 3.10]). *Let $(\mathcal{J}_i)_{i=1}^n$ be a family of sets of morphisms of each \mathcal{A}_i . Then*

$$\square^\perp(\mathcal{J}_1, \dots, \text{Cell}(\mathcal{J}_k), \dots, \mathcal{J}_n) \subseteq \text{Cell}(\square^\perp(\mathcal{J}_1, \dots, \mathcal{J}_k, \dots, \mathcal{J}_n)).$$

Proof. Again, by symmetry, it is enough to prove this for the case $k = 1$. Let f_1 be a map in \mathcal{A}_1 that belongs to $\text{Cell}(\mathcal{J}_1)$. Then this gives the data of a cocontinuous diagram $D : \alpha \rightarrow \mathcal{A}_1$ for an ordinal α with colimit C and structure maps $\phi_i : D(i) \rightarrow C$ such that $f_1 = \phi_0$. Additionally, for any $i < \alpha$, the maps $g_i : D(i) \rightarrow D(i+1)$ are pushouts of maps belonging to \mathcal{J}_1 . Let $D^+ : \alpha^\flat \rightarrow \mathcal{A}_1$ be the extension of D sending α to C and such that $D^+(i \rightarrow \alpha) = \phi_i$ for $i < \alpha$. Define $D^2 : \alpha^\flat \rightarrow \mathcal{A}_1^{[1]}$ by the rule $i \mapsto \phi_i$ for $i < \alpha$ and $\alpha \mapsto \text{id}_C$ and sending $i \rightarrow i+1$ to the commutative square $(g_i, \text{id}_C) : \phi_i \rightarrow \phi_{i+1}$ and sending the map $i \rightarrow \alpha$ to the commutative square $(\phi_i, \text{id}_C) : \phi_i \rightarrow \text{id}_C$. Then the colimit of D^2 is id_C as pictured below.

$$\begin{array}{ccccccc} D(0) & \xrightarrow{g_0} & D(1) & \xrightarrow{g_1} & \cdots & D(i) & \cdots \longrightarrow C \\ \downarrow \phi_0 & & \downarrow \phi_1 & & & \downarrow \phi_i & \parallel \\ C & \xlongequal{\quad} & C & \xlongequal{\quad} & \cdots & C & \cdots \xlongequal{\quad} C \end{array}.$$

Let $f_j : X_i \rightarrow Y_j \in \mathcal{J}_j$ for each $2 \leq j \leq n$. Denote $\square^\flat(\bullet, f_2, \dots, f_n)$ by $\square(\bullet, \mathbf{f})$ and the domain by $Q_{\bullet, \mathbf{f}}$. Then since \square^\flat preserves connected colimits argument by argument, it follows that

$$\text{colim } \square^\flat(D^2, \mathbf{f}) = \square^\flat(\text{id}_C, \mathbf{f}),$$

and so we have the structure maps of the colimiting cocone

$$(\phi_i^\flat, \text{id}_C) : \square^\flat(\phi_i, \mathbf{f}) \rightarrow \square^\flat(\text{id}_C, \mathbf{f}).$$

In particular, this demonstrates that the map

$$(\phi_0^\flat, \text{id}_C) : \square^\flat(\phi_0, \mathbf{f}) \rightarrow \square^\flat(\text{id}_C, \mathbf{f})$$

is the transfinite composite of the maps

$$(g_i^\flat, \text{id}_C) : \square^\flat(\phi_i, \mathbf{f}) \rightarrow \square^\flat(\phi_{i+1}, \mathbf{f})$$

But by commutativity of the square

$$\begin{array}{ccc} Q_{\phi_0, \mathbf{f}} & \xrightarrow{\phi_0^\flat} & Q_{\text{id}_C, \mathbf{f}} \\ \downarrow \square^\flat(\phi_0, \mathbf{f}) & & \downarrow \square^\flat(\text{id}_C, \mathbf{f}) \\ \square(C, \mathbf{Y}) & \xrightarrow{\text{id}} & \square(C, \mathbf{Y}) \end{array},$$

we see that

$$\square^\flat(\phi_0, \mathbf{f}) = \square^\flat(\text{id}_C, \mathbf{f}) \circ \phi_0^\flat,$$

but from Lemma 1.4.8, we see $\square^\flat(\text{id}_C, \mathbf{f}) = \text{id}$, so it immediately follows that $\square^\flat(\phi_0, \mathbf{f}) = \phi_0^\flat$. This exhibits $\square^\flat(\phi_0, \mathbf{f})$ as a transfinite composite of the g_i^\flat . Then it suffices to show that the g_i^\flat are pushouts of maps belonging to

$$\square^\flat(\mathcal{J}_1, \dots, \mathcal{J}_n).$$

Since each g_i was a pushout of a morphism $h_i \in \mathcal{J}_1$, and since we have a commutative square $(g_i, \text{id}) : \phi_i \rightarrow \phi_{i+1}$, we are exactly in the situation of Lemma 1.4.13, which implies that each g_i^\flat is a pushout of $\square^\flat(h_i, \mathbf{h})$, which proves the proposition. \square

Corollary 1.4.15. *There is an inclusion*

$$\square^\perp(\text{Cell}(\mathcal{J}_1), \dots, \text{Cell}(\mathcal{J}_n)) \subseteq \text{Cell}(\square^\perp(\mathcal{J}_1, \dots, \mathcal{J}_n)).$$

Proof. Apply the previous lemma n times, using the fact that Cell is idempotent, since it is a closure operator. \square

Definition 1.4.16. Let Rex_c denote the symmetric sub-multi-category of \mathbf{Cat} whose objects are the categories admitting all finite colimits and whose k -morphisms are the k -fold functors that preserve finite connected colimits.

Observation 1.4.17. The corner tensor construction, sending a multimorphism

$$F : \mathcal{A}_1 \times \dots \times \mathcal{A}_n \rightarrow \mathcal{D}$$

to its corner tensor

$$F^\perp : \mathcal{A}_1^{[1]} \times \dots \times \mathcal{A}_n^{[1]} \rightarrow \mathcal{D}^{[1]}$$

is a morphism of multicategories from Rex_c to itself. In particular, it is functorial with respect to the composition in Rex_c .

1.5 The regular Reedy structure of $\Theta[\mathcal{C}]$

In this section, we demonstrate that the class of monomorphisms is exactly the class of relative cell complexes for \mathcal{M} , which we will show coincides with the set of boundary maps for the regular Reedy category $\Theta[\mathcal{C}]$.

Proposition 1.5.1. *The category $\Theta[\mathcal{C}]$ is a regular skeletal Reedy category whenever \mathcal{C} is a regular Cartesian Reedy category. The dimension function of this regular Reedy category is given by*

$$\dim[n](c_1, \dots, c_n) \stackrel{\text{def}}{=} n + \dim_{\mathcal{C}} c_1 + \dots + \dim_{\mathcal{C}} c_n.$$

Proof. It follows by [BR11, Proposition 4.4] that $\Theta[\mathcal{C}]$ is normal skeletal Reedy with the desired dimension function (see [Cis06, 8.1]). To prove that $\Theta[\mathcal{C}]$ is regular, it suffices to show that any nondegenerate section

$$(\alpha, \mathbf{f}) : [n](c_1, \dots, c_n) \rightarrow [m](d_1, \dots, d_m)$$

is monic. If $n > m$, then the map α is a simplicial degeneracy and therefore the map factors through $[m](c_{i_1}, \dots, c_{i_m})$, which means that (α, \mathbf{f}) cannot be nondegenerate.

Then we have two cases, when $n = m$ or $n < m$, and therefore either α is the identity or a simplicial face map. If $n = m$, then clearly α, \mathbf{f} is monic, since each nondegenerate section $c_i \rightarrow d_i$ must be monic by the regularity of \mathcal{C} .

If α is a composite of outer face maps, then (α, \mathbf{f}) lands in $[n](d_1, \dots, d_n)$ or $[n](d_{m-n}, \dots, d_m)$, and then from the previous case together with the fact that those inclusions are monic. Otherwise, by induction on dimension, we can assume that α is the inclusion of a codimension 1 inner face map obtained by removing the k^{th} vertex. Then (α, \mathbf{f}) factors through the inclusion

$$V[n](d_1, \dots, d_{k-1} \times d_k, \dots, d_m) \subseteq [m](d_1, \dots, d_m)$$

by a map id, \mathbf{g} where each g_i is a nondegenerate section. Then each g_i must be monic since for $i \neq k$, g_i is a nondegenerate section of a representable, which is monic, and for $i = k$, g_i is a nondegenerate section $c \rightarrow d_{k-1} \times d_k$, which is monic by the fact that \mathcal{C} is regular Cartesian. \square

Corollary 1.5.2. *The boundary $\partial[n](c_1, \dots, c_n)$ can be computed using the corner-intertwiner (see 1.6)*

$$Q = \text{dom } \square_n^J(\delta^n, \delta^{c_1}, \dots, \delta^{c_n}).$$

Proof. It is clear that $Q \subseteq \partial[n](c_1, \dots, c_n)$. The calculations in the proof of the proposition prove the converse. \square

Corollary 1.5.3. *We define the set*

$$\mathcal{M} = \{\partial c \rightarrow c \mid c \in \text{Ob}(\Theta[\mathcal{C}])\}.$$

Then the class $\text{Cell}(\mathcal{M})$ is exactly the the class of monomorphisms of $\widehat{\Theta[\mathcal{C}]}$.

Proof. This is an immediate consequence of [Cis06, Proposition 8.1.37] or [BR11, 4.4]. \square

Lemma 1.5.4. *The category $\Theta[\mathcal{C}]$ is regular Cartesian Reedy when \mathcal{C} is.*

Proof. \square

1.6 The anodyne theorem for horizontal inner anodynes

In this section, following [Our10, 3.4.4], we will demonstrate that the horizontal inner anodynes are closed under corner products with monomorphisms. As a corollary of the analysis in this section, we will demonstrate that $\Theta[\mathcal{C}]$ is regular Cartesian Reedy. We make no claim to originality.

Lemma 1.6.1. *Given a simplicial set S , the category of labelings of S , that is, the fibre $\left(\widehat{\Delta} \int \widehat{\mathcal{C}}\right)_S$ is closed under finite Cartesian products.*

Proof. The product of two S -labeled simplicial sets $(S, \Omega), (S, \Omega')$ can be given by the formula

$$S \xrightarrow{\Delta} S \times S \xrightarrow{\Omega \times \Omega'} \widehat{\mathcal{C}}_{\bullet} \times \widehat{\mathcal{C}}_{\bullet} \xrightarrow{\times} \widehat{\mathcal{C}}_{\bullet}.$$

We leave the verification to the reader. \square

Definition 1.6.2. Given a simplicial set S define the functor

$$H_S : \left(\widehat{\Delta} \downarrow S\right) \times \left(\widehat{\Delta} \int \widehat{\mathcal{C}}\right)_S \rightarrow \widehat{\Delta} \int \widehat{\mathcal{C}}$$

by the rule

$$\left(S' \xrightarrow{f} S, (S, \Omega)\right) \mapsto (S', \Omega \circ f),$$

and we define the *relative intertwiner over S*

$$\square_S \stackrel{\text{def}}{=} \square \circ H_S.$$

Notice that when $S = \Delta^n$, the fibre decomposes as

$$\left(\widehat{\Delta} \int \widehat{\mathcal{C}}\right)_n \cong \widehat{\mathcal{C}}^n.$$

So we can write

$$\square_n : \left(\widehat{\Delta} \downarrow \Delta^n\right) \times \widehat{\mathcal{C}}^n \rightarrow \widehat{\Theta[\mathcal{C}]}. \quad \square$$

Observation 1.6.3. Given a labeled simplicial set (S, Ω) , a map of simplicial sets $Y \rightarrow S$, and an object $[t] = [n](c_1, \dots, c_n)$ of $\Theta[\mathcal{C}]$, we can compute $\square_S(f, \Omega)_t$ as follows: For any n -simplex $s \in S_n$, let $(W_{s,i})_{i=1}^n$ be the family of \mathcal{C} -sets obtained by evaluation of Ω on s . A map $[t] \rightarrow Y \square \Omega \circ f$ is by definition a map $[t] \rightarrow (Y, \Omega \circ f)$. Such a map is determined by giving an n -simplex $y \in Y_n$ together with a family of maps

$$(c_i \xrightarrow{\zeta_i} W_{fy,i})_{i=1}^n.$$

Then we can compute

$$\square_S(f, \Omega)_t \cong \coprod_{y \in Y_n} \prod_{i=1}^n W_{fy,i,c_i}.$$

Definition 1.6.4. Given a finite family of simplicial sets $\mathbf{S} = (S_i)_{i=1}^n$, we define a functor:

$$H_{\mathbf{S}} : \left(\widehat{\Delta} \downarrow \prod_{i=1}^n S_i \right) \times \prod_{i=1}^n \left(\widehat{\Delta} \int \widehat{\mathcal{C}} \right)_{S_i} \rightarrow \widehat{\Delta} \int \widehat{\mathcal{C}}$$

by the rule:

$$\left(S \xrightarrow{\prod_{i=1}^n f_i} \prod_{i=1}^n S_i, (\Omega_i)_{i=1}^n \right) \mapsto \left(S, \prod_{i=1}^n (\Omega_i \circ f_i) \right).$$

As in the previous definition, we define the *relative multi-intertwiner* by the formula

$$\square_{\mathbf{S}} \stackrel{\text{def}}{=} \square \circ H_{\mathbf{S}}.$$

Remark 1.6.5. Notice that given a finite family of labeled simplicial sets $(\mathbf{S}, \mathbf{\Omega}) = (S_i, \Omega_i)_{i=1}^n$ and a family $\mathbf{f} = \left(S \xrightarrow{f_i} S_i \right)_{i=1}^n$,

$$H_{\mathbf{S}}(\mathbf{f}, \mathbf{\Omega}) \cong H_{S_1}(f_1, \Omega_1) \times^S \cdots \times^S H_{S_n}(f_n, \Omega_n),$$

where \times^S denotes the product in $(\widehat{\Delta} \int \widehat{\mathcal{C}})_S$.

Observation 1.6.6. Let

$$(\mathbf{S}, \mathbf{\Omega}) = (\Delta^{m_i}, \Omega_i)_{i=1}^n$$

be a family of labeled simplices, and let

$$\mathbf{f} = \left(\Delta^r \xrightarrow{f_i} \Delta^{m_i} \right)_{i=1}^n,$$

be a family of maps defining an r -simplex of the product. We may identify the Ω_i with families of \mathcal{C} -presheaves $(X_{i,\ell})_{\ell=1}^{m_i}$, so we compute $H_{\Delta^{m_i}}(f_i, \Omega_i)$ as

$$\left(\prod_{k=f_i(j-1)+1}^{f_i(j)} X_{i,k} \right)_{j=1}^r,$$

and therefore, we can compute $H_{\mathbf{S}}(\mathbf{f}, \mathbf{\Omega})$ as

$$\left(\prod_{i=1}^n \left(\prod_{k=f_i(j-1)+1}^{f_i(j)} X_{i,k} \right) \right)_{j=1}^r.$$

Lemma 1.6.7. *The relative intertwiner \square_S preserves colimits in the first variable.*

Proof. Since colimits are computed objectwise in presheaves, it suffices to show that the functor $\square_S(\bullet, \Omega)_t$ preserves colimits for all $[t] = [n](c_1, \dots, c_n) \in \Theta[\mathcal{C}]$ and all labels Ω of S . Therefore, it suffices by Observation 1.6.3 to show this in the case where \mathcal{C} is the terminal category, since we may fix the family of objects (c_1, \dots, c_n) . For each $s \in S_n$ let $(W_{s,i})_{i=1}^n$ be the evaluation of Ω on s . Then given $f : Y \rightarrow S$, we have a Cartesian square

$$\begin{array}{ccc} \coprod_{y \in Y_n} \prod_{i=1}^n W_{fy,i} & \longrightarrow & Y_n \\ \tau^* f \downarrow & & \downarrow f \\ \coprod_{s \in S_n} \prod_{i=1}^n W_{s,i} & \xrightarrow{\tau} & S_n \end{array},$$

exhibiting $\coprod_{y \in Y_n} \prod_{i=1}^n W_{fy,i}$ as the pullback of f along τ , but by the universality of colimits in the category of sets, we are done. \square

Note 1.6.8. This is Oury's proof, but this statement can also be seen to follow immediately from 1.2.10.

Lemma 1.6.9. *The relative intertwiner \square_n preserves connected colimits in each variable.*

Proof. We saw from the previous lemma that it preserves colimits in the first variable, so representing Y as the colimit of its simplices, we immediately reduce to the case where Y is a simplex. But we know in this case that any map $[p] \rightarrow [n]$ factors as a degeneracy followed by a face map. In the case that f is a face map, we can compute the pullback of $V[n](X_1, \dots, X_n)$ along f to be

$$V[p] \left(\prod_{i=f(0)+1}^{f(1)} X_i, \dots, \prod_{i=f(p-1)}^{f(p)} X_i \right).$$

By universality of colimits in $\widehat{\mathcal{C}}$, we see that it suffices to show that

$$V[p](\bullet, \dots, \bullet) = \square_p(\text{id}_{\Delta^p}, \bullet, \dots, \bullet)$$

preserves connected colimits in each variable. In the case where f is a degeneracy map, we can compute the pullback over $[p]$ to be

$$V[p](*, \dots, X_1, \dots, *, \dots, X_n, \dots *),$$

where we fill in the terminal object of $\widehat{\mathcal{C}}$ in each argument i where $f(i-1) = f(i)$. In this case again, it again suffices to show that $V[p]$ preserves connected colimits in each variable, but this is precisely the content of [Rez10, Proposition 4.5], where the proof proceeds by first showing that if we set $X_k = \emptyset$, then

$$V[p+1+q](A_1, \dots, A_p, \emptyset, B_1, B_q) \cong V[p](A_1, \dots, A_p) \coprod V[q](B_1, \dots, B_q),$$

and then exhibiting the obvious parametric right adjoint

$$\left(\left(V[p](A_1, \dots, A_p) \coprod V[q](B_1, \dots, B_q) \right) \downarrow \widehat{\Theta[\mathcal{C}]} \right) \rightarrow \widehat{\mathcal{C}}.$$

□

Note 1.6.10. This proof is substantially easier than Oury's proof, which relies on a long direct computation.

Definition 1.6.11. Since the categories $\widehat{\mathcal{C}}$, $\widehat{\Theta[\mathcal{C}]}$, $\widehat{\mathcal{C}}^n$, and $(\widehat{\Delta} \downarrow \Delta^n)$ are all cocomplete (since they are all presheaf categories), and since the intertwiner preserves connected colimits argument-by-argument, we can use 1.4 to define the functor

$$\square_n^\sqcup : (\widehat{\Delta} \downarrow \Delta^n)^{[1]} \times \underbrace{\widehat{\mathcal{C}}^{[1]} \times \dots \times \widehat{\mathcal{C}}^{[1]}}_{n \text{ times}} \rightarrow \widehat{\Theta[\mathcal{C}]}^{[1]},$$

called the *corner intertwiner*.

More generally, for any finite family of simplices $(\Delta^{m_i})_{i=1}^n$, we can do the same trick and define the *corner-multi-intertwiner*

$$\square_{m_1, \dots, m_n}^\sqcup : (\widehat{\Delta} \downarrow \Delta^{m_1})^{[1]} \times \dots \times (\widehat{\Delta} \downarrow \Delta^{m_n})^{[1]} \times (\widehat{\mathcal{C}}^{[1]})^{m_1} \times \dots \times (\widehat{\mathcal{C}}^{[1]})^{m_n} \rightarrow \widehat{\Theta[\mathcal{C}]}^{[1]}.$$

Following [Our10, 3.85], we begin with the following observations:

Observation 1.6.12. We saw from the definition of \square and the definition of products in $\widehat{\Delta} f \widehat{\mathcal{C}}$ that the diagram

$$\begin{array}{ccc} \widehat{\Delta} f \widehat{\mathcal{C}} \times \widehat{\Delta} f \widehat{\mathcal{C}} & \xrightarrow{\square \times \square} & \widehat{\Theta[\mathcal{C}]} \times \widehat{\Theta[\mathcal{C}]} \\ \downarrow \times & & \downarrow \times \\ \widehat{\Delta} f \widehat{\mathcal{C}} & \xrightarrow{\square} & \widehat{\Theta[\mathcal{C}]} \end{array}$$

commutes. We also computed that the diagram

$$\begin{array}{ccc} (\widehat{\Delta} \downarrow \Delta^n) \times \widehat{\mathcal{C}}^n \times (\widehat{\Delta} \downarrow \Delta^m) \times \widehat{\mathcal{C}}^m & \xrightarrow{H_n \times H_m} & \widehat{\Delta} f \widehat{\mathcal{C}} \times \widehat{\Delta} f \widehat{\mathcal{C}} \\ \downarrow \varsigma & & \downarrow \times \\ (\widehat{\Delta} \downarrow \Delta^n) \times (\widehat{\Delta} \downarrow \Delta^m) \times \widehat{\mathcal{C}}^n \times \widehat{\mathcal{C}}^m & & \\ \downarrow P \times \text{id} \times \text{id} & & \\ (\widehat{\Delta} \downarrow \Delta^n \times \Delta^m) \times \widehat{\mathcal{C}}^n \times \widehat{\mathcal{C}}^m & \xrightarrow{H_{n,m}} & \widehat{\Delta} f \widehat{\mathcal{C}} \end{array}$$

commutes as well where ς permutes the factors and P is the functor sending a pair of labeled simplicial sets $f : S \rightarrow \Delta^n, \Omega$ and $g : S' \rightarrow \Delta^m, \Omega'$ over Δ^n and Δ^m respectively to the labeled simplicial set

$$f \times g : S \times S' \rightarrow \Delta^n \times \Delta^m, \Omega \times \Omega'$$

over the product. Taking these two diagrams together, we see that the diagram

$$\begin{array}{ccc} (\widehat{\Delta} \downarrow \Delta^n) \times \widehat{\mathcal{C}}^n \times (\widehat{\Delta} \downarrow \Delta^m) \times \widehat{\mathcal{C}}^m & \xrightarrow{\square_n \times \square_m} & \widehat{\Theta}[\widehat{\mathcal{C}}] \times \widehat{\Theta}[\widehat{\mathcal{C}}] \\ \downarrow \varsigma & & \downarrow \times \\ (\widehat{\Delta} \downarrow \Delta^n) \times (\widehat{\Delta} \downarrow \Delta^m) \times \widehat{\mathcal{C}}^n \times \widehat{\mathcal{C}}^m & & \\ \downarrow P \times \text{id} \times \text{id} & & \downarrow \\ (\widehat{\Delta} \downarrow \Delta^n \times \Delta^m) \times \widehat{\mathcal{C}}^n \times \widehat{\mathcal{C}}^m & \xrightarrow{\square_{n,m}} & \widehat{\Theta}[\widehat{\mathcal{C}}] \end{array}$$

also commutes.

Then by 1.6.9, every functor appearing in this diagram preserves connected colimits in each argument, the intertwiners by the lemma, and the functors P and \times , since they are products in presheaf categories and therefore preserve colimits in both arguments. Then by the functoriality of the corner tensor functor 1.4.17, we obtain a commutative diagram

$$\begin{array}{ccc} (\widehat{\Delta} \downarrow \Delta^n)^{[1]} \times (\widehat{\mathcal{C}}^{[1]})^n \times (\widehat{\Delta} \downarrow \Delta^m)^{[1]} \times (\widehat{\mathcal{C}}^{[1]})^m & \xrightarrow{\square_n^J \times \square_m^J} & \widehat{\Theta}[\widehat{\mathcal{C}}]^{[1]} \times \widehat{\Theta}[\widehat{\mathcal{C}}]^{[1]} \\ \downarrow \varsigma & & \downarrow \times^J \\ (\widehat{\Delta} \downarrow \Delta^n)^{[1]} \times (\widehat{\Delta} \downarrow \Delta^m)^{[1]} \times (\widehat{\mathcal{C}}^{[1]})^n \times (\widehat{\mathcal{C}}^{[1]})^m & & \\ \downarrow P^J \times \text{id} \times \text{id} & & \downarrow \\ (\widehat{\Delta} \downarrow \Delta^n \times \Delta^m)^{[1]} \times (\widehat{\mathcal{C}}^{[1]})^n \times (\widehat{\mathcal{C}}^{[1]})^m & \xrightarrow{\square_{n,m}^J} & \widehat{\Theta}[\widehat{\mathcal{C}}]^{[1]} \end{array}$$

also commutes, where $P^J = \times^J$.

Observation 1.6.13. Consider the corner product of a simplicial inner horn inclusion with a simplicial boundary inclusion

$$\lambda_j^n \times^J \delta^m : \Lambda_j^n \times \Delta^m \cup \Delta^n \times \partial \Delta^m \hookrightarrow \Delta^n \times \Delta^m.$$

Then it is a standard fact of quasicategory theory that we can factor this map as a sequence

$$\Lambda_j^n \times \Delta^m \cup \Delta^n \times \partial \Delta^m = X_0 \subseteq X_1 \subseteq \cdots \rightarrow X_{k-1} \subseteq X_k = \Delta^n \times \Delta^m$$

where each inclusion $X_{i-1} \hookrightarrow X_i$ is the pushout of an inner horn inclusion $\Lambda_{\ell_i}^{r_i} \rightarrow \Delta^{r_i}$ along an inclusion $\Lambda_{\ell_i}^{r_i} \hookrightarrow X_{i-1}$. By the construction of the sequence, each $[r_i] \rightarrow X_i \rightarrow \Delta^n \times \Delta^m$ is nondegenerate and does not factor through X_{i-1} , so in particular, it does not factor through X_0 , and therefore the maps $\alpha_i : \Delta^{r_i} \rightarrow \Delta^n$ and $\beta_i : [r_i] \rightarrow \Delta^m$ do not factor through Λ_j^n or $\partial\Delta^m$. In particular, the image of α_i is either $\partial_j\Delta^n$ or all of Δ^n , and the image of β_i must be all of Δ^m , so all three maps α_i , β_i , and $\alpha_i \times \beta_i$ send the initial and terminal vertices of Δ^{r_i} to the initial and terminal vertices of Δ^n , Δ^m , and $\Delta^n \times \Delta^m$ respectively.

Observation 1.6.14. Let $(\alpha \times \beta) : \Delta^r \rightarrow \Delta^n \times \Delta^m$ be an injective map preserving initial and terminal elements. Let $\mathbf{A} = (A_i)_{i=1}^n$ and $\mathbf{B} = (B_i)_{i=1}^m$ be objects of $\widehat{\mathcal{C}}^n$ and $\widehat{\mathcal{C}}^m$ respectively. Let

$$K_{\alpha,\beta} : \widehat{\mathcal{C}}^n \times \widehat{\mathcal{C}}^m \rightarrow \widehat{\mathcal{C}}^r$$

be the functor defined by the rule

$$(\mathbf{U}, \mathbf{V}) \mapsto \alpha^* \mathbf{U} \times \beta^* \mathbf{V},$$

taking the product of the pullbacks to the fibre over Δ^r . Then we have a diagram:

$$\begin{array}{ccc} (\widehat{\Delta} \downarrow \Delta^r) \times \widehat{\mathcal{C}}^n \times \widehat{\mathcal{C}}^m & \xrightarrow{\text{id} \times K_{\alpha,\beta}} & (\widehat{\Delta} \downarrow \Delta^r) \times \widehat{\mathcal{C}}^r \\ \downarrow ((\alpha,\beta)) \circ (-) \times \text{id} \times \text{id} & & \downarrow H_r \\ (\widehat{\Delta} \downarrow \Delta^n \times \Delta^m) \times \widehat{\mathcal{C}}^n \times \widehat{\mathcal{C}}^m & \xrightarrow{H_{n,m}} & \widehat{\Delta} \int \widehat{\mathcal{C}} \end{array}$$

To show that the diagram commutes, let $p : X \rightarrow \Delta^r$ be a map. Then evaluating $H_r(p, K_{\alpha,\beta}(\mathbf{U}, \mathbf{V})) = H_r(p, \alpha^* \mathbf{U} \times \beta^* \mathbf{V})$ on a simplex $x : \Delta^s \rightarrow X$ is

$$\begin{aligned} (px)^*(\alpha^* \mathbf{U} \times \beta^* \mathbf{V}) &= (px)^* \alpha^* \mathbf{U} \times (px)^* \beta^* \mathbf{V} \\ &= (\alpha px)^* \mathbf{U} \times (\beta px)^* \mathbf{V} \\ &= H_{n,m}((\alpha px, \beta px), \mathbf{U}, \mathbf{V}) \\ &= H_{n,m} \circ ((\alpha, \beta) \circ \text{id} \times \text{id})(px, \mathbf{U}, \mathbf{V}), \end{aligned}$$

which demonstrates that the diagram commutes. Let $(t_i)_{i=1}^r$ such that $t_i = \alpha(i) - \alpha(i-1) + \beta(i) - \beta(i-1)$. Note that the sum of the t_i is exactly $n + m$, since α and β preserve initial and terminal objects. We define a functor

$$\tau_i : \widehat{\mathcal{C}}^n \times \widehat{\mathcal{C}}^m \rightarrow \widehat{\mathcal{C}}^{t_i}$$

by the rule

$$(\mathbf{A}, \mathbf{B}) \mapsto (A_{\alpha(i-1)+1}, \dots, A_{\alpha(i)}, B_{\beta(i-1)+1}, \dots, B_{\beta(i)}).$$

Then define

$$\tau : \widehat{\mathcal{C}}^n \times \widehat{\mathcal{C}}^m \rightarrow \prod_{i=1}^r \widehat{\mathcal{C}}^{t_i}.$$

It is clear that τ is a permutation of variables and therefore an isomorphism. Then let

$$P_i : \widehat{\mathcal{C}}^{t_i} \rightarrow \widehat{\mathcal{C}}$$

be the functor defined by the rule

$$(X_1, \dots, X_{t_i}) \mapsto X_1 \times \dots \times X_{t_i}$$

Then the P_i assemble to a map (P_1, \dots, P_r) such that

$$(P_1, \dots, P_r) \circ \tau = K_{\alpha, \beta}.$$

Then the diagram

$$\begin{array}{ccc} (\widehat{\Delta} \downarrow \Delta^r) \times \widehat{\mathcal{C}}^n \times \widehat{\mathcal{C}}^m & \xrightarrow{\text{id} \times \tau} & (\widehat{\Delta} \downarrow \Delta^r) \times \prod_{i=1}^r \widehat{\mathcal{C}}^{t_i} \xrightarrow{\text{id} \times (P_i)_{i=1}^r} (\widehat{\Delta} \downarrow \Delta^r) \times \widehat{\mathcal{C}}^r \\ \downarrow ((\alpha, \beta)) \circ (-) \times \text{id} \times \text{id} & & \downarrow H_r \\ (\widehat{\Delta} \downarrow \Delta^n \times \Delta^m) \times \widehat{\mathcal{C}}^n \times \widehat{\mathcal{C}}^m & \xrightarrow{H_{n,m}} & \widehat{\Delta} \int \widehat{\mathcal{C}} \end{array}$$

commutes, and therefore, composing the bottom horizontal and right vertical maps with \square , we have another commutative diagram

$$\begin{array}{ccc} (\widehat{\Delta} \downarrow \Delta^r) \times \widehat{\mathcal{C}}^n \times \widehat{\mathcal{C}}^m & \xrightarrow{\text{id} \times \tau} & (\widehat{\Delta} \downarrow \Delta^r) \times \prod_{i=1}^r \widehat{\mathcal{C}}^{t_i} \xrightarrow{\text{id} \times (P_i)_{i=1}^r} (\widehat{\Delta} \downarrow \Delta^r) \times \widehat{\mathcal{C}}^r \\ \downarrow ((\alpha, \beta)) \circ (-) \times \text{id} \times \text{id} & & \downarrow \square_r \\ (\widehat{\Delta} \downarrow \Delta^n \times \Delta^m) \times \widehat{\mathcal{C}}^n \times \widehat{\mathcal{C}}^m & \xrightarrow{\square_{n,m}} & \widehat{\Theta}[\widehat{\mathcal{C}}] \end{array}$$

The bottom horizontal and right vertical maps preserve connected colimits, as we have seen. The left vertical map preserves connected colimits because colimits are computed in the domain for comma categories. The map $\prod_{i=1}^r P_i$ preserves colimits in each argument because colimits are universal in toposes. Then applying the corner tensor functor, we have the commutative diagram

$$\begin{array}{ccc} (\widehat{\Delta} \downarrow \Delta^r)^{[1]} \times (\widehat{\mathcal{C}}^{[1]})^n \times (\widehat{\mathcal{C}}^{[1]})^m & \xrightarrow{\text{id} \times \tau} & (\widehat{\Delta} \downarrow \Delta^r)^{[1]} \times \prod_{i=1}^r (\widehat{\mathcal{C}}^{[1]})^{t_i} \xrightarrow{\text{id} \times (P_i^J)} (\widehat{\Delta} \downarrow \Delta^r) \times (\widehat{\mathcal{C}}^{[1]})^r \\ \downarrow ((\alpha, \beta)) \circ (-)^J \times \text{id} \times \text{id} & & \downarrow \square_r^J \\ (\widehat{\Delta} \downarrow \Delta^n \times \Delta^m)^{[1]} \times (\widehat{\mathcal{C}}^{[1]})^{n+m} & \xrightarrow{\square_{n,m}^J} & \widehat{\Theta}[\widehat{\mathcal{C}}]^{[1]} \end{array}$$

Finally, we reach our destination. As we mentioned before, this entire section is due to David Oury [Our10], and the following theorem is the crescendo:

Theorem 1.6.15 (Anodyne Theorem [Our10, 3.88]). *The class of horizontal anodynes is closed under corner products with monomorphisms. In particular, if we let*

$$\mathcal{J} = \{\square_n^J(\lambda_k^n, \delta^{c_1}, \dots, \delta^{c_n}) \mid \text{for } n \geq 2, 0 < k < n\}.$$

Then we have

$$\mathcal{M} \times^\perp \mathcal{J} \subseteq \text{Cell}(\mathcal{J}).$$

Proof. Let $f_0 : \partial\Delta^n \hookrightarrow \Delta^n$, and let $\mathbf{f} = (f_i)_{i=1}^n$ be a family of boundary inclusions in $\widehat{\mathcal{C}}$. Let $g_0 : \Lambda_k^m \hookrightarrow \Delta^n$ be an inner horn inclusion, and let $\mathbf{g} = (g_i)_{i=1}^n$ be a family of boundary inclusions in $\widehat{\mathcal{C}}$. By 1.6.12, we have

$$\square_n^\perp(f_0, \dots, f_n) \times^\perp \square_m^\perp(g_0, \dots, g_m) \cong \square_{n,m}^\perp(f_0 \times^\perp g_0, f_1, \dots, f_n, g_1, \dots, g_n).$$

By 1.6.13, we know that $f_0 \times^\perp g_0$ can be factored as a finite sequence of pushouts of inner horn inclusions. By 1.4.11 and 1.4.14, it follows that

$$\square_{n,m}^\perp(f_0 \times^\perp g_0, f_1, \dots, f_n, g_1, \dots, g_n)$$

is a finite composites of pushouts of maps

$$\square_{n,m}^\perp(h_i, f_1, \dots, f_n, g_1, \dots, g_n)$$

where $(h_i : \Lambda_{\ell_i}^{r_i} \rightarrow \Delta^{r_i})_{i=1}^k$ are inner horn inclusions where h_i is a morphism over $\Delta^n \times \Delta^m$. Let h'_i denote this morphism as map in simplicial sets over Δ^{r_i} , and let $(\alpha_i, \beta_i) : \Delta^r \rightarrow \Delta^m \times \Delta^n$ be initial and terminal vertex preserving as in 1.6.13. Then by 1.6.14, we see that

$$\square_{n,m}^\perp(h_i, \mathbf{f}, \mathbf{g}) \cong \square_{r_i}^\perp(h'_i, (P_1^\perp, \dots, P_{r_i}^\perp) \circ \tau(\mathbf{f}, \mathbf{g})).$$

But the value of the argument in position $1 \leq j \leq r_i$ is

$$P_j^\perp \circ \tau_j(\mathbf{f}, \mathbf{g}) = f_{\alpha_i(j-1)+1} \times^\perp \dots \times^\perp f_{\alpha_i(j)} \times^\perp g_{\beta_i(j-1)+1} \times^\perp \dots \times^\perp g_{\beta_i(j)},$$

which belongs to $\text{Cell}(\mathcal{B})$, where \mathcal{B} is the set of boundary inclusions in $\widehat{\mathcal{C}}$. So we see that

$$\square_{r_i}^\perp(h'_i, (P_1^\perp, \dots, P_{r_i}^\perp) \circ \tau(\mathbf{f}, \mathbf{g}))$$

is in

$$\{\square_{r_i}^\perp(\mathcal{L}, \text{Cell}(\mathcal{B}), \dots, \text{Cell}(\mathcal{B}))\},$$

where \mathcal{L} is the set of inner horn inclusions, which by 1.4.14 is a subset of

$$\text{Cell}(\{\square_{r_i}^\perp(\mathcal{L}, \mathcal{B}, \dots, \mathcal{B})\}),$$

which belongs to $\text{Cell}(\mathcal{J})$, since $\{\square_{r_i}^\perp(\mathcal{L}, \mathcal{B}, \dots, \mathcal{B})\} \subseteq \mathcal{J}$. □

1.7 Comparison with Rezk's complete $\Theta[\mathcal{C}]$ -spaces

Since Rezk's complete Segal model structure on $\text{Psh}_\Delta(\Theta[\mathcal{C}])$ is Cartesian, and since $* \hookrightarrow E$ is one of the generators of the localization (see [Rez10]), and since $\Theta[\mathcal{C}]$ is regular skeletal Reedy, it follows by several results of Cisinski [Cis06, Proposition 8.2.9, Theorem 3.4.36, Proposition 2.3.30] that Rezk's localizer for complete $\Theta[\mathcal{C}]$ -Segal spaces is the simplicial completion of a localizer on $\Theta[\mathcal{C}]$.

Observation 1.7.1. To show that Rezk's localizer is the simplicial completion of the localizer generated by the horizontal inner anodynes, it suffices to show the following two properties hold:

- (i) The maps $\square_n^j(\lambda_i^n, \delta^{c_1}, \dots, \delta^{c_n}) \times \Delta^0$ belong to the localizer for complete $\Theta[\mathcal{C}]$ -Segal spaces for $n \geq 2$ and $0 < i < n$.
- (ii) The Segal maps $\text{Se}[n](c_1, \dots, c_n) : \text{Sp}[n](c_1, \dots, c_n) \hookrightarrow [n](c_1, \dots, c_n)$ are horizontal inner anodyne.

We will make use of the following lemma:

Lemma 1.7.2. *For any inner horn inclusion $\Lambda_k^n \hookrightarrow \Delta^n$, and any presheaves X_1, \dots, X_n on \mathcal{C} , the map $V_{\Lambda_k^n}(X_1, \dots, X_n) \times \Delta^0 \hookrightarrow V[n](X_1, \dots, X_n) \times \Delta^0$ belongs to the localizer for complete Segal- $\Theta[\mathcal{C}]$ -spaces.*

Proof. We will suppress the $\times \Delta^0$ factor denoting discrete simplicial presheaves for the duration of this proof. By [Rez10, 5.2], we know that the maps $\text{Se}[n](X_1, \dots, X_n)$ are already weak equivalences. Then we proceed following the argument of [JT07, Lemma 3.5]. Notice that trivial cofibrations have the right-cancellation property with respect to monomorphisms. Then we show that since the class of trivial cofibrations contains the class of maps $\text{Se}[n](X_1, \dots, X_n)$, it also contains the class of maps

$$V_{\partial_0 \Delta^n \cup \partial_n \Delta^n}(X_1, \dots, X_n) \hookrightarrow V[n](X_1, \dots, X_n)$$

by induction on n . Notice first that the map

$$V_{\Lambda_1^2}(X_1, X_2) \hookrightarrow V[2](X_1, X_2)$$

is automatically a trivial cofibration, since $\Lambda_1^2 = \text{Sp}[2]$. For the case of $n > 2$, notice that by cancellation, it suffices to show that the maps

$$V_{\text{Sp}[n]}(X_1, \dots, X_n) \xrightarrow{i_n} V_{\partial_0 \Delta^n \cup \text{Sp}[n]}(X_1, \dots, X_n) \xrightarrow{j_n} V_{\partial_0 \Delta^n \cup \partial_n \Delta^n}(X_1, \dots, X_n)$$

are trivial cofibrations. Then notice that

$$V_{\text{Sp}[n]}(X_1, \dots, X_n) \xrightarrow{i_n} V_{\partial_0 \Delta^n \cup \text{Sp}[n]}(X_1, \dots, X_n)$$

is a pushout of the map

$$V_{\text{Sp}[n-1]}(X_1, \dots, X_n) \hookrightarrow V_{\partial_0 \Delta^n}(X_1, \dots, X_n),$$

and is therefore a trivial cofibration. Notice that for $d_0 : [n-1] \rightarrow [n]$, $d_0^{-1}(\text{Sp}[n]) = \text{Sp}[n-1]$ and $d_0^{-1}(\partial_n \Delta^n) = \partial_{n-1} \Delta^{n-1}$. Then the square

$$\begin{array}{ccc} V_{\text{Sp}[n-1] \cup \partial_{n-1} \Delta^{n-1}}(X_1, \dots, X_n) & \longrightarrow & V_{\text{Sp}[n-1] \cup \partial_n \Delta^n}(X_1, \dots, X_n) \\ \downarrow k_{n-1} & & \downarrow j_n \\ V_{\partial_0 \Delta^n}(X_1, \dots, X_n) & \xrightarrow{d'_0} & V_{\partial_0 \Delta^n \cup \partial_n \Delta^n}(X_1, \dots, X_n) \end{array}$$

is coCartesian, and k_{n-1} is a trivial cofibration by using the cancellation property with the map j_{n-1} . Therefore, it follows that j_n is also a trivial cofibration.

We now prove the lemma: By the cancellation property, it suffices to show that

$$V_{\text{Sp}[n]}(X_1, \dots, X_n) \hookrightarrow V_{\Lambda_k^n}(X_1, \dots, X_n)$$

is a trivial cofibration for $n \geq 2$ and $0 < i < n$. The case $n = 2$ is obvious, so it suffices to show for the case $n > 2$. Given a set $S \subseteq [n]$, let

$$\Lambda_S^n = \bigcup_{i \notin S} \partial_i \Delta^n.$$

We will show that for $n > 2$ and S a nonempty subset of $[1, \dots, n-1]$, the map

$$V_{\text{Sp}[n]}(X_1, \dots, X_n) \hookrightarrow V_{\Lambda_S^n}(X_1, \dots, X_n)$$

is a trivial cofibration. We argue by induction on n and $s = n - \text{Card}(S)$. If $s = 1$, $\Lambda_S^n = \partial_0 \Delta^n \cup \partial_n \Delta^n$, in which case we are done by the previous argument. If $s > 1$, let $T = S \cup \{b\}$ for some $b \in [1, \dots, n-1] \setminus S$. Then by the inductive hypothesis,

$$V_{\text{Sp}[n]}(X_1, \dots, X_n) \hookrightarrow V_{\Lambda_T^n}(X_1, \dots, X_n)$$

is a trivial cofibration. Then it suffices to show that

$$V_{\Lambda_T^n}(X_1, \dots, X_n) \hookrightarrow V_{\Lambda_S^n}(X_1, \dots, X_n)$$

is a trivial cofibration. We see that the diagram

$$\begin{array}{ccc} V_{\Lambda_T^n \cap \partial_b \Delta^n}(X_1, \dots, X_n) & \longrightarrow & V_{\Lambda_T^n}(X_1, \dots, X_n) \\ \downarrow & & \downarrow \\ V_{\partial_b \Delta^n}(X_1, \dots, X_n) & \longrightarrow & V_{\Lambda_S^n}(X_1, \dots, X_n) \end{array}$$

is a pushout, and therefore, it suffices to show that

$$V_{\Lambda_T^n \cap \partial_b \Delta^n}(X_1, \dots, X_n) \hookrightarrow V_{\partial_b \Delta^n}(X_1, \dots, X_n)$$

is a trivial cofibration. But this is true by the inductive hypothesis on n . Therefore, we are done. \square

Proposition 1.7.3. *The map*

$$\square_n^\perp(\lambda_k^n, \delta^{c_1}, \dots, \delta^{c_n}) \times \Delta^0$$

is a trivial cofibration for the model structure on complete $\Theta[\mathcal{C}]$ -Segal spaces.

Proof. We again suppress the $\times \Delta^0$ factor. Let $Q = \square_n^\perp(\lambda_k^n, \delta^{c_1}, \dots, \delta^{c_n})$. Evaluation of Q on 0 is the source and evaluation on 1 is the target. We must show that the monomorphism $Q : Q(0) \hookrightarrow Q(1)$ is a trivial cofibration. Notice first that

$$\square_n(\Lambda_k^n, c_1, \dots, c_n) \hookrightarrow Q(0) \hookrightarrow Q(1) = [n](c_1, \dots, c_n)$$

is a weak equivalence by the lemma. Then by right-cancellation, it suffices to show that

$$\square_n(\Lambda_k^n, c_1, \dots, c_n) \hookrightarrow Q(0)$$

is a trivial cofibration. Let

$$U(s, t) = \int^{u_0, \dots, u_n} ([1](u_0, s) \times [1](u_1 \wedge \dots \wedge u_n, t)) \cdot \square(\lambda_k^n(u_0), \delta^{c_1}(u_1), \dots, \delta^{c_n}(u_n)),$$

where evaluation on $u_i \in [1]$ denotes taking the source or target. Then we see by coend reduction that

$$\int^{s, t} [1](s \wedge t, x) \times [1](u_0, s) \times [1](u_1 \wedge \dots \wedge u_n, t) = [1](u_0 \wedge u_1 \wedge \dots \wedge u_n, x),$$

so by commutation of coends, we see that

$$Q(x) = \int^{s, t} [1](s \wedge t, x) U(s, t),$$

which proves that

$$Q(0) = U(1, 0) \coprod_{U(0, 0)} U(0, 1),$$

but $U(0, 1) = \square_n(\Lambda_k^n, c_1, \dots, c_n)$, so the map

$$\square_n(\Lambda_k^n, c_1, \dots, c_n) \hookrightarrow Q(0)$$

is a pushout of $U(0, 0) \rightarrow U(1, 0)$, which we will show is a trivial cofibration. Notice that in $U(0, 0)$, everything vanishes when $u_0 = 1$, so we have that

$$U(0, 0) \cong \int^{u_1, \dots, u_n} [1](u_1 \wedge \dots \wedge u_n, 0) \cdot \square(\lambda_k^n(0), \delta^{c_1}(u_1), \dots, \delta^{c_n}(u_n)).$$

Notice also that by cofinality, we have that

$$U(1, 0) = \int^{u_0, \dots, u_n} ([1](u_0, 1) \times [1](u_1 \wedge \dots \wedge u_n, 0)) \cdot \square(\lambda_k^n(u_0), \delta^{c_1}(u_1), \dots, \delta^{c_n}(u_n)),$$

is isomorphic to

$$\int^{u_1, \dots, u_n} [1](u_1 \wedge \dots \wedge u_n, t) \cdot \square(\lambda_k^n(1), \delta^{c_1}(u_1), \dots, \delta^{c_n}(u_n)).$$

Then the map $U(0, 0) \hookrightarrow U(1, 0)$ is induced by the natural maps

$$\square(\lambda_k^n(0), \delta^{c_1}(u_1), \dots, \delta^{c_n}(u_n)) \hookrightarrow \square(\lambda_k^n(1), \delta^{c_1}(u_1), \dots, \delta^{c_n}(u_n)).$$

But $\lambda_k^n(0) \hookrightarrow \lambda_k^n(1)$ is the inner horn inclusion $\Lambda_k^n \hookrightarrow \Delta^n$, and therefore, by the lemma, these are all trivial cofibrations. But $U(0, 0)$ and $U(0, 1)$ are clearly homotopy coends, and therefore, the map $U(0, 0) \hookrightarrow U(0, 1)$ is a monic weak equivalence and therefore a trivial cofibration, which proves the proposition. \square

This proves one direction of the theorem; now we prove the converse.

Proposition 1.7.4. *The maps*

$$\mathrm{Se}[n](c_1, \dots, c_n) : \mathrm{Sp}[n](c_1, \dots, c_n) \hookrightarrow [n](c_1, \dots, c_n)$$

are horizontal inner anodyne.

Proof. Since the map

$$\lambda^n : \Lambda^n = \partial_0 \Delta^n \cup \partial_n \Delta^n \hookrightarrow \Delta^n$$

is inner anodyne, and since the empty maps $e^{c_i} : \emptyset \hookrightarrow c_i$ are monic, it follows by 1.4.14 that the corner-intertwiner

$$\square_n^J(\lambda^n, e^{c_1}, \dots, e^{c_n})$$

is horizontal inner anodyne. However, it is easy to see that this map is exactly

$$\square_n(\Lambda^n, c_1, \dots, c_n) \hookrightarrow [n](c_1, \dots, c_n).$$

Therefore, it suffices to show that the map

$$\square_n(\mathrm{Sp}[n], c_1, \dots, c_n) \hookrightarrow \square_n(\Lambda^n, c_1, \dots, c_n)$$

is a horizontal inner anodyne. We will first show that the map

$$\square_n(\mathrm{Sp}[n] \cup \partial_0 \Delta^n, c_1, \dots, c_n) \hookrightarrow \square_n(\Lambda^n, c_1, \dots, c_n)$$

is horizontal inner anodyne. To see this, we proceed by induction on n . This is immediate for $n \leq 2$. Suppose $n > 2$. Then the map

$$\square_n(\mathrm{Sp}[n] \cup \partial_0 \Delta^n, c_1, \dots, c_n) \hookrightarrow \square_n(\Lambda^n, c_1, \dots, c_n)$$

is horizontal inner anodyne, as it is a pushout of the map

$$\square_n(\mathrm{Sp}[n-1] \cup \partial_0 \Delta^{n-1}, c_1, \dots, c_n) \hookrightarrow \square_n(\partial_n \Delta^n, c_1, \dots, c_n),$$

which is horizontal inner anodyne by the induction hypothesis. Then it suffices to show that

$$\square_n(\mathrm{Sp}[n], c_1, \dots, c_n) \hookrightarrow \square_n(\mathrm{Sp}[n] \cup \partial_0 \Delta^n, c_1, \dots, c_n)$$

is horizontal inner anodyne. Again, we proceed by induction on n and notice that this is clear for $n \leq 2$, but we see immediately that

$$\square_n(\mathrm{Sp}[n], c_1, \dots, c_n) \hookrightarrow \square_n(\mathrm{Sp}[n] \cup \partial_0 \Delta^n, c_1, \dots, c_n)$$

is a pushout of

$$\square_n(\mathrm{Sp}[n-1], c_1, \dots, c_n) \hookrightarrow \square_n(\partial_0 \Delta^n, c_1, \dots, c_n),$$

which is horizontal inner anodyne by the induction hypothesis, which concludes the proof. \square

Corollary 1.7.5. *The left Kan extension of the functor*

$$Y \times E^\bullet : \Theta[\mathcal{C}] \times \Delta \rightarrow \widehat{\Theta[\mathcal{C}]}$$

(where Y is the Yoneda embedding) defined by the rule

$$[n](c_1, \dots, c_n) \times \Delta^m \mapsto [n](c_1, \dots, c_n) \times E^m$$

induces a Quillen equivalence

$$\mathrm{Psh}_\Delta(\Theta[\mathcal{C}])_{\mathrm{CSS}} \xrightleftharpoons[\mathrm{Sing}_E]{\mathrm{Real}_E} \widehat{\Theta[\mathcal{C}]}_{\mathrm{hJoyal}}$$

between the model structure for complete $\Theta[\mathcal{C}]$ -Segal spaces and the horizontal Joyal model structure, and the left Kan extension of the functor

$$d : \Theta[\mathcal{C}] \rightarrow \mathrm{Psh}_\Delta(\Theta[\mathcal{C}])$$

defined by the rule

$$[n](c_1, \dots, c_n) \mapsto [n](c_1, \dots, c_n) \times \Delta^0$$

induces a Quillen equivalence

$$\widehat{\Theta[\mathcal{C}]}_{\mathrm{hJoyal}} \xrightleftharpoons[d^*]{d_!} \mathrm{Psh}_\Delta(\Theta[\mathcal{C}])_{\mathrm{CSS}}.$$

That is to say, the two model categories are Quillen bi-equivalent.

Proof. This is an immediate consequence of [Cis06, Proposition 2.3.27]. \square

1.8 Recognition of horizontal Joyal fibrations

In this section, we will prove the analogue of Joyal's pseudofibration theorem for the horizontal model structure on $\Theta[\mathcal{C}]$. We will need to set up some definitions.

Definition 1.8.1. Recall from Definition 1.6.4, we defined the functor $H_{\mathbf{S}}$ for a finite family of simplicial sets \mathbf{S} . Consider the case of the functor $H_{S,n}$, where the family is made up of two simplicial sets Δ^n and some simplicial set S . Then we define the functor

$$\mathcal{H}_{S,n} : \left(\widehat{\Delta} \downarrow S \times \Delta^n \right) \times \left(\widehat{\Delta} \int \widehat{\mathcal{C}} \right)_n \rightarrow \widehat{\Delta} \int \widehat{\mathcal{C}},$$

to be the restriction of $H_{S,n}$ to the terminal labeling of S , the unique labeling of S that sends all edges of S to the terminal object of $\widehat{\mathcal{C}}$. Composing \square with $\mathcal{H}_{S,n}$ is denoted by $\square_{S,n}$. For the remainder of this section, we also, by abuse of notation, define

$$\square_S : \left(\widehat{\Delta} \downarrow S \right) \rightarrow \widehat{\Theta[\mathcal{C}]}$$

to be the composite of \square with the restriction of H_S to the terminal labeling of S .

Observation 1.8.2. Observe that from 1.6.9 the functor

$$\square_{S,n} : (\widehat{\Delta} \downarrow \Delta^n \times S) \times \widehat{\mathcal{C}}^n \rightarrow \widehat{\Theta}[\mathcal{C}]$$

preserves connected colimits in each argument and therefore can be corner tensored. By the same argument as 1.6.12, given $h : Y \rightarrow S$ a map of simplicial sets, we can compute

$$\begin{aligned} \square_n^\perp(f_0, \dots, f_n) \times^\perp h &= \square_n^\perp(f_0, \dots, f_n) \square_S^\perp(h) \\ &= \square_{S,n}^\perp(h \times^\perp f_0, f_1, \dots, f_n). \end{aligned}$$

In what follows, we will use a very nice observation of Danny Stevenson [Ste]. Consider the case of the simplicial set E^1 and the map $e : \Delta^0 \hookrightarrow E^1$.

Theorem 1.8.3 ([Ste, Lemma 2.19]). *The map of simplicial sets*

$$e \times^\perp \delta^n : E^1 \times \partial \Delta^n \cup \Delta^0 \times \Delta^n \hookrightarrow E \times \Delta^n,$$

is inner anodyne for all $n > 0$.

Observation 1.8.4. It follows from the small object argument that we can factor $e \times^\perp \delta^n$ as a composite

$$E^1 \times \partial \Delta^n \coprod_{\Delta^0 \times \partial \Delta^n} \Delta^0 \times \Delta^n \xrightarrow{\iota} E \xrightarrow{\varepsilon} E^1 \times \Delta^n,$$

where ι is a relative cell complex for the inner horn inclusions and where ε is an inner fibration between quasicategories. Since $e \times^\perp \delta^n$ is inner anodyne, it follows by the lifting property that it is a retract of ι .

Observation 1.8.5. A simplex $(\alpha, \beta) : \Delta^r \rightarrow E^1 \times \Delta^n$ is determined by the destination of its vertices. We label the vertices by (a, i) and (b, i) for $0 \leq i \leq n$, where e is the inclusion of $a \in E^1$. Given a simplex $(\alpha, \beta) : \Delta^r \rightarrow E^1 \times \Delta^n$, and a family of presheaves $\mathbf{U} = (U_i)_{i=1}^n$ on \mathcal{C} .

$$K_\beta : \widehat{\mathcal{C}}^n \rightarrow \widehat{\mathcal{C}}^r$$

to be the functor sending

$$\mathbf{U} \mapsto \beta^* \mathbf{U}.$$

Similar to 1.6.14, we have a diagram

$$\begin{array}{ccc} (\widehat{\Delta} \downarrow \Delta^r) \times \widehat{\mathcal{C}}^n & \xrightarrow{\text{id} \times K_\beta} & (\widehat{\Delta} \downarrow \Delta^r) \times \widehat{\mathcal{C}}^r \\ \downarrow ((\alpha, \beta)) \circ (-) \times \text{id} \times \text{id} & & \downarrow H_r \\ (\widehat{\Delta} \downarrow E^1 \times \Delta^n) \times \widehat{\mathcal{C}}^n & \xrightarrow{H_{E^1, n}} & \widehat{\Delta} \int \widehat{\mathcal{C}} \end{array},$$

and the diagram commutes by a direct computation. By abuse of notation, set $\beta(-1) = 0$ and $\beta(r+1) = n$. Then we define the family

$$(t_i = \beta(i) - \beta(i-1))_{i=0}^{r+1}$$

and let

$$\tau_i : \widehat{\mathcal{C}}^n \rightarrow \widehat{\mathcal{C}}^{t_i}$$

be the map defined by sending

$$\mathbf{U} \mapsto (U_{\beta(i-1)+1}, \dots, U_{\beta(i)}).$$

Then this family of maps defines a map

$$\widehat{\mathcal{C}}^n \rightarrow \prod_{i=1}^r \widehat{\mathcal{C}}^{t_i}$$

which is a permutation and therefore an isomorphism. Then for each $0 < i < r + 1$ let

$$P_i : \widehat{\mathcal{C}}^{t_i} \rightarrow \widehat{\mathcal{C}}$$

be the map defined by the rule

$$(X_1, \dots, X_{t_i}) \mapsto X_1 \times \dots \times X_{t_i},$$

and for $i = 0$ or $i = r + 1$, define

$$P_i : \widehat{\mathcal{C}}^{t_i} \rightarrow *$$

to be the terminal functor. Then these P_i assemble to a map

$$(P_0, \dots, P_{r+1}) : \prod_{i=0}^{r+1} \widehat{\mathcal{C}}^{t_i} \rightarrow \widehat{\mathcal{C}}^r,$$

such that $(P_0, \dots, P_{r+1}) \circ \tau = K_\beta$. Then we have a commutative diagram

$$\begin{array}{ccc} (\widehat{\Delta} \downarrow \Delta^r) \times \widehat{\mathcal{C}}^n & \xrightarrow{\text{id} \times \tau} & (\widehat{\Delta} \downarrow \Delta^r) \times \prod_{i=0}^{r+1} \widehat{\mathcal{C}}^{t_i} \xrightarrow{\text{id} \times (P_i)_{i=0}^{r+1}} (\widehat{\Delta} \downarrow \Delta^r) \times \widehat{\mathcal{C}}^r \\ \downarrow ((\alpha, \beta)) \circ (-) \times \text{id} \times \text{id} & & \downarrow H_r \\ (\widehat{\Delta} \downarrow E^1 \times \Delta^n) \times \widehat{\mathcal{C}}^n & \xrightarrow{H_{E^1, n}} & \widehat{\Delta} \int \widehat{\mathcal{C}} \end{array},$$

and therefore, composing the bottom horizontal and right vertical maps with \square , we have another commutative diagram

$$\begin{array}{ccc} (\widehat{\Delta} \downarrow \Delta^r) \times \widehat{\mathcal{C}}^n & \xrightarrow{\text{id} \times \tau} & (\widehat{\Delta} \downarrow \Delta^r) \times \prod_{i=0}^{r+1} \widehat{\mathcal{C}}^{t_i} \xrightarrow{\text{id} \times (P_i)_{i=0}^{r+1}} (\widehat{\Delta} \downarrow \Delta^r) \times \widehat{\mathcal{C}}^r \\ \downarrow ((\alpha, \beta)) \circ (-) \times \text{id} \times \text{id} & & \downarrow \square_r \\ (\widehat{\Delta} \downarrow E^1 \times \Delta^n) \times \widehat{\mathcal{C}}^n & \xrightarrow{\square_{E^1, n}} & \widehat{\Theta}[\widehat{\mathcal{C}}] \end{array}.$$

We see that each of the arrows in this picture preserves connected colimits argument-by-argument, so applying the corner tensor functor, we obtain a commutative diagram

$$\begin{array}{ccc}
(\widehat{\Delta} \downarrow \Delta^r)^{[1]} \times (\widehat{\mathcal{C}}^{[1]})^n & \xrightarrow{\text{id} \times \tau} & (\widehat{\Delta} \downarrow \Delta^r)^{[1]} \times \prod_{i=0}^{r+1} (\widehat{\mathcal{C}}^{[1]})^{t_i \text{id} \times (P_i^\perp)} \xrightarrow{\quad} (\widehat{\Delta} \downarrow \Delta^r) \times (\widehat{\mathcal{C}}^{[1]})^r \\
\downarrow ((\alpha, \beta)) \circ (-)^\perp \times \text{id} \times \text{id} & & \downarrow \square_r^\perp \\
(\widehat{\Delta} \downarrow E^1 \times \Delta^n)^{[1]} \times (\widehat{\mathcal{C}}^{[1]})^n & \xrightarrow{\square_{E^1, n}^\perp} & \widehat{\Theta}[\mathcal{C}]^{[1]}
\end{array}$$

Theorem 1.8.6. *Set $e : \Delta^0 \rightarrow E^1$. Then for any boundary inclusion*

$$\square_n^\perp(\delta^n, \delta^{c_1}, \dots, \delta^{c_n}),$$

with $n > 0$, the map

$$e \times^\perp \square_n^\perp(\delta^n, \delta^{c_1}, \dots, \delta^{c_n})$$

is a horizontal inner anodyne.

Proof. From 1.8.2, we see that

$$e \times^\perp \square_n^\perp(\delta^n, \delta^{c_1}, \dots, \delta^{c_n}) \cong \square_{E^1, n}^\perp(e \times^\perp \delta^n, \delta^{c_1}, \dots, \delta^{c_n}),$$

and by 1.8.4 we see that $e \times^\perp \delta^n$ can be factored as $\varepsilon \circ \iota$ such that it is a retract of ι , which is a relative cell complex of inner horn inclusions. By a similar argument to 1.4.11 and 1.4.14, it follows that

$$\square_{E^1, n}^\perp(e \times^\perp \delta^n, \delta^{c_1}, \dots, \delta^{c_n})$$

is a retract of the transfinite composite of inner horn inclusions

$$\square_{E^1, n}^\perp(h_i, \delta^{c_1}, \dots, \delta^{c_n}),$$

where the $h_i : \Lambda_{\ell_i}^{r_i} \rightarrow \Delta^{r_i}$ are inner horn inclusions and the h_i are maps in $(\widehat{\Delta} \downarrow E^1 \times \Delta^n)$. Let h'_i denote the same map in the category $(\widehat{\Delta} \downarrow \Delta^{r_i})$, and let (α, β) denote the map $\Delta^{r_i} \rightarrow E^1 \times \Delta^n$, with β surjective as we saw above. Then by 1.8.5,

$$\square_{E^1, n}^\perp(h_i, \delta^{c_1}, \dots, \delta^{c_n}) \cong \square_{r_i}^\perp(h'_i, (P_0^\perp, \dots, P_{r_i+1}^\perp) \circ \tau(\delta^{c_1}, \dots, \delta^{c_n})),$$

but we see that the P_i^\perp are corner products of boundary inclusions as in 1.6.15, and therefore, the maps

$$\square_{r_i}^\perp(h'_i, (P_1^\perp, \dots, P_{r_i}^\perp) \circ \tau(\delta^{c_1}, \dots, \delta^{c_n}))$$

are all horizontal inner anodyne, demonstrating the claim that

$$e \times^\perp \square_n^\perp(\delta^n, \delta^{c_1}, \dots, \delta^{c_n})$$

is inner anodyne. □

Corollary 1.8.7. *The formal \mathcal{C} -quasicategories are the fibrant objects of the horizontal Joyal model structure.*

Proof. Since the formal \mathcal{C} -quasicategories have the right lifting property with respect to all inner anodynes, it suffices to check that they have the right lifting property with respect to $e : \Delta^0 \hookrightarrow E^1$, but such a lift always exists by choosing the obvious degenerate lift. \square

Corollary 1.8.8. *The fibrations between fibrant objects in the horizontal Joyal model structure are horizontal inner fibrations having the right lifting property with respect to the map $e : \Delta^0 \hookrightarrow E^1$.*

Proof. A fibration between fibrant objects must have the right lifting property with respect to all horizontal inner anodynes and all maps of the form $e \times \lrcorner \square_n^\lrcorner(\delta^n, \delta^{c_1}, \dots, \delta^{c_n})$. Since every inner fibration has the right lifting property with respect to all of those maps for $n > 0$, it follows that an inner fibration between fibrant objects need only have the right lifting property with respect to the case where $n = 0$, which is exactly the map e . \square

2 The Coherent Nerve, Horizontal case

2.1 The coherent realization for $\Theta[\mathcal{C}]$

There is an adjoint pair that should be familiar to those who have experience with $(\infty, 1)$ -categories, namely the coherent realization and nerve pair,

$$\widehat{\Theta[*]} \xrightleftharpoons[\mathfrak{N}_\Delta]{\mathfrak{C}_\Delta} \mathbf{Cat}_{\mathbf{Psh}_\Delta(*)},$$

where we have obvious isomorphisms $\Theta[*] \cong \Delta$ and $\mathbf{Psh}_\Delta(*) \cong \widehat{\Delta}$

An important theorem early in *Higher Topos Theory* tells us that this adjunction gives a Quillen equivalence when the lefthand side is equipped with the Joyal model structure and the righthand side is equipped with the Bergner model structure. The goal of this section is to show that for any \mathcal{C} with the aforementioned properties, we can construct an analogous adjunction:

$$\widehat{\Theta[\mathcal{C}]} \xrightleftharpoons[\mathfrak{N}]{\mathfrak{C}} \mathbf{Cat}_{\mathbf{Psh}_\Delta(\mathcal{C})}$$

Definition 2.1.1. The category of \mathcal{C} -precategories is the full subcategory of $\mathbf{Psh}_\Delta(\mathcal{C})$ spanned by the *precategory objects*, namely those simplicial presheaves F on \mathcal{C} such that F_0 is a constant presheaf on \mathcal{C} . We denote this category by $\mathbf{PCat}(\mathcal{C})$.

Definition 2.1.2. The functor $k : \Delta \times \mathcal{C} \rightarrow \Theta[\mathcal{C}]$ defined by the rule

$$([n], c) \mapsto [n](c, \dots, c).$$

Induces a colimit-preserving functor $k^* : \widehat{\Theta[\mathcal{C}]} \rightarrow \mathbf{Psh}_\Delta(\mathcal{C})$ that lands in $\mathbf{PCat}(\mathcal{C})$ called the *associated precategory*.

Definition 2.1.3. The *pointwise realization* functor $\mathfrak{C}_{\Delta, \bullet} : \mathbf{Psh}_\Delta(\mathcal{C}) \rightarrow \mathbf{Cat}_\Delta^{\text{cop}}$ defined by the rule

$$\mathfrak{C}_{\Delta, \bullet}(X)_c \stackrel{\text{def}}{=} \mathfrak{C}_\Delta(X_c)$$

restricts to colimit preserving functor $\mathfrak{C}_{\Delta, \bullet} : \mathbf{PCat}(\mathcal{C}) \rightarrow \mathbf{Cat}_{\mathbf{Psh}_\Delta(\mathcal{C})}$ in the obvious way, which by abuse of notation, we also refer to as the *pointwise realization*.

Definition 2.1.4. The *coherent realization* $\mathfrak{C}_{\Theta[\mathcal{C}]} : \widehat{\Theta[\mathcal{C}]} \rightarrow \mathbf{Cat}_{\mathbf{Psh}_\Delta(\mathcal{C})}$, also denoted by \mathfrak{C} by abuse of notation when \mathcal{C} is fixed is defined as the composite:

$$\widehat{\Theta[\mathcal{C}]} \xrightarrow{k^*} \mathbf{PCat}(\mathcal{C}) \xrightarrow{\mathfrak{C}_{\Delta, \bullet}} \mathbf{Cat}_{\mathbf{Psh}_\Delta(\mathcal{C})}.$$

Note 2.1.5. It is immediate from the cocontinuity of each functor in this composite that the functor $\mathfrak{C}_{\Theta[\mathcal{C}]}$ is cocontinuous and therefore determined on representables. In what follows, we will give an explicit computation of its values on representables and in fact more generally on cellular sets of the form $V[n](A_1, \dots, A_n)$.

Remark 2.1.6. It is easy to see that if we substitute the terminal category for \mathcal{C} , this specializes precisely to the usual coherent realization $\mathfrak{C}_\Delta = \mathfrak{C}_{\Theta[*]}$.

We will extensively abuse notation in what follows by identifying a simplicial set with its associated constant simplicial presheaf on \mathcal{C} and identifying a presheaf on \mathcal{C} with its associated discrete simplicial presheaf.

Definition 2.1.7. We define a construction on objects

$$Q : \Delta \int \widehat{\mathcal{C}} \rightarrow \mathbf{Cat}_{\mathbf{Psh}_\Delta(\mathcal{C})}.$$

Suppose $[n](X_1, \dots, X_n)$ is any object of $\Delta \int \widehat{\mathcal{C}}$. Then we define $Q([n](X_1, \dots, X_n))$ as follows:

- The objects are the vertices $\{0, \dots, n\}$
- The Hom-object

$$\mathrm{Hom}(i, j) = \begin{cases} \emptyset & \text{for } i > j \\ c\Delta^0 & \text{for } i = j \\ X_{i+1} \times \Delta^1 \times \dots \times \Delta^1 \times X_j & \text{for } i < j \end{cases}$$

- The associative composition law, $\mathrm{Hom}(i, j) \times \mathrm{Hom}(j, k) \rightarrow \mathrm{Hom}(i, k)$ which is the inclusion on the bottom face with respect to j :

$$\begin{array}{c} X_{i+1} \times \Delta^1 \times \dots \times \Delta^1 \times X_j \times \{1\} \times X_{j+1} \times \Delta^1 \times \dots \times \Delta^1 \times X_k \\ \downarrow \\ X_{i+1} \times \Delta^1 \times \dots \times \Delta^1 \times X_j \times \Delta^1 \times X_{j+1} \times \Delta^1 \times \dots \times \Delta^1 \times X_k \end{array}$$

Proposition 2.1.8. *The construction Q is functorial.*

Proof. Recall that a map

$$[n](X_1, \dots, X_n) \rightarrow [m](Y_1, \dots, Y_m)$$

in $\Delta \int \widehat{\mathcal{C}}$ is given by a pair (γ, \mathbf{f}) , where $\gamma : [n] \rightarrow [m]$ is a map of simplices together with a family of maps

$$\mathbf{f} = \left(f_i : X_i \rightarrow \prod_{j=\gamma(i-1)}^{\gamma(i)} Y_j \right)_{i=1}^n.$$

If for $0 < i \leq n$, we have $\gamma(i-1) = \gamma(i)$, we can see easily that γ factors through the codegeneracy map

$$[n](X_1, \dots, X_n) \rightarrow [n-1](X_1, \dots, \widehat{X_i}, \dots, X_n).$$

Applying this factorization repeatedly, we factor (γ, \mathbf{f}) as a codegeneracy followed by a map (γ', f') such that γ' is the inclusion of a coface $[n'] \hookrightarrow [m]$.

Since $\Delta \int \widehat{\mathcal{C}}$ is fibred over Δ , we may take the Cartesian lift of γ' , which is the map

$$\overline{\gamma'} = (\gamma', \mathbf{id}) : [n'] \left(\prod_{j=\gamma'(0)}^{\gamma'(1)} Y_j, \dots, \prod_{j=\gamma'(n'-1)}^{\gamma'(n')} Y_j \right) \hookrightarrow [m](Y_1, \dots, Y_m).$$

By Cartesianness, we have a unique factorization of (γ', \mathbf{f}') by this map, yielding a map

$$(\text{id}, \overline{f'}) : [n'](X'_1, \dots, X'_{n'}) \rightarrow [n'] \left(\prod_{j=\gamma'(0)}^{\gamma'(1)} Y_j, \dots, \prod_{j=\gamma'(n'-1)}^{\gamma'(n')} Y_j \right).$$

Then to prove the proposition, we need to show functoriality in three cases:

- If the map (γ, \mathbf{f}) is a codegeneracy of codimension 1, suppose $\gamma = \sigma^i : [n+1] \rightarrow [n]$ for $0 \leq i \leq n$. Then $Q((\sigma^i, \mathbf{id}))_{ab} : \text{Hom}(a, b) \rightarrow \text{Hom}(\sigma^i(a), \sigma^i(b))$ is defined on the homs as follows:

$$Q(\sigma^i)_{ab} = \begin{cases} \text{id} \times \min \circ \tau_{X_i} \times \text{id} & \text{if } a < i \leq b \\ \text{id} & \text{otherwise} \end{cases}$$

where $\min : \Delta^1 \times \Delta^1 \rightarrow \Delta^1$ is induced by the map of posets sending $(x, y) \mapsto \min(x, y)$ and $\tau_{X_i} \rightarrow *$ is the terminal map. Specifically, in the case where $a < i \leq b$, the map is given by the composite:

$$\begin{array}{c} X_{a+1} \times \Delta^1 \times \dots \times \Delta^1 \times X_i \times \Delta^1 \times \dots \times \Delta^1 \times X_b \\ \downarrow \\ X_{a+1} \times \Delta^1 \times \dots \times \Delta^1 \times * \times \Delta^1 \times \dots \times \Delta^1 \times X_b \\ \parallel \\ X_{a+1} \times \Delta^1 \times \dots \times \Delta^1 \times \Delta^1 \times \dots \times \Delta^1 \times X_b \\ \downarrow \\ X_{a+1} \times \Delta^1 \times \dots \times \Delta^1 \times \dots \times \Delta^1 \times X_b \end{array}$$

If $i = 0$ or $i = n$, we consider $\Delta^0 = *$ to be $\{0\}$.

- If the map (γ, \mathbf{f}) is a pure coface of codimension 1, we have two subcases: If it is an outer coface, the map is just the obvious inclusion. If it is an inner coface, it has a term that looks like $X_i \times X_{i+1}$, and this is included in all of the Hom objects as $X_i \times \{0\} \times X_{i+1}$
- If the map (γ, \mathbf{f}) is such that $\gamma = \text{id}$, since each of the Hom objects is given as a product of the X_i with Δ^1 of the same length, just map them by $f_a \times \Delta^1 \times \dots \times \Delta^1 \times f_{b-1}$ using the functoriality of the Cartesian product.

It is an easy exercise to see that this assignment is functorial and completely analogous to the un-enriched case. \square

Note 2.1.9. We specify the following abuse of notation: Given an ordered family of sets S_1, \dots, S_n , we denote by

$$[n](S_1, \dots, S_n)$$

the nerve of the free category on the directed graph specified as follows:

- The set of vertices is $\{0, \dots, n\}$, and
- The set of arrows from the vertex $i - 1$ to i is S_i .

Proposition 2.1.10. *The functor $Q : \Delta \int \widehat{\mathcal{C}} \rightarrow \mathbf{Cat}_{\text{Psh}_\Delta(\mathcal{C})}$ factors as the composite*

$$\Delta \int \widehat{\mathcal{C}} \xrightarrow{V} \Theta[\widehat{\mathcal{C}}] \xrightarrow{\mathfrak{C}} \mathbf{Cat}_{\text{Psh}_\Delta(\mathcal{C})}$$

Proof. Since for any $[m](X_1, \dots, X_m) \in \Delta \int \widehat{\mathcal{C}}$, the enriched categories $Q([m](X_1, \dots, X_m))$ and $\mathfrak{C}(V[m](X_1, \dots, X_m))$ have object sets in natural bijection with the set of vertices of $[m]$ it suffices to produce, for each pair of vertices $i, j \in [m]$ a natural isomorphism

$$\mathfrak{C}(V[m](X_1, \dots, X_m))(i, j) \rightarrow Q([m](X_1, \dots, X_m))(i, j),$$

which amounts to the data of a natural (in c) isomorphism of simplicial sets

$$\mathfrak{C}(V[m](X_1, \dots, X_m))(x, y)_c \rightarrow Q([m](X_1, \dots, X_m))(x, y)_c.$$

First, observe that specifying a simplex

$$\Delta^n \rightarrow Q([m](X_1, \dots, X_m))(i, j)_c$$

is equivalent to specifying a morphism

$$\begin{aligned} (\Delta^n, c) &\rightarrow X_{i+1} \times \Delta^1 \times \dots \times \Delta^1 \times X_j \\ &= (\Delta^1)^{j-i-1} \times \prod_{k=i+1}^j X_k \end{aligned}$$

which is itself equivalent to specifying a simplex

$$\Delta^n \rightarrow (\Delta^1)^{j-i-1} \times \prod_{k=i+1}^j X_k(c),$$

so in particular, we may make the identification

$$Q([m](X_1, \dots, X_m))(i, j)_c \cong \mathfrak{C}_\Delta([m](X_1(c), \dots, X_m(c)))(i, j),$$

and therefore it follows that we have an isomorphism, natural in c :

$$Q([m](X_1, \dots, X_m))_c \cong \mathfrak{C}_\Delta([m](X_1(c), \dots, X_m(c))).$$

It therefore suffices to demonstrate a natural isomorphism of simplicial sets

$$k^*(V[m](X_1, \dots, X_m))_c \cong [m](X_1(c), \dots, X_m(c)),$$

but it can be observed that specifying a simplex

$$\Delta^n \rightarrow k^*(V[m](X_1, \dots, X_m))_c$$

corresponds to specifying a map

$$[n](c, \dots, c) \rightarrow V[m](X_1, \dots, X_m),$$

which in turn is given by the data of a map

$$\gamma : [n] \rightarrow [m]$$

together with maps

$$c \rightarrow \prod_{k=\gamma(0)+1}^{\gamma(n)} X_k,$$

or equivalently, a family

$$(f_{\gamma(0)+1}, \dots, f_{\gamma(n)}) \in \prod_{k=\gamma(0)+1}^{\gamma(n)} X_k(c),$$

which specifies a unique simplex

$$\Delta^n \rightarrow [m](X_1(c), \dots, X_m(c)),$$

which proves the claim. \square

2.2 Necklaces and the coherent realization

Necklaces were introduced by Dugger and Spivak in [DS11a] in order to understand the mapping objects $\mathfrak{C}_\Delta(X)(x, y)$. They prove a useful theorem that allows one to compute the coherent realization up to homotopy as a simplicially-enriched category whose hom-objects are the nerves of ordinary categories. We will demonstrate here how their theory generalizes to our setting. We begin by recalling the definition of a necklace:

Definition 2.2.1. A *necklace* is a bi-pointed simplicial set $(T, (\alpha, \omega))$ of the form

$$\Delta^{m_1} \vee \dots \vee \Delta^{m_n},$$

with specified vertices

$$(\alpha, \omega) : \Delta^0 \amalg \Delta^0 \xrightarrow{\{0\} \amalg \{m_n\}} \Delta^{m_1} \amalg \Delta^{m_n} \xrightarrow{\iota_1 \amalg \iota_n} S.$$

By abuse of notation, we will simply refer to such necklaces by their underlying simplicial sets.

Dugger and Spivak construct a functor

$$\mathfrak{C}_{\Delta}^{\mathcal{N}ec} : \widehat{\Delta} \rightarrow \mathbf{Cat}_{\widehat{\Delta}}$$

whose evaluation on a simplicial set X is given as follows:

- The set of objects of $\mathfrak{C}_{\Delta}^{\mathcal{N}ec}(X)$ is X_0 .
- Given any two vertices $x, x' \in X_0$, the simplicial set of morphisms from x to x' is given by the formula

$$\mathfrak{C}_{\Delta}^{\mathcal{N}ec}(X)(x, x') \stackrel{\text{def}}{=} N(\mathcal{N}ec \downarrow X_{x, x'}),$$

where $(\mathcal{N}ec \downarrow X_{x, x'})$ denotes the slice over the bi-pointed simplicial set $X_{x, x'}$ in the category of bi-pointed simplicial sets.

- The composition map is induced by concatenation of necklaces. That is, given a pair of necklaces $T \rightarrow X_{x, x'}$ and $T' \rightarrow X_{x', x''}$, their composite is given by the necklace

$$T \omega_T \vee_{\alpha_{T'}} T' \rightarrow X_{x, x''}.$$

In order to compare $\mathfrak{C}_{\Delta}^{\mathcal{N}ec}$ and \mathfrak{C}_{Δ} , Dugger and Spivak introduce an auxiliary functor $\mathfrak{C}_{\Delta}^{\mathcal{H}oc}$ that admits specified natural transformations to both. This leads to the main theorem:

Theorem 2.2.2 ([DS11a, Theorem 5.2]). *There is a specified natural zig-zag of weak equivalences of functors valued in simplicially-enriched categories:*

$$\mathfrak{C}_{\Delta}^{\mathcal{N}ec} \leftarrow \mathfrak{C}_{\Delta}^{\mathcal{H}oc} \rightarrow \mathfrak{C}_{\Delta}.$$

The functors $\mathfrak{C}_{\Delta}^{\mathcal{N}ec}$ and $\mathfrak{C}_{\Delta}^{\mathcal{H}oc}$, much like the functor \mathfrak{C}_{Δ} , send simplicial sets to simplicially enriched categories with set of objects equal to the set of 0-simplices. Ergo, they induce useful functor when applied pointwise to precategories.

Definition 2.2.3. As in Definition 2.1.3, the *pointwise necklace realization*

$$\mathfrak{C}_{\Delta, \bullet}^{\mathcal{N}ec} : \mathbf{PCat}(\mathcal{C}) \rightarrow \mathbf{Cat}_{\mathbf{Psh}_{\Delta}(\mathcal{C})}$$

is defined by the rule

$$\mathfrak{C}_{\Delta, \bullet}^{\mathcal{N}ec}(X)_c \stackrel{\text{def}}{=} \mathfrak{C}_{\Delta}^{\mathcal{N}ec}(X_c).$$

The *pointwise homotopy colimit realization* $\mathfrak{C}^{\mathcal{H}oc}$ is defined similarly.

Definition 2.2.4. The *necklace realization* functor $\mathfrak{C}^{\mathcal{N}ec} : \widehat{\Theta}[\mathcal{C}] \rightarrow \mathbf{Cat}_{\mathbf{Psh}_{\Delta}(\mathcal{C})}$ is the composite

$$\widehat{\Theta}[\mathcal{C}] \xrightarrow{k^*} \mathbf{PCat}(\mathcal{C}) \xrightarrow{\mathfrak{C}_{\Delta, \bullet}^{\mathcal{N}ec}} \mathbf{Cat}_{\mathbf{Psh}_{\Delta}(\mathcal{C})}.$$

The *homotopy colimit realization* functor $\mathfrak{C}^{\mathcal{H}oc}$ is defined similarly.

From these definitions and the theorem above, we deduce the following useful corollary.

Corollary 2.2.5. *There is a specified natural zig-zag of weak equivalences of functors valued in $\mathbf{Cat}_{\mathbf{Psh}_{\Delta}(\mathcal{C})}$*

$$\mathfrak{C}^{\mathcal{N}ec} \leftarrow \mathfrak{C}^{\mathcal{H}oc} \rightarrow \mathfrak{C}$$

2.3 Gadgets

To prove the equivalence between quasicategories and simplicially-enriched categories, Dugger and Spivak make use of another intermediate construction called a category of Gadgets. These are subcategories of bi-pointed simplicial sets that generalize necklaces while still retaining many of their useful properties. We begin by recalling the definition of a category of gadgets.

Definition 2.3.1 ([DS11a]). A *Category of Gadgets* is a subcategory \mathcal{G} of the category $\widehat{\Delta}_{*,*}$ satisfying the following properties:

- The category \mathcal{G} contains *Nec*.
- For all $G \in \mathcal{G}$ and all necklaces T , there is an equality

$$\mathcal{G}(T, G) = \widehat{\Delta}(T, G).$$

- For any $G \in \mathcal{G}$, the simplicial set $\mathfrak{C}(G)(\alpha, \omega)$ is contractible.

The category \mathcal{G} is moreover said to be *closed under wedges* if

- For any $G, G' \in \mathcal{G}$, the wedge $G \vee G'$ is as well.

For any bi-pointed simplicial set $X_{x,x'}$ and any category of gadgets \mathcal{G} , we functorially define a simplicial set

$$\mathfrak{C}_{\Delta}^{\mathcal{G}}(X)(x, x') \stackrel{\text{def}}{=} N(\mathcal{G} \downarrow X_{x,x'}).$$

Moreover, if \mathcal{G} is closed under wedges, the collection of simplicial sets

$$(\mathfrak{C}_{\Delta}^{\mathcal{G}}(X)(x, x'))_{x, x' \in X}$$

assembles to a simplicially enriched category $\mathfrak{C}_{\Delta}^{\mathcal{G}}(X)$ with composition induced by concatenation of gadgets.

Dugger and Spivak then prove the following useful proposition:

Proposition 2.3.2 ([DS11a, Proposition 5.5]). *For any category of gadgets \mathcal{G} , the natural map*

$$\mathfrak{C}_{\Delta}^{\mathcal{Nec}}(X)(x, x') \rightarrow \mathfrak{C}_{\Delta}^{\mathcal{G}}(X)(x, x')$$

induced by the inclusion $\mathcal{Nec} \hookrightarrow \mathcal{G}$ is a weak homotopy equivalence. Moreover, if \mathcal{G} is closed under wedges, the natural transformation of functors valued in simplicially-enriched categories

$$\mathfrak{C}_{\Delta}^{\mathcal{Nec}} \rightarrow \mathfrak{C}_{\Delta}^{\mathcal{G}}$$

is a weak equivalence.

As in the previous section, given a category \mathcal{G} of gadgets, we can extend the realization functor to a functor

$$\mathfrak{C}^{\mathcal{G}} : \widehat{\Theta}[\mathcal{C}]_{*,*} \rightarrow \mathbf{Psh}_{\Delta}(\mathcal{C}),$$

and when \mathcal{G} is closed under wedges these specify a functor

$$\mathfrak{C}^{\mathcal{G}} : \widehat{\Theta}[\mathcal{C}] \rightarrow \mathbf{Cat}_{\mathbf{Psh}_{\Delta}(\mathcal{C})},$$

from which we obtain the following corollary:

Corollary 2.3.3. *The inclusion $\mathcal{Nec} \hookrightarrow \mathcal{G}$ induces a natural equivalence of functors valued in $\mathbf{Psh}_\Delta(\mathcal{C})$*

$$\mathfrak{C}^{\mathcal{Nec}}(X)(x, x') \rightarrow \mathfrak{C}^{\mathcal{G}}(X)(x, x'),$$

and when \mathcal{G} is closed under wedges, these assemble to a natural equivalence of functors valued in $\mathbf{Cat}_{\mathbf{Psh}_\Delta(\mathcal{C})}$:

$$\mathfrak{C}^{\mathcal{Nec}} \rightarrow \mathfrak{C}^{\mathcal{G}}.$$

2.4 Quillen functoriality

In this section, we show that the adjunction

$$\widehat{\Theta[\mathcal{C}]}_{\mathbf{hJoyal}} \xrightleftharpoons[\mathfrak{N}]{\mathfrak{C}} \mathbf{Cat}_{\mathbf{Psh}_\Delta(\mathcal{C})_{\text{inj}}}$$

is a Quillen pair. We will extensively use the characterization of \mathfrak{C} given in Proposition 2.1.10. We begin with the following observation:

Proposition 2.4.1. *For any $n > 0$, let $K \subseteq \{1, \dots, n-1\}$ and define*

$$\Lambda_K^n = \bigcup_{i \notin K} \partial_i \Delta^n,$$

and let

$$\lambda_K^n : \Lambda_K^n \hookrightarrow \Delta^n$$

denote the inclusion map. Then

$$\mathfrak{C}(\square_n^\perp(\lambda_K^n, \delta^{c_1}, \dots, \delta^{c_n}))(i, j)$$

is an isomorphism whenever $i \neq 0$ or $j \neq n$. Moreover, the map

$$\mathfrak{C}(\square_n^\perp(\lambda_K^n, \delta^{c_1}, \dots, \delta^{c_n}))(0, n)$$

is exactly

$$\delta^{c_1} \times^\perp h_K^1 \times^\perp \dots \times^\perp h_K^{n-1} \times^\perp \delta^{c_n},$$

where

$$h_K^k = \begin{cases} \lambda_1^1 & \text{if } k \in K \\ \delta^1 & \text{otherwise} \end{cases}.$$

Proof. Let X denote the domain of $\square_n^\perp(\lambda_K^n, \delta^{c_1}, \dots, \delta^{c_n})$. If $f : T \rightarrow [n](c_1, \dots, c_n)_{i,j}$ is a bi-pointed map from a necklace T , with $i \neq 0$, then f factors through the inclusion of the subobject $[n-1](c_2, \dots, c_n) \subseteq V_{\Lambda_K^n}(c_1, \dots, c_n)$, so $\mathfrak{C}(X)(i, j) = \mathfrak{C}([n](c_1, \dots, c_n))$. The case where $j \neq n$ follows by symmetry.

The second part comes from the observation that when $K = \{1, \dots, n-1\}$,

$$\mathfrak{C}(V_{\Lambda_K^n}(c_1, \dots, c_n))(0, n) = \bigcup_{i=1}^{n-1} c_1 \times \Gamma_i^1 \times \dots \times \Gamma_i^{n-1} \times c_n,$$

where

$$\Gamma_i^\ell = \begin{cases} \Lambda_1^1 & \text{for } \ell = i \\ \Delta^1 & \text{otherwise} \end{cases}.$$

To see this, notice that Λ_K^n is the union of the two outer faces, and attaching them along their common face gives a colimit in $\mathbf{Cat}_{\mathbf{Psh}_\Delta(\mathcal{C})}$ where $\mathfrak{C}(V_{\Lambda_K^n}(c_1, \dots, c_n)(0, n)$ is freely generated by compositions

$$\mathfrak{C}([n-1](c_1, \dots, c_{n-1}))(0, \ell) \times \{1\} \times \mathfrak{C}([n-1](c_2, \dots, c_n))(\ell, n).$$

For when K is otherwise, each additional inner face gives the factor

$$\mathfrak{C}([n-1](c_1, \dots, c_{n-1}))(0, \ell) \times \{0\} \times \mathfrak{C}([n-1](c_2, \dots, c_n))(\ell, n),$$

so in general,

$$\mathfrak{C}(V_{\Lambda_K^n}(c_1, \dots, c_n))(0, n) = \bigcup_{i=1}^{n-1} c_1 \times \Gamma_{i,K}^1 \times \dots \times \Gamma_{i,K}^{n-1} \times c_n,$$

where

$$\Gamma_{i,K}^\ell = \begin{cases} \partial\Delta^1 & \text{for } \ell = i \text{ and } i \in K \\ \Lambda_1^1 & \text{for } \ell = i \text{ and } i \notin K \\ \Delta^1 & \text{otherwise} \end{cases}.$$

Each factor

$$V[n](c_1, \dots, \partial c_j, \dots, c_n)$$

contributes

$$\mathfrak{C}(V[n](c_1, \dots, \partial c_j, \dots, c_n))(0, n) = c_1 \times \Delta^1 \times \dots \times \Delta^1 \times \partial c_j \times \Delta^1 \times \dots \times \Delta^1 \times c_n,$$

and taking the union of all of the factors gives exactly the domain of the inclusion

$$\delta^{c_1} \times^\perp h_K^1 \times^\perp \dots \times^\perp h_K^{n-1} \times^\perp \delta^{c_n}.$$

□

Proposition 2.4.2. *The functor \mathfrak{C} sends monomorphisms to cofibrations and horizontal inner anodynes to trivial cofibrations.*

Proof. Let

$$\mathbf{2} : \mathbf{Psh}_\Delta(\mathcal{C}) \rightarrow \mathbf{Cat}_{\mathbf{Psh}_\Delta(\mathcal{C})}$$

be the functor sending a simplicial presheaf X to the enriched category with objects $\{0, 1\}$ with $\mathbf{2}(X)(0, 0) = \mathbf{2}(X)(1, 1) = *$, $\mathbf{2}(X)(1, 0) = \emptyset$, and $\mathbf{2}(X)(0, 1) = X$.

When $K = \emptyset$, $\lambda_K^n = \delta^n$, so the lemma tells us that

$$\mathfrak{C}(\square_n^\perp(\delta^n, \delta^{c_1}, \dots, \delta^{c_n}))$$

is a pushout of the map

$$\mathbf{2}(\delta^{c_1} \times^\perp \delta^1 \times^\perp \dots \times^\perp \delta^1 \times^\perp \delta^{c_n}),$$

which is a cofibration, which proves the claim.

Similarly, when K is a singleton, $\lambda_K^n = \lambda_k^n$ is the inclusion of an inner horn, so

$$\mathfrak{C}(\square_n^\perp(\lambda_k^n, \delta^{c_1}, \dots, \delta^{c_n}))$$

is the pushout of the map

$$\mathbf{2}(\delta^{c_1} \times^\perp h_k^1 \times^\perp \dots \times^\perp h_k^{n-1} \times^\perp \delta^{c_n}),$$

where $h_k^k = \lambda_1^1$. This is a corner map where one factor is a trivial cofibration (because it is Kan anodyne), and therefore its image under $\mathbf{2}$ is a trivial cofibration. Since the pushout of a trivial cofibration is a trivial cofibration, we are done. \square

Corollary 2.4.3. *The coherent nerve of a fibrant $\text{Psh}_\Delta(\mathcal{C})_{\text{inj}}$ -enriched category is a formal \mathcal{C} -quasicategory.*

Lemma 2.4.4. *The object $\mathfrak{C}(E^n)$ is weakly contractible for all n .*

Proof. We notice immediately that $\mathfrak{C}(E^n)(i, j)_\bullet$ is a constant simplicial presheaf for all i, j , so it suffices to show that $\mathfrak{C}(E^n)(i, j)_*$ is contractible for all i, j , but then it follows immediately from the classical case. \square

Proposition 2.4.5. *The coherent nerve sends fibrations between fibrant $\text{Psh}_\Delta(\mathcal{C})$ -enriched categories to fibrations for the horizontal Joyal model structure.*

Proof. Given a fibration between two fibrant $\text{Psh}_\Delta(\mathcal{C})$ -enriched categories, $p : \mathcal{D} \rightarrow \mathcal{D}'$, we see immediately that the coherent nerve takes this fibration to a horizontal inner fibration between formal \mathcal{C} -quasicategories by Proposition 2.4.2. To show that it is a fibration for the horizontal Joyal model structure, it suffices by Theorem 1.3.8 to show that it has the right lifting property with respect to the inclusion $e : \Delta^0 \hookrightarrow E^1$. By Proposition 2.4.2, we see that \mathfrak{C} takes the monomorphism e to a cofibration, and by the previous lemma, we see that $\mathfrak{C}(e)$ is a weak equivalence. It follows that $\mathfrak{N}(p)$ is a fibration for the horizontal Joyal model structure. \square

Corollary 2.4.6. *The adjunction*

$$\widehat{\Theta[\mathcal{C}]_{\text{hJoyal}}} \xrightleftharpoons[\mathfrak{N}]{\mathfrak{C}} \mathbf{Cat}_{\text{Psh}_\Delta(\mathcal{C})_{\text{inj}}}$$

is a Quillen pair.

Proof. If \mathfrak{C} takes cofibrations to cofibrations, and \mathfrak{N} takes fibrations between fibrant objects to fibrations between fibrant objects, then the adjunction is a Quillen pair, but this is exactly what we proved in this section. \square

The following lemma will be useful for later.

Lemma 2.4.7. *If $A \hookrightarrow B$ is an inner anodyne map of simplicial sets, the induced map*

$$k_!(A \times c) \hookrightarrow k_!(B \times c)$$

is horizontal inner anodyne. Consequently, if X is a horizontal \mathcal{C} -quasicategory, each simplicial set $k^(X)_c$ is a quasicategory.*

Proof. This follows directly by Lemma 1.7.2 applied to the maps

$$V_{\Lambda_i^n}(c, \dots, c) \hookrightarrow V_{\Delta^n}(c, \dots, c),$$

as these maps are exactly the maps

$$k_!(\Lambda_i^n \times c) \hookrightarrow k_!(\Delta^n \times c).$$

□

2.5 The Hom by cosimplicial resolutions for \mathcal{C} -precategories

In this section, we show that if X is a \mathcal{C} -precategory such that X_c is a quasicategory for all $c \in \mathcal{C}$, the simplicial presheaves $\mathfrak{C}_{\Delta, \bullet}(X)(x, x')$ can be computed by resolutions. To show this, we will make use of some helpful results in [DS11b].

We define the following four cosimplicial bi-pointed simplicial sets:

$$C_{\text{cyl}}^\bullet \stackrel{\text{def}}{=} \text{colim} \left(\partial \Delta^1 \leftarrow \Delta^\bullet \times \partial \Delta^1 \hookrightarrow \Delta^\bullet \times \Delta^1 \right)$$

$$C_E^\bullet \stackrel{\text{def}}{=} \text{colim} \left(\partial \Delta^1 \leftarrow \Delta^\bullet \times \partial \Delta^1 \hookrightarrow \Delta^\bullet \times E^1 \right)$$

$$C_R^\bullet \stackrel{\text{def}}{=} \text{colim} \left(* \leftarrow \Delta^\bullet \xrightarrow{\partial^{\bullet+1}} \Delta^{\bullet+1} \right)$$

$$C_L^\bullet \stackrel{\text{def}}{=} \text{colim} \left(* \leftarrow \Delta^\bullet \xrightarrow{\partial^0} \Delta^{\bullet+1} \right).$$

These cosimplicial bi-pointed simplicial sets fit in a natural diagram

$$\begin{array}{ccc} C_R^\bullet & & \\ & \searrow & \\ & C_{\text{cyl}}^\bullet & \longrightarrow C_E^\bullet \\ & \nearrow & \\ C_L^\bullet & & \end{array}.$$

As these are cosimplicial objects in a cocomplete category, they induce adjunctions admit right adjoints

$$\text{Map}^{(-)} : \widehat{\Delta}_{*,*} \rightarrow \widehat{\Delta},$$

where

$$\text{Map}_X^{(-)}(x, x')_n \stackrel{\text{def}}{=} \text{Hom}(C_{(-)}^n, X_{x, x'})$$

Lemma 2.5.1 ([DS11b, Proposition 9.4]). *Each of the cosimplicial bi-pointed simplicial sets above is a Reedy-cofibrant cosimplicial resolution of Δ^1 with respect to the bi-pointed Joyal model structure.*

The following corollary is immediate from the fact that quasicategories are the fibrant objects of the Joyal model structure together with the general theory of cosimplicial resolutions:

Corollary 2.5.2. *If $X_{x,x'}$ is a bi-pointed quasicategory, then there is a natural isomorphism in the homotopy category*

$$\mathrm{Map}_X^{(-)}(x, x') \cong \mathrm{hMap}_X(x, x'),$$

where the $(-)$ on the lefthand side can take any of the values cyl , E , R or L , and where the righthand side denotes the homotopy function complex of maps of bi-pointed simplicial sets

$$\Delta^1 \rightarrow X_{x,x'}.$$

Since the choice of resolution doesn't matter, we abuse notation and let Map and C^\bullet denote whichever choice of resolution and adjoint that is convenient. The following theorem is a key result in [DS11b].

Theorem 2.5.3 ([DS11b, Corollary 5.3]). *There is a zig-zag, natural in $X_{x,x'}$*

$$\mathfrak{C}_\Delta(X)(x, x') \rightleftarrows \mathrm{Map}_X(x, x'),$$

which becomes a zig-zag of natural weak homotopy equivalences upon restriction to bi-pointed quasicategories.

From the naturality of this result, we obtain the following corollary:

Corollary 2.5.4. *If $X_{x,x'}$ is a bi-pointed \mathcal{C} -precategory such that each X_c is a quasicategory, the component of the natural zig-zag of maps of simplicial presheaves on \mathcal{C} at $X_{x,x'}$ is a zig-zag of weak equivalences in $\mathrm{Psh}_\Delta(\mathcal{C})$:*

$$\mathfrak{C}_{\Delta, \bullet}(X)(x, x')_c \rightleftarrows \mathrm{Map}_{X_c}(x, x')$$

Combining this corollary with Lemma 2.4.7, we obtain the following:

Corollary 2.5.5. *Upon restriction to bi-pointed horizontal quasicategories, we have a natural zig-zag of weak equivalences in $\mathrm{Psh}_\Delta(\mathcal{C})$*

$$\mathfrak{C}(X)(x, x')_c \rightleftarrows \mathrm{Map}_{k^*(X)_c}(x, x').$$

Unwinding the definitions, we note the following corollary:

Corollary 2.5.6. *If $X_{x,x'}$ is a bi-pointed horizontal quasicategory, we have zig-zags of weak equivalences in $\mathrm{Psh}_\Delta(\mathcal{C})$*

$$\mathfrak{C}(X)(x, x')_c \rightleftarrows \mathrm{Hom}(k_!(C_R^\bullet \times c), X_{x,x'}),$$

and

$$\mathfrak{C}(X)(x, x')_c \rightleftarrows \mathrm{Hom}(k_!(C_L^\bullet \times c), X_{x,x'}).$$

We also make note of one more useful fact:

Lemma 2.5.7. *The cosimplicial bi-pointed \mathcal{C} -cellular sets $k_!(C_R^\bullet \times c)$ and $k_!(C_L^\bullet \times c)$ are Reedy-cofibrant cosimplicial resolutions of $[1](c) = k_!(\Delta^1 \times c)$ in $\widehat{\Theta[\mathcal{C}]_{*,*}}$.*

Proof. From [DS11b, Proposition 9.4], we see that the maps

$$C_R^n \rightarrow \Delta^1$$

and

$$C_L^n \rightarrow \Delta^1$$

are retracts of the inner anodynes inclusions

$$\Delta^1 \hookrightarrow C_R^n$$

and

$$\Delta^1 \hookrightarrow C_L^n$$

respectively. It then follows immediately from Lemma 2.4.7 that the maps

$$k_!(C_R^n \times c) \rightarrow k_!(\Delta^1 \times c) = [1](c)$$

and

$$k_!(C_L^n \times c) \rightarrow k_!(\Delta^1 \times c) = [1](c)$$

are retracts of horizontal inner anodynes and ergo equivalences. Reedy cofibrancy is clear. \square

2.6 The Hom by cosimplicial resolution

For every object $[1](c)$ in $\Theta[\mathcal{C}]$, we introduce four canonical cosimplicial resolutions, which we can use to define simplicial presheaves that represent the mapping space between two vertices of a $\Theta[C]$ -set.

First, we define the functor

$$(\bullet)^\triangleright(c), \text{ resp. } (\bullet)^\triangleleft(c) : \Delta \rightarrow \Theta[\mathcal{C}],$$

which sends

$$[n] \mapsto [n+1](*, \dots, *, c), \text{ resp. } [n] \mapsto [n+1](c, *, \dots, *).$$

We see immediately that there are natural embeddings

$$([n])^{\triangleright, c} \hookrightarrow \Delta^n \times [1](c) \hookleftarrow ([n])^{\triangleleft, c}$$

where each map embeds along the respective outer shuffle. We also have an obvious natural embedding

$$\mathcal{H}(\Delta^n) \times [1](c) \hookrightarrow E^n \times [1](c).$$

Then we define the following four cosimplicial objects:

$$C_{\text{cyl}}^\bullet(c) \stackrel{\text{def}}{=} \text{colim} (V[1](\emptyset) \leftarrow \Delta^\bullet \times V[1](\emptyset) \hookrightarrow \Delta^\bullet \times [1](c))$$

$$C_E^\bullet(c) \stackrel{\text{def}}{=} \text{colim} (V[1](\emptyset) \leftarrow E^\bullet \times V[1](\emptyset) \hookrightarrow E^\bullet \times [1](c))$$

$$C_R^\bullet(c) \stackrel{\text{def}}{=} \text{colim} (* \leftarrow \Delta^\bullet \hookrightarrow (\bullet)^\triangleright(c))$$

$$C_L^\bullet(c) \stackrel{\text{def}}{=} \text{colim} (* \leftarrow \Delta^\bullet \hookrightarrow (\bullet)^\triangleleft(c)).$$

These cosimplicial objects fit in a natural diagram

$$\begin{array}{ccc}
C_R^\bullet(c) & & \\
& \searrow & \\
& C_{\text{cyl}}^\bullet(c) & \longrightarrow C_E^\bullet(c) \\
& \nearrow & \\
C_L^\bullet(c) & &
\end{array}$$

induced by the inclusions we described above.

Proposition 2.6.1. *Each of the cosimplicial objects described above is a Reedy-cofibrant object of bipointed presheaves on $\Theta[\mathcal{C}]$, and each is objectwise horizontal-Joyal equivalent to the constant cosimplicial object $[1](c)$. That is to say, each of these cosimplicial objects is a cosimplicial resolution of $[1](c)$ in $\Theta[\widehat{\mathcal{C}}]_{*,*}$.*

Proof. That they are Reedy-cofibrant is obvious, and it is also obvious that $C_E^\bullet(c)$ is objectwise horizontal-Joyal equivalent to $[1](c)$. To treat the case of $C_L^\bullet(c)$, we follow the proof of [DS11b, Lemma 9.3]. We begin by defining a filtration on $C_L^n(c)$ as follows:

$$\begin{aligned}
X_0 &= [0, 1](c), \\
X_1 &= \bigcup_{1 < i \leq n} [0, 1, i](c, *), \\
X_2 &= \bigcup_{1 < i < j \leq n} [0, 1, i, j](c, *, *),
\end{aligned}$$

and so on. We see that $X_{n-1} = C_L^n(c)$, so it suffices to show that each inclusion $X_{k-1} \subseteq X_k$ is obtained by horizontal inner anodyne attachments. Notice first of all that

$$h_1^\ell : \square_\ell(\Lambda_1^\ell, c, *, \dots, *) \hookrightarrow \square_\ell(\Delta^\ell, c, *, \dots, *)$$

is horizontal inner anodyne, since it is exactly

$$\square_k^\perp(\lambda_1^\ell, e^c, e^*, \dots, e^*),$$

where e^x denotes the empty map $\emptyset \rightarrow x$. Then choose $1 < i_1 < \dots < i_k \leq n$ in X_k . Then X_{k-1} contains all of the faces $[0, 1, i_1, \dots, \hat{i}_{k_j}, \dots, i_k](c, *, \dots, *)$, but it also contains the degenerate face $[1, i_1, \dots, i_k](*, \dots, *)$, so therefore the map $X_{k-1} \subseteq X_k$ is obtained by pushing out along copies of h_1^{k+1} , and therefore it is horizontal inner anodyne. Therefore, the composite of the filtration $[1](c) = X_0 \subseteq \dots \subseteq X_{n-1}$ is horizontal inner anodyne, and by 3-for-2, we see that the retraction $C_L^n(c) \rightarrow [1](c)$ is a horizontal Joyal equivalence. The proof for C_R^\bullet follows by symmetry. The proof for $C_{\text{cyl}}^\bullet(c)$ is similar and can be obtained by applying the scheme for $C_L^\bullet(c)$ to the proof of that case in [DS11b, Proposition 9.4]. We omit it because we do not make use of that particular resolution. \square

Definition 2.6.2. Given a bipointed \mathcal{C} -cellular set $X_{x,y}$ we define the *mapping object* from x to y to the simplicial presheaf obtained by taking homotopy function complexes

$$\text{Map}_X(x, y)_c \stackrel{\text{def}}{=} h\widehat{\Theta[\mathcal{C}]}_{*,*}([1](c), X).$$

Remark 2.6.3. It is a general fact of abstract homotopy theory that if X is a fibrant object, we can compute the homotopy function complex with any Reedy-cofibrant cosimplicial resolution of $[1](c)$. Any two Reedy-cofibrant cosimplicial resolutions are related by a natural zig-zag of weak equivalences, so any one will do.

Proposition 2.6.4. *For any formal \mathcal{C} -quasicategory X and any pair of vertices x, y , there is a natural zig-zag of weak equivalences between $\mathrm{Map}_X(x, y)$ and $\mathfrak{C}(X)(x, y)$.*

Proof. By Lemma 2.5.7, we see that we can equivalently compute $\mathrm{Map}_X(x, y)$ using the cosimplicial resolutions $k_!(C_R^\bullet \times c)$ of $[1](c)$, from which the result follows by Corollary 2.5.6. \square

Lemma 2.6.5. *For any bi-pointed formal \mathcal{C} -quasicategory $X_{x,y}$, the functor*

$$\left(\Delta \downarrow \widehat{\Theta}[\mathcal{C}]_{*,*}(C_R^\bullet(c), X_{x,y}) \right) \rightarrow \left(\Delta \downarrow \widehat{\Theta}[\mathcal{C}]_{*,*}(k_!(C_R^\bullet \times c), X_{x,y}) \right)$$

induced by the map of cosimplicial resolutions of $[1](c)$

$$k_!(C_R^\bullet \times c) \rightarrow C_R^\bullet(c)$$

induces a weak homotopy equivalence on nerves. Moreover, as we have a natural isomorphism

$$\left(\Delta \downarrow \widehat{\Theta}[\mathcal{C}]_{*,*}(k_!(C_R^\bullet \times c), X_{x,y}) \right) \cong \left(\Delta \downarrow \widehat{\Delta}_{*,*}(C_R^\bullet, k^*(X_{x,y})_c) \right),$$

we have a commutative diagram in which the specified maps are weak homotopy equivalences

$$\begin{array}{ccc} \mathrm{colim}_{[n], C_R^n \rightarrow k^*(X_{x,y})_c} \mathfrak{C}_\Delta(C_R^m)(\alpha, \omega) & \xleftarrow{\sim} & \mathrm{colim}_{[n], C_R^n(c) \rightarrow X_{x,y}} \mathfrak{C}_\Delta(C_R^m)(\alpha, \omega) \\ \uparrow & & \uparrow \\ \mathrm{hocolim}_{[n], C_R^n \rightarrow k^*(X_{x,y})_c} \mathfrak{C}_\Delta(C_R^m)(\alpha, \omega) & \xleftarrow{\sim} & \mathrm{hocolim}_{[n], C_R^n(c) \rightarrow X_{x,y}} \mathfrak{C}_\Delta(C_R^m)(\alpha, \omega) \\ \downarrow \sim & & \downarrow \sim \\ \mathrm{hocolim}_{[n], C_R^n \rightarrow k^*(X_{x,y})_c} * & \xleftarrow{\sim} & \mathrm{hocolim}_{[n], C_R^n(c) \rightarrow X_{x,y}} * \end{array},$$

Proof. Any weak homotopy equivalence of simplicial sets induces a weak homotopy equivalence on nerves of their categories of elements. As the map $k_!(C_R^\bullet \times c) \rightarrow C_R^\bullet(c)$ is a map between Reedy-cofibrant resolutions of $[1](c)$, it follows that the induced map

$$\widehat{\Theta}[\mathcal{C}]_{*,*}(C_R^\bullet(c), X_{x,y}) \rightarrow \widehat{\Theta}[\mathcal{C}]_{*,*}(k_!(C_R^\bullet \times c), X_{x,y})$$

is a weak homotopy equivalence whenever $X_{x,y}$ is fibrant. Taken together, these observations prove the first claim. To obtain the diagram in the statement, notice that the vertical maps are equivalences as each of the simplicial sets $\mathfrak{C}_\Delta(C_R^m)(\alpha, \omega)$ is weakly contractible and that the bottom horizontal map is an equivalence, since each of these homotopy colimits is naturally weakly equivalent to the nerve of the diagram category. The fact that the middle horizontal map is a weak homotopy equivalence follows by 3-for-2. \square

Note 2.6.6. In what follows, we will make use of a special category of gadgets denoted by \mathcal{Y} . It is the full subcategory of $\widehat{\Delta}_{*,*}$ spanned by those bi-pointed simplicial sets $Y_{\alpha,\omega}$ such that $\mathfrak{C}_{\Delta}(Y)(\alpha,\omega)$ is contractible. The properties of this category of gadgets are spelled out in [DS11b, Section 5].

Lemma 2.6.7. *Given a $\text{Psh}_{\Delta}(\mathcal{C})$ -enriched category \mathcal{D} and any pair of objects $x, y \in \mathcal{D}$, there is a natural commutative diagram*

$$\begin{array}{ccc} \mathfrak{C}(\mathfrak{N}\mathcal{D})(x, y)_c & \xrightarrow{\quad\quad\quad} & \mathcal{D}(x, y)_c \\ \uparrow \cong & & \uparrow \\ \text{colim}_{T \in (\text{Nec} \downarrow k^*(\mathfrak{N}\mathcal{D}_{x,y})_c)} \mathfrak{C}_{\Delta}(T)(\alpha, \omega) & \longrightarrow & \text{colim}_{Y \in (\mathcal{Y} \downarrow k^*(\mathfrak{N}\mathcal{D}_{x,y})_c)} \mathfrak{C}_{\Delta}(Y)(\alpha, \omega) \end{array},$$

in which the specified map is an isomorphism.

Proof. Recall that $\mathfrak{N} = k_* \circ \mathfrak{N}_{\Delta, \bullet}$, so given a map

$$f : Y \rightarrow k^*(\mathfrak{N}\mathcal{D}_{x,y})_c,$$

we obtain the map $\mathfrak{C}(Y) \rightarrow \mathcal{D}_c$ as the transpose of the composite of f with the whiskered counit of the adjunction $k^*k_* \rightarrow \text{id}$. In particular, for each such map, we obtain a universal factorization

$$\mathfrak{C}_{\Delta}(Y) \rightarrow \mathfrak{C}\mathfrak{N}\mathcal{D}_c \rightarrow \mathcal{D}_c.$$

This determines the righthand vertical map after taking colimits. Upon restriction to

$$(\text{Nec} \downarrow k^*(\mathfrak{N}\mathcal{D}_{x,y})_c),$$

we see that this factorization gives rise to the lefthand vertical map, which is an isomorphism by [DS11a, Proposition 4.3]. \square

Theorem 2.6.8. *For any fibrant $\text{Psh}_{\Delta}(\mathcal{C})_{\text{inj}}$ -enriched category \mathcal{D} , the counit map*

$$\epsilon_{\mathcal{D}} : \mathfrak{C}(\mathfrak{N}\mathcal{D}) \rightarrow \mathcal{D}$$

is a weak equivalence of $\text{Psh}_{\Delta}(\mathcal{C})_{\text{inj}}$ -enriched categories.

Proof. Let $C^{\bullet} = C_R^{\bullet}$. Then consider the following commutative diagram:

$$\begin{array}{ccccc}
\mathfrak{C}(\mathfrak{N}\mathcal{D})(x, y)_c & \longrightarrow & \mathcal{D}(x, y)_c & & \\
\uparrow \cong & & \uparrow & \nwarrow & \\
\operatorname{colim}_{T \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} \mathfrak{C}_\Delta(T)(\alpha, \omega) & \longrightarrow & \operatorname{colim}_{Y \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} \mathfrak{C}_\Delta(Y)(\alpha, \omega) & \longleftarrow & \operatorname{colim}_{[n], C^n(c) \rightarrow \mathfrak{N}\mathcal{D}} \mathfrak{C}_\Delta(C^n)(\alpha, \omega) \\
\uparrow \sim & & \uparrow & & \uparrow \\
\operatorname{hocolim}_{T \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} \mathfrak{C}_\Delta(T)(\alpha, \omega) & \longrightarrow & \operatorname{hocolim}_{Y \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} \mathfrak{C}_\Delta(Y)(\alpha, \omega) & \longleftarrow & \operatorname{hocolim}_{[n], C^n(c) \rightarrow \mathfrak{N}\mathcal{D}} \mathfrak{C}_\Delta(C^n)(\alpha, \omega) \\
\downarrow \sim & & \downarrow \sim & & \downarrow \sim \\
\operatorname{hocolim}_{T \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} * & \xrightarrow{\sim} & \operatorname{hocolim}_{Y \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} * & \xleftarrow{\sim} & \operatorname{hocolim}_{[n], C^n(c) \rightarrow \mathfrak{N}\mathcal{D}} *
\end{array}$$

This diagram is obtained by composing the diagram from [DS11b, Proposition 5.2] reproduced below with the diagrams appearing in Lemmas 2.6.5 and 2.6.7

$$\begin{array}{ccccc}
\operatorname{colim}_{T \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} \mathfrak{C}_\Delta(T)(\alpha, \omega) & \longrightarrow & \operatorname{colim}_{Y \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} \mathfrak{C}_\Delta(Y)(\alpha, \omega) & \longleftarrow & \operatorname{colim}_{[n], C^n \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} \mathfrak{C}_\Delta(C^n)(\alpha, \omega) \\
\uparrow \sim & & \uparrow & & \uparrow \\
\operatorname{hocolim}_{T \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} \mathfrak{C}_\Delta(T)(\alpha, \omega) & \longrightarrow & \operatorname{hocolim}_{Y \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} \mathfrak{C}_\Delta(Y)(\alpha, \omega) & \longleftarrow & \operatorname{hocolim}_{[n], C^n \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} \mathfrak{C}_\Delta(C^n)(\alpha, \omega) \\
\downarrow \sim & & \downarrow \sim & & \downarrow \sim \\
\operatorname{hocolim}_{T \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} * & \xrightarrow{\sim} & \operatorname{hocolim}_{Y \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} * & \xleftarrow{\sim} & \operatorname{hocolim}_{[n], C^n \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} *
\end{array}$$

The colimits in the left column are indexed over $(\mathcal{N}ec \downarrow k^*(\mathfrak{N}\mathcal{D}_{x,y})_c)$, the colimits in the middle column are indexed by $(\mathcal{Y} \downarrow k^*(\mathfrak{N}\mathcal{D}_{x,y})_c)$, and the colimits in the righthand column are indexed by $(\Delta \downarrow \widehat{\Delta}_{*,*}(C_R^\bullet, k^*(X_{x,y})_c))$. The two horizontal maps in the bottom row are weak equivalences by [DS11b, Proposition 5.2], as are the downward-oriented vertical maps, using the fact that $k^*(\mathfrak{N}\mathcal{D})_c$ is a quasicategory. The indicated upward-oriented vertical map is a weak equivalence by [DS11a, Theorem 5.2]. By 3-for-2, it follows that the horizontal maps in the third row are all weak equivalences. We therefore reduce the composite diagram to a smaller diagram:

$$\begin{array}{ccccc}
\mathfrak{C}(\mathfrak{N}\mathcal{D})(x, y)_c & \longrightarrow & \mathcal{D}(x, y)_c & & \\
\uparrow \sim & & \uparrow & \nwarrow \gamma_c & \\
\operatorname{hocolim}_{T \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} \mathfrak{C}_\Delta(T)(\alpha, \omega) & \xrightarrow{\sim} & \operatorname{hocolim}_{Y \rightarrow k^*(\mathfrak{N}\mathcal{D})_c} \mathfrak{C}_\Delta(Y)(\alpha, \omega) & \xleftarrow{\sim} & \operatorname{hocolim}_{[n], C^n(c) \rightarrow \mathfrak{N}\mathcal{D}} \mathfrak{C}_\Delta(C^n)(\alpha, \omega)
\end{array}$$

By 3-for-2, we can see that it suffices to show that the map

$$\gamma_c : \operatorname{hocolim}_{[n], C^n(c) \rightarrow \mathfrak{N}\mathcal{D}} \mathfrak{C}_\Delta(C^n)(\alpha, \omega) \rightarrow \operatorname{colim}_{[n], C^n(c) \rightarrow \mathfrak{N}\mathcal{D}} \mathfrak{C}_\Delta(C^n)(\alpha, \omega) \rightarrow \mathcal{D}(x, y), c$$

is a weak equivalence. To do this, notice that

$$\begin{aligned} \widehat{\Theta[\mathcal{C}]}_{*,*}(C^n(c), \mathfrak{N}\mathcal{D}_{x,y}) &\cong (\partial[1] \downarrow \mathbf{Cat}_{\operatorname{Psh}_\Delta(\mathcal{C})})(\mathfrak{C}(C^n(c))_{\alpha, \omega}, \mathcal{D}_{x,y}) \\ &\cong \operatorname{Psh}_\Delta(\mathcal{C})(\mathfrak{C}(C^n(c))(\alpha, \omega), \mathcal{D}(x, y)). \end{aligned}$$

As in the proof of [DS11b, Proposition 5.8], we define the cosimplicial simplicial set $Q^\bullet = \mathfrak{C}_\Delta(C^\bullet)(\alpha, \omega)$, which is obviously isomorphic to the Q^\bullet defined in [DS11b]. Moreover, by direct computation, we see that

$$\mathfrak{C}(C^\bullet(c))(\alpha, \omega) \cong Q^\bullet \times c.$$

Then we see immediately that

$$\operatorname{Psh}_\Delta(\mathcal{C})(\mathfrak{C}(C^n(c))(\alpha, \omega), \mathcal{D}(x, y)) \cong \operatorname{Psh}_\Delta(\mathcal{C})(Q^n \times c, \mathcal{D}(x, y)) \cong \widehat{\Delta}(Q^n, \mathcal{D}(x, y)_c),$$

so γ_c is precisely the map obtained by composing

$$\left(\operatorname{hocolim}_{[n], Q^n \rightarrow \mathcal{D}(x, y)_c} Q^n \right) \rightarrow \left(\operatorname{colim}_{[n], Q^n \rightarrow \mathcal{D}(x, y)_c} Q^n \right) \rightarrow \mathcal{D}(x, y)_c.$$

The result then follows immediately by application of [DS11b, Lemma 5.9]. \square

Remark 2.6.9. This result is even stronger than it first appears, because it implies that the counit map is a weak equivalence for fibrant categories enriched in any Cartesian-closed left-Bousfield localization of $\operatorname{Psh}_\Delta(\mathcal{C})_{inj}$. It reduces proving comparison theorems for such localizations to showing that \mathfrak{C} is a left-Quillen functor (something we already know for the horizontal Joyal model structure by Corollary 2.4.6) and reflects weak equivalences.

2.7 The horizontal comparison theorem

Dugger and Spivak introduce a definition of a Dwyer-Kan equivalence as a stepping stone to proving the comparison theorem. They use the definition of DK-equivalence as an intermediate step to proving that \mathfrak{C}_Δ is homotopy-conservative. We give an analogous definition as follows:

Definition 2.7.1. A map $f : X \rightarrow Y$ of presheaves on $\Theta[\mathcal{C}]$ is called a *horizontal Dwyer-Kan equivalence* if the following two properties hold:

- The induced map

$$f_* : \operatorname{Ho}(\widehat{\Theta[\mathcal{C}]}_{\operatorname{hJoyal}}(*, X) \rightarrow \operatorname{Ho}(\widehat{\Theta[\mathcal{C}]}_{\operatorname{hJoyal}}(*, Y)$$

is bijective, and

- For any two vertices $x, x' \in X_0$, the induced map

$$\operatorname{Map}_X(x, x') \rightarrow \operatorname{Map}_Y(f(x), f(x'))$$

is a weak equivalence of simplicial presheaves on \mathcal{C} .

Proposition 2.7.2. *A map $f : X \rightarrow Y$ of presheaves on $\Theta[\mathcal{C}]$ is a horizontal weak equivalence if and only if it is a horizontal Dwyer-Kan equivalence.*

Proof. It is clear that any horizontal Joyal equivalence is automatically horizontally Dwyer-Kan since our constructions are all homotopy-invariant, so we prove that all horizontal Dwyer-Kan equivalences are horizontal Joyal equivalences. We notice immediately that if X and Y are fibrant, the horizontal Dwyer-Kan condition implies that the associated map $\mathcal{Q}(f) : \mathcal{Q}(X) \rightarrow \mathcal{Q}(Y)$ between complete $\Theta[\mathcal{C}]$ Segal spaces is an equivalence, where

$$\mathcal{Q} : \widehat{\Theta[\mathcal{C}]} \rightarrow \widehat{\Theta[\mathcal{C}]} \times \Delta$$

is defined by the rule

$$\mathcal{Q}(X)_{[n](c_1, \dots, c_n), m} \stackrel{\text{def}}{=} \text{Hom}([n](c_1, \dots, c_n) \times E^m, X).$$

Since \mathcal{Q} is the right adjoint of a Quillen equivalence by Section 1.7, a map f between fibrant objects is a weak equivalence if and only if its image under \mathcal{Q} is. Since Dwyer-Kan equivalences between Complete $\Theta[\mathcal{C}]$ Segal spaces are exactly the weak equivalences by [Rez10], the claim holds for X and Y fibrant.

In general, given a horizontal Dwyer-Kan equivalence $f : X \rightarrow Y$ where X and Y are no longer assumed to be fibrant, we can take a fibrant replacement \tilde{Y} of Y such that $Y \rightarrow Y'$ is a trivial cofibration for the horizontal Joyal model structure. Then we can also factor $X \rightarrow Y \rightarrow \tilde{Y}$ into a trivial Joyal cofibration $X \rightarrow \tilde{X}$ followed by a fibration $\tilde{X} \rightarrow \tilde{Y}$. But notice now that the condition of being horizontally DK-equivalent is homotopy invariant, so the map $\tilde{X} \rightarrow \tilde{Y}$ is also a horizontal DK-equivalence. Since \tilde{Y} is fibrant and $\tilde{X} \rightarrow \tilde{Y}$ is a horizontal Joyal fibration, this is a horizontal Joyal equivalence. Then by 3-for-2 we see that f is also a horizontal Joyal equivalence, which concludes the proof. \square

Proposition 2.7.3. *A map $f : X \rightarrow Y$ of presheaves on $\Theta[\mathcal{C}]$ is a horizontal Joyal equivalence if and only if $\mathfrak{C}(f)$ is a weak equivalence of $\text{Psh}_\Delta(\mathcal{C})_{\text{inj}}$ -enriched categories.*

Proof. We only need to check one direction, since the other direction is immediate by the fact that \mathfrak{C} is left-Quillen. Assume $f : X \rightarrow Y$ has the property that $\mathfrak{C}(f)$ is an equivalence. Then as in the previous proposition, we can reduce to the case where X and Y are fibrant, but in this case, we know from Proposition 2.6.4 that $\mathfrak{C}(X)(x, x')$ is connected by a natural zig-zag of weak equivalences to $\text{Map}_X(x, y)$, so if the map $\mathfrak{C}(X)(x, x') \rightarrow \mathfrak{C}(Y)(f(x), f(x'))$ is a weak equivalence, it follows that the map $\text{Map}_X(x, x') \rightarrow \text{Map}_Y(f(x), f(x'))$ is also a weak equivalence.

Then it suffices to show that when $\mathfrak{C}(f)$ is an equivalence, the induced map on sets of homotopy classes

$$[* , X]_{E^1} \rightarrow [* , Y]_{E^1}$$

is a bijection. Notice that

$$[* , X]_{E^1} \cong \pi_0 \widehat{\Theta[\mathcal{C}]}(E^\bullet, X),$$

and since $E^n = \mathcal{H} \text{cosk}_0 \Delta^n$. By abuse of notation, we also denote the simplicial set $\text{cosk}_0 \Delta^n$ by E^n . We noticed earlier that \mathcal{H} has a right adjoint, which we now denote by \mathcal{N} . Using this, we can rewrite the question as asking for the induced map to give a bijection

$$\pi_0 \widehat{\Delta}(E^n, \mathcal{N}X) \rightarrow \pi_0 \widehat{\Delta}(E^n, \mathcal{N}X),$$

which is the same as giving a bijection

$$[\Delta^0, \mathcal{N}X]_{E^1} \rightarrow [\Delta^0, \mathcal{N}X]_{E^1}$$

Notice also that the data classifying an equivalence in $\mathfrak{C}(X)$ all factor through the simplicial category $\mathfrak{C}(X)_{*_{\mathfrak{C}}}$ obtained by evaluating each of the Hom objects at the terminal object $*_{\mathfrak{C}}$ of \mathfrak{C} . It is an easy exercise to see that

$$\mathfrak{C}(X)_{*_{\mathfrak{C}}} \cong \mathfrak{C}_{\Delta}(\mathcal{N}X),$$

but since $\mathcal{N}X$ is quite clearly a quasicategory, the claim follows immediately from the ordinary case. This implies that f is a horizontal Dwyer-Kan equivalence, and therefore by the previous proposition, a horizontal Joyal equivalence, which concludes the proof. \square

Theorem 2.7.4. *The Quillen pair*

$$\widehat{\Theta[\mathfrak{C}]}_{\mathfrak{hJoyal}} \xrightleftharpoons[\mathfrak{N}]{\mathfrak{C}} \mathbf{Cat}_{\mathbf{Psh}_{\Delta}(\mathfrak{C})_{\text{inj}}}$$

is a Quillen equivalence.

Proof. All we have left to show is that the derived unit of the adjunction,

$$X \rightarrow \mathfrak{N}\mathfrak{C}(X) \rightarrow \mathfrak{N}\mathcal{D}$$

is a weak equivalence for all presheaves X on $\Theta[\mathfrak{C}]$, where $\mathfrak{C}(X) \rightarrow \mathcal{D}$ is a weak equivalence and \mathcal{D} is fibrant. However, by the previous proposition, we see that it suffices to show that the map

$$\mathfrak{C}(X) \rightarrow \mathfrak{C}\mathfrak{N}\mathfrak{C}(X) \rightarrow \mathfrak{C}\mathfrak{N}\mathcal{D}$$

is a weak equivalence. We obtain a naturality square from the counit

$$\begin{array}{ccc} \mathfrak{C}\mathfrak{N}\mathfrak{C}(X) & \longrightarrow & \mathfrak{C}\mathfrak{N}\mathcal{D} \\ \downarrow & & \downarrow \sim \\ \mathfrak{C}(X) & \xrightarrow{\sim} & \mathcal{D} \end{array},$$

in which the indicated arrows are equivalences (for the bottom horizontal, this was by choice, and for the righthand vertical, it comes from Theorem 2.6.8). But if we precompose with the unit map $\mathfrak{C}\eta_X : \mathfrak{C}(X) \rightarrow \mathfrak{C}\mathfrak{N}\mathfrak{C}(X)$, the lefthand arrow becomes the identity by the triangle identities, which proves the claim by applying 3-for-2 to the commutative diagram

$$\begin{array}{ccccc} \mathfrak{C}(X) & \longrightarrow & \mathfrak{C}\mathfrak{N}\mathfrak{C}(X) & \longrightarrow & \mathfrak{C}\mathfrak{N}\mathcal{D} \\ & \searrow & \downarrow & & \downarrow \sim \\ & & \mathfrak{C}(X) & \xrightarrow{\sim} & \mathcal{D} \end{array}.$$

\square

3 The Coherent Nerve, Local case

3.1 The $(\mathcal{C}, \mathcal{S})$ -enriched model structure

While our presentation of the horizontal Joyal model structure comes mainly from David Oury's thesis [Our10], what follows is independent, making use of the resolution technology we developed in the previous section to give a simple and satisfying story. Suppose $\mathcal{M} = (\mathcal{C}, \mathcal{S})$ is a Cartesian presentation in the sense of Rezk, where \mathcal{S} is a set of monomorphisms of $\text{Psh}_\Delta(\mathcal{C})$ such that the left-Bousfield localization of $\text{Psh}_\Delta(\mathcal{C})_{\text{inj}}$ at \mathcal{S} is a Cartesian model category.

Recall that we had a number of functorial cosimplicial objects

$$C_{(-)}^\bullet(\bullet) : \Delta \times \mathcal{C} \rightarrow \widehat{\Theta[\mathcal{C}]_{*,*}},$$

such that $C_{(-)}^\bullet(\bullet)$ was a cosimplicial resolution of $[1](\bullet)$, which is a Reedy cofibrant diagram $\mathcal{C} \rightarrow \widehat{\Theta[\mathcal{C}]_{*,*}}$. Since $\widehat{\Theta[\mathcal{C}]_{*,*}}$ is cocomplete, $C_{(-)}^\bullet(\bullet)$ extends to a cocontinuous functor

$$\Sigma : \text{Psh}_\Delta(\mathcal{C}) \rightarrow \widehat{\Theta[\mathcal{C}]_{*,*}}.$$

Proposition 3.1.1. *The functor $\Sigma_{(-)}$ is left-Quillen when $\widehat{\Theta[\mathcal{C}]_{*,*}}$ is equipped with the horizontal Joyal model structure.*

Proof. It clearly preserves cofibrations, so it suffices to show that its right adjoint preserves fibrations between fibrant objects. However, this is clear, since the right adjoint sends a bi-pointed formal \mathcal{C} quasicategory $X_{x,y}$ to $\text{Map}_X(x, y)$, which we saw sends horizontal Joyal fibrations to injective fibrations of simplicial presheaves on \mathcal{C} . \square

Corollary 3.1.2. *The functor $\Sigma_{(-)}$ is independent up-to-homotopy of choice of resolution $C_{(-)}^\bullet(\bullet)$.*

Proof. Since simplicial presheaves are always canonically the homotopy-colimit of their representables, and since left-Quillen functors send homotopy-colimits to homotopy-colimits, it suffices to show that $\Sigma_{(-)}(\Delta^n \times c)$ is independent up-to-homotopy. But this is clear since all $C_{(-)}^\bullet(\bullet)$ are connected by natural zig-zags of natural weak equivalences, since they are all cosimplicial resolutions of the same functor $[1](\bullet)$. \square

We can therefore, without any worry, denote $\Sigma_{(-)}$ simply by Σ . Then we define the following model structure:

Definition 3.1.3. If $\mathcal{M} = (\mathcal{C}, \mathcal{S})$ is a Cartesian presentation, we define the model category $\widehat{\Theta[\mathcal{C}]_{\mathcal{M}}}$ to be the left-Bousfield localization of $\widehat{\Theta[\mathcal{C}]_{\text{hJoyal}}}$ at the set $\Sigma(\mathcal{S})$, where we call the fibrant objects \mathcal{M} -enriched quasicategories or simply \mathcal{M} -quasicategories.

Proposition 3.1.4. *Let \mathcal{B} denote the set of simplicial boundary inclusions. Then a formal \mathcal{C} -quasicategory is an \mathcal{M} -quasicategory if and only if it has the right lifting property with respect to*

$$\Sigma(\mathcal{B} \times^\perp \mathcal{S}).$$

Proof. Let $E : \widehat{\Delta} \rightarrow \widehat{\Theta[\mathcal{C}]}$ be the left Kan extension of the cosimplicial object E^\bullet . We know by the construction of the left-Bousfield localization of Cisinski model categories that a formal \mathcal{C} -quasicategory

X is $\Sigma(\mathcal{S})$ -local if and only if it has the right-lifting property with respect to $E(\mathcal{B}) \times^\perp \Sigma(\mathcal{S})$, but this occurs only when for every $s : A \rightarrow B$ in \mathcal{S} , the map $X^{\Sigma(f)} : X^{\Sigma(B)} \rightarrow X^{\Sigma(A)}$ has the right lifting property with respect to $E(\mathcal{B})$.

By we claim that by adjunction, this happens if and only if $\mathcal{G}(X^{\Sigma(f)})$ is a trivial fibration, where \mathcal{G} is the functor sending a formal \mathcal{C} -quasicategory X to the Kan core of the underlying quasicategory $\mathcal{N}(X)$. To see this, notice that since all of the maps $f \in \mathcal{S}$ are monic, and since Σ preserves cofibrations, it follows that the maps $X^{\Sigma(f)}$ are all horizontal Joyal fibrations because the horizontal Joyal model structure is Cartesian. Then, since \mathcal{N} sends horizontal Joyal fibrations between formal \mathcal{C} -quasicategories to Joyal fibrations of quasicategories, and since the Kan core of a Joyal fibration between quasicategories is a Kan fibration of Kan complexes, it suffices to show that the Kan fibration $\mathcal{G}(X^f)$ is a trivial fibration as we claimed.

Notice that $\mathcal{G}(X^\bullet)$ is exactly the simplicial mapping space $\mathrm{hMap}^\Delta(\bullet, X)$. For any two vertices x, y of X , it suffices therefore to show that $\mathrm{hMap}_{*,*}^\Delta(\Sigma(f), X_{x,y})$ is a trivial fibration, since a Kan fibration is a trivial fibration if and only if it has contractible fibres. But $\mathrm{hMap}_{*,*}^\Delta(\Sigma(f), X_{x,y})$ is exactly $\mathrm{hMap}^\Delta(f, \mathrm{Map}_X(x, y))$, which is a trivial fibration if and only if $\mathrm{Map}_X(x, y)$ has the right lifting property with respect to $b \times^\perp f$ where b is a simplicial boundary inclusion, which proves the proposition. \square

Corollary 3.1.5. *A formal \mathcal{C} -quasicategory X is an \mathcal{M} -quasicategory if and only if $\mathrm{Map}_X(x, y)$ is \mathcal{S} -local for all pairs of vertices x, y in X .*

Theorem 3.1.6. *For any Cartesian presentation $\mathcal{M} = (\mathcal{C}, \mathcal{S})$, the model category $\widehat{\Theta[\mathcal{C}]}_{\mathcal{M}}$ is Cartesian-closed.*

Proof. This is exactly [Rez10, Proposition 8.5]. \square

In what follows, let Σ be Σ_R .

Proposition 3.1.7. *The pair $\widehat{\Theta[\mathcal{C}]}_{\mathcal{M}} \xrightleftharpoons[\mathfrak{N}]{\mathfrak{C}} \mathbf{Cat}_{\mathrm{Psh}_\Delta(\mathcal{C})_{\mathcal{S}}}$ is a Quillen pair.*

Proof. It suffices to show that \mathfrak{N} preserves fibrant objects by the properties of the left-Bousfield localization. Since the coherent nerve of any fibrant $\mathrm{Psh}_\Delta(\mathcal{C})$ -enriched category \mathcal{D} is already a formal \mathcal{C} -quasicategory, it suffices to show that $\mathfrak{N}\mathcal{D}$ has the right-lifting property with respect to $\Sigma(\mathcal{B} \times^\perp \mathcal{S})$. This will be true so long as the maps $\mathfrak{C}(\Sigma(\mathcal{B} \times^\perp \mathcal{S}))$ are all weak equivalences. To see this, let $\mathbf{2}(A)$ for any simplicial presheaf A on \mathcal{C} denote the $\mathrm{Psh}_\Delta(\mathcal{C})$ -enriched category whose objects are $\{0, 1\}$ and where

$$\mathbf{2}(A)(x, y) = \begin{cases} * & \text{if } x = y \\ A & \text{if } x < y \\ \emptyset & \text{otherwise} \end{cases}.$$

For all $n \geq 0$ and $c \in \mathcal{C}$, there is a natural weak equivalence

$$\mathfrak{C}(\Sigma(\Delta^n \times c))(\alpha, \omega) \cong Q^n \times c \xrightarrow{\sim} \Delta^n \times c \cong \mathbf{2}(c \times \Delta^n)(0, 1).$$

Following [Lur09, Proposition 2.2.2.7], We define a realization

$$|\bullet|_Q : \mathrm{Psh}_\Delta(\mathcal{C}) \rightarrow \mathrm{Psh}_\Delta(\mathcal{C})$$

by left Kan extension of the functor $\Delta^n \times c \mapsto Q^n \times c$ along the Yoneda embedding. Let \mathcal{A} denote the class of simplicial presheaves A on \mathcal{C} such that the map

$$|A|_Q \rightarrow A$$

is an injective equivalence. This class is closed under filtered colimits, since injective weak equivalences are closed under filtered colimits, so it suffices to consider the case where A has finitely many nondegenerate representable cells $[n] \times c$. Since Δ and \mathcal{C} are regular Cartesian Reedy, so is their product by [Cis06, 8.2.7], and the boundary of a representable cell is given by the formula

$$\partial(\Delta^n \times c) = \partial\Delta^n \times c \cup \Delta^n \times \partial c.$$

We work by induction on Reedy dimension and number of cells. If $A = \emptyset$, we are done, since the map in question is the identity. Otherwise, suppose

$$A = A' \coprod_{\partial(\Delta^n \times c)} \Delta^n \times c.$$

This is a homotopy pushout since $\partial(\Delta^n \times c) \rightarrow \Delta^n \times c$ is an injective cofibration. Similarly,

$$|A|_Q = |A'| \coprod_{|\partial(\Delta^n \times c)|_Q} |\Delta^n \times c|$$

is also a homotopy-pushout since $|\bullet|_Q$ preserves monomorphisms. Then we see that the map

$$|\Delta^n \times c|_Q = Q^n \times c \rightarrow \Delta^n \times c$$

is already a weak equivalence since $Q^n \rightarrow \Delta^n$ is a weak equivalence and the injective model structure is Cartesian. The map

$$|\partial(\Delta^n \times c)|_Q \rightarrow \partial(\Delta^n \times c)$$

is a weak equivalence by the induction hypothesis, since the Reedy dimension of $\partial(\Delta^n \times c)$ is less than the dimension of $\Delta^n \times c$. Finally, we see that

$$|A'|_Q \rightarrow A'$$

is a weak equivalence since A' has one fewer nondegenerate cell than A and is therefore also covered in the induction hypothesis.

Therefore, the natural map

$$\mathfrak{C}(\Sigma(A)) \cong \mathbf{2}(|A|_Q) \xrightarrow{\sim} \mathbf{2}(A)$$

is a weak equivalence in $\mathbf{Cat}_{\mathbf{Psh}_\Delta(\mathcal{C})_{\text{inj}}}$ for all simplicial presheaves A on \mathcal{C} . From this, it follows that since $\mathbf{2}(b \times^\cdot f)$ is an \mathcal{M} -equivalence for any $f \in \mathcal{J}$, and since we have a natural equivalence of arrows

$$\mathfrak{C}(\Sigma(b \times^\cdot f)) \xrightarrow{\sim} \mathbf{2}(b \times^\cdot f),$$

then by 3-for-2, $\mathfrak{C}(\Sigma(b \times^\cdot f))$ is a weak equivalence, which proves left-Quillen functoriality. □

Theorem 3.1.8. *The Quillen pair $\widehat{\Theta}[\widehat{\mathcal{C}}]_{\mathcal{M}} \xrightleftharpoons[\mathfrak{N}]{\mathfrak{C}} \mathbf{Cat}_{\mathbf{Psh}_\Delta(\mathcal{C})_{\mathcal{J}}}$ is a Quillen equivalence.*

Proof. It suffices to show that \mathfrak{C} is homotopy-conservative, so let $f : X \rightarrow Y$ be a map in $\widehat{\Theta[\mathfrak{C}]}$ such that $\mathfrak{C}(f)$ is an equivalence in $\mathbf{Cat}_{\mathbf{Psh}_\Delta(\mathfrak{C})_{\mathcal{M}}}$. Using the same argument as in Section 2.7, we reduce to the case where $f : X \rightarrow Y$ is a map between \mathcal{M} -quasicategories. Since \mathcal{M} -quasicategories are also formal \mathfrak{C} -quasicategories, we can apply Proposition 2.6.4 to obtain a natural zig-zag of weak equivalences between $\mathrm{Map}_X(x, y)$ and $\mathfrak{C}(X)(x, y)$ for any pair of vertices x, y of X . By 3-for-2 and since

$$\mathfrak{C}(X)(x, y) \rightarrow \mathfrak{C}(Y)(fx, fy)$$

was assumed to be an \mathcal{M} -equivalence, we see that the map $\mathrm{Map}_X(x, y) \rightarrow \mathrm{Map}_Y(fx, fy)$ must also be an \mathcal{M} -equivalence. In fact, since both $\mathrm{Map}_X(x, y)$ and $\mathrm{Map}_Y(fx, fy)$ are local, this map is actually an equivalence for $\mathbf{Cat}_{\mathbf{Psh}_\Delta(\mathfrak{C})_{\mathrm{inj}}}$. The argument showing that f is bijective on iso-components is the same as in the proof of Proposition 2.7.3 by passing to the underlying quasicategory. Therefore, it follows that f is a horizontal Dwyer-Kan equivalence, which concludes the proof. \square

3.2 The Yoneda embedding and Yoneda's lemma

We need the following easy lemma:

Lemma 3.2.1. *There is a natural isomorphism $\mathfrak{C}(X^{\mathrm{op}}) \cong \mathfrak{C}(X)^{\mathrm{op}}$.*

Proof. It suffices to check on representables, and this is left as an easy exercise to the reader. \square

Before we give a construction of the Yoneda embedding and a proof of Yoneda's lemma for \mathcal{M} -quasicategories, we need two lemmas from [Lur09]. We fix a Cartesian presentation $\mathcal{M} = (\mathfrak{C}, \mathcal{S})$ for the remainder of this section.

Proposition 3.2.2. [Lur09, 4.2.4.4] *Let $X \in \widehat{\Theta[\mathfrak{C}]}$ be a cellular \mathfrak{C} -set, \mathcal{D} a small $\mathbf{Psh}_\Delta(\mathfrak{C})$ -enriched category, and let $\mathfrak{C}(X) \rightarrow \mathcal{D}$ be an equivalence of $\mathbf{Psh}_\Delta(\mathfrak{C})_{\mathcal{S}}$ -enriched categories. Suppose \mathbf{A} is a $\mathbf{Psh}_\Delta(\mathfrak{C})_{\mathcal{S}}$ -enriched model category, and let \mathcal{U} be \mathcal{D} -chunk (see [Lur09, A.3.4.9] for the definition). Then the induced map*

$$\mathfrak{N}((\mathcal{U}^{\mathcal{D}})^{\circ}) \rightarrow \mathfrak{N}(\mathcal{U}^{\circ})^X$$

is an equivalence of \mathcal{M} -quasicategories.

Proof. Although we have altered the statement slightly, the only result used in the proof in [Lur09] that doesn't hold for all excellent monoidal model categories is [Lur09, 2.2.5.1], but the analogue of this is exactly Theorem 3.1.8. \square

Proposition 3.2.3. [Lur09, 4.2.4.7] *Let \mathcal{J} be a fibrant $\mathbf{Psh}_\Delta(\mathfrak{C})_{\mathcal{S}}$ -enriched category, X an object of $\widehat{\Theta[\mathfrak{C}]}$, and $p : \mathfrak{N} \rightarrow X$ be any map. Then we can find the following:*

- A fibrant $\mathbf{Psh}_\Delta(\mathfrak{C})_{\mathcal{S}}$ -enriched category \mathcal{D} .
- An enriched functor $P : \mathcal{J} \rightarrow \mathcal{D}$.
- A map $j : X \rightarrow \mathfrak{N}(\mathcal{D})$ that is a weak equivalence in $\widehat{\Theta[\mathfrak{C}]}_{\mathcal{M}}$.
- An equivalence between $j \circ p$ and $\mathfrak{N}(P)$ as objects of the \mathcal{M} -quasicategory $\mathfrak{N}(\mathcal{D})^{\mathfrak{N}(\mathcal{J})}$.

Proof. No change to the proof of [Lur09, 4.2.4.7] is needed. \square

We begin by constructing the Yoneda embedding:

Definition 3.2.4. Let X be a \mathcal{C} -cellular set, and let $\Phi : \mathfrak{C}(X) \xrightarrow{\sim} \mathcal{D}$ be an \mathcal{M} -enriched fibrant replacement. Since \mathcal{D} is fibrant, the functor

$$\mathrm{Hom}_{\mathcal{D}} : \mathcal{D} \times \mathcal{D}^{\mathrm{op}} \rightarrow \mathrm{Psh}_{\Delta}(\mathcal{C})$$

factors through the full subcategory $\mathrm{Psh}_{\Delta}(\mathcal{C})^{\circ}$. By [Lur09, Corollary A.3.4.14], this gives rise up to homotopy to a universal map

$$J_{\mathcal{D}} : \mathcal{D} \rightarrow \left(\mathrm{Psh}_{\Delta}(\mathcal{C})^{\mathcal{D}^{\mathrm{op}}} \right)_{\mathrm{proj}}^{\circ}$$

that is fully faithful up to homotopy, since it is homotopic to the enriched Yoneda embedding. Then we have a map

$$\mathfrak{C}(X \times X^{\mathrm{op}}) \xrightarrow{\mathfrak{C}(p_1) \times \mathfrak{C}(p_2)} \mathfrak{C}(X) \times \mathfrak{C}(X)^{\mathrm{op}} \xrightarrow{\Phi \times \Phi^{\mathrm{op}}} \mathcal{D} \times \mathcal{D}^{\mathrm{op}} \xrightarrow{\mathrm{Hom}_{\mathcal{D}}, \mathrm{Psh}_{\Delta}(\mathcal{C})_{\mathcal{F}}^{\circ}}$$

which yields by adjunction

$$X \rightarrow \mathfrak{N}(\mathrm{Psh}_{\Delta}(\mathcal{C})_{\mathcal{F}}^{\circ})^{X^{\mathrm{op}}}.$$

We denote $\mathfrak{N}(\mathrm{Psh}_{\Delta}(\mathcal{C})_{\mathcal{F}}^{\circ})$ simply by \mathcal{M} , and finally, we obtain *the Yoneda embedding* for \mathcal{M} -quasicategories:

$$j : X \rightarrow \mathcal{M}^{X^{\mathrm{op}}}.$$

Let $\mathcal{P}(X)$ denote the large \mathcal{M} -quasicategory $\mathcal{M}^{X^{\mathrm{op}}}$, and let \mathcal{M}^+ denote the coherent nerve of the huge enriched category of not-necessarily-small \mathcal{S} -local injectively fibrant simplicial presheaves on \mathcal{C} .

Definition 3.2.5. We say that a functor $F : X^{\mathrm{op}} \rightarrow \mathcal{M}$ is *representable* if the object it classifies belongs to the essential image of the Yoneda embedding $j : X \rightarrow \mathcal{P}(X)$. If $x : * \rightarrow X$ is a vertex of X , we denote the associated representable functor by h_x .

Proposition 3.2.6 (Yoneda embedding). [Lur09, 5.1.3.1] *The Yoneda embedding is fully faithful.*

Proof. First, let $\Phi : \mathfrak{C}(X^{\mathrm{op}}) \xrightarrow{\sim} \mathcal{D}$ be a fibrant replacement. We have an equivalence by Proposition 3.2.2

$$j'' : \mathfrak{N} \left(\left((\mathrm{Psh}_{\Delta}(\mathcal{C})_{\mathcal{F}})^{\mathcal{D}} \right)_{\mathrm{proj}}^{\circ} \right) \xrightarrow{\sim} \mathcal{M}^{\mathfrak{N}(\mathcal{D})},$$

and since \mathcal{M} is fibrant and $\Psi : X^{\mathrm{op}} \rightarrow \mathfrak{N}\mathcal{D}$, the adjunct of Φ is an equivalence between cofibrant objects, it follows that the induced map $\mathcal{M}^{\mathfrak{N}\mathcal{D}} \rightarrow \mathcal{P}(X)$ is an equivalence between fibrant objects, since the model structure is Cartesian. Since X is cofibrant, it follows that there is a map $h : X \rightarrow \mathcal{M}^{\mathfrak{N}\mathcal{D}}$ such that the composite

$$X \xrightarrow{h} \mathcal{M}^{\mathfrak{N}\mathcal{D}} \xrightarrow{\mathcal{M}^{\Psi}} \mathcal{P}(X)$$

is equivalent to j , and again, since $\mathfrak{N} \left(\left((\mathrm{Psh}_{\Delta}(\mathcal{C})_{\mathcal{F}})^{\mathcal{D}} \right)_{\mathrm{proj}}^{\circ} \right)$ is fibrant and j'' is a weak equivalence between fibrant objects, we can find a map

$$j' : X \rightarrow \mathfrak{N} \left(\left((\mathrm{Psh}_{\Delta}(\mathcal{C})_{\mathcal{F}})^{\mathcal{D}} \right)_{\mathrm{proj}}^{\circ} \right)$$

such that the composite

$$X \xrightarrow{j'} \mathfrak{N} \left(\left((\mathrm{Psh}_{\Delta}(\mathcal{C})_{\mathcal{F}})^{\mathcal{D}} \right)_{\mathrm{proj}}^{\circ} \right) \xrightarrow{j''} \mathcal{M}^{\mathfrak{N}\mathcal{D}} \xrightarrow{\sim} \mathcal{P}(X)$$

is equivalent to j . It suffices to show that the map j' is fully faithful. Let

$$J : \mathfrak{C}(X) \rightarrow ((\mathrm{Psh}_\Delta(\mathcal{C})_{\mathcal{J}})^{\mathcal{D}}_{\mathrm{proj}})^{\circ}$$

be the adjunct of j' . Then we see that J is equivalent to the composite $J_{\mathcal{D}^{\mathrm{op}}} \circ \Phi^{\mathrm{op}}$, where $J_{\mathcal{D}^{\mathrm{op}}}$ is fully-faithful and Φ^{op} is an equivalence, which concludes the proof. \square

Proposition 3.2.7 (Yoneda's Lemma). [Lur09, 5.5.2.1] *Let X be a small \mathcal{C} -cellular set, and let $f : X^{\mathrm{op}} \rightarrow \mathcal{M}$ be an object of $\mathcal{P}(X)$. Then let $F : \mathcal{P}(X)^{\mathrm{op}} \rightarrow \mathcal{M}^+$ be the functor represented by f . Then the composite*

$$X^{\mathrm{op}} \xrightarrow{j_X^{\mathrm{op}}} \mathcal{P}(X)^{\mathrm{op}} \xrightarrow{F} \mathcal{M}^+$$

is equivalent to f .

Proof. By Proposition 3.2.3, we can choose a small fibrant \mathcal{M} -enriched category \mathcal{D} and an equivalence $\Phi : X^{\mathrm{op}} \rightarrow \mathfrak{N}(\mathcal{D})$ such that $f \sim \mathfrak{N}(f') \circ \Phi$ for some $f' : \mathcal{D} \rightarrow \mathrm{Psh}_\Delta(\mathcal{C})_{\mathcal{J}}^{\circ}$. Without loss of generality, we can assume that f' is a projectively cofibrant diagram. Using Proposition 3.2.2, we have an equivalence of \mathcal{M} -quasicategories

$$\Psi : \mathfrak{N}\left(\left((\mathrm{Psh}_\Delta(\mathcal{C})_{\mathcal{J}})^{\mathcal{D}}_{\mathrm{proj}}\right)^{\circ}\right) \xrightarrow{\sim} \mathcal{P}(X).$$

We observe that $F \circ \Psi^{\mathrm{op}}$ can be identified with the coherent nerve of the the map

$$G : \left(\left((\mathrm{Psh}_\Delta(\mathcal{C})_{\mathcal{J}})^{\mathcal{D}}_{\mathrm{proj}}\right)^{\circ}\right)^{\mathrm{op}} \rightarrow (\mathrm{Psh}_\Delta(\mathcal{C})_{\mathcal{J}}^+)^{\circ}$$

represented by f' . The Yoneda embedding factors through Ψ by the adjunct of the composite

$$j' : \mathfrak{C}(X) \xrightarrow{\Phi^{\mathrm{op}}} \mathcal{D}^{\mathrm{op}} \hookrightarrow ((\mathrm{Psh}_\Delta(\mathcal{C})_{\mathcal{J}})^{\mathcal{D}}_{\mathrm{proj}})^{\circ},$$

so it follows that $F \circ j^{\mathrm{op}}$ can be identified with the adjunct of

$$\mathfrak{C}(X)^{\mathrm{op}} \xrightarrow{(j')^{\mathrm{op}}} \left(\left((\mathrm{Psh}_\Delta(\mathcal{C})_{\mathcal{J}})^{\mathcal{D}}_{\mathrm{proj}}\right)^{\circ}\right)^{\mathrm{op}} \xrightarrow{G} (\mathrm{Psh}_\Delta(\mathcal{C})_{\mathcal{J}}^+)^{\circ}.$$

This composite is equal to the f' we started with, so its coherent nerve is equivalent to f . \square

3.3 Weighted limits and colimits

We fix a Cartesian presentation $\mathcal{M} = (\mathcal{C}, \mathcal{J})$. Our presentation here follows the one given in [na18].

Definition 3.3.1. Let D be a small \mathcal{C} -cellular set, that is, an object of $\widehat{\Theta[\mathcal{C}]}$, and suppose we have a diagram $f : D \rightarrow \mathcal{M}$ and a *weight* $W : D \rightarrow \mathcal{M}$. If X is an \mathcal{M} -quasicategory and $f : D \rightarrow X$ is a diagram, we define the map $h_f : X^{\mathrm{op}} \rightarrow \mathcal{M}^D$ to be the adjunct of the composite

$$D \rightarrow X \xrightarrow{j} \mathcal{M}^{X^{\mathrm{op}}},$$

and we define the *weight representation* $h^W : \mathcal{M}^D \rightarrow \mathcal{M}^+$ be the map co-representing W as an object of \mathcal{M}^D . Then if the composite $h^W \circ h_f : X^{\mathrm{op}} \rightarrow \mathcal{M}^+$ is representable, we call its representing object *the limit of f weighted by W* , denoted by $\lim^W f$.

Dually, coweights for colimits are diagrams $D^{\text{op}} \rightarrow \mathcal{M}$, and their coweighting functors are their associated corepresentable functors $\mathcal{M}^{D^{\text{op}}} \rightarrow \mathcal{M}^+$. Given a diagram $f : D \rightarrow X$, the *colimit of f weighted by W* is defined to be an object of X that corepresents the weighted limit of f^{op} , which is denoted by $\text{colim}^W f$.

Proposition 3.3.2. *The large \mathcal{M} -quasicategory \mathcal{M} has all small weighted limits and colimits.*

Proof. Given $f : D \rightarrow \mathcal{M}$ and a weight $W : D \rightarrow \mathcal{M}$, it is straightforward to see that a representing object for the weighted limit is exactly

$$\text{Map}_{\mathcal{M}^D}(W, f),$$

by unwinding the definitions. □

For now, we don't have much to say for this particular application. If other definitions are proposed for weighted limits and colimits, they should be equivalent to this one.

3.4 Examples

The only examples we really care about are the cases where $\mathcal{C} = \Theta_n$ for $0 \leq n \leq \omega$ and where \mathcal{S} is the set of generating anodynes for the model structure on weak n -categories. We invite the reader to consider other applications. We expect that a simple application would be to consider the left-Bousfield localization of spaces at homology equivalences, but we aren't certain if this is a Cartesian model structure.

Also note that our definitions of weighted limits and colimits do not work for computing lax and oplax weighted limits and colimits in weak ω -categories (taking \mathcal{C} to be $\Theta = \Theta_\omega$ and \mathcal{S} to be the union of all of the generating weak equivalences for the Rezk model structure). The problem is that the lax Gray tensor product has not yet been shown to be homotopy-invariant (in particular, a left-Quillen bifunctor), so the function complexes of lax and oplax natural transformations are themselves not yet known to be homotopy invariant. This is why we encourage people in the future to refer to weighted limits and colimits in these cases as *weighted pseudolimits* and *weighted pseudocolimits* respectively.

References

- [Ber07] C. Berger, *Iterated wreath product of the simplex category and iterated loop spaces*, Adv. Math. **213** (2007), no. 1, 230–270. MR2331244 (2008f:55010)
- [BR11] J. E. Bergner and C. Rezk, *Reedy categories and the Θ -Construction*, Mathematische Zeitschrift **1-2** (2011).
- [Cis06] D.-C. Cisinski, *Les préfaisceaux comme modèles des types d'homotopie*, Astérisque, vol. 308, Soc. Math. France, 2006.
- [DS11a] D. Dugger and D. Spivak, *Rigidification of quasi-categories*, Algebr. Geom. Topol. **11** (2011), no. 1, 225–261. MR2764042
- [DS11b] ———, *Mapping spaces in quasi-categories*, Algebr. Geom. Topol. **11** (2011), no. 1, 263–325. MR2764043
- [Lur09] Jacob Lurie, *Higher Topos Theory*, Princeton University Press, 2009.
- [JT07] A. Joyal and M. Tierney, *Quasi-categories vs Segal spaces*, Contemporary Mathematics **431** (2007), 277–326.
- [na18] nLab authors, *Weighted Limit* (2018), <http://ncatlab.org/nlab/show/weighted%20limit>. Revision 29.
- [Our10] D. Oury, *Duality for Joyal's category Θ and homotopy concepts for Θ_2 -sets*, Macquarie University, 2010.
- [Rez10] C. Rezk, *A Cartesian presentation of weak n -categories*, Geom. Topol. **14** (2010), no. 1, 521–571, DOI 10.2140/gt.2010.14.521. MR2578310
- [Ste] D. Stevenson, *Model Structures for Correspondences and Bifibrations*, 41 pp., available at [arXiv:1807.08226v1](https://arxiv.org/abs/1807.08226).