

Affine descents and the Steinberg torus

T. Kyle Petersen
University of Michigan

(joint with Kevin Dilks and John Stembridge, [arXiv:0709.4291](https://arxiv.org/abs/0709.4291))

FPSAC 08

Affine descents and the Steinberg torus

Eulerian polynomials

Coxeter complexes

Affine Eulerian polynomials

The Steinberg torus

Eulerian polynomials

The Eulerian polynomials, $A_n(t) = \sum_{k=0}^n a_{n,k} t^k$ (classical):

$$A_1(t) = 1 + t$$

$$A_2(t) = 1 + 4t + t^2$$

$$A_3(t) = 1 + 11t + 11t^2 + t^3$$

$$A_4(t) = 1 + 26t + 66t^2 + 26t^3 + t^4$$

$$\vdots$$

Eulerian polynomials

The Eulerian polynomials, $A_n(t) = \sum_{k=0}^n a_{n,k} t^k$ (classical):

$$A_1(t) = 1 + t$$

$$A_2(t) = 1 + 4t + t^2$$

$$A_3(t) = 1 + 11t + 11t^2 + t^3$$

$$A_4(t) = 1 + 26t + 66t^2 + 26t^3 + t^4$$

$$\vdots$$

- symmetric, unimodal coefficients

Eulerian polynomials

The Eulerian polynomials, $A_n(t) = \sum_{k=0}^n a_{n,k} t^k$ (classical):

$$A_1(t) = 1 + t$$

$$A_2(t) = 1 + 4t + t^2$$

$$A_3(t) = 1 + 11t + 11t^2 + t^3$$

$$A_4(t) = 1 + 26t + 66t^2 + 26t^3 + t^4$$

$$\vdots$$

- ▶ symmetric, unimodal coefficients
- ▶ real-rooted (Harper '67)

Combinatorial interpretation

Let $d(w) := \#\{i : w_i > w_{i+1}\}$. Then,

$$A_n(t) = \sum_{w \in S_{n+1}} t^{d(w)}$$

Combinatorial interpretation

Let $d(w) := \#\{i : w_i > w_{i+1}\}$. Then,

$$A_n(t) = \sum_{w \in S_{n+1}} t^{d(w)}$$

w	$d(w)$	$t^{d(w)}$
123		
132		
213		
231		
312		
321		

Combinatorial interpretation

Let $d(w) := \#\{i : w_i > w_{i+1}\}$. Then,

$$A_n(t) = \sum_{w \in S_{n+1}} t^{d(w)}$$

w	$d(w)$	$t^{d(w)}$
123	0	1
132		
213		
231		
312		
321		

Combinatorial interpretation

Let $d(w) := \#\{i : w_i > w_{i+1}\}$. Then,

$$A_n(t) = \sum_{w \in S_{n+1}} t^{d(w)}$$

w	$d(w)$	$t^{d(w)}$
123	0	1
132	1	t
213		
231		
312		
321		

Combinatorial interpretation

Let $d(w) := \#\{i : w_i > w_{i+1}\}$. Then,

$$A_n(t) = \sum_{w \in S_{n+1}} t^{d(w)}$$

w	$d(w)$	$t^{d(w)}$
123	0	1
132	1	t
213	1	t
231		
312		
321		

Combinatorial interpretation

Let $d(w) := \#\{i : w_i > w_{i+1}\}$. Then,

$$A_n(t) = \sum_{w \in S_{n+1}} t^{d(w)}$$

w	$d(w)$	$t^{d(w)}$
123	0	1
132	1	t
213	1	t
231	1	t
312		
321		

Combinatorial interpretation

Let $d(w) := \#\{i : w_i > w_{i+1}\}$. Then,

$$A_n(t) = \sum_{w \in S_{n+1}} t^{d(w)}$$

w	$d(w)$	$t^{d(w)}$
123	0	1
132	1	t
213	1	t
231	1	t
312	1	t
321		

Combinatorial interpretation

Let $d(w) := \#\{i : w_i > w_{i+1}\}$. Then,

$$A_n(t) = \sum_{w \in S_{n+1}} t^{d(w)}$$

w	$d(w)$	$t^{d(w)}$
123	0	1
132	1	t
213	1	t
231	1	t
312	1	t
321	2	t^2

Combinatorial interpretation

Let $d(w) := \#\{i : w_i > w_{i+1}\}$. Then,

$$A_n(t) = \sum_{w \in S_{n+1}} t^{d(w)}$$

w	$d(w)$	$t^{d(w)}$
123	0	1
132	1	t
213	1	t
231	1	t
312	1	t
321	2	t^2

$$A_2(t) = 1 + 4t + t^2$$

A generalization

The notion of descent makes sense in any Coxeter system (W, S) (and simple roots Δ):

$$\begin{aligned} d(w) &:= \#\{s \in S : \ell(ws) < \ell(w)\} \\ &= \#\{\alpha \in \Delta : w(\alpha) < 0\} \end{aligned}$$

Define the W -Eulerian polynomial:

$$W(t) := \sum_{w \in W} t^{d(w)}$$

A generalization

The notion of descent makes sense in any Coxeter system (W, S) (and simple roots Δ):

$$\begin{aligned} d(w) &:= \#\{s \in S : \ell(ws) < \ell(w)\} \\ &= \#\{\alpha \in \Delta : w(\alpha) < 0\} \end{aligned}$$

Define the W -Eulerian polynomial:

$$W(t) := \sum_{w \in W} t^{d(w)}$$

- The W -Eulerian polynomials are symmetric, unimodal (Brenti '94) (in fact, γ -nonnegative)

A generalization

The notion of descent makes sense in any Coxeter system (W, S) (and simple roots Δ):

$$\begin{aligned} d(w) &:= \#\{s \in S : \ell(ws) < \ell(w)\} \\ &= \#\{\alpha \in \Delta : w(\alpha) < 0\} \end{aligned}$$

Define the W -Eulerian polynomial:

$$W(t) := \sum_{w \in W} t^{d(w)}$$

- ▶ The W -Eulerian polynomials are symmetric, unimodal (Brenti '94) (in fact, γ -nonnegative)
- ▶ Brenti has conjectured real-rootedness as well (D_n remains unproved)

Affine descents and the Steinberg torus

Eulerian polynomials

Coxeter complexes

Affine Eulerian polynomials

The Steinberg torus

The f - and h -polynomials

Let Σ be a finite set of simplices, $f_k(\Sigma)$ = number of faces of dimension $k - 1$


$$f(\Sigma; t) := \sum_{k=0}^n f_k(\Sigma) t^k$$

(f_0, f_1, \dots, f_n) is the f -vector


$$h(\Sigma; t) := (1 - t)^n f(\Sigma; t/(1 - t)) = \sum_{k=0}^n h_k(\Sigma) t^k$$

(h_0, h_1, \dots, h_n) is the h -vector

The f - and h -polynomials




The f - and h -polynomials



► $f_0 = 1$


The f - and h -polynomials



► $f_0 = 1$


► $f_1 = 6$

The f - and h -polynomials



- ▶ $f_0 = 1$
- ▶ $f_1 = 6$
- ▶ $f_2 = 6$

The f - and h -polynomials




► $f_0 = 1$

► $f_1 = 6$

► $f_2 = 6$

$$f(\Sigma; t) = 1 + 6t + 6t^2$$

The f - and h -polynomials



► $f_0 = 1$


► $f_1 = 6$

► $f_2 = 6$

$$f(\Sigma; t) = 1 + 6t + 6t^2$$

$$h(\Sigma; t) = 1 + 4t + t^2$$

The f - and h -polynomials



► $f_0 = 1$


► $f_1 = 6$

► $f_2 = 6$

$$f(\Sigma; t) = 1 + 6t + 6t^2$$


$$h(\Sigma; t) = 1 + 4t + t^2 = A_2(t) \text{ (hmm...)}$$

The Coxeter complex



For a Coxeter system (W, S) , the reflecting hyperplanes partition the ambient vector space


The Coxeter complex



For a Coxeter system (W, S) , the reflecting hyperplanes partition the ambient vector space

By intersecting the hyperplanes with the unit sphere we achieve a topological realization of the *Coxeter complex*, $\Sigma(W)$

The W -Eulerian polynomial



Theorem (Björner '84, Brenti '94)

For any finite Coxeter group W ,

$$h(\Sigma(W); t) = \sum_{w \in W} t^{d(w)} = W(t)$$

Affine descents and the Steinberg torus

Eulerian polynomials

Coxeter complexes

Affine Eulerian polynomials

The Steinberg torus

Affine descents

If W is crystallographic, it has a unique lowest root $\alpha_0 = -\tilde{\alpha}$

Let s_0 be the corresponding reflection, $\Delta_0 = \Delta \cup \{\alpha_0\}$, and

$$\begin{aligned}\tilde{d}(w) &:= d(w) + \chi(\ell(ws_0) > \ell(w)) \\ &= \#\{\alpha \in \Delta_0 : w(\alpha) < 0\}\end{aligned}$$

Affine descents

If W is crystallographic, it has a unique lowest root $\alpha_0 = -\tilde{\alpha}$

Let s_0 be the corresponding reflection, $\Delta_0 = \Delta \cup \{\alpha_0\}$, and

$$\begin{aligned}\tilde{d}(w) &:= d(w) + \chi(\ell(ws_0) > \ell(w)) \\ &= \#\{\alpha \in \Delta_0 : w(\alpha) < 0\}\end{aligned}$$

Definition (Dilks-Petersen-Stembridge '07)

The affine W -Eulerian polynomial is

$$\widetilde{W}(t) := \sum_{w \in W} t^{\tilde{d}(w)}$$

Affine descents

$$\widetilde{W}(t) := \sum_{w \in W} t^{\widetilde{d}(w)}$$

Results of D-P-S, $\widetilde{W}(t)$ is:

Affine descents

$$\widetilde{W}(t) := \sum_{w \in W} t^{\widetilde{d}(w)}$$

Results of D-P-S, $\widetilde{W}(t)$ is:

- ▶ γ -nonnegative (\Rightarrow symmetric, unimodal)

Affine descents

$$\widetilde{W}(t) := \sum_{w \in W} t^{\widetilde{d}(w)}$$

Results of D-P-S, $\widetilde{W}(t)$ is:

- ▶ γ -nonnegative (\Rightarrow symmetric, unimodal)
- ▶ conjecturally real-rooted (\widetilde{A}_n , \widetilde{C}_n , exceptional groups are proved; \widetilde{B}_n and \widetilde{D}_n are verified for $n \leq 100$)

Affine descents

In type A, $\tilde{d}(w)$ is the number of cyclic descents

Affine descents

In type A, $\tilde{d}(w)$ is the number of cyclic descents

w	$\tilde{d}(w)$	$t^{\tilde{d}(w)}$
123		
132		
213		
231		
312		
321		

Affine descents

In type A, $\tilde{d}(w)$ is the number of cyclic descents

w	$\tilde{d}(w)$	$t^{\tilde{d}(w)}$
123	1	t
132		
213		
231		
312		
321		

Affine descents

In type A, $\tilde{d}(w)$ is the number of cyclic descents

w	$\tilde{d}(w)$	$t^{\tilde{d}(w)}$
123	1	t
132	2	t^2
213		
231		
312		
321		

Affine descents

In type A, $\tilde{d}(w)$ is the number of cyclic descents

w	$\tilde{d}(w)$	$t^{\tilde{d}(w)}$
123	1	t
132	2	t^2
213	2	t^2
231		
312		
321		

Affine descents

In type A, $\tilde{d}(w)$ is the number of cyclic descents

w	$\tilde{d}(w)$	$t^{\tilde{d}(w)}$
123	1	t
132	2	t^2
213	2	t^2
231	1	t
312		
321		

Affine descents

In type A, $\tilde{d}(w)$ is the number of cyclic descents

w	$\tilde{d}(w)$	$t^{\tilde{d}(w)}$
123	1	t
132	2	t^2
213	2	t^2
231	1	t
312	1	t
321		

Affine descents

In type A, $\tilde{d}(w)$ is the number of cyclic descents

w	$\tilde{d}(w)$	$t^{\tilde{d}(w)}$
123	1	t
132	2	t^2
213	2	t^2
231	1	t
312	1	t
321	2	t^2

Affine descents

In type A, $\tilde{d}(w)$ is the number of cyclic descents

w	$\tilde{d}(w)$	$t^{\tilde{d}(w)}$
123	1	t
132	2	t^2
213	2	t^2
231	1	t
312	1	t
321	2	t^2

$$\tilde{A}_2(t) = 3t + 3t^2$$

Jeopardy!

The answer is ...

Jeopardy!

The answer is ...

It is the topological construction whose h -vector is encoded by the affine Eulerian polynomial.

Jeopardy!

The answer is ...

It is the topological construction whose h -vector is encoded by the affine Eulerian polynomial.

What is the Steinberg torus? (Correct!)

Affine descents and the Steinberg torus


Eulerian polynomials

Coxeter complexes

Affine Eulerian polynomials

The Steinberg torus

Affine Coxeter complexes



The affine Weyl group \tilde{W} is generated by S along with the reflection through $H_{\tilde{\alpha},1} := \{\lambda : \langle \tilde{\alpha}, \lambda \rangle = 1\}$, drawing all hyperplanes gives $\Sigma(\tilde{W})$ (... if W is irreducible...)

The Steinberg torus

Standard fact: the coroot lattice is a translation subgroup;

$$\widetilde{W} \cong W \ltimes \mathbb{Z}\Phi^\vee$$

Thus \widetilde{W} -action on V restricts to a W -action on the torus $V/\mathbb{Z}\Phi^\vee$
(Steinberg '68 - Bott's formula for Poincaré series of \widetilde{W})

Definition (D-P-S)

The **Steinberg torus** of \widetilde{W} is

$$\Sigma_T(\widetilde{W}) := \Sigma(\widetilde{W})/\mathbb{Z}\Phi^\vee$$

The Steinberg torus

Standard fact: the coroot lattice is a translation subgroup;

$$\widetilde{W} \cong W \ltimes \mathbb{Z}\Phi^\vee$$

Thus \widetilde{W} -action on V restricts to a W -action on the torus $V/\mathbb{Z}\Phi^\vee$
(Steinberg '68 - Bott's formula for Poincaré series of \widetilde{W})

Definition (D-P-S)

The **Steinberg torus** of \widetilde{W} is

$$\Sigma_T(\widetilde{W}) := \Sigma(\widetilde{W})/\mathbb{Z}\Phi^\vee$$

- ▶ $\Sigma_T(\widetilde{W})$ is a finite complex (boolean complex, or simplicial poset)

The Steinberg torus

Standard fact: the coroot lattice is a translation subgroup;

$$\widetilde{W} \cong W \ltimes \mathbb{Z}\Phi^\vee$$

Thus \widetilde{W} -action on V restricts to a W -action on the torus $V/\mathbb{Z}\Phi^\vee$
(Steinberg '68 - Bott's formula for Poincaré series of \widetilde{W})


Definition (D-P-S)

The **Steinberg torus** of \widetilde{W} is

$$\Sigma_T(\widetilde{W}) := \Sigma(\widetilde{W})/\mathbb{Z}\Phi^\vee$$


- ▶ $\Sigma_T(\widetilde{W})$ is a finite complex (boolean complex, or simplicial poset)
- ▶ maximal cells of in bijection with elements of W

The Steinberg torus



$$\Sigma_T(\tilde{A}_2) := \Sigma(\tilde{A}_2)/\mathbb{Z}\{\alpha_1^\vee, \alpha_2^\vee\}$$

The Steinberg torus




Equivalently, observe that exactly one vertex of every alcove is in $\mathbb{Z}\Phi^\vee$, so we translate can translate to the origin

The Steinberg torus

The union of (closures of) the alcoves neighboring the origin is a convex, W -invariant simplicial polytope:


$$P_\Phi := \{\lambda \in V : -1 \leq \langle \lambda, \beta \rangle \leq 1 \text{ for all } \beta \in \Phi\}$$

We obtain the Steinberg torus by identifying opposite faces of P_Φ




The Steinberg torus

$\Sigma_T(\tilde{A}_2)$:




The Steinberg torus



- $f_0 = 0$ (...if we ignore the empty face, things work out nicer...)


The Steinberg torus



- ▶ $f_0 = 0$ (...if we ignore the empty face, things work out nicer...)
- ▶ $f_1 = 3$

The Steinberg torus


$\Sigma_T(\tilde{A}_2)$:



- ▶ $f_0 = 0$ (... if we ignore the empty face, things work out nicer...)
- ▶ $f_1 = 3$
- ▶ $f_2 = 9$

The Steinberg torus


$\Sigma_T(\tilde{A}_2)$:



- ▶ $f_0 = 0$ (... if we ignore the empty face, things work out nicer...)
- ▶ $f_1 = 3$
- ▶ $f_2 = 9$
- ▶ $f_3 = 6$

The Steinberg torus


$\Sigma_T(\tilde{A}_2)$:



$$f(\Sigma_T; t) = 3t + 9t^2 + 6t^3$$

The Steinberg torus

$\Sigma_T(\tilde{A}_2)$:




$$f(\Sigma_T; t) = 3t + 9t^2 + 6t^3$$

$$h(\Sigma_T; t) = 3t + 3t^2$$

The Steinberg torus

$\Sigma_T(\tilde{A}_2)$:



$$f(\Sigma_T; t) = 3t + 9t^2 + 6t^3$$

$$h(\Sigma_T; t) = 3t + 3t^2 = \tilde{A}_2(t)$$

The Steinberg torus

Theorem (D-P-S)

For any irreducible affine Weyl group $\widetilde{W} = W \ltimes \mathbb{Z}\Phi^\vee$,

$$h(\Sigma_T(\widetilde{W}); t) = \sum_{w \in W} t^{\widetilde{d}(w)} = \widetilde{W}(t)$$

The Steinberg torus

Theorem (D-P-S)

For any irreducible affine Weyl group $\widetilde{W} = W \ltimes \mathbb{Z}\Phi^\vee$,

$$h(\Sigma_T(\widetilde{W}); t) = \sum_{w \in \widetilde{W}} t^{\widetilde{d}(w)} = \widetilde{W}(t)$$

Dehn-Sommerville relations for the torus now imply symmetry of $\widetilde{W}(t)$

Next

- Real-rootedness: The remaining cases, $D_n, \tilde{B}_n, \tilde{D}_n$ are all γ -nonnegative (a necessary condition for real roots in this situation)

Next

- Real-rootedness: The remaining cases, $D_n, \tilde{B}_n, \tilde{D}_n$ are all γ -nonnegative (a necessary condition for real roots in this situation)

Each case boils down to type A peak combinatorics:

$$w_{i-1} < w_i > w_{i+1}$$

Next

- ▶ Real-rootedness: The remaining cases, $D_n, \tilde{B}_n, \tilde{D}_n$ are all γ -nonnegative (a necessary condition for real roots in this situation)

Each case boils down to type A peak combinatorics:

$$w_{i-1} < w_i > w_{i+1}$$

- ▶ What does γ -nonnegativity mean topologically? (related to Gal's conjecture in spherical case?)

Next

- ▶ Real-rootedness: The remaining cases, $D_n, \tilde{B}_n, \tilde{D}_n$ are all γ -nonnegative (a necessary condition for real roots in this situation)

Each case boils down to type A peak combinatorics:

$$w_{i-1} < w_i > w_{i+1}$$

- ▶ What does γ -nonnegativity mean topologically? (related to Gal's conjecture in spherical case?)
- ▶ General topological reasons to expect $h_i \geq 0$ here? unimodality? (Novik, Swartz)

Next

- ▶ Real-rootedness: The remaining cases, $D_n, \tilde{B}_n, \tilde{D}_n$ are all γ -nonnegative (a necessary condition for real roots in this situation)

Each case boils down to type A peak combinatorics:

$$w_{i-1} < w_i > w_{i+1}$$

- ▶ What does γ -nonnegativity mean topologically? (related to Gal's conjecture in spherical case?)
- ▶ General topological reasons to expect $h_i \geq 0$ here? unimodality? (Novik, Swartz)
 g -theorem for boundary complex of a torus?

Next

- ▶ Real-rootedness: The remaining cases, $D_n, \tilde{B}_n, \tilde{D}_n$ are all γ -nonnegative (a necessary condition for real roots in this situation)

Each case boils down to type A peak combinatorics:

$$w_{i-1} < w_i > w_{i+1}$$

- ▶ What does γ -nonnegativity mean topologically? (related to Gal's conjecture in spherical case?)
- ▶ General topological reasons to expect $h_i \geq 0$ here?
unimodality? (Novik, Swartz)
 g -theorem for boundary complex of a torus?
details: this is a *boolean* complex (not a simplicial complex)

Next

- ▶ Real-rootedness: The remaining cases, $D_n, \tilde{B}_n, \tilde{D}_n$ are all γ -nonnegative (a necessary condition for real roots in this situation)


Each case boils down to type A peak combinatorics:

$$w_{i-1} < w_i > w_{i+1}$$

- ▶ What does γ -nonnegativity mean topologically? (related to Gal's conjecture in spherical case?)
- ▶ General topological reasons to expect $h_i \geq 0$ here?
unimodality? (Novik, Swartz)
 g -theorem for boundary complex of a torus?
details: this is a *boolean* complex (not a simplicial complex)
we ignore the empty face (compare with reduced/unreduced homology)

Questions?

Art gallery:



<http://www.math.lsa.umich.edu/~tkpeters/steinberg>