




Departamento de
Economía
Universidad de Concepción

The Impact of Environmental Variability on Fishers' Harvest Decisions in Chile

Using a Multi-Species Approach

Felipe J. Quezada-Escalona
Encuentro EfD Chile 2025

- Marine resource distribution and abundance is changing due to climate variability, with heterogenous spatial effects (Poloczanska, Brown, Sydeman et al., 2013; Sumaila, Cheung, Lam et al., 2011).
- Harvest levels would be affected (Quezada, Tommasi, Frawley et al., 2023),
 - Prices and value of catches, fishing costs, fishers' incomes, among others (Sumaila, Cheung, Lam et al., 2011)



Source: Cheung, Lam, Sarmiento et al. (2010)

Research question

How will fishing decisions, aggregate catch levels, and the price of marine resources be affected under different climatic scenarios in the multispecies small pelagic fishery (SPF) in Chile?

- How do fishers **substitute between species**?
- Contribute to the limited local literature on multi-species economic modeling and climate variability in Chile (e.g., [Peña-Torres, Dresdner, and Vasquez \(2017\)](#))
- Understand fishers' adaptive capacity helps to inform climate-resilient fisheries policies
- We will focus in **climate variability** to estimate short-run responses.
 - *i.e.*, climate change effect without adaptation ([Auffhammer, 2018](#))
 - Upper-bound estimates

Chile's Small Pelagic Fishery

Some facts

- Mainly composed by anchoveta, Jack mackerel, Sardine
- ~94% of national catch ([SUBPESCA, 2020](#))
- Mostly harvested with purse-seiners
- In the Central-South (CS) region (Valparaíso-Los Lagos) all three species play a major role.
- SPF have been used primarily for fishmeal and fish oil production ([Peña-Torres, Dresdner, and Vasquez, 2017](#)) (~85% of jack mackerel for reduction)

Regulations

- Quota (TAC); divided between the small-scale and industrial sector
- Industrial sector operates under ITQ
- RAE (*Regimen Artesanal de Extracción*) in some areas for small-scale fishery
 - Allocates regional quota to area or fishermen organization (i.e., catch shares)
- Anchoveta and sardine are regulated as a mixed-species fishery

Status of the stocks (CS)

- Anchoveta:
 - Collapsed until 2018,
 - Overexploited in 2019,
 - Since 2020, within MSY limits.
- Sardine:
 - Within MSY levels, except in 2021 and 2023 (overexploited)
- Jack mackerel:
 - Overexploited until 2018, then within MSY limits.

Methods and data

Based on [Kasperski \(2015\)](#):

- Econometrics models
 - Estimate stock dynamics
 - Estimate trip-level costs
 - Estimate annual trips
 - Estimate inverse demand
- Use parameters to:
 - Obtain the optimal **harvest** and **quota** conditional on climate variables.
 - Simulate the effects on harvest, prices and **profits**, and **species substitution** of different climate scenarios.

Harvest and biomass data


From IFOP (2012-2025):

- Trip-level data:
 - ID, departure and arrival times, vessel capacity, fleet and gear type, ports of departure and landing, haul timing and location, species, retained catch.
- Annual stock biomass by macro region and species (e.g. south-central Chile) -- (1999-2025)
- Monthly landings by port/species.
- Prices paid by processing plants (IFOP surveys; month-region)

Environmental covariates

For the period 2000-2025:

- Daily salinity, sea surface temperature, and current speed and direction (E.U. Copernicus Marine Service Information, 2025a)
- Hourly wind speed and direction at the surface (E.U. Copernicus Marine Service Information, 2025b)
- Chlorophyll-a concentration (E.U. Copernicus Marine Service Information, 2025c)



To be requested

- Average wage pay to crew member per hour (available?)
- Diesel cost.
- Permits by vessels
- Quota prices (auction or secondary markets, if available)
 - Captures forward-looking behavior and information ([Birkenbach, Lee, and Smith, 2024](#)).
 - Simplify the dynamic model to a static one ([Reimer, Abbott, and Haynie, 2022](#)).
- Quota by area/fishing organization for small-scale sector, and ITQ for industrial (by vessel?)

Data for projections

Bio-ORACLE

- Advantages:
 - Different climate scenarios
 - SST, salinity, currents and chlorophyll (4km resolution)
- Disadvantages
 - Only decadal (e.g., 2040–2050) projections
 - No winds; CMIP6 for winds? (~100 km).



Econometrics models

$$x_{i,y+1} + h_{iy} = \underbrace{(1 + r_i)x_{iy} + \eta_i x_{iy}^2}_{R_i(x_{iy})} + \underbrace{\sum_{j \neq i}^{n-1} \alpha_{ij} x_{iy} x_{jy}}_{I_i(x_y)} + \rho_i Env_{iy} + \varepsilon_{iy} \quad i = 1, \dots, n$$

where:

- x_{iy} is the fish stock by species $i = 1, \dots, n$ in year y , n is the total number of species,
- h_{iy} is the annual harvest of species i on year y ,
- r_i is the intrinsic growth rate of the resource i ,
- η_i is a density-dependent factor related to the carrying capacity,
- α_{ij} are the interaction parameters between species.
- Env_{iy} includes year–macroregion averages of **environmental covariates** (SST and chlorophyll-a).

The system of n growth equations can be estimated simultaneously using SUR

Model 2: Trip-Level Costs



$$C_{vg} = \sum_{i=1}^{2n+M+k} \alpha_{g, \mathbf{X}_i} \mathbf{X}_{ivg} + \frac{1}{2} \sum_{i=1}^{2n+M+k} \sum_{j=1}^{2n+M+k} \alpha_{g, \mathbf{X}_i \mathbf{X}_j} \mathbf{X}_{ivg} \mathbf{X}_{jvg}$$

where $C_{vg} = w z_{vg}^*$ is the total cost incurred by vessel $v = 1, \dots, V_g$ conditional on gear used $g = 1, \dots, G$ -- Mostly purse seiners!

- w is a matrix of variable input prices.
- z_{vg}^* is the optimal quantity of input used
 - Crew members?
 - Time spent at sea?
 - Distance traveled?
- $\mathbf{X}'_{vg} = [w; h_{vg}; x; Z_v; Env]$ is a matrix of explanatory variables:
 - Env is a matrix of **environmental covariates** (e.g., wind intensity and wave conditions in each trip)

Model 3: Total Annual Trips



The number of trips a vessel will take in a given year is assumed to follow a Poisson distribution:

$$Pr [T_{vgy}^* = t_v] = \frac{\exp^{-\exp(U'_{vg}\beta_g)} \exp(U'_{vg}\beta_g)^{t_v}}{t_v!}$$

where $U_{vg} = [p, w, h_{vg}, \bar{q}, Z_{vg}, Env]$ is a matrix of explanatory variables,

- t_v is the number of trips taken by vessel v using gear type g in year y
- Env include variables that reflect **annual weather conditions**
 - Accumulation of *bad weather days*?
 - *Number of storms*?

The price of each species is modeled using an Inverse Almost Ideal Demand System (IAIDS). The log of the price p_{iy} of a species i in year y is the following:

$$\ln p_{iy} = \sum_j^n \gamma_j \ln h_{j,y} + \gamma_H \ln H_y + \gamma_{FM} \ln P_y^{\text{FishMeal}} + \epsilon_{iy}$$

where:

- $H_y = \sum h_{j,y}$
- P_y^{FishMeal} is the fish meal world price

Harvest may be endogenous


- Three Stage Least Squares (3SLS) procedure,
- $h_{j,y}$ instrumented by variables that affect supply function such as *SST*, *Chl*, and fuel prices.

Preliminary results

Is there any substitution?




Small-scale vessels



Is there any substitution?



Industrial vessels



Seemingly Unrelated Regression (SUR) estimates of biomass dynamics for small pelagic species in Central–Southern Chile.

	Species		
	<i>Sardine</i>	<i>Anchoveta</i>	<i>Jack mackerel</i>
Biomass dynamics			
Constant	23.781***	10.355***	25.891***
Biomass (t)	0.353*	1.260***	1.035***
Biomass sq (t)	-0.011	-0.209***	-0.022**
Environmental effects			
SST	-12.377	-4.985	5.168
(SST) sq	52.233*	-2.526	-46.039
Chlorophyll-a	70.441**	-6.189	-15.631
Chlorophyll-a sq	155.277	59.220	438.550
Cross-species interactions			
Sardine × Jack mackerel	0.012	-	-0.008
Sardine × Anchoveta	0.017	-0.010	-
Anchoveta × Jack mackerel	-	0.011	0.045
Model fit			
R-squared (Adj.)	0.516	0.279	0.732
Entries are coefficients; significance: * p<0.10, ** p<0.05, *** p<0.01. Quadratic terms shown with “sq”; interactions use ×.			

- SST and CHL improves model performance ($F = 1.908$; p -value = 0.07).

- Finish biomass estimations
- Start soon with total annual trips
 - Hopefully, a short paper can come out of that work
- Two undergraduate students are working on the inverse demand (i.e., price) module for their theses
 - Results expected by July 2026
 - If time allows, they will also analyze long-run and short-run dynamics using a VEC model
 - Plan to write a paper based on their dissertations

¡Muchas gracias!

¿Preguntas?

Felipe J. Quezada-Escalona



Departamento de
Economía
Universidad de Concepción

 felipequezada.com

I gratefully acknowledge financial support for this research from ANID-Chile, under project Fondecyt
Iniciación No. 11250223.

Auffhammer, M. (2018). "Quantifying economic damages from climate change". In: *Journal of Economic Perspectives* 32.4, pp. 33-52.

Birkenbach, A. M., M. Lee, and M. D. Smith (2024). "Counterfactual Modeling of Multispecies Fisheries Outcomes under Market-Based Regulation". In: *Journal of the Association of Environmental and Resource Economists* 11.3, pp. 755-796. DOI: [10.1086/727356](https://doi.org/10.1086/727356). eprint: <https://doi.org/10.1086/727356>.

Cheung, W. W., V. W. Lam, J. L. Sarmiento, et al. (2010). "Large-scale redistribution of maximum fisheries catch potential in the global ocean under climate change". In: *Global change biology* 16.1, pp. 24-35.

E.U. Copernicus Marine Service Information (2025c). *Global Ocean Colour (Copernicus-GlobColour)*. Accessed on 23-SEP-2025. DOI: [10.48670/moi-00281](https://doi.org/10.48670/moi-00281).

E.U. Copernicus Marine Service Information (2025b). *Global Ocean Hourly Reprocessed Sea Surface Wind and Stress from Scatterometer and Model*. Accessed on 23-SEP-2025. DOI: [10.48670/moi-00185](https://doi.org/10.48670/moi-00185).

E.U. Copernicus Marine Service Information (2025a). *Global Ocean Physics Reanalysis*. Accessed on 23-SEP-2025. DOI: [10.48670/moi-00021](https://doi.org/10.48670/moi-00021).

Finkbeiner, E. M. (2015). "The role of diversification in dynamic small-scale fisheries: Lessons from Baja California Sur, Mexico". In: *Glob. Environ. Change* 32, pp. 139-152.

Gonzalez-Mon, B., Ö. Bodin, E. Lindkvist, et al. (2021). "Spatial diversification as a mechanism to adapt to environmental changes in small-scale fisheries". In: *Environ. Sci. Policy* 116, pp. 246-257.

Jardine, S. L., M. C. Fisher, S. K. Moore, et al. (2020). "Inequality in the Economic Impacts from Climate Shocks in Fisheries: The Case of Harmful Algal Blooms". In: *Ecol. Econ.* 176, p. 106691.

Kasperski, S. (2015). "Optimal Multi-species Harvesting in Ecologically and Economically Interdependent Fisheries". In: *Environ. Resour. Econ.* 61.4, pp. 517-557.

Kasperski, S. and D. S. Holland (2013). "Income diversification and risk for fishermen". En. In: *Proc. Natl. Acad. Sci. U. S. A.* 110.6, pp. 2076-2081.

Peña-Torres, J., J. Dresdner, and F. Vasquez (2017). "El Niño and Fishing Location Decisions: The Chilean Straddling Jack Mackerel Fishery". In: *Mar. Resour. Econ.* 32.3, pp. 249-275.

Poloczanska, E. S., C. J. Brown, W. J. Sydeman, et al. (2013). "Global imprint of climate change on marine life". En. In: *Nat. Clim. Chang.* 3.10, pp. 919-925.

Powell, F., A. Levine, and L. Ordonez-Gauger (2022). "Climate adaptation in the market squid fishery: fishermen responses to past variability associated with El Niño Southern Oscillation cycles inform our understanding of adaptive capacity in the face of future climate change". En. In: *Clim. Change* 173.1-2, p. 1.

Quezada, F. J., D. Tommasi, T. H. Frawley, et al. (2023). "Catch as catch can: markets, availability, and fishery closures drive distinct responses among the U.S. West Coast coastal pelagic species fleet segments". En. In: *Can. J. Fish. Aquat. Sci.*. Just-IN.

Reimer, M. N., J. K. Abbott, and A. C. Haynie (2022). "Structural behavioral models for rights-based fisheries". In: *Resource and Energy Economics* 68, p. 101294.

SUBPESCA (2020). *Informe Sectorial de Pesca y Acuicultura 2019*. Accessed: 02-04-2025.
URL: https://www.subpesca.cl/portal/618/articles-106845_documento.pdf.

Sumaila, U. R., W. W. Cheung, V. W. Lam, et al. (2011). "Climate change impacts on the biophysics and economics of world fisheries". In: *Nature climate change* 1.9, pp. 449-456.

Zhang, J. and M. D. Smith (2011). "Heterogeneous Response to Marine Reserve Formation: A Sorting Model approach". In: *Environ. Resour. Econ.* 49.3, pp. 311-325.

Appendix A: Numerical optimization

Vessel maximization problem

In each year, a vessel maximizes profits by choosing their optimal number of trips T_g and harvest levels per trip $h_{g\tau}$ given a gear type:

$$\begin{aligned} \max_{h_{gt}, T_g} \quad & \pi_{vgt} = \sum_{\tau=t}^{T_g} \rho^\tau \{P(h)h_{g\tau} - C_g(h_{g\tau}|w, x, Z, Env)\} \quad \tau = t, \dots, T_g \\ \text{s.t} \quad & q_{g,t+1} = \omega * \bar{q} - \sum_{t=1}^t h_{gt} \geq 0, \quad t = 1, \dots, T-1, \quad g = 1, \dots, G \end{aligned}$$

- where:
 - ρ is the intra-annual discount factor,
 - ω is a vector of shares of \bar{q} , and
 - $h_{lt} = 0$ for all $l \neq g$.