

# Phase transition in the exclusion process with slowed boundary reservoirs on resistance spaces

Joe P. Chen

Department of Mathematics  
Colgate University  
Hamilton, NY


2019 Finger Lakes Probability Seminar  
Cornell University  
April 19, 2019

Joint work with Patrícia Gonçalves (IST Lisboa)  
[arXiv:1904.08789](https://arxiv.org/abs/1904.08789)



# Exclusion process on a finite connected weighted graph

A system of random walkers subject to the **exclusion rule**: no 2 walkers occupy the same vertex at any time.



Let  $G = (V, E)$  be a finite connected graph endowed with positive edge weights  $\mathbf{c} = \{c_{xy}\}_{xy \in E}$ .


The **symmetric exclusion process** on  $(G, \mathbf{c})$  is a continuous-time Markov process on  $\{0, 1\}^V$  with generator

$$(\mathcal{L}^{\text{EX}} f)(\eta) = \sum_{xy \in E} c_{xy} [f(\eta^{xy}) - f(\eta)], \quad f : \{0, 1\}^V \rightarrow \mathbb{R},$$

$$\text{where } (\eta^{xy})(z) = \begin{cases} \eta(y), & \text{if } z = x, \\ \eta(x), & \text{if } z = y, \\ \eta(z), & \text{otherwise.} \end{cases}$$

- **Conserved quantity**: Total # of particles.
- Each product Bernoulli measure  $\nu_\alpha$ ,  $\alpha \in [0, 1]$ , with marginal  $\nu_\alpha\{\eta : \eta(x) = 1\} = \alpha$  for each  $x \in V$ , is an **invariant measure**.
- **Dirichlet energy**:  $\mathcal{E}^{\text{EX}}(f) = \langle f, -\mathcal{L}^{\text{EX}} f \rangle_{\nu_\alpha} = \frac{1}{2} \sum_{zw \in E} c_{zw} \int_{\{0, 1\}^V} [f(\eta^{zw}) - f(\eta)]^2 d\nu_\alpha(\eta).$

# Adding reservoirs (Kawasaki dynamics) to the exclusion process



Designate a finite boundary set  $\partial V \subset V$ . For each  $a \in \partial V$ :


- At rate  $\lambda_+(a)$ ,  $\eta(a) = 0 \rightarrow \eta(a) = 1$  (birth).
- At rate  $\lambda_-(a)$ ,  $\eta(a) = 1 \rightarrow \eta(a) = 0$  (death).

Formally,  $(\mathcal{L}_{\partial V}^{\text{boun}} f)(\eta) = \sum_{a \in \partial V} [\lambda_+(a)(1 - \eta(a)) + \lambda_-(a)\eta(a)][f(\eta^a) - f(\eta)]$ ,  $f : \{0, 1\}^V \rightarrow \mathbb{R}$ , where


$$\eta^a(z) = \begin{cases} 1 - \eta(a), & \text{if } z = a, \\ \eta(z), & \text{otherwise.} \end{cases}$$

- **1D boundary-driven simple exclusion process:** generator  $N^2 \left( \mathcal{L}_{\{1,2,\dots,N-1\}}^{\text{EX}} + \mathcal{L}_{\{1,N-1\}}^{\text{boun}} \right)$ .
- Has been studied extensively for the past  $\sim 15$  years:  
Hydrodynamic limits, fluctuations, large deviations, etc.
- **Difficulties:** # of particles is no longer conserved; the invariant measure is in general not explicit.

# Extending the analysis to higher dims & with $> 2$ reservoirs?



# Boundary-driven exclusion process on the Sierpinski gasket




- Construction of Brownian motion with invariant measure  $m$  (the standard self-similar measure) as scaling limit of RWs accelerated by  $\mathcal{T}_N = 5^N$ .  
[Goldstein '87, Kusuoka '88, Barlow-Perkins '88]
- A robust notion of calculus on SG which in some sense mimics (but in many other senses differs from) calculus in 1D: Laplacian, Dirichlet form, integration by parts, boundary-value problems, etc.  
[See the books by Kigami '01, Strichartz '06]

What is the analog of " $\int_K |\nabla f|^2 dx$ " in the fractal setting?

- A good model for rigorously studying (non)equilibrium stochastic dynamics with  $\geq 3$  boundary reservoirs.

# Boundary-driven exclusion process on the Sierpinski gasket




- Construction of Brownian motion with invariant measure  $m$  (the standard self-similar measure) as scaling limit of RWs accelerated by  $\mathcal{T}_N = 5^N$ .  
[Goldstein '87, Kusuoka '88, Barlow-Perkins '88]
- A robust notion of calculus on SG which in some sense mimics (but in many other senses differs from) calculus in 1D: Laplacian, Dirichlet form, integration by parts, boundary-value problems, etc.  
[See the books by Kigami '01, Strichartz '06]

$$\mathcal{E}(f) = \lim_{N \rightarrow \infty} \frac{5^N}{3^N} \sum_{xy \in E_N} (f(x) - f(y))^2, \quad f \in L^2(K, m).$$

- A good model for rigorously studying (non)equilibrium stochastic dynamics with  $\geq 3$  boundary reservoirs.

# Boundary-driven exclusion process on the Sierpinski gasket



$$5^N \mathcal{L}_N^{\text{bEX}} = 5^N \left( \mathcal{L}_N^{\text{EX}} + \frac{1}{b^N} \mathcal{L}_N^{\text{boun}} \right).$$

Parameter  $b > 0$  governs the inverse speed (relative to the bulk jump rate) at which the reservoir injects/extracts particles into/from the boundary vertices  $V_0$ .

Our main result in a nutshell [C.–Gonçalves '19]

A **phase transition** in the scaling limit of the particle density depending on the value of  $b$ , reflected by the different **boundary conditions**. The critical value of  $b$  is  $\frac{5}{3}$ .

Dirichlet ( $b < \frac{5}{3}$ ), Robin ( $b = \frac{5}{3}$ ), Neumann ( $b > \frac{5}{3}$ )

## Hydrodynamic limit: a LLN result

Assume that sequence of probability measures  $\{\mu_N\}_{N \geq 1}$  on  $\{0, 1\}^{V_N}$  is associated to a density profile  $\varrho : K \rightarrow [0, 1]$ : for any continuous function  $F : K \rightarrow \mathbb{R}$  and any  $\delta > 0$ ,

$$\lim_{N \rightarrow \infty} \mu_N \left\{ \eta \in \{0, 1\}^{V_N} : \left| \frac{1}{|V_N|} \sum_{x \in V_N} F(x) \eta(x) - \int_K F(x) \varrho(x) dm(x) \right| > \delta \right\} = 0.$$

Given the process  $\{\eta_t^N : t \geq 0\}$  generated by  $5^N \mathcal{L}_N^{\text{bEX}}$ , the **empirical density measure**  $\pi_t^N$  given by

$$\pi_t^N = \frac{1}{|V_N|} \sum_{x \in V_N} \eta_t^N(x) \delta_{\{x\}}$$

and for any test function  $F : K \rightarrow \mathbb{R}$ , we denote the integral of  $F$  wrt  $\pi_t^N$  by  $\pi_t^N(F)$  which equals

$$\pi_t^N(F) = \frac{1}{|V_N|} \sum_{x \in V_N} \eta_t^N(x) F(x).$$


**Claim.** The sequence  $\{\pi_t^N\}_N$  converges in the Skorokhod topology on  $D([0, T], \mathcal{M}_+)$  to the unique measure  $\pi$ . with  $d\pi(x) = \rho(\cdot, x) dm(x)$ .

For any  $t \in [0, T]$ , any continuous  $F : K \rightarrow \mathbb{R}$  and any  $\delta > 0$ ,

$$\lim_{N \rightarrow \infty} \mu_N \left\{ \eta_t^N : \left| \frac{1}{|V_N|} \sum_{x \in V_N} \eta_t^N(x) F(x) - \int_K F(x) \rho(t, x) dm(x) \right| > \delta \right\} = 0,$$



# Hydrodynamic limit: a LLN result



$$5^N \mathcal{L}_N^{\text{bEX}} = 5^N \left( \mathcal{L}_N^{\text{EX}} + \frac{1}{b^N} \mathcal{L}_N^{\text{boun}} \right).$$

$$\lambda_{\Sigma}(a) = \lambda_{+}(a) + \lambda_{-}(a)$$

$$\bar{\rho}(a) = \frac{\lambda_{+}(a)}{\lambda_{\Sigma}(a)}$$

## Theorem (Density hydrodynamic limit)


For any  $t \in [0, T]$ , any continuous  $F : K \rightarrow \mathbb{R}$  and any  $\delta > 0$ ,

$$\lim_{N \rightarrow \infty} \mu_N \left\{ \eta_t^N : \left| \frac{1}{|V_N|} \sum_{x \in V_N} \eta_t^N(x) F(x) - \int_K F(x) \rho(t, x) dm(x) \right| > \delta \right\} = 0,$$

where  $\rho$  is the unique weak solution of the heat equation  
with Dirichlet boundary condition if  $b < \frac{5}{3}$ :

$$\begin{cases} \partial_t \rho(t, x) = \frac{2}{3} \Delta \rho(t, x), & t \in [0, T], x \in K \setminus V_0, \\ \rho(t, a) = \bar{\rho}(a), & t \in (0, T], a \in V_0, \\ \rho(0, x) = \varrho(x), & x \in K. \end{cases}$$

# Hydrodynamic limit: a LLN result



$$5^N \mathcal{L}_N^{\text{bEX}} = 5^N \left( \mathcal{L}_N^{\text{EX}} + \frac{1}{b^N} \mathcal{L}_N^{\text{boun}} \right).$$

$$\lambda_{\Sigma}(a) = \lambda_{+}(a) + \lambda_{-}(a)$$

$$\bar{\rho}(a) = \frac{\lambda_{+}(a)}{\lambda_{\Sigma}(a)}$$

## Theorem (Density hydrodynamic limit)


For any  $t \in [0, T]$ , any continuous  $F : K \rightarrow \mathbb{R}$  and any  $\delta > 0$ ,

$$\lim_{N \rightarrow \infty} \mu_N \left\{ \eta_t^N : \left| \frac{1}{|V_N|} \sum_{x \in V_N} \eta_t^N(x) F(x) - \int_K F(x) \rho(t, x) dm(x) \right| > \delta \right\} = 0,$$

where  $\rho$  is the unique weak solution of the heat equation with Neumann boundary condition if  $b > \frac{5}{3}$ :

$$\begin{cases} \partial_t \rho(t, x) = \frac{2}{3} \Delta \rho(t, x), & t \in [0, T], x \in K \setminus V_0, \\ (\partial^\perp \rho)(t, a) = 0, & t \in (0, T], a \in V_0, \\ \rho(0, x) = \varrho(x), & x \in K. \end{cases}$$

# Hydrodynamic limit: a LLN result



$$5^N \mathcal{L}_N^{\text{bEX}} = 5^N \left( \mathcal{L}_N^{\text{EX}} + \frac{1}{b^N} \mathcal{L}_N^{\text{boun}} \right).$$

$$\lambda_{\Sigma}(a) = \lambda_{+}(a) + \lambda_{-}(a)$$

$$\bar{\rho}(a) = \frac{\lambda_{+}(a)}{\lambda_{\Sigma}(a)}$$

## Theorem (Density hydrodynamic limit)

For any  $t \in [0, T]$ , any continuous  $F : K \rightarrow \mathbb{R}$  and any  $\delta > 0$ ,

$$\lim_{N \rightarrow \infty} \mu_N \left\{ \eta_t^N : \left| \frac{1}{|V_N|} \sum_{x \in V_N} \eta_t^N(x) F(x) - \int_K F(x) \rho(t, x) dm(x) \right| > \delta \right\} = 0,$$

where  $\rho$  is the unique weak solution of the heat equation with linear Robin boundary condition if  $b = \frac{5}{3}$ :

$$\begin{cases} \partial_t \rho(t, x) = \frac{2}{3} \Delta \rho(t, x), & t \in [0, T], x \in K \setminus V_0, \\ (\partial^\perp \rho)(t, a) = -\lambda_{\Sigma}(a)(\rho(t, a) - \bar{\rho}(a)), & t \in (0, T], a \in V_0, \\ \rho(0, x) = \varrho(x), & x \in K. \end{cases}$$

# Heuristics for hydrodynamics

Analysis of Dynkin's martingale (which has QV tending to 0 as  $N \rightarrow \infty$ ):

$$\begin{aligned} M_t^N(F) &:= \pi_t^N(F_t) - \pi_0^N(F_0) - \int_0^t \pi_s^N \left( \left( \frac{2}{3} \Delta + \partial_s \right) F_s \right) ds \\ &\quad + \int_0^t \frac{3^N}{|V_N|} \sum_{a \in V_0} \left[ \eta_s^N(a) (\partial^\perp F_s)(a) + \frac{5^N}{3^N b^N} \lambda_\Sigma(a) (\eta_s^N(a) - \bar{\rho}(a)) F_s(a) \right] ds + o_N(1). \end{aligned}$$

---

## [Ingredient #1] Analysis on fractals

Convergence of discrete Laplacian to the continuous counterpart; normal derivatives at the boundary; integration by parts formula ... [Kigami '01, Strichartz '06].

# Heuristics for hydrodynamics

Analysis of Dynkin's martingale (which has QV tending to 0 as  $N \rightarrow \infty$ ):

$$\begin{aligned} M_t^N(F) &:= \pi_t^N(F_t) - \pi_0^N(F_0) - \int_0^t \pi_s^N \left( \left( \frac{2}{3} \Delta + \partial_s \right) F_s \right) ds \\ &\quad + \int_0^t \frac{3^N}{|V_N|} \sum_{a \in V_0} \left[ \eta_s^N(a) (\partial^\perp F_s)(a) + \frac{5^N}{3^N b^N} \lambda_\Sigma(a) (\eta_s^N(a) - \bar{\rho}(a)) F_s(a) \right] ds + o_N(1). \end{aligned}$$

---

**[Ingredient #1] Analysis on fractals**

This part will produce the weak formulation of the heat equation.

# Heuristics for hydrodynamics

Analysis of Dynkin's martingale (which has QV tending to 0 as  $N \rightarrow \infty$ ):

$$M_t^N(F) := \pi_t^N(F_t) - \pi_0^N(F_0) - \int_0^t \pi_s^N \left( \left( \frac{2}{3} \Delta + \partial_s \right) F_s \right) ds \\ + \int_0^t \frac{3^N}{|V_N|} \sum_{a \in V_0} \left[ \eta_s^N(a) (\partial^\perp F_s)(a) + \frac{5^N}{3^N b^N} \lambda_\Sigma(a) (\eta_s^N(a) - \bar{\rho}(a)) F_s(a) \right] ds + o_N(1).$$

## [Ingredient #2] Treatment of the boundary term

- $b > 5/3$ : The first term dominates, should converge to  $\int_0^t \frac{2}{3} \sum_{a \in V_0} \rho_s(a) (\partial^\perp F_s)(a) ds$
- $b = 5/3$ : Both terms contribute equally, should converge to  $\int_0^t \frac{2}{3} \sum_{a \in V_0} \left[ \rho_s(a) (\partial^\perp F_s)(a) + \lambda_\Sigma(a) (\rho_s(a) - \bar{\rho}(a)) F_s(a) \right] ds$
- $b < 5/3$ : Impose  $\rho_t(a) = \bar{\rho}(a)$  for all  $a \in V_0$ , should converge to  $\int_0^t \frac{2}{3} \sum_{a \in V_0} \bar{\rho}(a) (\partial^\perp F_s)(a) ds$

Require a series of **replacement lemmas** — not trivial on state spaces without translational invariance!  
→ I will come back to address this at the end.

Analysis of Dynkin's martingale (which has QV tending to 0 as  $N \rightarrow \infty$ ):

$$M_t^N(F) := \pi_t^N(F_t) - \pi_0^N(F_0) - \int_0^t \pi_s^N \left( \left( \frac{2}{3} \Delta + \partial_s \right) F_s \right) ds \\ + \int_0^t \frac{3^N}{|V_N|} \sum_{a \in V_0} \left[ \eta_s^N(a) (\partial^\perp F_s)(a) + \frac{5^N}{3^N b^N} \lambda_\Sigma(a) (\eta_s^N(a) - \bar{\rho}(a)) F_s(a) \right] ds + o_N(1).$$

$\downarrow N \rightarrow \infty$

$$0 = \pi_t(F_t) - \pi_0(F_0) - \int_0^t \pi_s \left( \left( \frac{2}{3} \Delta + \partial_s \right) F_s \right) ds + (\text{boundary term})$$

---

## [Ingredient #3] Convergence of stochastic processes

- Show that  $\{\pi_t^N\}_N$  is tight in the Skorokhod topology on  $D([0, T], \mathcal{M}_+)$  via Aldous' criterion.
- Prove that any limit point  $\pi_\cdot$  is absolutely continuous w.r.t. the self-similar measure  $m$ , with  $\pi_t(dx) = \rho(t, x) dm(x)$ , and  $\rho \in L^2(0, T, \text{dom} \mathcal{E})$ .
- Finally, prove ! of the weak solution to the heat equation to conclude ! of the limit point.

# Density fluctuation field (at equilibrium): Heuristics

**Equilibrium**  $\Leftrightarrow \lambda_+(a) = \lambda_+$  and  $\lambda_-(a) = \lambda_-$  for all  $a \in V_0$ . (Otherwise, **nonequilibrium**.)

The product Bernoulli measure  $\nu_\rho^N$  with  $\rho = \lambda_+ / (\lambda_+ + \lambda_-)$  is stationary for the process.

**Density fluctuation field (DFF)**

$$\mathcal{Y}_t^N(F) = \frac{1}{\sqrt{|V_N|}} \sum_{x \in V_N} \left( \eta_t^N(x) - \rho \right) F(x)$$

The corresponding Dynkin's martingale is

$$\begin{aligned} \mathcal{M}_t^N(F) &= \mathcal{Y}_t^N(F) - \mathcal{Y}_0^N(F) - \int_0^t \mathcal{Y}_s^N(\Delta_N F) ds + o_N(1) \\ &\quad + \frac{3^N}{\sqrt{|V_N|}} \int_0^t \sum_{a \in V_0} \bar{\eta}_s^N(a) \left[ (\partial_N^\perp F)(a) + \frac{5^N}{b^N 3^N} \lambda_\Sigma F(a) \right] ds, \end{aligned}$$

which has QV

$$\begin{aligned} \langle M^N(F) \rangle_t &= \int_0^t \frac{5^N}{|V_N|^2} \sum_{x \in V_N} \sum_{\substack{y \in V_N \\ y \sim x}} (\eta_s^N(x) - \eta_s^N(y))^2 (F(x) - F(y))^2 ds \\ &\quad + \int_0^t \sum_{a \in V_0} \frac{5^N}{b^N |V_N|^2} \{ \lambda_-(a) \eta_s^N(a) + \lambda_+(a) (1 - \eta_s^N(a)) \} F^2(a) ds. \end{aligned}$$



# Density fluctuation field (at equilibrium): Heuristics

**Equilibrium**  $\Leftrightarrow \lambda_+(a) = \lambda_+$  and  $\lambda_-(a) = \lambda_-$  for all  $a \in V_0$ . (Otherwise, **nonequilibrium**.)  
The product Bernoulli measure  $\nu_\rho^N$  with  $\rho = \lambda_+ / (\lambda_+ + \lambda_-)$  is stationary for the process.

**Density fluctuation field (DFF)**

$$\mathcal{Y}_t^N(F) = \frac{1}{\sqrt{|V_N|}} \sum_{x \in V_N} \left( \eta_t^N(x) - \rho \right) F(x)$$

The corresponding Dynkin's martingale is

$$\begin{aligned} \mathcal{M}_t^N(F) &= \mathcal{Y}_t^N(F) - \mathcal{Y}_0^N(F) - \int_0^t \mathcal{Y}_s^N(\Delta_N F) ds + o_N(1) \\ &\quad + \frac{3^N}{\sqrt{|V_N|}} \int_0^t \sum_{a \in V_0} \bar{\eta}_s^N(a) \left[ (\partial_N^\perp F)(a) + \frac{5^N}{b^N 3^N} \lambda_\Sigma F(a) \right] ds, \end{aligned}$$

which, as  $N \rightarrow \infty$ , has the QV of a space-time white noise (with boundary condition)

$$\frac{2}{3} \cdot 2\rho(1-\rho)t\mathcal{E}_b(F), \quad \text{where } \mathcal{E}_b(F) = \mathcal{E}(F) + \lambda_\Sigma \sum_{a \in V_0} F^2(a) \mathbf{1}_{\{b=5/3\}}$$

# Density fluctuation field (at equilibrium): Heuristics

**Equilibrium**  $\Leftrightarrow \lambda_+(a) = \lambda_+$  and  $\lambda_-(a) = \lambda_-$  for all  $a \in V_0$ . (Otherwise, **nonequilibrium**.)

The product Bernoulli measure  $\nu_\rho^N$  with  $\rho = \lambda_+ / (\lambda_+ + \lambda_-)$  is stationary for the process.

**Density fluctuation field (DFF)**

$$\mathcal{Y}_t^N(F) = \frac{1}{\sqrt{|V_N|}} \sum_{x \in V_N} \left( \eta_t^N(x) - \rho \right) F(x)$$

The corresponding Dynkin's martingale is

$$\begin{aligned} \mathcal{M}_t^N(F) &= \mathcal{Y}_t^N(F) - \mathcal{Y}_0^N(F) - \int_0^t \mathcal{Y}_s^N(\Delta_N F) ds + o_N(1) \\ &\quad + \frac{3^N}{\sqrt{|V_N|}} \int_0^t \sum_{a \in V_0} \bar{\eta}_s^N(a) \left[ (\partial_N^\perp F)(a) + \frac{5^N}{b^N 3^N} \lambda_\Sigma F(a) \right] ds, \end{aligned}$$

We then argue that the test function  $F \in \text{dom} \Delta_b$  be chosen appropriate to each boundary condition such that **the boundary term vanishes** as  $N \rightarrow \infty$ .

$$\text{dom} \Delta_b := \begin{cases} \{F \in \text{dom} \Delta : F|_{V_0} = 0\}, & \text{if } b < 5/3, \\ \{F \in \text{dom} \Delta : (\partial^\perp F)|_{V_0} = -\lambda_\Sigma F|_{V_0}\}, & \text{if } b = 5/3, \\ \{F \in \text{dom} \Delta : (\partial^\perp F)|_{V_0} = 0\}, & \text{if } b > 5/3. \end{cases}$$

For technical reasons (in order to use Mitoma's tightness criterion) we use a smaller test function space

$S_b := \{F \in \text{dom} \Delta_b : \Delta_b F \in \text{dom} \Delta_b\}$ , which can be made into a Frechét space.

Let  $S'_b$  be the topological dual of  $S_b$ .

## Definition (Ornstein-Uhlenbeck equation)

We say that a random element  $\mathcal{Y}$  taking values in  $C([0, T], S'_b)$  is a solution to the **Ornstein-Uhlenbeck equation** on  $K$  with parameter  $b$  if:

- ① For every  $F \in S_b$ ,

$$\mathcal{M}_t(F) = \mathcal{Y}_t(F) - \mathcal{Y}_0(F) - \int_0^t \mathcal{Y}_s\left(\frac{2}{3}\Delta_b F\right) ds$$

$$\text{and } \mathcal{N}_t(F) = (\mathcal{M}_t(F))^2 - \frac{2}{3} \cdot 2\rho(1 - \rho)t\mathcal{E}_b(F)$$


are  $\mathcal{F}_t$ -martingales, where  $\mathcal{F}_t := \sigma\{\mathcal{Y}_s(F) : s \leq t\}$  for each  $t \in [0, T]$ .

- ②  $\mathcal{Y}_0$  is a centered Gaussian  $S'_b$ -valued random variable with covariance

$$\mathbb{E}_\rho^b[\mathcal{Y}_0(F)\mathcal{Y}_0(G)] = \rho(1 - \rho) \int_K F(x)G(x) dm(x), \quad \forall F, G \in S_b.$$

Moreover, for every  $F \in S_b$ , the process  $\{\mathcal{Y}_t(F) : t \geq 0\}$  is Gaussian: the distribution of  $\mathcal{Y}_t(F)$  conditional upon  $\mathcal{F}_s$ ,  $s < t$ , is Gaussian with mean  $\mathcal{Y}_s(\tilde{T}_{t-s}^b F)$  and variance  $\int_0^{t-s} \frac{2}{3} \cdot 2\rho(1 - \rho)\mathcal{E}_b(\tilde{T}_r^b F) dr$ , where  $\{\tilde{T}_t^b : t > 0\}$  is the heat semigroup associated with  $\frac{2}{3}\mathcal{E}_b$ .

# O-U limit of equilibrium density fluctuations: a CLT result



$$5^N \mathcal{L}_N^{\text{bEX}} = 5^N \left( \mathcal{L}_N^{\text{EX}} + \frac{1}{b^N} \mathcal{L}_N^{\text{boun}} \right).$$

Dirichlet ( $b < \frac{5}{3}$ ), Robin ( $b = \frac{5}{3}$ ), Neumann ( $b > \frac{5}{3}$ )

**Equilibrium**  $\Leftrightarrow \lambda_+(a) = \lambda_+$  and  $\lambda_-(a) = \lambda_-$  for all  $a \in V_0$ .

Let  $\mathbb{Q}_\rho^{N,b}$  be the probability measure on  $D([0, T], S'_b)$  induced by the DFF  $\mathcal{Y}^N$  started from  $\nu_\rho^N$  and boundary parameter  $b$ .


## Theorem (CLT)

*The sequence  $\{\mathbb{Q}_\rho^{N,b}\}_N$  converges in distribution, as  $N \rightarrow \infty$ , to a unique solution of the Ornstein-Uhlenbeck equation with parameter  $b$  (as defined previously).*

**Key Lemma.**  $\tilde{T}_t^b(S_b) \subset S_b$  for any  $t > 0$ . Enough to verify that  $\tilde{T}_t^b(L^1(K, m)) \subset \text{dom} \Delta_b$ , which can be shown using e.g. the Nash inequality (heat kernel upper bound).

The rest of the argument follows a martingale approach of Kipnis–Landim.

# A quick word on the replacement lemma



For finite  $\Lambda \subset V$ , denote  $\text{Av}_\Lambda[\eta] := |\Lambda|^{-1} \sum_{z \in \Lambda} \eta(z)$ .

We need to prove that for every  $t > 0$ ,

## Replacement lemma

$$\overline{\lim}_{j \rightarrow \infty} \overline{\lim}_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[ \left| \int_0^t \left( \eta_s^N(a) - \text{Av}_{V_N \cap K_j(a)}[\eta_s^N] \right) ds \right| \right] = 0, \quad a \in V_0.$$

where  $K_j(a)$  is the cell of depth level  $j$  which contains  $a$ .

Proving the replacement lemma without using translational invariance is difficult!

What other mechanism can be used to carry out this replacement?

## A quick word on the replacement lemma

For finite  $\Lambda \subset V$ , denote  $A_{V_\Lambda}[\eta] := |\Lambda|^{-1} \sum_{z \in \Lambda} \eta(z)$ .

We need to prove that for every  $t > 0$ ,

### Replacement lemma

$$\overline{\lim}_{j \rightarrow \infty} \overline{\lim}_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[ \left| \int_0^t \left( \eta_s^N(a) - A_{V_{V_N \cap K_j(a)}}[\eta_s^N] \right) ds \right| \right] = 0, \quad a \in V_0.$$

where  $K_j(a)$  is the cell of depth level  $j$  which contains  $a$ .


For resistance spaces, we can use the **moving particle lemma** [C., ECP '17]:

$$\frac{1}{2} \int [f(\eta^{xy}) - f(\eta)]^2 d\nu_\alpha(\eta) \leq R_{\text{eff}}(x, y) \mathcal{E}^{\text{EX}}(f), \quad f : \{0, 1\}^V \rightarrow \mathbb{R},$$

where  $R_{\text{eff}}(x, y)$  is the **effective resistance** between  $x$  and  $y$  in the *random walk process* on the same weighted graph  $(G, c)$ .

For more details on the MPL and its connection to the octopus inequality of [Caputo–Liggett–Richthammer JAMS '10], see my ECP paper.

# A sneak preview of upcoming series of works, and **Thank you!**



$$5^N \mathcal{L}_N^{\text{bEX}} = 5^N \left( \mathcal{L}_N^{\text{EX}} + \frac{1}{b^N} \mathcal{L}_N^{\text{boun}} \right).$$

**Symmetric** exclusion process with **slowed** boundary on the Sierpinski gasket

Dirichlet ( $b < \frac{5}{3}$ ), Robin ( $b = \frac{5}{3}$ ), Neumann ( $b > \frac{5}{3}$ )

**Equilibrium**  $\Leftrightarrow \lambda_+(a) = \lambda_+$  and  $\lambda_-(a) = \lambda_-$  for all  $a \in V_0$ . (Otherwise, **nonequilibrium**.)

- (Non)equilibrium density hydrodynamic limit (DRN✓) [This talk](#)
- Ornstein-Uhlenbeck limit of equilibrium density fluctuations (DRN✓). [This talk](#)
- Large deviations principle for the (non)equilibrium density (D✓) [[C.–Hinz '19](#)]
- Hydrostatic limit, scaling limit of nonequilibrium density fluctuations (D in progress). [[C.–Franceschini–Gonçalves–Menezes '19+](#)]  $\rightarrow$  careful study of two-particle correlations

- **More in the pipeline:**

Motion of the tagged particle (a fractional BM on the gasket?).

Add (suitably rescaled) weak asymmetry to the jump rate, prove that the equilibrium density fluctuations converges (subsequentially) to a stochastic Burgers equation [[C. '19+](#)]