

Mixing Times of Card Shuffling and Road Traffic

Rodrigo Marinho and Joe P. Chen



IST, University of Lisbon



Colgate University

October 09, 2020

Card shuffling: some history

- ▶ Choose your favourite way to shuffle a deck of cards.

- Choose your favourite way to shuffle a deck of cards.

How many times do you have
to shuffle the deck like that to
mix it up?

Card shuffling: some history

- What is the meaning of mixing a deck?

Card shuffling: some history

- ▶ What is the meaning of mixing a deck?
- ▶ Deck with n cards $\Rightarrow n!$ permutations

Card shuffling: some history

- ▶ What is the meaning of mixing a deck?
- ▶ Deck with n cards $\Rightarrow n!$ permutations
- ▶ Deck with 52 cards \Rightarrow order 10^{68} permutations

Card shuffling: some history

- ▶ What is the meaning of mixing a deck?
- ▶ Deck with n cards $\Rightarrow n!$ permutations
- ▶ Deck with 52 cards \Rightarrow order 10^{68} permutations

This is much more than the
number of particles on Earth!

Card shuffling: some history

- ▶ What is the meaning of mixing a deck?
- ▶ Deck with n cards $\Rightarrow n!$ permutations
- ▶ Deck with 52 cards \Rightarrow order 10^{68} permutations

This is much more than the
number of particles on Earth!

- ▶ A deck will be considered mixed whenever all permutations are equiprobable.

Card shuffling as maths

- ▶ μ_t : Probability distribution of your shuffle at time t

Card shuffling as maths

- μ_t : Probability distribution of your shuffle at time t , that is,

$$\mu_t(\sigma) = \mathbb{P}(\sigma_t = \sigma);$$

Card shuffling as maths

- ▶ μ_t : Probability distribution of your shuffle at time t , that is,

$$\mu_t(\sigma) = \mathbb{P}(\sigma_t = \sigma);$$

- ▶ U_n : Uniform distribution on S_n ;

Card shuffling as maths

- ▶ μ_t : Probability distribution of your shuffle at time t , that is,

$$\mu_t(\sigma) = \mathbb{P}(\sigma_t = \sigma);$$

- ▶ U_n : Uniform distribution on S_n ;

- ▶ $d_n(t) := \|\mu_t - U_n\|_{TV} = \frac{1}{2} \sum_{\sigma \in S_n} |\mu_t(\sigma) - \frac{1}{n!}|$
(distance to equilibrium);

Card shuffling as maths

- ▶ μ_t : Probability distribution of your shuffle at time t , that is,

$$\mu_t(\sigma) = \mathbb{P}(\sigma_t = \sigma);$$

- ▶ U_n : Uniform distribution on S_n ;

- ▶ $d_n(t) := \|\mu_t - U_n\|_{TV} = \frac{1}{2} \sum_{\sigma \in S_n} |\mu_t(\sigma) - \frac{1}{n!}|$
(distance to equilibrium);

- ▶ $t_{\text{mix}}^n(\varepsilon) := \inf\{t \geq 0; d_n(t) < \varepsilon\};$

Card shuffling as maths

- ▶ μ_t : Probability distribution of your shuffle at time t , that is,

$$\mu_t(\sigma) = \mathbb{P}(\sigma_t = \sigma);$$

- ▶ U_n : Uniform distribution on S_n ;

- ▶ $d_n(t) := \|\mu_t - U_n\|_{TV} = \frac{1}{2} \sum_{\sigma \in S_n} |\mu_t(\sigma) - \frac{1}{n!}|$
(distance to equilibrium);

- ▶ $t_{\text{mix}}^n(\varepsilon) := \inf\{t \geq 0; d_n(t) < \varepsilon\};$

- ▶ $t_{\text{mix}}^n := t_{\text{mix}}^n(1/4).$

Riffle shuffle



Riffle shuffle



- Bayer and Diaconis: $t_{\text{mix}}^n(\epsilon) = \frac{3}{2} \log_2 n$.

Riffle shuffle



► Bayer and Diaconis: $t_{\text{mix}}^n(\epsilon) = \frac{3}{2} \log_2 n$.

► $n = 52 \Rightarrow t_{\text{mix}}^{52} = 7$.

The Annals of Applied Probability
1992, Vol. 2, No. 2, 294–313

TRAILING THE DOVETAIL SHUFFLE TO ITS LAIR

BY DAVE BAYER¹ AND PERSI DIACONIS²

Columbia University and Harvard University

2. A card trick. Rising sequences, the basic invariant of riffle shuffling, were discovered by magicians Williams and Jordan at the beginning of this



TRAILING THE DOVETAIL SHUFFLE TO ITS LAIR

297

century. A rising sequence is a maximal subset of an arrangement of cards,

Overhand shuffle



Overhand shuffle



- $t_{\text{mix}}^n = \Theta(n^2 \log n)$. The correct constant is unknown.


Overhand shuffle




► $t_{\text{mix}}^n = \Theta(n^2 \log n)$. The correct constant is unknown.

► $n = 52 \Rightarrow t_{\text{mix}}^{52} \approx 10000$.

Top-to-random shuffle




Top-to-random shuffle



► Aldous and Diaconis: $t_{\text{mix}}^n(\epsilon) = n \log n$.


Top-to-random shuffle




► Aldous and Diaconis: $t_{\text{mix}}^n(\epsilon) = n \log n$.

► $n = 52 \Rightarrow t_{\text{mix}}^{52} \approx 200$.

Rudvalis shuffle




Rudvalis shuffle




- **Hildebrand** and **Wilson**: $t_{\text{mix}}^n(\varepsilon) = \Theta(n^3 \log n)$. The correct constant is unknown.

Rudvalis shuffle



- Hildebrand and Wilson: $t_{\text{mix}}^n(\varepsilon) = \Theta(n^3 \log n)$. The correct constant is unknown. Work in progress with P. Gonçalves, M. Jara and D. Moreira.

Rudvalis shuffle



- Hildebrand and Wilson: $t_{\text{mix}}^n(\varepsilon) = \Theta(n^3 \log n)$. The correct constant is unknown. Work in progress with P. Gonçalves, M. Jara and D. Moreira.
- $n = 52 \Rightarrow t_{\text{mix}}^{52} \approx 14000$.

Random transposition shuffle

Example

1	2	3	4	5	6	7
1	5	3	4	2	6	7

Random transposition shuffle

Example

1	2	3	4	5	6	7
1	5	3	4	2	6	7

- Diaconis and Shahshahani, and Matthews

Random transposition shuffle

Example

1	2	3	4	5	6	7
1	5	3	4	2	6	7

- ▶ Diaconis and Shahshahani, and Matthews
- ▶ $t_{\text{mix}}^n(\varepsilon) = \frac{1}{2}n \log n$.

Random transposition shuffle

Example

1	2	3	4	5	6	7
1	5	3	4	2	6	7

- ▶ Diaconis and Shahshahani, and Matthews
- ▶ $t_{\text{mix}}^n(\varepsilon) = \frac{1}{2}n \log n.$
- ▶ $n = 52 \Rightarrow t_{\text{mix}}^{52} \approx 100.$

Random adjacent transposition shuffle

Example

1	2	3	4	5	6	7
1	2	4	3	5	6	7

Random adjacent transposition shuffle

Example

1	2	3	4	5	6	7
1	2	4	3	5	6	7

- Wilson (2004) and Lacoïn (2016).

Random adjacent transposition shuffle


Example

1	2	3	4	5	6	7
1	2	4	3	5	6	7


► Wilson (2004) and Lacoïn (2016).

► $t_{\text{mix}}^n(\varepsilon) = \frac{1}{\pi^2} n^3 \log n$.

Cutoff phenomenon



Cutoff phenomenon



$$\lim_{n \rightarrow \infty} d_n(\kappa t_{\text{mix}}^n) = \begin{cases} 1, & \text{if } \kappa < 1; \\ 0, & \text{if } \kappa > 1. \end{cases}$$


Shuffling cards on graphs

Shuffling cards on graphs

- ▶ $G = (V, E)$: finite connected graph;


Shuffling cards on graphs

- $G = (V, E)$: finite connected graph;




Shuffling cards on graphs

- ▶ $G = (V, E)$: finite connected graph;




- ▶ At each vertex of G we put a card and at each edge of G we attach an exponential clock of rate 1.

Shuffling cards on graphs




Shuffling cards on graphs



- Flip edges, exchanging the cards on the two incident vertices.


Shuffling cards on graphs



- Flip edges, exchanging the cards on the two incident vertices.
- Interchange process $\{\sigma_t : t \geq 0\}$.

Shuffling cards on graphs

- ▶ When G is the complete graph $K_n \Rightarrow$ Random transposition shuffle;




Shuffling cards on graphs

- ▶ When G is the complete graph $K_n \Rightarrow$ Random transposition shuffle;



- ▶ When G is the graph $P_n \Rightarrow$ Adjacent transposition shuffle;



Shuffling cards on graphs

- ▶ When G is the complete graph $K_n \Rightarrow$ Random transposition shuffle;



- ▶ When G is the graph $P_n \Rightarrow$ Adjacent transposition shuffle;



- ▶ The cutoff has been proven only for: K_n , P_n and star graphs ($n \log n$)



until now...

Shuffling cards on graphs

- We proved that the interchange process on any graphs G_n presents cutoff at times

$$\frac{1}{2\lambda_1^n} \log |V_n|,$$

whenever the G_n is a discretization of a compact set on \mathbb{R}^d .

Shuffling cards on graphs

- ▶ We proved that the interchange process on any graphs G_n presents cutoff at times

$$\frac{1}{2\lambda_1^n} \log |V_n|,$$

whenever the G_n is a discretization of a compact set on \mathbb{R}^d .

- ▶ Above $-\lambda_1^n$ stands for the largest-nonzero eigenvalue of the Laplacian operator Δ_n on G_n which is given by

$$\Delta_n f(x) = \sum_{y \sim x} (f(y) - f(x))$$

for any $f : V_n \rightarrow \mathbb{R}$.

Shuffling cards on graphs

- ▶ We proved that the interchange process on any graphs G_n presents cutoff at times

$$\frac{1}{2\lambda_1^n} \log |V_n|,$$

whenever the G_n is a discretization of a compact set on \mathbb{R}^d .

- ▶ Above $-\lambda_1^n$ stands for the largest-nonzero eigenvalue of the Laplacian operator Δ_n on G_n which is given by

$$\Delta_n f(x) = \sum_{y \sim x} (f(y) - f(x))$$

for any $f : V_n \rightarrow \mathbb{R}$.

- ▶ We call λ_1^n by *spectral gap*.

Shuffling cards on graphs

- ▶ We proved that the interchange process on any graphs G_n presents cutoff at times

$$\frac{1}{2\lambda_1^n} \log |V_n|,$$

whenever the G_n is a discretization of a compact set on \mathbb{R}^d .

- ▶ Above $-\lambda_1^n$ stands for the largest-nonzero eigenvalue of the Laplacian operator Δ_n on G_n which is given by


$$\Delta_n f(x) = \sum_{y \sim x} (f(y) - f(x))$$

for any $f : V_n \rightarrow \mathbb{R}$.

- ▶ We call λ_1^n by *spectral gap*. In spectral graph theory it is called *algebraic connectivity*.

This is more or less what we are assuming on G_n


Σ



\int



Example: adjacent transposition shuffle on a grid



Example: L -adjacent transposition shuffle



The exclusion process on a graph

- ▶ Let us colour ℓ cards in black and $|V_n| - \ell$ cards in red;

The exclusion process on a graph

- ▶ Let us colour ℓ cards in black and $|V_n| - \ell$ cards in red;
- ▶ Define $\eta_t^n : V \rightarrow \{0, 1\}$ as the function that assigns value 1 to vertices with black cards and value 0 to vertices with red cards of permutation σ_t^n .

The exclusion process on a graph


- ▶ Let us colour ℓ cards in black and $|V_n| - \ell$ cards in red;
- ▶ Define $\eta_t^n : V \rightarrow \{0, 1\}$ as the function that assigns value 1 to vertices with black cards and value 0 to vertices with red cards of permutation σ_t^n .

The exclusion process on a graph

- ▶ Let us colour ℓ cards in black and $|V_n| - \ell$ cards in red;
- ▶ Define $\eta_t^n : V \rightarrow \{0, 1\}$ as the function that assigns value 1 to vertices with black cards and value 0 to vertices with red cards of permutation σ_t^n .

$$\eta(x) = \begin{cases} 1, & \text{if the card at vertex } x \text{ is black;} \\ 0, & \text{if the card at vertex } x \text{ is red.} \end{cases}$$

The exclusion process on a graph



The Projection

- ▶ $d_{n,\ell}(t)$ and $t_{\text{mix}}^{n,\ell}(\varepsilon)$: Distance to equilibrium and ε -mixing time of the exclusion process with ℓ particles;

The Projection


- ▶ $d_{n,\ell}(t)$ and $t_{\text{mix}}^{n,\ell}(\varepsilon)$: Distance to equilibrium and ε -mixing time of the exclusion process with ℓ particles;
- ▶ $d_n(t)$ and $t_{\text{mix}}^n(\varepsilon)$: Distance to equilibrium and ε -mixing time of the interchange process.

The Projection

- ▶ $d_{n,\ell}(t)$ and $t_{\text{mix}}^{n,\ell}(\varepsilon)$: Distance to equilibrium and ε -mixing time of the exclusion process with ℓ particles;
- ▶ $d_n(t)$ and $t_{\text{mix}}^n(\varepsilon)$: Distance to equilibrium and ε -mixing time of the interchange process.
- ▶ Observe that a projection can not increase distances.


The Projection

- ▶ $d_{n,\ell}(t)$ and $t_{\text{mix}}^{n,\ell}(\varepsilon)$: Distance to equilibrium and ε -mixing time of the exclusion process with ℓ particles;
- ▶ $d_n(t)$ and $t_{\text{mix}}^n(\varepsilon)$: Distance to equilibrium and ε -mixing time of the interchange process.
- ▶ Observe that a projection can not increase distances.



The Projection

- ▶ $d_{n,\ell}(t)$ and $t_{\text{mix}}^{n,\ell}(\varepsilon)$: Distance to equilibrium and ε -mixing time of the exclusion process with ℓ particles;
- ▶ $d_n(t)$ and $t_{\text{mix}}^n(\varepsilon)$: Distance to equilibrium and ε -mixing time of the interchange process.
- ▶ Observe that a projection can not increase distances.



Thus, $d_{n,\ell}(t) \leq d_n(t)$ and

$$t_{\text{mix}}^{n,\ell}(\varepsilon) \leq t_{\text{mix}}^n(\varepsilon).$$