Bayesian and Frequentist Perspectives in Three Statistical Inverse Problems

Bayesian and Frequentist Perspectives in Three Statistical Inverse Problems

Dissertation

for the purpose of obtaining the degree of doctor
at Delft University of Technology
by the authority of the Rector Magnificus, prof. dr. ir. H. Bijl,
chair of the Board for Doctorates
to be defended publicly on
Friday, 4 December 2026 at 15:00 o'clock

by

Francesco GILI

This dissertation has been approved by the promotors.

Composition of the doctoral committee:

Rector Magnificus,	Voorzitter
Prof. dr. A. W. van der Vaart,	Technische Universiteit Delft, *promotor*
Prof. dr. ir. G. Jongbloed,	Technische Universiteit Delft, *promotor*
Independent members:	
Prof. dr. S. Ghoshal,	North Carolina State University
Prof. dr. K. Ray,	Imperial College London
Prof. dr. D. Anevski,	Lund University
Dr. S. van der Pas,	Vrije Universiteit Amsterdam
Prof. dr. A.J. Cabo,	Technische Universiteit Delft
Reserve:	
Prof. dr. F. Redig,	Technische Universiteit Delft

Dit onderzoek werd gedeeltelijk gefinancierd door de Spinozapremie, toegekend aan A. W. van der Vaart door de Nederlandse Organisatie voor Wetenschappelijk Onderzoek

COLOPHON:
Bayesian and Frequentist Perspectives in Three Statistical Inverse Problems

An electronic copy of this dissertation is available at
`https://repository.tudelft.nl/`.

Contents

1 Introduction **1**
- 1.1 Statistical inverse problems and regularisation . . . 2
- 1.2 Efficiency and minimax lower bounds . . . 2
- 1.3 Uncertainty quantification and BvM theorem . . . 3
- 1.4 How is this thesis built? . . . 6

2 Wicksell's problem: the frequentist nonparametric approach **11**
- 2.1 Introduction . . . 11
- 2.2 Adaptive and Efficient Isotonic Estimation . . . 12
  - 2.2.1 Asymptotics of the Isotonic Inverse Estimator . . . 15
  - 2.2.2 Local Asymptotic Minimax Theorem . . . 24
  - 2.2.3 Conclusion . . . 30
- 2.3 Efficient estimation under local constraint . . . 30
  - 2.3.1 Construction of the informed estimators . . . 31
  - 2.3.2 Asymptotic distributions of the informed estimators . . . 33
  - 2.3.3 Local asymptotic minimax risk under constraint . . . 34
  - 2.3.4 Simulation study . . . 36
  - 2.3.5 Conclusion . . . 38
- 2.4 Appendix . . . 39
  - 2.4.1 Proofs of complemental statements for Theorem 2.2.3. . . . 39
  - 2.4.2 Proofs complemental statements for the LAN expansion. . . . 52
  - 2.4.3 Proofs of Section 2.3. . . . 59
  - 2.4.4 Efficiency Theory Constrained Space . . . 73
- 2.5 Bibliographic remarks . . . 76

3 Wicksell's problem: the Bayesian nonparametric approach **77**
- 3.1 Semiparametric BvM via Isotonized Posterior . . . 77
  - 3.1.1 Introduction . . . 77
  - 3.1.2 Main results . . . 80
  - 3.1.3 Simulation Study and Discussion about UQ . . . 87
  - 3.1.4 Conclusion . . . 92
- 3.2 Posterior Contraction via Hierarchical Modelling . . . 93
  - 3.2.1 Introduction . . . 93
  - 3.2.2 Main results . . . 93
  - 3.2.3 Simulation study . . . 95
  - 3.2.4 Conclusion . . . 98

3.3 Appendix . . . 99
3.3.1 Proofs of complemental statements for Semiparametric BvM . . . 99
3.3.2 Proofs of complemental statements for Hierarchical Modelling . . . 115
3.4 Bibliographic remarks . . . 135

4 Deconvolution: semiparametric UQ via isotonized posterior 139
4.1 Introduction . . . 139
4.2 Main results . . . 142
4.3 Simulation study . . . 148
4.4 Conclusion . . . 153
4.5 Appendix . . . 154
4.5.1 Complementary results . . . 154
4.5.2 Review of the integral equation . . . 163
4.6 Bibliographic Remarks . . . 165

5 Proximal inference based on mixtures of linear models 167
5.1 Introduction . . . 167
5.2 Mixture of linear models . . . 169
5.3 Contraction . . . 178
5.3.1 *Priors* . . . 178
5.3.2 *Results* . . . 179
5.4 Conclusion . . . 182
5.5 Appendix . . . 182
5.5.1 Definitions of function spaces and topologies used in proofs . . . 182
5.5.2 Proofs . . . 184
5.5.3 Consistency by entropy . . . 199

Summary 201

Samenvatting 203

Acknowledgements 205

Curriculum Vitæ 209

List of Publications 211

Chapter 1

Introduction

A recurring theme in statistics is that the quantity of scientific interest is not directly observable. Instead, one observes a transformed or corrupted version of it, and the task is to invert the transformation. In a *statistical inverse problem*, the data $Z_1, \ldots, Z_n$ are drawn from a distribution G_0 that is linked to an unknown distribution F_0 through a forward operator K:

$$G_0 = K(F_0).$$

The goal is to recover F_0, or functionals thereof, from the observations. Problems of this form arise across a wide range of scientific disciplines. In astronomy, one cannot observe the full three-dimensional structure of a globular cluster — a spherical aggregation of thousands of stars bound by gravity — but only its two-dimensional projection on the sky. The apparent positions of stars are known functions of their true spatial positions, and recovering the three-dimensional radial distribution of stars from these projected observations requires inverting that geometric transform [2, 115]. In signal processing and epidemiology, a measured signal is the convolution of an unknown source distribution with a known noise kernel (arising when signal and noise are independent additive components), and deconvolution is required to recover the source [33, 98]. In causal inference from observational data, unobserved confounders create a systematic distortion between the observed data distribution and the causal distribution of interest; identifying and estimating the causal effect requires inverting this confounding mechanism [21, 75].



Figure 1.1: Schematic diagram of a statistical inverse problem. The forward operator K maps the latent law F_0 to the observable law G_0. Statistical inference inverts this map from data $Z_1, \ldots, Z_n$. The thesis treats Wicksell's problem (K = stereological transform, Chapters 2–3), nonparametric deconvolution (K = convolution, Chapters 4), and proximal causal inference (K = unobserved confounding mechanism, Chapter 5).

Despite their apparent diversity, these problems share a common mathematical skeleton: a known forward map K, an unknown object F_0, and noisy indirect observations (see Fig. 1.1), and they share thus a common set of statistical challenges.

1.1 Statistical inverse problems and regularisation

The fundamental difficulty of statistical inverse problems is their *ill-posedness*: small perturbations in G_0 can correspond to large perturbations in F_0, because the operator K typically smooths or compresses information. A classical illustration is deconvolution: in the frequency domain, K multiplies by the characteristic function of the noise distribution, which may decay to zero. Inverting K then amplifies high-frequency components, turning small noise into large estimation errors. As a consequence, unconstrained nonparametric estimators of F_0 tend to be erratic: non-monotone, irregular, or with infinite variance at rates too slow for statistical use and thus *regularisation* is essential.

A natural and particularly powerful form of regularisation, which runs through this entire thesis, is the effective use of intrinsically present shape constraints. Distribution functions are monotone; their integral transforms inherit convexity or monotonicity properties; and estimators that respect these constraints are better conditioned. The analytical device used here is *isotonisation*: replacing a naive unconstrained estimator by its projection onto the set of monotone functions, computed as the derivative of the least concave majorant of a primitive. This operation is computationally efficient (it reduces to the Pool-Adjacent-Violators Algorithm [4]), tuning-parameter-free, and, in some cases, as established in this thesis, *asymptotically efficient*.

1.2 Efficiency and minimax lower bounds

A central question, once an estimator has been constructed, is whether it makes optimal use of the information in the data. The theory of statistical efficiency provides the language to answer this question precisely, and minimax theory provides lower bounds that characterise the fundamental difficulty of an estimation problem independently of any particular method. Let us start by briefly recalling efficiency in parametric models versus nonparametric models.

Efficiency. In a parametric model $\{P_\theta : \theta \in \Theta \subset \mathbb{R}^d\}$ with Fisher information matrix $I(\theta)$, the Cramér–Rao inequality asserts that any unbiased estimator T_n of θ satisfies $\mathrm{Var}_\theta(T_n) \geq I(\theta)^{-1}$, in the matrix sense. The classical asymptotic version, due to Hájek and Le Cam (cf. [102], Chapter 8), is sharper and more general: for any *regular* estimator sequence T_n of θ, the limiting distribution L in $\sqrt{n}(T_n - \theta) \to L$ must be a convolution $N(0, I(\theta)^{-1}) * M$ for some probability measure M. The Gaussian component $N(0, I(\theta)^{-1})$ cannot be removed; it is the irreducible cost of estimation, set by the Fisher information. An estimator is *asymptotically efficient* if $L = N(0, I(\theta)^{-1})$, i.e. if $M = \delta_0$: there is no superfluous noise beyond what the data unavoidably carry.

In a nonparametric model, where F_0 ranges over an infinite-dimensional class, there is no single Fisher information matrix. The relevant notion is instead the *efficient asymptotic variance* for estimating a real-valued functional $T(F_0)$. Fix a smooth one-dimensional submodel $\{F_t\}$ passing through F_0 at $t = 0$, differentiable in quadratic mean with score $h \in L_0^2(F_0)$. Suppose the functional T is *pathwise differentiable* at F_0 relative to such paths: the map $t \mapsto T(F_t)$ is differentiable at $t = 0$, with derivative $\dot{T}_{F_0}(h)$ depending linearly and continuously on the score h. The Riesz representation theorem then gives a unique element $\tilde{\ell}_{F_0} \in L_0^2(F_0)$ — the *efficient influence function* — such that $\dot{T}_{F_0}(h) = \langle \tilde{\ell}_{F_0}, h \rangle_{F_0}$ for all smooth directions h. The variance $\sigma^2_{\mathrm{eff}}(F_0) = \mathrm{E}_{F_0}[\tilde{\ell}_{F_0}(Z)^2]$ is the *semiparametric efficiency bound*: no regular estimator of $T(F_0)$ can have asymptotic variance smaller than $\sigma^2_{\mathrm{eff}}(F_0)$ (cf. [7], [102], Chapter 25). An

estimator achieving this bound is *semiparametrically efficient*. The bound is intrinsic: it depends only on F_0, the model and the functional T, not on the estimation method.

Minimax lower bounds. Efficiency bounds are local, attached to a fixed F_0. Minimax lower bounds certify that, over a class of distributions, no estimator can do better than a certain rate or variance; the class may be a fixed nonparametric family (a *global* bound) or a shrinking neighbourhood of F_0 (a *local* bound), and in general we allow it to depend on the sample size, writing $\mathcal{F}_n$. The prototypical statement is

$$\liminf_{n\to\infty} \inf_{T_n} \sup_{F\in\mathcal{F}_n} r_n^2 \, \mathrm{E}_F\big[\, T_n - T(F)\,\big]^2 \geq c > 0,$$

for some sequence $r_n \to \infty$ and constant $c > 0$, where the infimum is over all estimator sequences. Taking $\mathcal{F}_n \equiv \mathcal{F}$ fixed gives a *global* rate lower bound of order r_n^{-1}, uniformly over $\mathcal{F}$; taking $\mathcal{F}_n$ to shrink to F_0 at the appropriate rate gives a *local asymptotic minimax* bound, which recovers the semiparametric efficiency bound of the previous paragraph as its sharp constant. This says that no estimator can estimate $T(F_0)$ at a rate faster than r_n^{-1}, uniformly over $\mathcal{F}$. The proof technique is typically a reduction to a hypothesis testing problem: if the estimation problem were easy, one could distinguish between hypotheses that are hard to separate, a contradiction. When an estimator T_n satisfies $r_n(T_n - T(F_0)) \to N(0, \sigma^2(F_0))$ and $\sigma^2(F_0) = c$ matches the minimax lower bound, the estimator is simultaneously *rate-optimal* and *asymptotically efficient*: it is the best possible in both rate and constant.

A distinctive feature of inverse problems is that optimal estimation rates are often slower than the parametric $\sqrt{n}$. In Wicksell's problem, the optimal rate for estimating $F_0(x)$ is $\sqrt{n}/\sqrt{\log n}$, slower by a logarithmic factor. This logarithmic loss is a direct consequence of the ill-posedness of the Abel inversion: the operator K loses information at a rate proportional to $(\log n)^{1/2}$, and this loss is irrecoverable. In deconvolution, rates can be polynomial in n when the kernel is smooth [33]. These non-standard rates mean that the classical $\sqrt{n}$-asymptotic theory does not apply directly; one must carry out the minimax analysis afresh for each problem.

1.3 How UQ works in the frequentist versus Bayesian world

A point estimator $\hat{F}_n(x)$ answers the question of where $F_0(x)$ likely is, but for scientific inference one also needs to know how reliably. We now give precise definitions of the two main tools for this.

A *confidence set* at level $1 - \alpha$ is a data-dependent set $C_n = C_n(Z_1, \ldots, Z_n)$ satisfying

$$\inf_{F_0\in\mathcal{F}} \mathrm{P}_{F_0}(T(F_0) \in C_n) \geq 1 - \alpha,$$

for all n (or asymptotically as $n \to \infty$), where $T(F_0)$ is the functional of interest and the infimum is over all F_0 in the model class $\mathcal{F}$. The probability is taken over the randomness in the data for fixed F_0: the set C_n is random, and the parameter $T(F_0)$ is fixed. A $(1 - \alpha)$ confidence set contains the true value in at least $1 - \alpha$ of all hypothetical repetitions of the experiment.

Bayesian inference proceeds from a prior distribution Π on the parameter F and is based on the posterior distribution $\Pi(F \in \cdot \mid Z_1, \ldots, Z_n)$, the conditional distribution of the parameter

given the data. A *credible set* at level $1-\alpha$ is a set $D_n = D_n(Z_1, \dots, Z_n)$ satisfying

$$\Pi(F \in \mathcal{F} : T(F) \in D_n \mid Z_1, \dots, Z_n) \geq 1 - \alpha,$$

Here the data are fixed and F is random: the credible set concentrates $1-\alpha$ of the posterior mass on the event that the functional $T(F)$ takes a value in D_n. The Bernstein-von Mises (BvM) theorem is the classical result that connects these two objects in the parametric setting.

Building intuition: UQ in the parametric world. Consider a parametric model $\{P_\theta : \theta \in \Theta \subset \mathbb{R}^d\}$ with Fisher information I_θ and a prior Π that is absolutely continuous with a positive density near θ_0. The BvM theorem (cf. [102], Chapter 10) states that the rescaled posterior distribution of $\sqrt{n}(\theta - \theta_0)$ converges in total variation to a Gaussian:

$$\Big\| \Pi\Big(\sqrt{n}(\theta - \theta_0) \in \cdot \mid X_1, \dots, X_n \Big) - N\Big(I_{\theta_0}^{-1} \Delta_{n,\theta_0},\, I_{\theta_0}^{-1} \Big) \Big\|_{\mathrm{TV}} \xrightarrow{P^n_{\theta_0}} 0,$$

where $\Delta_{n,\theta_0} = n^{-1/2} \sum_{i=1}^n \dot{\ell}_{\theta_0}(X_i)$ is the rescaled score. The key consequences are: (i) the posterior concentrates at the rate $1/\sqrt{n}$, regardless of the prior; (ii) the asymptotic shape is Gaussian with variance $I_{\theta_0}^{-1}$, the Cramér–Rao bound — so the posterior automatically concentrates at the rate and variance of an efficient estimator; (iii) the prior Π disappears in the limit, meaning the result is robust to the choice of prior. Intuitively, the likelihood dominates the prior for large n, and the local shape of the likelihood is always approximately Gaussian around its maximum. For this reason, the BvM theorem is sometimes called the *Bayesian central limit theorem*: just as the sample mean concentrates and Gaussianises by the CLT, the posterior concentrates and Gaussianises by the BvM. The practical upshot is that any $(1-\alpha)$ credible set is asymptotically a valid $(1-\alpha)$ frequentist confidence set, since both are determined by the same limiting Gaussian.

A useful reformulation, sometimes stated as a corollary, shifts the centring from θ_0 to the maximum likelihood estimator $\hat{\theta}_n$. Under the same conditions, if $\hat{\theta}_n$ satisfies $\sqrt{n}(\hat{\theta}_n - \theta_0) - I_{\theta_0}^{-1}\Delta_{n,\theta_0} \to 0$ in $P^n_{\theta_0}$-probability — a condition met by the MLE under standard regularity conditions — then

$$\left\| \Pi(\theta \in \cdot \mid X_1, \dots, X_n) - N\left(\hat{\theta}_n,\, \frac{1}{n} I_{\theta_0}^{-1} \right) \right\|_{\mathrm{TV}} \xrightarrow{P^n_{\theta_0}} 0.$$

In words: the posterior is approximately Gaussian centred at the MLE with covariance $\frac{1}{n} I_{\theta_0}^{-1}$, matching the MLE's own sampling distribution exactly in the limit. This makes the alignment between the Bayesian posterior and the frequentist sampling distribution particularly transparent.

The *frequentist sampling distribution* of an estimator $\hat{\theta}_n$ is the distribution of $\hat{\theta}_n$ induced by imagining the experiment repeated many times: each time, a fresh sample $Z_1, \dots, Z_n \sim P_{\theta_0}$ is drawn and $\hat{\theta}_n$ is computed. The histogram of all those estimates converges to the sampling distribution, which is centred at the truth θ_0 for consistent estimators and narrows at rate $1/\sqrt{n}$ essentially by the central limit theorem. It is a statement about hypothetical replications of the experiment and does not depend on the observed data.

The *Bayesian posterior* $\Pi(\theta \mid Z_1, \dots, Z_n)$, by contrast, is a distribution over the parameter θ given the *actually observed* data. It is usually centred near the maximum likelihood estimate $\hat{\theta}_n$ and encodes the uncertainty about θ from the Bayesian perspective. It describes our belief about θ after seeing the data, and it changes every time new data arrive.



Figure 1.2: The BvM phenomenon as n increases. The red dashed curve is the sampling law of $\hat{\theta}_n$ (centred at θ_0); blue curves are posteriors from separately sampled datasets, each centred at its own $\hat{\theta}_n$. As n grows all curves narrow, but each posterior stays centred at $\hat{\theta}_n$. The posterior width approximately matches the spread of the frequentist sampling distribution, so the 95% credible set (shaded) achieves asymptotically correct frequentist coverage.

These two objects answer different questions: one describes what would happen across many hypothetical experiments, the other describes uncertainty given the single observed dataset. Crucially, a 95% credible set C_n satisfies $\Pi(\theta \in C_n \mid Z_1, \ldots, Z_n) \geq 0.95$ by definition, but there is no guarantee that it will contain the underlying true parameter θ_0. In parametric models, however, the BvM theorem guarantees that the posterior converges in total variation to the same limiting Gaussian as the sampling distribution of $\hat{\theta}_n$, so a $(1-\alpha)$ credible set is asymptotically a valid $(1-\alpha)$ frequentist confidence set.

Why UQ is harder in nonparametric inverse problems. In nonparametric inverse problems the situation is fundamentally harder, for two reasons. First, the rates of convergence are non-standard: in Wicksell's problem the optimal rate for estimating $F_0(x)$ is $\sqrt{n}/\sqrt{\log n}$, slower than $\sqrt{n}$ by a logarithmic factor, so the rescaling in the BvM display above is no longer $\sqrt{n}$. Second, the limiting distributions involve nuisance parameters, like the local density $g_0(x)$ and the local Hölder smoothness γ of F_0, that are themselves difficult to estimate consistently. For a frequentist, constructing a confidence interval therefore requires estimating these nuisance quantities, which introduces additional uncertainty and may lead to under-coverage in practice. Bootstrap procedures are an attractive alternative; while they can fail in non-standard-rate settings [93], a consistent bootstrap does exist for Wicksell's problem.

The Bayesian approach offers a way out. One places a nonparametric prior Π on F_0 and studies the posterior $\Pi(\cdot \mid Z_1, \ldots, Z_n)$. Credible sets are produced automatically as a by-product of posterior computation and do not require the user to estimate nuisance parameters. Whether these credible sets are valid from a frequentist perspective depends on whether a nonparametric analogue of the BvM theorem holds. The rigorous justification is provided by *semiparametric BvM theorems*: under suitable conditions, the posterior for a functional $T(F_0)$, after centring at an efficient estimator and rescaling at the correct rate, converges to the same Gaussian as the sampling distribution of that estimator. While such results require considerably more work than in the parametric case [13], in the inverse-problem settings treated in this thesis, a BvM theorem takes the form: there exists an efficient estimator $\hat{T}_n$ and

a variance σ^2 such that

$$\sup_B \left| \Pi\Big(r_n \left(T(F) - \hat{T}_n\right) \in B \;\Big|\; Z_1, \ldots, Z_n \Big) - \Phi_{\sigma^2}(B) \right| \xrightarrow{\mathrm{P}_{F_0}} 0,$$

where r_n is a rate, Φ_{σ^2} is the centred Gaussian with variance σ^2 and the supremum is over Borel sets B. The variance σ^2 is the asymptotic variance of the efficient estimator and does not involve nuisance parameters beyond those already determined by F_0 itself. Figure 1.2 illustrates schematically how the BvM theorem aligns the posterior with the frequentist sampling distribution, making Bayesian credible sets into asymptotically valid frequentist confidence sets.

1.4 How is this thesis built?

The organisation of the this is as follows.

Chapter 2 — Frequentist nonparametric estimation in Wicksell's problem. Wicksell's problem originates in the medical setting: it was introduced to model the size distribution of spherical tumours in the spleen from post-mortem cross-sections. More generally, in materials science and stereology [115] we have the following setting: spherical particles of unknown radii are embedded in an opaque medium, and the only observable data are the radii of the circles produced when a plane cuts through the medium. Very interestingly, the same underlying mathematical structure appears in astronomy [2, 92]: one observes only the two-dimensional projected positions of stars in a globular cluster and wishes to recover the three-dimensional radial distribution. In both settings, the forward operator maps the distribution F_0 of squared sphere radii to the distribution G_0 of squared circle radii through an Abel-type integral equation, and the inversion formula involves the functional $V_0(x) = \int_x^\infty g_0(z)(z-x)^{-1/2}\,dz$. The nonparametric estimation of F_0 in this setting was pioneered by Groeneboom and Jongbloed [52], who established rates of convergence for an isotonic estimator; the question of asymptotic efficiency was subsequently studied by Golubev and Levit [49], who identified the optimal asymptotic variance. The naive empirical plug-in estimator of V_0 is non-monotone and erratic; its isotonised version, the *isotonic inverse estimator* (IIE), is monotone by construction. Chapter 2 establishes that the IIE adapts automatically to the *local* Hölder smoothness $\gamma > 1/2$ of F_0 at the estimation point and at zero, without requiring knowledge of γ, and that it attains the minimax lower bound, making it the uniquely efficient nonparametric estimator in this problem. A striking consequence is that when F_0 is flat near the estimation point, the IIE automatically attains the faster $\sqrt{n}$ rate — a phenomenon with no analogue in standard nonparametric density estimation.

Chapter 3 — Bayesian nonparametric approaches to Wicksell's problem. In this chapter, we develop two distinct Bayesian nonparametric approaches to Wicksell's problem. The first, and main, approach places a Dirichlet Process prior directly on the *observable* distribution G_0, which is conjugate and computationally explicit. The key innovation is to isotonise the posterior draws, projecting each draw through the inversion and isotonisation pipeline, to obtain the *Isotonized Inverse Posterior* (IIP), following the projection-posterior methodology of Chakraborty and Ghosal [17, 18]. The main results are semiparametric BvM theorems: the IIP concentrates at the optimal rate $\sqrt{n}/\sqrt{\log n}$ with a variance that adapts to the local smoothness $\gamma > 1/2$, without estimating any nuisance parameter, and the resulting credible sets have asymptotically correct frequentist coverage. These results were published in the

Annals of Statistics [45]. The second approach places a Dirichlet Process prior directly on *the unknown* F_0 itself, yielding a hierarchical Dirichlet-Beta model. Here the posterior is a mixture of Dirichlet processes, and a BvM theorem is currently out of reach; instead, posterior consistency and contraction rates are established for this more classical Bayesian formulation.

Chapter 4 — Semiparametric UQ via isotonized posterior for deconvolutions. The deconvolution problem on the half-line arises when one observes $Z_i = X_i + Y_i$, where $0 \leq X_i \sim F_0$ is the unknown quantity of interest and Y_i follows a known distribution with density k on $[0, \infty)$. Applications include back-calculation of historical HIV infection rates in epidemiology [4], image restoration [73, 87], astronomical spectroscopy, and seismology. Chapter 4 develops the isotonized-posterior methodology for this setting. A Dirichlet Process prior is placed on the observable distribution G_0 and the posterior draws are isotonised, producing the IIP $\hat{F}_G$. The main result is an analogue of the asymptotic distribution theorem of van Es, Jongbloed, and van Zuijlen [109] for the Bayesian procedure: $n^{1/3}(\hat{F}_G(x) - F_0(x))$ converges, conditionally on the data, to the *Bayesian Chernoff distribution* introduced by Chakraborty and Ghosal [17] — the same distribution that governs isotonic regression posteriors. This parallels exactly the cube-root Chernoff asymptotics of the frequentist IIE, establishing a precise frequentist-Bayesian correspondence in a non-Gaussian, non-$\sqrt{n}$ regime. Because the limiting distribution is non-Gaussian, classical BvM theory does not apply; valid credible sets require a known recalibration of the type developed by Chakraborty and Ghosal in [17], and the question of efficiency in any rigorous sense remains open.

Chapter 5 — Proximal causal inference via mixtures of linear models. The final chapter addresses causal inference in the presence of unmeasured confounding, a challenge that arises in observational studies in medicine, economics, and the social sciences. When treatment and outcome are both affected by a latent confounder U, standard regression does not recover the causal effect. *Proximal inference* [21, 75] provides a solution by exploiting proxy variables: a negative-control exposure Z (not causing the outcome Y) and a negative-control outcome W (not causally affected by treatment A). Under suitable completeness conditions, the average causal effect of A on Y is identifiable from the observed distribution of (A, W, Y, Z) via a *bridge function*. Existing identification results rely on strong linearity or categoricity assumptions. Chapter 5 introduces a causal model based on mixtures of linear models for the regression of Y on the unmeasured confounder, a strategy also adopted concurrently and independently in the related work of Hu, Zorzetto, and Dominici [58]. Under this model the bridge function exists and has a closed-form expression, extending identifiability to a substantially richer class of outcome models. A Bayesian nonparametric estimator is proposed, based on a probit stick-breaking process with covariate-dependent mixing weights, and a posterior consistency theorem for the average treatment effect is proved by establishing Kullback-Leibler support and sieve entropy conditions for the dependent mixture prior.

In an ideal world: a programme for efficiency and UQ. The ideal approach to each inverse problem would follow five steps.

(i) **Construct a shape-constrained frequentist estimator.** Compose the empirical plug-in with an isotonisation or regularisation step to obtain a monotone, tuning-parameter-free estimator of the functional of interest.

(ii) **Derive the limiting distribution.** Establish that the estimator, after appropriate centring and rescaling by a rate r_n, converges in distribution to a known limit law. In the Gaussian

case this yields an asymptotic variance $\sigma^2(F_0)$ that may depend on local properties of F_0 — such as the local Hölder smoothness γ or the density $g_0(x)$ — but the estimator should adapt to these automatically, without knowing γ or g_0.

(iii) **Prove a matching minimax lower bound.** Show that no estimator sequence can have a smaller asymptotic mean squared error than $r_n^{-2}\,\sigma^2(F_0)$, uniformly over a suitable class $\mathcal{F}$. Together with step (ii), this certifies the estimator as asymptotically efficient: it is the best possible in both rate and constant.

(iv) **Construct a Bayesian counterpart.** Place a nonparametric prior on the observable distribution G_0 and apply the same isotonisation pipeline to the posterior draws, obtaining an *Isotonized Inverse Posterior* (IIP) that is computationally no harder than the frequentist estimator.

(v) **Prove a Bernstein-von Mises theorem.** Show that the IIP concentrates at the same rate r_n and with the same variance $\sigma^2(F_0)$ as the efficient frequentist estimator. By step (iii), the posterior is then *Bayes-efficient*: its $(1-\alpha)$ credible sets are asymptotically valid $(1-\alpha)$ frequentist confidence sets at the minimax-optimal level, free of nuisance parameters.

Unfortunately we do not live in an ideal world and bringing this program to completion is a considerable undertaking that not always is possible, also because some mathematical concepts are not yet clear (!), e.g. the concept of efficiency for non-Gaussian limiting distributions.

Wicksell's problem (Chapters 2–3). This is the setting where the complete chain is established. Chapter 2 constructs the IIE, derives its limiting distribution, and proves the matching minimax lower bound, establishing the IIE as asymptotically efficient. Chapter 3 then constructs the Bayesian counterpart (the IIP) and proves a semiparametric BvM theorem showing that the posterior concentrates with the same asymptotic variance, making the credible sets asymptotically valid confidence sets at the minimax-optimal level.

Deconvolution (Chapter 4). For the deconvolution problem on the half-line, the situation is different. The asymptotic distribution of the isotonic inverse estimator was already established by van Es, Jongbloed, and van Zuijlen [109], so step (ii) of the programme was known prior to this thesis. Crucially, however, the limiting distributions in the deconvolution setting are in general *not Gaussian*: depending on the kernel and the smoothness of F_0, one encounters non-standard limit laws. This means that the standard theory of semiparametric efficiency — which is built around Gaussian limits and Fisher information — does not directly apply, and the question of whether the IIE is efficient in any rigorous sense remains open in the literature. The contribution of this thesis, developed in Chapter 4, is to show that the Bayesian IIP behaves asymptotically in the same way as the frequentist IIE. The two procedures do not agree exactly in distribution: the posterior requires a known calibration in order for its credible sets to achieve correct frequentist coverage, of the type studied by Chakraborty and Ghosal in the related monotone regression setting [17, 18]. Steps (iii) (minimax lower bound matching the Bayesian variance) and a full treatment of efficiency thus remain open.

Causal inference (Chapter 5). Proximal causal inference is a relatively nascent field, and many fundamental questions like identification conditions, optimal estimation rates and efficiency bounds remain unsettled in the literature [21, 75]. Chapter 5 takes a more modest first step: it introduces a flexible causal model based on mixtures of linear models, establishes identifiability of the average treatment effect in closed form, proposes a Bayesian nonparametric estimator,

and proves that this estimator is consistent. The questions of efficient estimation, asymptotic distributions, and UQ in the proximal causal setting are, to the best of my knowledge, still to these days open problems.

Scope of this thesis. The common thread of this thesis is the study of Bayesian nonparametric procedures for statistical inverse problems. For each problem considered, we design a procedure, typically by placing a nonparametric prior on the observable distribution and isotonising the posterior draws, and then investigate its theoretical properties: posterior consistency, asymptotic distribution, and the validity of the resulting uncertainty quantification. Where possible, we compare the Bayesian procedure to its frequentist counterpart and ask whether the two are asymptotically equivalent; where a notion of efficiency is available, we ask whether the procedure is optimal. The answers vary by problem. For Wicksell's problem the analysis can be carried to completion: we obtain an efficient frequentist estimator, a matching minimax lower bound, and a Bayesian procedure that is asymptotically equivalent to the efficient estimator, giving valid and nuisance-parameter-free credible sets. For deconvolution, the Bayesian procedure can be shown to match the frequentist one asymptotically, but efficiency in the classical sense is not defined because the limiting distributions are non-Gaussian. For the causal inference problem, consistency of the Bayesian estimator is the main result, and the deeper questions of asymptotic distribution and efficiency are left open.

Chapter 2

Wicksell's problem: the frequentist nonparametric approach

2.1 Introduction

In the field of stereology, scientists are usually concerned with the study of three-dimensional properties of materials and objects by interpreting their two-dimensional cross-sections. This approach differs from three-dimensional reconstructions, as it allows for the estimation of certain quantities without the need for costly 3D reconstructions. Consider now these two stereological problems. The first one: astronomers are interested in the study of the way stars are distributed in the universe. In this setting, globular clusters — spherical aggregations of stars held together by gravity (cf. [2, 92]) — are a topic of particular focus. But how to determine the distribution of stars within these three-dimensional structures when given only a 2D photo of the clusters? The second one: having material that presents a globular microstructure and that cannot be scanned (e.g. steel) but only sectioned, how can we determine the distribution of particles' size inside the material when given only a limited number of 2D cross sections (cf. [22, 72])?



Figure 2.1: Two examples of sampling in stereological inverse problems. Left panel: The Messier 4 (also designated NGC 6121) is a globular cluster in the constellation of Scorpius. Center and right panel: simulation 3D configuration with spheres and the intersecting plane in material microstructures (from [43]).

Despite their seemingly distant nature, both of these problems share a common thread: the inherent structure of an inverse problem, originally conceptualized by Wicksell (cf. [43, 52, 114, 115]). There, the setting of the opaque material having globular microstructure is

formalized by assuming the embedded particles' radii are i.i.d. realization from a random variable $X^* \sim F^*$ and our observations consist of cross sections of the medium (i.e. "planes cutting the medium").

To formalize further the model note first the following two aspects: the observable radii in the cross sections cannot exceed the radii of the associated sphere; the spheres with bigger radii are more likely to be sampled. From this follows that the distribution of the squared radii of the spheres actually cut is not F^* but a biased version of it; thus we write for the squared radius of the sphere actually cut: $X \sim F$. Since we assume that the samples are gathered uniformly, by Pythagoras Theorem, the observable squared circle radius Z is related to X via: $Z = (1 - U^2)X = YX$, where $U \sim U[0, 1]$ and $Y \sim \text{Be}(1, 1/2)$. In the astronomical application, let $\mathbf{Z} = (Z_{p_1}, Z_{p_2}, Z_{p_3})$ be a spherically symmetric random vector (the three-dimensional position of a star in a galaxy) of which only (Z_{p_1}, Z_{p_2}) can be observed. Under the assumption that the joint of $\mathbf{Z}$ is a function of the distance to the center of the galaxy, astronomers are interested in estimating the distribution of the squared distance of a star to the center of the globular cluster or galaxy $X = Z_{p_1}^2 + Z_{p_2}^2 + Z_{p_3}^2 \sim F$ using $Z = Z_{p_1}^2 + Z_{p_2}^2 \sim G$.

From [52] and [92], it follows that both for the application in materials science and the one in astronomy, the density of the observable squared circle radii $Z \sim G$ is:

$$g(z) = \frac{1}{2m_0} \int_z^\infty \frac{dF^*(x)}{\sqrt{x - z}} \quad \left(= \int_z^\infty \frac{dF(x)}{2\sqrt{x^2 - xz}} \right), \tag{2.1.1}$$

where $0 < m_0 = \int_0^\infty \sqrt{y}\, dF^*(y) < \infty$ is the expected sphere radius under F^*. We can invert the above expression and obtain for $x > 0$:

$$F^*(x) = 1 - \frac{V(x)}{V(0)}, \qquad \text{and} \qquad F(x) = 1 + \frac{2}{\pi} \int_x^\infty \sqrt{s}\, dV(s) \tag{2.1.2}$$

where, for all $x \geq 0$:

$$V(x) := \int_x^\infty \frac{g(z)}{\sqrt{z - x}}\, dz \quad \left(= \int_x^\infty \frac{\pi}{2\sqrt{s}}\, dF(s) \right) \tag{2.1.3}$$

and $F^*(x) = 0$ for all $x \leq 0$. In the following sections, we will often focus on the estimation of V, being the common denominator for the estimation of both F and F^*. We will also show formally how asymptotic results for the estimation of V carry on to the estimation of F^*. Finally it is also possible to see that asymptotic results for the estimation of V at x carry on to the estimation of F due to the following inverse formula (c.f. [92]):

$$F(x) = \frac{2}{\pi} \int_0^\infty \sin^{-1} \left\{ \sqrt{\frac{x \wedge z}{z}} \right\} g(z)\, dz - \frac{2}{\pi} \sqrt{x}\, V(x),$$

where the functional $\int_0^\infty \sin^{-1} \sqrt{\frac{x \wedge z}{z}} g(z)\, dz$ is generally estimable at a faster rate than V, as its variance is finite.

2.2 Adaptive and Efficient Isotonic Estimation

Given a sample $Z_1, \ldots, Z_n$ from the density g and for $\mathbb{G}_n$ the empirical distribution function based on this sample, a naive (empirical plug-in) estimator of V is:

$$V_n(x) := \int_x^\infty \frac{d\mathbb{G}_n(z)}{\sqrt{z - x}}. \tag{2.2.1}$$

The estimator for F^* is then defined by $F_n^*(x) = 1 - V_n(x)/V_n(0)$. As a function of x, this estimator has some undesirable properties, being non-monotone and having infinite discontinuities at a set of points becoming dense in the support of the density g as $n \to \infty$. This led in [52] to the definition of the so-called "isotonic inverse estimator" (IIE) $\hat{F}_n^*$ for F^*, which is obtained by replacing V_n by its isotonic projection. This can be calculated as the right continuous derivative of the least concave majorant of the primitive function of V_n, which is:

$$U_n(x) = \int_0^x V_n(y)\,dy = 2\int_0^\infty \sqrt{z}\,d\mathbb{G}_n(z) - 2\int_x^\infty \sqrt{z-x}\,d\mathbb{G}_n(z).$$

Note that $x \mapsto U(x) = \int_0^x V(y)\,dy$ is a concave and increasing function on $[0,\infty)$, whereas U_n is not. Taking the least concave majorant of U_n on $[0,\infty)$ yields an increasing concave function. Denote by $\hat{V}_n$ the right derivative of this least concave majorant of U_n, and note that this will be a decreasing function on $[0,\infty)$. Then $\hat{F}_n^*$ is given by:

$$\hat{F}_n^*(x) := 1 - \hat{V}_n(x)/\hat{V}_n(0). \tag{2.2.2}$$

This estimator has the important properties of being monotone (see Figure 2.2), right-continuous and does not depend on tuning parameters, unlike other estimators in the literature (cf. [49]).



Figure 2.2: 200 observed draws from g and isotonic estimator for $F \sim \text{Ga}(2, 0.5)$.

The work in this section is closely related to [49] and [52]. In [52], the authors observed a peculiar behavior of the isotonic estimator (IIE) compared to the naive estimator $F_n^*(x) = 1 - V_n(x)/V_n(0)$. Both estimators exhibit the same rate of estimation, $(\log n)^{-1/2}\sqrt{n}$, and normal limiting behavior. However, the naive estimator's asymptotic variance $\sigma^2(F^*) = \frac{4m_0^2}{\pi^2}(g(x) + g(0)(1 - F^*(x))^2)$ is exactly twice that of the IIE. The authors of [49] propose an explanation, showing that for F^* in the Hölder class Λ_γ with $\gamma > \frac{1}{2}$, any efficient estimator attains the variance $\sigma^2(F^*)/(2\gamma)$. In [52], it is assumed that F^* has a bounded density f, which corresponds to $\gamma = 1$, thus explaining the variance reduction by half.

Moreover, the authors of [49] propose an estimator based on Kernel Density Estimation (KDE) that achieves the variance $\sigma^2(F^*)/(2\gamma)$ and which is, to the best of our knowledge, the only other efficient estimator in the literature. However, the estimator they propose depends directly on the degree γ of smoothness, which should be guesstimated before using the estimator, making it non-adaptive. Furthermore, under local conditions at x and 0 (cf. (24) and (25)), they derive an asymptotic minimax lower bound assuming F is in the Hölder class Λ_γ, for $\gamma > \frac{1}{2}$.

This section has two main contributions. On the one hand, we show that the IIE $\hat{F}_n^*$ automatically adapts to the *local* smoothness of F^*, without requiring F^* to belong to a *global*

Hölder class. It suffices to consider the *local* smoothness at points x and 0, measured by constants $\gamma_x > \frac{1}{2}$ and $\gamma_0 > \frac{1}{2}$ (see conditions (2.2.3) and (2.2.36)). These can be interpreted as the tight degree of Hölder smoothness at the points x and 0. For F^* with density f (i.e. differentiable at x and 0) we retrieve the $\gamma_x = \gamma_0 = 1$ that explains the behavior observed in [52]. On the other hand, we prove a lower bound on the minimax risk for any estimator sequence. The minimax theorem holds as well under local conditions of smoothness of F^* around x and 0. If all conditions are met, this shows that the isotonic estimator is asymptotically efficient. No other estimator in the literature simultaneously has these properties. Consequently, the IIE emerges as the unequivocal choice, surpassing alternative estimators previously proposed in Wicksell's problem.

First, we prove that the IIE is asymptotically normal and the smoother the cdf F^* is at x and 0, the smaller the asymptotic variance automatically attained by the IIE (Theorem 2.2.1.a.). Here, the smoothness of F^* at a point p is measured by considering constants $K_p > 0$ and $\gamma_p > \frac{1}{2}$ such that:

$$H_p(\delta) := \int_0^1 (F^*(p + u\delta) - F^*(p))\, du \sim \operatorname{sgn}(\delta)|\delta|^{\gamma_p} K_p, \qquad \text{as } \delta \to 0. \tag{2.2.3}$$

Note that for $p = 0$ we can only allow $\delta > 0$, because $\delta < 0$ would lead to $K_0 = 0$.

The adaptivity extends also to the situation where F^* is constant around x and 0. Then a faster $\sqrt{n}$-rate of convergence is attained (Theorem 2.2.1.b.). Note that previously there were no results in the literature for nonparametric estimators that attained $\sqrt{n}$-rate in Wicksell's problem. In the case in which F^* is constant only around x but not around 0, or vice versa, the rate will be again $(\log n)^{-1/2}\ \sqrt{n}$. We now state the first main result and its proof is given in Appendix 2.4.

Theorem 2.2.1. *Let F^* be the distribution function of the squared sphere radii and g the corresponding density of the squared circle radii according to* (2.1.1)*. Assume $\int_0^\infty y^{\frac{3}{2}}\, dF^*(y) < \infty$ and choose $x > 0$.*

a. If F^* satisfies condition (2.2.3) *both at* 0 *and at x for $\gamma_0 > \frac{1}{2}$ and $\gamma_x > \frac{1}{2}$ respectively and $g(x) < \infty$ holds true, then, as $n \to \infty$:*

$$\sqrt{\frac{n}{\log n}}\left(\hat{F}_n^*(x) - F^*(x)\right) \rightsquigarrow N\left(0, \frac{4m_0^2}{\pi^2}\left(\frac{g(x)}{2\gamma_x} + (1 - F^*(x))^2 \frac{g(0)}{2\gamma_0}\right)\right).$$

b. If F^* is constant around x and 0*, with K_x and K_0 the biggest compact intervals in $[0, \infty)$ that contain x and* 0*, respectively, so that F^* is constant, on K_x and K_0, with $K_x \cap K_0 = \emptyset$, then, as $n \to \infty$:*

$$\sqrt{n}\left(\hat{F}_n^*(x) - F^*(x)\right) \rightsquigarrow \frac{2m_0}{\pi}\Big(L_x + (1 - F^*(x))L_0\Big),$$

where, for any $a, b \in \mathbb{R}$, the joint distribution of L_x and L_0 is given by:

$$\mathrm{P}\left(L_x \le a, L_0 \le b\right) = \mathrm{P}\Big(\operatorname*{arg\,max}_{s \in K_x}\{\mathbb{Z}_x(s) - as\} \le x, \operatorname*{arg\,max}_{s \in K_0}\{\mathbb{Z}_0(s) - bs\} \le 0\Big).$$

Here $\mathbb{Z}_p(s) = \mathbb{Z}(s) - \mathbb{Z}(p)$, for $s \ge 0$ and $p \in \{0, x\}$. $\mathbb{Z}$ is a continuous centered Gaussian Process with covariance structure given by:

$$\operatorname{Cov}\left(\mathbb{Z}(t), \mathbb{Z}(s)\right) = \operatorname{Cov}\Big(2(\sqrt{Z} - \sqrt{(Z_t)_+}), 2(\sqrt{Z} - \sqrt{(Z_s)_+})\Big), \tag{2.2.4}$$

where $Z_t = Z - t$ and $Z_s = Z - s$.

The second main result of this section consists of showing that the IIE is asymptotically efficient in this setting, i.e. the attained asymptotic variance is the smallest attainable. This kind of result is obtained by proving a minimax lower bound for any estimator sequence estimating $F^*(x)$. Here we give a weaker (but perhaps more insightful) version of this result and its proof is given in Appendix 2.4.

Theorem 2.2.2. *Let the conditions of Theorem 2.2.1.a. hold, assume that F^* for such γ_0, γ_x satisfies* (2.2.36) *and that g is continuous at x and 0. Let ℓ be a bowl-shaped loss function, i.e., ℓ is non-negative, symmetric and subconvex, then for any estimator sequence $(F_n^*(x))_{n\in\mathbb{N}}$:*

$$\lim_{\delta\downarrow 0} \liminf_{n\to\infty} \sup_{\|F'-F^*\|_\infty<\delta} \mathrm{E}_{G_{F'}}\, \ell\left(\sqrt{\frac{n}{\log n}}\,(F_n^*(x) - F'(x))\right) \geq \mathrm{E}\ell(L), \tag{2.2.5}$$

where L possesses the normal distribution in Theorem 2.2.1.a.. $G_{F'}$ is the probability measure associated to the density g in (2.1.1) *with F^* replaced by F'.*

Combining the above results shows that the IIE is an easy-to-compute, adaptive, and efficient estimator for F^*.

2.2.1 Asymptotics of the Isotonic Inverse Estimator

In this subsection, we prove two main theorems that will imply Theorem 2.2.1. In order to prove Theorem 2.2.1.a., it is enough to look at the behavior of $\hat{V}_n(x)$ and then apply the same result also to $\hat{V}_n(0)$. We now state the conditions needed for Theorem 2.2.1 and 2.2.3. First, a natural assumption is a moment constraint for Z, which can be translated into a moment condition on X^* (for $0 < m_0 < \infty$):

$$\int_0^\infty z\, g(z)\, dz = \frac{2}{3m_0} \int_0^\infty y^{\frac{3}{2}}\, dF^*(y) < \infty. \tag{2.2.6}$$

Additionally, for $x \geq 0$, we need our version of g to be well defined, i.e.:

$$\frac{1}{2m_0} \int_x^\infty \frac{dF^*(s)}{\sqrt{s-x}} < \infty. \tag{2.2.7}$$

Furthermore, we measure the smoothness of F^* at x; $\exists\, K > 0$ and $\gamma_x > \frac{1}{2}$ such that (2.2.3) is satisfied.

Before proving Theorem 2.2.1.a., we show that condition (2.2.3) entails the local smoothness of the distribution function F by looking at the following examples 2.2.1 and 2.2.2 below.

Example 2.2.1. For $\gamma_x > \frac{1}{2}$ consider the class of distribution functions locally behaved as follows. Let F^* be so that there exists a neighborhood U_x of $x > 0$ and some $K > 0$:

$$F^*(u) = \begin{cases} F^*(x) + K(u-x)^{\gamma_x}, & \text{for } u \geq x,\ u \in U_x \\ F^*(x) - K(x-u)^{\gamma_x}, & \text{for } u < x,\ u \in U_x \end{cases}$$

One can think of such distribution functions as tightly Hölder smooth of degree γ_x at the point x. This class is of particular importance because it illustrates what type of functions we are thinking of when we derive results under smoothness conditions at the point x and 0. Such

functions satisfy both conditions (2.2.3) and (2.2.20). Condition (2.2.20) follows immediately. For condition (2.2.3) note that as $|\delta| \downarrow 0$, small enough so that $[x - |\delta|, x + |\delta|] \subset U_x$:

$$H_x(\delta) = \int_0^1 (F^*(x + u\delta) - F^*(x))\, du = \operatorname{sgn}(\delta)|\delta|^{\gamma_x} K \int_0^1 u^{\gamma_x}\, du + o(\delta).$$

Example 2.2.2. For any $x > 0$, let μ_x be the discrete probability measure given by:

$$\mu_x = \frac{1}{2} \sum_{i=1}^{\infty} p_i \left(\delta_{x+t_i} + \delta_{x-t_i}\right),$$

for sequences p_i of probability weights and $t_i \downarrow 0$. In particular set: $p_i = \frac{6}{\pi^2} i^{-2}$ and $t_i = i^{-\frac{1}{\gamma_x}}$ where $\gamma_x > 1/2$ and $\delta_x(\cdot)$ indicates the Dirac measure at x. Denote by F^* the associated cdf, so that, because $\gamma_x > 1/2$, we have:

$$\int_x^{\infty} \frac{dF^*(s)}{\sqrt{s - x}} = \frac{3}{\pi^2} \sum_{i=1}^{\infty} \frac{1}{i^2} \frac{1}{\sqrt{t_i}} = \frac{3}{\pi^2} \sum_{i=1}^{\infty} \frac{1}{i^{2-\frac{1}{2\gamma_x}}} < \infty.$$

Moreover, we have for $\delta \downarrow 0$ (similar reasonings hold also for a generic sequence $|\delta| \downarrow 0$, we consider $\delta > 0$ for simplicity):

$$H_x(\delta) = \int_0^1 (F^*(x + u\delta) - F^*(x))\, du = \sum_i \frac{p_i}{2} \int_0^1 \mathbf{1}_{t_i \le u\delta}\, du = \sum_i \frac{p_i(\delta - t_i)_+}{2\delta},$$

which means $t_i < \delta$, thus $i > \frac{1}{\delta^{\gamma_x}}$. Therefore $H_x(\delta) = \sum_{i > 1/\delta^{\gamma_x}} \frac{p_i(\delta - i^{-1/\gamma_x})}{2\delta}$. Now we prove the following fact, as $\delta \downarrow 0$ and $k \ge 2$:

$$\frac{1}{(\delta^{\gamma_x})^{k-1}} \sum_{i > \frac{1}{\delta^{\gamma_x}}} \frac{1}{i^k} \xrightarrow{\delta \downarrow 0} \frac{1}{k-1}$$

This follows directly from the fact that for $\alpha \to \infty$ (e.g. $\alpha = 1/\delta^{\gamma_x}$) we have:

$$\frac{1}{k-1} \frac{1}{(\alpha+1)^{k-1}} = \int_{\alpha+1}^{\infty} \frac{1}{x^k} dx \le \sum_{i > \alpha} \frac{1}{i^k} \le \int_{\alpha-1}^{\infty} \frac{1}{x^k} dx = \frac{1}{k-1} \frac{1}{(\alpha-1)^{k-1}}$$

Moreover $(\alpha/(\alpha \pm 1))^{k-1} \to 1$. Therefore as $\delta \downarrow 0$ we have:

$$H_x(\delta) = \frac{6}{\pi^2} \left(\sum_{i > 1/\delta^{\gamma_x}} \frac{1}{i^2} - \frac{1}{\delta} \sum_{i > 1/\delta^{\gamma_x}} \frac{1}{i^{2+\frac{1}{\gamma_x}}} \right) \sim \frac{6}{\pi^2(1+\gamma)} \delta^{\gamma_x}.$$

This demonstrates that the required condition for Theorem 2.2.3 can also be satisfied for a discrete cumulative distribution function.

Remark 2.2.1. Example 2.2.1 shows that *tight* Hölder smoothness of degree γ_x at the point x implies (2.2.3). The implication in the other direction is not true as it is shown by Example 2.2.2, where a distribution that is not Hölder continuous at x satisfies (2.2.3). This example also shows that (2.2.3) does not imply Hölder continuity of F^* at x (i.e. $|F^*(x+\delta) - F^*(x)| = O(|\delta|^{\gamma_x})$ as $\delta \to 0$). The condition $|F^*(x + \delta) - F^*(x)| = O(|\delta|^{\gamma_x})$ as $\delta \to 0$ is also not sufficient for Theorem 2.2.3 because $|F^*(x + \delta) - F^*(x)| = O(|\delta|^{\gamma_x})$ only gives information about the upper bound of the difference, but not about the lower bound, which is also needed to be tight Hölder of degree γ_x, as can be seen by inspection of the proof of Theorem 2.2.3.

Theorem 2.2.3. *Let $x \geq 0$, suppose that*(2.2.3)*,* (2.2.6)*, and*(2.2.7) *hold true and $g(x)$ is given by* (2.1.1)*. Then:*

$$\sqrt{\frac{n}{\log n}}\left(\hat{V}_n(x) - V(x)\right) \rightsquigarrow N\left(0, \frac{1}{2\gamma_x} g(x)\right) \quad \text{as } n \to \infty.$$

Proof. We generalize the proof of [52], for $\gamma_x \neq 1$.

First note that (2.2.3), implies that F^* is not constant in a neighborhood of x (c.f. Lemma 2.4.2). Fix $x \geq 0$ such that x satisfies the conditions stated above. Define, for $a > 0$, the process T_n as follows:

$$T_n(a) = \inf\{t \geq 0 : U_n(t) - at \text{ is maximal }\}.$$

The relation between T_n and $\hat{V}_n$ is given by the switch relation (see [53]):

$$T_n(a) \leq t \quad \Leftrightarrow \quad \hat{V}_n(t) \leq a. \tag{2.2.8}$$

From this relation (Lemma 3.2, [53]), it follows that for each vanishing sequence (δ_n) of positive numbers, for $a \in \mathbb{R}$, for $\delta_n^* = \delta_n^{1/\gamma_x}$ (γ_x as in (2.2.3)), and all n sufficiently large:

$$\begin{aligned} \delta_n^{-1}\left(\hat{V}_n(x) - V(x)\right) \leq a \quad &\Leftrightarrow \quad \delta_n^{-1}\left(T_n\left(a_0 + \delta_n a\right) - x\right) \leq 0 \\ &\Leftrightarrow \quad (\delta_n^*)^{-1}\left(T_n\left(a_0 + \delta_n a\right) - x\right) \leq 0. \end{aligned}$$

We aim to study the probability of the event on the left-hand side. We write a_0 for $V(x)$, which is fixed. The introduction of $\delta_n^* = \delta_n^{1/\gamma_x}$ is one of the main steps that differentiates the current proof from the one in [52]. Now denote by:

$$I_x := \begin{cases} [0, \infty), & \text{if } x = 0 \\ \mathbb{R}, & \text{if } x > 0 \end{cases}$$

Define the localized process $\tilde{Z}_n$ in $C(I_x)$, the space of continuous functions on I_x, as follows (recall $a_0 = V(x)$):

$$\tilde{Z}_n(t) = \delta_n^{-1}(\delta_n^*)^{-1}\left[U_n\left(x + \delta_n^* t\right) - U_n(x) - a_0 \delta_n^* t\right].$$

For each fixed $a \in \mathbb{R}$ we can, for all n sufficiently large, write

$$\begin{aligned} &(\delta_n^*)^{-1}\left(T_n\left(a_0 + \delta_n a\right) - x\right) \\ &\quad = (\delta_n^*)^{-1}\big(\inf\{x + t \geq 0 : U_n(x+t) - a_0 x + \\ &\qquad\qquad - a_0 t - \delta_n a x - \delta_n a t \text{ is maximal }\} - x\big) \\ &\quad = (\delta_n^*)^{-1} \inf\{t \geq -x : U_n(x+t) - U_n(x) - a_0 t - \delta_n a t \text{ is maximal }\} \\ &\quad = \inf\left\{t \geq -(\delta_n^*)^{-1} x : U_n\left(x + \delta_n^* t\right) - U_n(x) - a_0 \delta_n^* t - \delta_n \delta_n^* a t \text{ is maximal }\right\} \\ &\quad = \inf\left\{t \geq -(\delta_n^*)^{-1} x : \tilde{Z}_n(t) - at \text{ is maximal }\right\}, \end{aligned}$$

where we use that the location of a maximum of a function is equivariant under translations and invariant under multiplication by a positive number and addition of a constant. Thus we analyze the asymptotic behavior of the process:

$$\tilde{Z}_n(t) - at = \delta_n^{-1}(\delta_n^*)^{-1}\left\{U_n\left(x + \delta_n^* t\right) - U_n(x) - U\left(x + \delta_n^* t\right) + U(x)\right\} + \tag{2.2.9}$$

$$+ \delta_n^{-1}(\delta_n^*)^{-1}\left\{U\left(x+\delta_n^* t\right) - U(x) - V(x)\delta_n^* t\right\} - at. \tag{2.2.10}$$

We will now analyze the behavior of the stochastic and deterministic terms (2.2.9) and (2.2.10) separately. Let us start with (2.2.10). The Taylor expansion of U with integral form of the remainder yields:

$$\begin{aligned} U(x+\delta_n^* t) - U(x) - V(x)\delta_n^* t &= \delta_n^* t \int_0^1 V(x+u\delta_n^* t) - V(x)\, du \\ &= -\frac{\delta_n^* t\pi}{2m_0}\int_0^1 F^*(x+u\delta_n^* t) - F^*(x)\, du. \end{aligned}$$

When we look at the limit of $\tilde{Z}_n(t) - at$ we multiply the above expression by $\delta_n^{-1}(\delta_n^*)^{-1} = (\delta_n^*)^{-\gamma_x - 1}$. Therefore we have to look at the limit behavior of:

$$-(\delta_n^*)^{-\gamma_x-1}\frac{\delta_n^* t\pi}{2m_0}\int_0^1 F^*(x+u\delta_n^* t) - F^*(x)\, du = -\frac{|t|^{\gamma_x+1}\pi}{2m_0}(K_x + o(1)), \tag{2.2.11}$$

where we use assumption (2.2.3).

Now we consider (2.2.9). Defining the processes (Z_n) in $C(I_x)$ as

$$\begin{aligned} Z_n(t) &= \delta_n^{-1}(\delta_n^*)^{-1}\left\{U_n\left(x+\delta_n^* t\right) - U_n(x) - U\left(x+\delta_n^* t\right) + U(x)\right\} \\ &= 2\delta_n^{-1}(\delta_n^*)^{-1}\int\left\{\sqrt{z-x}\mathbf{1}_{[x,\infty)}(z) - \sqrt{z-x-\delta_n^* t}\mathbf{1}_{[x+\delta_n^* t,\infty)}(z)\right\} d\left(\mathbb{G}_n - G\right)(z) \end{aligned} \tag{2.2.12}$$

we will now establish convergence in distribution of the process Z_n. In order to do so, we need that $g(x) < \infty$. This follows directly from (2.2.7). Moreover, we need the asymptotic covariance structure of the process Z_n, which is given by:

$$\mathrm{Cov}\left(Z_n(s), Z_n(t)\right) = \frac{1}{2\gamma_x}g(x)st\left(1 - \frac{\log\log n}{\log n}\right) + O\left(\frac{1}{\log n}\right), \tag{2.2.13}$$

for each fixed $s, t \in I_x$, provided that $\delta_n = n^{-1/2}\sqrt{\log n}$. Therefore, using (2.2.13) and the Lindeberg-Feller Central Limit theorem we have that:

$$Z_n(1) \rightsquigarrow N(0, g(x)/(2\gamma_x)), \tag{2.2.14}$$

(the proof of (2.2.13) and the proof of (2.2.14) are provided in Appendix 2.4). On the other hand, by Chebyshev's inequality and (2.2.13), we have, $\forall\ \varepsilon > 0$:

$$\mathrm{P}\left(|Z_n(t) - tZ_n(1)| > \varepsilon\right) \leq \frac{\mathrm{Var}(Z_n(t) - tZ_n(1))}{\varepsilon^2} \to 0.$$

Therefore, the finite-dimensional distributions of Z_n converge weakly to the finite-dimensional distributions of the process: $Z(t) = tX$, in $C(I_x)$, where $X \sim N(0, g(x)/(2\gamma_x))$. We now need a stochastic equicontinuity condition to apply Theorem 2.3 from [65]. Take the class of functions for $\varepsilon > 0,\ M > 0$:

$$\begin{aligned} \mathscr{F}_{n,\varepsilon}^M = \{ &\sqrt{z-x-\delta_n^* t}\mathbf{1}_{[x+\delta_n^* t,\infty)}(z) - \sqrt{z-x-\delta_n^* s}\mathbf{1}_{[x+\delta_n^* s,\infty)}(z) : \\ &: s, t \in I_x, |s-t| < \varepsilon, \max(|s|,|t|) < M\}. \end{aligned} \tag{2.2.15}$$

This class has square integrable natural envelope H_ε that satisfies for all $\varepsilon > 0$ small (cf. Lemma 2.4.1): $\int H_\varepsilon^2(z)g(z)dz \leq g(x+\delta_n^* s \wedge t)\,(\varepsilon\delta_n^*)^2 \log \frac{1}{(\varepsilon\delta_n^*)^2}$ (as proved in Lemma 2.2.1 this class of functions is also VC). Therefore, by using maximal inequality 3.1 (ii) from [65] we obtain:

$$\limsup_{n\to\infty} \mathrm{P}\left\{\sup_{\mathscr{F}_{n,\varepsilon}^M} |Z_n(s) - Z_n(t)| > \eta\right\} \leq \limsup_{n\to\infty} \frac{\mathrm{E}\sup_{\mathscr{F}_{n,\varepsilon}^M} |Z_n(s) - Z_n(t)|^2}{\eta^2}$$
$$\leq \limsup_{n\to\infty} \frac{J(1)^2 g(x+\delta_n^* s \wedge t)}{(\delta_n^*)^2 \log n} (\varepsilon\delta_n^*)^2 \log (\varepsilon\delta_n^*)^{-2} = J(1)^2 \gamma_x^{-1} \eta^{-2} g(x) \varepsilon^2 \overset{\varepsilon\to 0}{\longrightarrow} 0.$$

where the reason why the uniform entropy integral is finite is as in the proof of Lemma 2.2.1. The stochastic equicontinuity condition, together with the convergence in distribution of the finite-dimensional distributions, implies convergence in distribution of Z_n to Z in $(C(I_x), d)$ where d is the metric of uniform convergence on compacta. Next, we need the stochastic boundedness of the sequence T_n given below in (2.2.16) (the proof of which is given in Appendix 2.4). For $a_0 = V(x)$ and $n \to \infty$:

$$T_n\left(a_0 + \delta_n a\right) = V^{-1}\left(a_0\right) + O_p\left(\delta_n^*\right). \tag{2.2.16}$$

One can now apply Theorem 2.7 from [65] to the process $\tilde{Z}_n$.

$$\left(\delta_n^*\right)^{-1}\left(T_n\left(a_0 + \delta_n a\right) - x\right) \rightsquigarrow \underset{t\in I_x}{\arg\max}\left\{tX - \frac{|t|^{\gamma_x+1}\pi}{2m_0} K_x - at\right\}. \tag{2.2.17}$$

But it is immediate to see that, almost surely (Lemma 2.4.4):

$$\underset{t\in I_x}{\arg\max}\left\{tX - \frac{|t|^{\gamma_x+1}\pi}{2m_0} K_x - at\right\} \leq 0 \quad \Longleftrightarrow \quad X \leq a. \tag{2.2.18}$$

Hence, using the two relations above:

$$\mathrm{P}\left\{\delta_n^{-1}\left(\hat{V}_n(x) - V(x)\right) \leq a\right\} = \mathrm{P}\left\{\left(\delta_n^*\right)^{-1}\left(T_n\left(a_0 + \delta_n a\right) - x\right) \leq 0\right\} \overset{n\to\infty}{\longrightarrow}$$
$$\overset{(2.2.17)}{\longrightarrow} \mathrm{P}\left\{\underset{t\in I_x}{\arg\max}\left\{tX - \frac{|t|^{\gamma_x+1}\pi}{2m_0} K_x - at\right\} \leq 0\right\} \overset{(2.2.18)}{=} \mathrm{P}\{X \leq a\}.$$

□

Example 2.2.3. As in [49], it is possible to conduct the analysis in accordance with the global smoothness properties of the underlying distribution function F^*. Consider, for arbitrary $\gamma = \alpha + \beta, \alpha = 0, 1, \ldots, 0 < \beta \leq 1$, the Hölder class C^γ of functions $f(y), y \geq 0$ having finite norm:

$$\|f\|_\gamma = \sup_{y\geq 0} |f(y)| + \sup_{x,y\geq 0} \frac{\left|f^{(\alpha)}(x) - f^{(\alpha)}(y)\right|}{|x-y|^\beta}. \tag{2.2.19}$$

Then assume $F \in C^\gamma$ for $\gamma > \frac{1}{2}$. Moreover, assume:

$$\lim_{t\to 0} \frac{|F^*(x+t) - F^*(x)|}{|t|^\gamma} = K > 0. \tag{2.2.20}$$

Note that it is possible to prove that if $F^* \in C^\gamma$ for $\gamma > 1/2$, then the density of the observations g is bounded and continuous (see Lemma 2.4.3). We show that the combination of (2.2.19) and (2.2.20) implies (2.2.3). Using the Dominated Convergence Theorem in combination with the assumption of the Hölder condition, and (2.2.20), we obtain as $\delta \to 0$:

$$\int_0^1 \frac{F^*(x+u\delta) - F^*(x)}{\operatorname{sgn}(\delta)|\delta|^\gamma u^\gamma} u^\gamma \, du \to K. \tag{2.2.21}$$

Also for the case of Theorem 2.2.1.b., we look first at the behavior of $\hat{V}_n(x)$.

Theorem 2.2.4. *Let $x \geq 0$, and $K_x := [\underline{x}, \bar{x}] \subset [0, \infty)$ the biggest interval that contains x on which F^* is constant. Assume that* (2.2.6) *holds true. Then:*

$$\sqrt{n}\left(\hat{V}_n(x) - V(x)\right) \rightsquigarrow L_x,$$

where, for any $a \in \mathbb{R}$:

$$\mathrm{P}\left(L_x \leq a\right) = \mathrm{P}\left(\arg\max_{s \in K_x} \{\mathbb{Z}_x(s) - as\} \leq x\right).$$

for $\mathbb{Z}_x$ a centered continuous Gaussian Process with covariance structure given by:

$$\mathrm{Cov}\left(\mathbb{Z}_x(t), \mathbb{Z}_x(s)\right) = \mathrm{Cov}\left(2(\sqrt{(Z_x)_+} - \sqrt{(Z_t)_+}), 2(\sqrt{(Z_x)_+} - \sqrt{(Z_s)_+})\right). \tag{2.2.22}$$

where $Z_x := Z - x$ and analogously Z_t and Z_s.

Proof. Using (2.2.8) we obtain for each fixed $a \in \mathbb{R}$, for all n sufficiently large:

$$\begin{aligned}
& \sqrt{n}\left(\hat{V}_n(x) - V(x)\right) \leq a \\
\Leftrightarrow \quad & \inf\{s \geq 0 : U_n(s) - \left(V(x) + a/\sqrt{n}\right)s \text{ is maximal}\} \leq x \\
\Leftrightarrow \quad & \inf\{s \geq 0 : (U_n(s) - U_n(x) - U(s) + U(x)) + \\
& \qquad + (U(s) - U(x) - V(x)(s-x)) - (as)/\sqrt{n} \text{ is maximal}\} \leq x.
\end{aligned}$$

where we use that the location of a maximum of a function is equivariant under translations. Now using the invariance under multiplication by a positive number and addition of a constant, and defining:

$$Z_n(s) := \sqrt{n}(U_n(s) - U_n(x) - U(s) + U(x)) \quad \text{and} \tag{2.2.23}$$

$$h_x(s) := U(x) - U(s) - V(x)(x - s). \tag{2.2.24}$$

we obtain that the above event can be rewritten as:

$$\inf\{s \geq 0 : Z_n(s) - as - \sqrt{n}h_x(s) \text{ is maximal}\} \leq x,$$

Lemma 2.2.1. *Let ℓ^∞ be endowed with uniform norm. Define:*

$$Z_n^*(s) := \sqrt{n}\left(U_n(s) - U(s)\right)$$

then:

$$Z_n^* \rightsquigarrow \mathbb{Z} \quad \text{in } \ell^\infty[0, \infty), \tag{2.2.25}$$

where $\mathbb{Z}$ is a Gaussian Process with a.s. continuous sample paths relative to the Euclidean metric. In particular:

$$\|U_n - U\|_\infty = \sup_{s \geq 0} |U_n(s) - U(s)| \xrightarrow{\mathrm{P}} 0. \tag{2.2.26}$$

The proof of this lemma is given in Appendix 2.4. Consequently:

$$Z_n = Z_n^* - Z_n^*(x) \rightsquigarrow \mathbb{Z} - \mathbb{Z}(x) =: \mathbb{Z}_x \quad \text{in } \ell^\infty[0,\infty), \tag{2.2.27}$$

where also $\mathbb{Z}_x$ is a Gaussian Process with a.s. continuous sample paths relative to the Euclidean metric. By the above computations, we conclude that the limiting behavior of:

$$\widetilde{Z}_n(s) := Z_n(s) - as - \sqrt{n}h_x(s),$$

is determined by the process $\mathbb{W}$ which is defined by:

$$\mathbb{W}(s) = \begin{cases} \mathbb{Z}_x(s) - as, & \text{for } s \in [\underline{x}, \bar{x}] \\ -\infty, & \text{for } s \notin [\underline{x}, \bar{x}] \end{cases}$$

This is because: $h_x(s) \geq 0$ for all $s \in (0, \underline{x}) \cup (\bar{x}, \infty)$, and therefore multiplied by $-\sqrt{n}$ will go to $-\infty$ in case the inequality is strict; however $h_x(s) = 0, \ \forall\ s \in [\underline{x}, \bar{x}]$ and thus on that interval the limiting behavior is completely determined by $\mathbb{Z}_x(s) - as$ (using the limiting behavior of Z_n derived above). Let:

$$\hat{s}_n := \inf\left\{ s \geq 0 \ : \ Z_n(s) - as - \sqrt{n}h_x(s) \ \text{ is maximal} \right\}, \tag{2.2.28}$$

$$\hat{s} := \arg\max_{s \geq 0} \{\mathbb{W}(s)\} = \arg\max_{s \in [\underline{x}, \bar{x}]} \{\mathbb{Z}_x(s) - as\}. \tag{2.2.29}$$

To prove convergence in distribution we use the Portmanteau Lemma and show for every closed subset F of $[0,\infty)$: $\limsup_{n\to\infty} \mathrm{P}(\hat{s}_n \in F) \leq \mathrm{P}(\hat{s} \in F)$.

Let us start by choosing a sequence $\varepsilon_n \downarrow 0$ such that $\sqrt{n}h_x(\underline{x}-\varepsilon_n) \to \infty$ and $\sqrt{n}h_x(\bar{x}+\varepsilon_n) \to \infty$. This exists since $\forall\ \varepsilon > 0$, we have $\sqrt{n}h_x(\underline{x} - \varepsilon) \wedge h_x(\bar{x} + \varepsilon) \to \infty$. Moreover, $h_x(\bar{x} + \varepsilon)$ is an increasing function in ε that has a zero in $\varepsilon = 0$. The behavior is analogous for $h_x(\underline{x} - \varepsilon)$. Therefore it is sufficient to construct a sequence of ε_n such that $h_x(\underline{x}-\varepsilon_n) \wedge h_x(\bar{x}+\varepsilon_n) = O(1/n^\alpha)$ for some $\alpha < 1/2$. For $K_n := [\underline{x}-\varepsilon_n, \bar{x}+\varepsilon_n]$ and for $K := [\underline{x}, \bar{x}]$ (denoted by K_x in the statement):

$$\begin{aligned} \limsup_{n\to\infty} \mathrm{P}(\hat{s}_n \in F) &\leq \limsup_{n\to\infty} \mathrm{P}(\hat{s}_n \in F \cap K_n) + \limsup_{n\to\infty} \mathrm{P}(\hat{s}_n \in K_n^c) \\ &\leq \underbrace{\limsup_{n\to\infty} \mathrm{P}\left(\sup_{s \in F \cap K_n} \widetilde{Z}_n(s) \geq \sup_{s \in K_n} \widetilde{Z}_n(s) \right)}_{(1)} + \underbrace{\limsup_{n\to\infty} \mathrm{P}(\hat{s}_n \in K_n^c)}_{(2)}. \end{aligned} \tag{2.2.30}$$

We look at (2) in (2.2.30) and show it goes to zero. To do this, we need relation (2.2.31) and its proof is given Appendix 2.4. Thus:

$$\sup_{s \in K_n^c} \widetilde{Z}_n(s) = \sup_{s \in K_n^c} \{Z_n(s) - as - \sqrt{n}h_x(s)\} \xrightarrow{\mathrm{P}} -\infty, \tag{2.2.31}$$

implies that:

$$\begin{aligned} \mathrm{P}(\hat{s}_n \in K_n^c) &\leq \mathrm{P}\Big(\sup_{s \in K_n^c} \widetilde{Z}_n(s) \geq \sup_{s \in K_n} \widetilde{Z}_n(s) \Big) \\ &\leq \mathrm{P}\Big(\sup_{s \in K_n^c} \widetilde{Z}_n(s) \geq \sup_{s \in K} \{Z_n(s) - as\} \Big) \to 0. \end{aligned} \tag{2.2.32}$$

The last convergence is a consequence of (2.2.31) and that $Z_n \rightsquigarrow \mathbb{Z}$ on $\ell^\infty(K)$, combined with the continuous mapping theorem (i.e. the term on the left-hand side of the inequality is $O_p(1)$).

Now we argue the behavior of (1). For that, we need an additional convergence in probability given in (2.2.33), and its proof is provided Appendix 2.4.

$$\sup_{s\in F\cap K_n}\left\{Z_n(s) - \sqrt{n}h_x(s) - as\right\} - \sup_{s\in F\cap K}\{Z_n(s) - as\} \xrightarrow{\mathrm{P}} 0. \tag{2.2.33}$$

Using that:

$$\sup_{s\in F\cap K}\{Z_n(s) - as\} \rightsquigarrow \sup_{s\in F\cap K}\{\mathbb{Z}_x(s) - as\},$$

we conclude using Theorem 2.7 (iv) from [102] and (2.2.33):

$$\sup_{s\in F\cap K_n}\left\{Z_n(s) - \sqrt{n}h_x(s) - as\right\} \rightsquigarrow \sup_{s\in F\cap K}\{\mathbb{Z}_x(s) - as\}.$$

We derive the behavior of (1) in (2.2.30). This term can be upper bounded by:

$$\begin{aligned}
&\limsup \mathrm{P}\left(\sup_{s\in F\cap K_n}\left\{Z_n(s) - \sqrt{n}h_x(s) - as\right\} \geq \sup_{s\in K}\{Z_n(s) - as\}\right) \\
&\leq \mathrm{P}\left(\sup_{F\cap K}\{\mathbb{Z}_x(s) - as\} \geq \sup_{s\in K}\{\mathbb{Z}_x(s) - as\}\right) \leq \underbrace{\mathrm{P}(\hat{s}\in K^c)}_{=0} + \mathrm{P}(\hat{s}\in F)
\end{aligned} \tag{2.2.34}$$

where we used the above derivations, together with Lemma 2.2.1 and Theorem 2.7 (v) from [102]. Therefore by (2.2.30) combined with (2.2.32), (2.2.34) and Portmanteau Lemma we obtain the desired convergence in distribution, which can be restated:

$$\begin{aligned}
&\mathrm{P}\left(\sqrt{n}(\hat{V}_n(x) - V(x)) \leq a\right) = \mathrm{P}\Big(\inf_{s\geq 0}\{Z_n(s) - as - \sqrt{n}h_x(s) \text{ is maximal}\} \leq x\Big) \\
&\xrightarrow{n\to\infty} \mathrm{P}\Big(\arg\max_{s\in K}\{\mathbb{Z}_x(s) - as\} \leq x\Big).
\end{aligned}$$

Note that the last expression defines the cdf of a random variable L_x, which is the right derivative of the least concave majorant of the process $\mathbb{Z}_x$ evaluated at x. Indeed, by the switch relation and Lemma 2.4.5, we have $\mathrm{P}\left(\arg\max_{s\in K}\{\mathbb{Z}_x(s) - as\} \leq x\right) = \mathrm{P}\left(L_x \leq a\right)$. □

Note that the proof of Theorem 2.2.1.a. (included in Appendix 2.4) is a consequence of Slutsky's Lemma, Theorem 2.2.3 and 2.2.4, the fact that $1 - F^*(x) = V(x)/V(0)$ together with the asymptotic independence of $\hat{V}_n(x)$ and $\hat{V}_n(0)$. The proof of Theorem 2.2.1.b. follows the same steps, using Theorem 2.2.4.

Remark 2.2.2. Let $x > 0$, and suppose that F^* satisfies condition (2.2.3) both at x and at 0 for a common $\gamma_x > \frac{1}{2}$ and (2.2.6) holds true. Then, by Theorem 2.2.1.a.:

$$\sqrt{\frac{n}{\log n}}\left(\hat{F}_n^*(x) - F^*(x)\right) \rightsquigarrow N\left(0, \frac{g(x)V(0)^2 + g(0)V(x)^2}{2\gamma_x V(0)^4}\right). \tag{2.2.35}$$

The asymptotic variance obtained here coincides with the one obtained in obtained in [49] where it is given a unique γ and the one in [52] with $\gamma_x = 1$.

Simulation study limiting process Theorem 2.2.4

In this subsection, we present some simulations about the distribution of L_x of Theorem 2.2.4; in particular we show how to constructively sample from L_x and that, even if L_x is not normal, its behavior is close to being normal. First, we consider an underlying distribution function F^* of the squared radii as displayed in the left panel of Figure 2.3, which is constant on $[2, 3]$; observe in light blue a histogram of 1000 gathered samples from the associated density of the observations g (note that we obtain observations also on the interval where F^* is constant).



Figure 2.3: Left panel: underlying cdf F^* with samples from g. Right panel: representation of covariance structure of the limiting Gaussian Process $\mathbb{Z}_x$.



Figure 2.4: Left panel: 200 samples from limiting Gaussian Process $\mathbb{Z}_x$. Right panel: zoom in on 15 samples paths, putting into evidence the interval $[2, 3]$.

In this setting, we show some simulations of the distribution of the limiting random variable of Theorem 2.2.4, above denoted by L_x. To do that, we compute the kernel given in (2.2.4) of the limiting Gaussian process $\mathbb{Z}_x$. For the F^* given in the left panel of Figure 2.3, the corresponding kernel is given in the right panel of Figure 2.3. Subsequently, we simulate 200 sample paths from the limiting Gaussian Process $\mathbb{Z}_x$ with the given kernel. These sample paths are shown in Figure 2.4. In the right panel of Figure 2.4 we zoom in on the interval $[2, 3]$ of interest and look at the behavior of 15 selected sample paths. As we would expect, all sample paths pass through $(2.5, 0)$.

In order to simulate the distribution of L_x, we use the fact that such random variable, by the proof of the theorem, is the right derivative of the least concave majorant of the Gaussian Process $\mathbb{Z}_x$ evaluated at $x = 2.5$. With all the gathered samples from L_x we looked at its estimated distribution and below we show some tests about the normality of the distribution of



Figure 2.5: Left panel: ecdf of limiting random variable L_x compared to a suitably fitted normal. Right panel: qq-plot of L_x compared to the fitted normal and p-values of the standard tests.

L_x. With the gathered samples, we see in Figure 2.5 that we could not reject the null hypothesis that L_x is normally distributed.

2.2.2 Local Asymptotic Minimax Theorem

In this subsection, we show that the isotonic estimator (IIE) is asymptotically efficient, in the sense that its asymptotic variance is the smallest attainable. For this reason, we establish a statement about a lower bound of the minimax risk for any estimator sequence. We do this by constructing a parametric perturbation along the least favorable direction in the proximity of the true distribution function F^*.

Let $x > 0$ be the point at which we want to estimate F^*. The existence of a least favorable path is a non-trivial matter. We will be guided by intuition in the construction of this path. We want to estimate is: $F^*(x) = 1 - V(x)/V(0)$. The local behavior at x and 0 of the kernels in the definitions of $V(x)$ and $V(0)$ is what essentially characterizes the estimation in Wicksell's problem and leads to the unusual rate $n^{-1/2}\sqrt{\log n}$. Thus, it should be enough to introduce a local perturbation of F^* around x and 0. However, such perturbation must be tailored according to the local level of smoothness of the true F^* near x and 0. Therefore, we need to identify such a level, and this is done in the following condition (2.2.36).

The level of locality of the condition must be tied to the smoothness itself, because if the true F^* is very rough around x or 0, then we expect a more stringent level of locality (and vice versa). Therefore we define the following intervals close to $p \in \{0, x\}$ for constants $\gamma_0 > \frac{1}{2}, \gamma_x > \frac{1}{2}$ to be determined later:

$$
\begin{aligned}
{}^{r}\mathrm{K}_n^0 &:= \left[n^{-\frac{1}{2\gamma_0}}, \eta_n\right], \\
{}^{l}\mathrm{K}_n^x := \left[x - \eta_n, x - n^{-\frac{1}{2\gamma_x}}\right], \quad {}^{r}\mathrm{K}_n^x &:= \left[x + n^{-\frac{1}{2\gamma_x}}, x + \eta_n\right],
\end{aligned}
$$

where $\eta_n \downarrow 0$, $0 < \eta_n \leq 1$, $\forall\, n$ and $\exists\, \alpha \in (0,1)$ such that $\eta_n \gtrsim \frac{1}{\log^{\alpha}(n)}$. Now, let γ_0, γ_x be the smallest constants greater than $1/2$, such that the true F^*, for $\mathrm{K}_n^p \in \{{}^{r}\mathrm{K}_n^x, {}^{l}\mathrm{K}_n^x, {}^{r}\mathrm{K}_n^0\}$ and therefore $p \in \{0, x\}$ satisfies:

$$
\liminf_{n\to\infty} \inf_{x_1, x_2 \in \mathrm{K}_n^p} \sqrt{n \log n} \frac{|F^*(x_2) - F^*(x_1)|}{\left|\log\left\{\frac{p - x_1}{p - x_2}\right\}\right|} = \infty. \tag{2.2.36}
$$

With condition (2.2.36), our path can be shown to be well-defined and it is the least restrictive we can require. As we naturally would expect, the γ_0, γ_x introduced above contain information about the local roughness of F^* near x and 0. We illustrate this with our previous example 2.2.1 and an additional example.

Example 2.2.4. Let us consider the same class of functions as in Example 2.2.1. These functions can be thought to be tightly Hölder smooth at x and satisfy (2.2.36). Let us consider the interval ${}^r\mathrm{K}_n^x = \left[x + n^{-1/(2\gamma_x)}, x + \log(n)^{-\alpha}\right]$. If $F^*(u) = F^*(x) + K(u-x)^{\gamma_x}$ for $u \in {}^r\mathrm{K}_n^x$, then

$$\begin{aligned}
&\inf_{x_1,x_2\in{}^r\mathrm{K}_n^x} \frac{|F^*(x_2) - F^*(x_1)|}{\left|\log\{(x-x_1)/(x-x_2)\}\right|} = K \inf_{\substack{x_1,x_2\in{}^r\mathrm{K}_n^x \\ x_1<x_2}} \frac{(x_2-x)^{\gamma_x} - (x_1-x)^{\gamma_x}}{\log\{1 + (x_2-x_1)/(x_1-x)\}} \\
&= K \inf_{\substack{x_1,x_2\in{}^r\mathrm{K}_n^x \\ x_1<x_2}} (x_1-x)^{\gamma_x} \frac{[1 + (x_2-x_1)/(x_1-x)]^{\gamma_x} - 1}{\log[1 + (x_2-x_1)/(x_1-x)]} = K\gamma_x \inf_{x_1\in{}^r\mathrm{K}_n^x} (x_1-x)^{\gamma_x} \\
&= K\gamma_x n^{-1/2},
\end{aligned}$$

because $\frac{(1+y)^\gamma - 1}{\log(1+y)}$ is increasing in $y > 0$ with limit γ as $y \downarrow 0$.

Example 2.2.5. Condition (2.2.36), which is similar to the conditions of [49] (cf. (24) and (25)), is a consequence of a stronger on, namely:

$$\liminf_{n\to\infty} \inf \frac{|F^*(x_2) - F^*(x_1)|}{|x_2 - x_1|^{\gamma_x}} > 0,$$

where the infimum is as in (2.2.36). We show that it implies (2.2.36) on ${}^r\mathrm{K}_n^x$:

$$\inf_{x_1,x_2\in{}^r\mathrm{K}_n^x} \frac{F^*(x_2) - F^*(x_1)}{\log\{\frac{x-x_1}{x-x_2}\}} = \inf_{x_1,x_2\in{}^r\mathrm{K}_n^x} \frac{|F^*(x_2) - F^*(x_1)|}{|x_2 - x_1|^{\gamma_x}} \frac{|x_2 - x_1|^{\gamma_x}}{\log\left(1 - \frac{x_2-x_1}{x_2-x}\right)}$$

where, $x_1 < x_2$, both $x_1, x_2 \in {}^r\mathrm{K}_n^x$ and thus $0 < \frac{x_2-x_1}{x_2-x} < 1$ going to zero as $n \to \infty$. The above expression, for all n big enough behaves like in the previous example. Multiplied by $\sqrt{n \log n}$ (2.2.36) diverges to ∞.

The second fundamental assumption to obtain our minimax lower bound is that g is continuous in a neighborhood of x and in a right neighborhood of 0. We will assume this all along. A sufficient condition to obtain this is that F^* is locally Hölder continuous around x and 0 of degrees greater than $\frac{1}{2}$ (see Lemma 2.4.3 in Appendix 2.4). We also note that from (2.2.36) we have $F^*(x) < 1$.

We will now proceed with the construction of the path on the space of distribution functions and verify that it is well defined. The construction of our path is based on ideas from [49]. Condition (2.2.36) gives us the appropriate γ_0, γ_x that we use for the definition of the perturbation. For $u \geq 0$ define:

$$F_{h_n}^*(u) = \frac{1}{D_{h_n}} \left(F^*(u) + h_{1,n} \int_0^u \chi_{1,n}(v)\, dv + h_{2,n} \int_0^u \chi_{2,n}(v-x)\, dv \right), \tag{2.2.37}$$

where for $\gamma_0 > \frac{1}{2}$, $\gamma_x > \frac{1}{2}$, using that $\int_0^\infty \chi_{2,n}(v-x)\, dv = 0$:

$$\begin{aligned}
\chi_{1,n}(v) &= (v)^{-1} \mathbf{1}_{\{n^{-1/(2\gamma_0)} \leq v \leq \eta_n\}}, \\
\chi_{2,n}(v-x) &= (v-x)^{-1} \mathbf{1}_{\{n^{-1/(2\gamma_x)} \leq |v-x| \leq \eta_n\}},
\end{aligned}$$



Figure 2.6: Visualization of $\int_0^u \chi_{2,n}(v-x)dv$ for $x = 2$, $n = 50$ and $\gamma_x = 0.52$, $\eta_n = 1/\sqrt{\log n}$.

$$\begin{aligned} D_{h_n} &= 1 + h_{1,n} \int_0^\infty \chi_{1,n}(v)\, dv + h_{2,n} \int_0^\infty \chi_{2,n}(v-x)\, dv \\ &= 1 + h_{1,n} \int_0^\infty \chi_{1,n}(v)\, dv. \end{aligned}$$

We will see that the truncation levels of the function $1/(v-x)$ proposed here above work, but they may not be unique. The function $\chi_{i,n}$, with similar truncation levels, was introduced for the first time in [49], however there it was not used directly for the perturbation as done here and the assumption $\gamma_0 = \gamma_x$ was used.

We will check that the perturbation $F^*_{h_n}$ is well defined for:

$$h_{i,n} = \frac{h_i}{\sqrt{n \log n}}, \quad h_i \in \mathbb{R}, \quad i = 1, 2. \tag{2.2.38}$$

First, we compute the case of $\chi_{2,n}$, to show how the perturbation looks like:

$$\begin{aligned} \int_0^u \chi_{2,n}(v-x)\, dv &= \log\left(\eta_n^{-1}(x-u)\right)\mathbf{1}_{\left\{x-\eta_n \le u \le x - n^{-\frac{1}{2\gamma_x}}\right\}} + \\ &+ \log\left(\eta_n^{-1}\, n^{-\frac{1}{2\gamma_x}}\right)\left(\mathbf{1}_{\left\{u > x - n^{-\frac{1}{2\gamma_x}}\right\}} - \mathbf{1}_{\{u > x+\eta_n\}}\right) + \\ &+ \log\left(\eta_n^{-1}(u-x)\right)\mathbf{1}_{\left\{x + n^{-\frac{1}{2\gamma_x}} \le u \le x+\eta_n\right\}}. \end{aligned} \tag{2.2.39}$$

Figure 2.6 visualizes the perturbation $\int_0^u \chi_{2,n}(v-x)dv$ in (2.2.39). It is immediate from (2.2.39) that this perturbation is continuous and therefore the summation with F^* results in a function $F^*_{h_n}$ that is right-continuous. Moreover, thanks to the normalization D_{h_n}, $\lim_{u\to+\infty} F^*_{h_n}(u) = 1$ and, from the definition of $F^*_{h_n}$, note that $\lim_{u\to 0} F^*_{h_n}(u) = 0$. What is not evident is if $F^*_{h_n}$ is non-decreasing. Intuitively, for this to be true, we need F^* to be "increasing enough" to compensate for the negative drop introduced by the perturbation. Condition (2.2.36) ensures that this is the case.

Lemma 2.2.2. *For all n sufficiently large, $F^*_{h_n}$ is non decreasing.*

The proof of this lemma is provided in Appendix 2.4. We start by stating the most interesting result of this section which is the so-called LAM (locally asymptotically minimax) theorem and it entails the minimax risk for any estimator sequence. The proof of this theorem requires a few additional technical statements (the LAN expansion and the expression of the derivative of the functional of interest), which will be provided right after.

Theorem 2.2.5 (LAM)**.** *Let γ_0, γ_x be the smallest constants above $\frac{1}{2}$ such that F^* satisfies* (2.2.36)*. Let $J = \frac{\pi^2}{8m_0^2}\operatorname{diag}\left(\frac{1}{\gamma_0 g(0)}, \frac{1}{\gamma_x g(x)}\right)$, $\dot\psi = \left[\frac{1}{2\gamma_0}(1-F^*(x)) \quad -\frac{1}{2\gamma_x}\right]^\top$, g as in* (2.1.1)*, and ℓ be a bowl-shaped loss function, i.e., ℓ is non-negative, symmetric and subconvex. Then for every estimator sequence $(F_n^*(x))_{n\in\mathbb{N}}$:*

$$\sup_{\substack{I\subset\mathbb{R}^2\\ I \text{ finite}}} \liminf_{n\to\infty} \sup_{h\in I} \mathrm{E}_h \ell\left(\sqrt{\frac{n}{\log n}}\left(F_n^*(x) - F_{h_n}^*(x)\right)\right) \geq \mathrm{E}\ell(L). \tag{2.2.40}$$

The first supremum is taken over all finite subsets I of $\mathbb{R}^2$, the expectation E_h is under $G_{F^*_{h_n}}$, the probability measures associated to the perturbed densities g_n in (2.2.41)*, and $L \sim N(0, \dot\psi^\top J^{-1}\dot\psi)$;*

$$\dot\psi^\top J^{-1}\dot\psi = \frac{4m_0^2}{\pi^2}\left(\frac{g(x)}{2\gamma_x} + (1-F^*(x))^2\frac{g(0)}{2\gamma_0}\right).$$

Proof. The proof is an immediate application of Theorem 3.11.5 from [103], using Theorem 2.2.6 and Lemma 2.2.4 given below. □

Now we proceed by proving the technical results needed for the proof of Theorem 2.2.5. First, we show that given the path (2.2.37), we can obtain a LAN expansion (see [102], Chapter 7). In the statement of the LAN Theorem, the perturbed density of the observations Z_i appears. Therefore, using (2.2.37), we derive it before proceeding:

$$\begin{aligned} g_n(z) &= \frac{1}{2m_{h_n}}\int_z^\infty \frac{dF^*_{h_n}(v)}{\sqrt{v-z}} \\ &= \frac{1}{2m_{h_n}D_{h_n}}\left\{\int_z^\infty \frac{dF^*(v)}{\sqrt{v-z}} + h_{1,n}\int_z^\infty \frac{\chi_{1,n}(v)}{\sqrt{v-z}}\,dv + h_{2,n}\int_z^\infty \frac{\chi_{2,n}(v-x)}{\sqrt{v-z}}\,dv\right\}. \end{aligned} \tag{2.2.41}$$

where $m_{h_n} = \int_0^\infty \sqrt{s}\,dF^*_{h_n}(s)$, analogously to m_0.

Theorem 2.2.6 (LAN expansion)**.** *Let γ_0, γ_x be constants above $\frac{1}{2}$ such that F^* satisfies* (2.2.36)*. Let g be continuous in a neighborhood of x and in a right-neighborhood of 0. Then for $h^\top = [h_1, h_2] \in \mathbb{R}^2$, $h_n = \frac{h}{\sqrt{n\log n}}$, $F^*_{h_n}$ as in* (2.2.37) *and accordingly g_n in* (2.2.41)*:*

$$\sum_{i=1}^n \log\frac{g_n(Z_i)}{g(Z_i)} = h^\top\Delta_n - \frac{1}{2}h^\top Jh + o_p(1),$$

where the $o_p(1)$ is under $Z_1, \cdots, Z_n \overset{i.i.d.}{\sim} g$, for $J = \frac{\pi^2}{8m_0^2}\operatorname{diag}\left(\frac{1}{\gamma_0 g(0)}, \frac{1}{\gamma_x g(x)}\right)$ and $\Delta_n \rightsquigarrow N(0, J)$, where:

$$\Delta_n = \begin{bmatrix} \sum_{i=1}^n \frac{1}{m_{h_n}D_{h_n}\sqrt{n\log n}}\left(\frac{1}{2g(Z_i)}\int_{Z_i}^\infty \frac{\chi_{1,n}(v)}{\sqrt{v-Z_i}}\,dv - \int_0^\infty \sqrt{v}\chi_{1,n}(v)\,dv\right) \\ \sum_{i=1}^n \frac{1}{m_{h_n}D_{h_n}\sqrt{n\log n}}\left(\frac{1}{2g(Z_i)}\int_{Z_i}^\infty \frac{\chi_{2,n}(v-x)}{\sqrt{v-Z_i}}\,dv - \int_0^\infty \sqrt{v}\chi_{2,n}(v-x)\,dv\right) \end{bmatrix}.$$

Proof. The structure of the proof of this result is as follows. First, we give the expansions of the functions $\int_z^\infty \frac{\chi_{2,n}(v-x)}{\sqrt{v-z}}\,dv$, $\int_z^\infty \frac{\chi_{1,n}(v)}{\sqrt{v-z}}\,dv$ that will highlight the leading terms that contribute to the limiting behavior.

Next, we use the obtained approximations to write out the Taylor approximation of the logarithm and prove some additional relations that will eventually give the LAN. Finally, we show that the remaining terms are negligible. First of all, using the definition of $F^*_{h_n}$ in (2.2.37), we obtain:

$$\frac{g_n(z)}{g(z)} = \frac{1}{m_{h_n} D_{h_n}} \left\{ m_0 + \frac{h_{1,n}}{2g(z)} \int_z^\infty \frac{\chi_{1,n}(v)}{\sqrt{v-z}}\, dv + \frac{h_{2,n}}{2g(z)} \int_z^\infty \frac{\chi_{2,n}(v-x)}{\sqrt{v-z}}\, dv \right\}.$$

Then using (2.4.18) (see Lemma 2.4.6 proved in Appendix 2.4):

$$\int_z^\infty \frac{\chi_{2,n}(v-x)}{\sqrt{v-z}}\, dv = \left\{ \frac{\pi}{\sqrt{z-x}} + O\left(\eta_n^{-\frac{1}{2}}\right) \right\} \mathbf{1}_{\left\{ n^{-\frac{1}{2\gamma_x}} \le z-x \le \eta_n \right\}} + \tag{2.2.42}$$

$$+ O\left(n^{\frac{1}{4\gamma_x}}\right) \mathbf{1}_{\left\{ -n^{-\frac{1}{2\gamma_x}} \le z-x \le n^{-\frac{1}{2\gamma_x}} \right\}} + O\left(\eta_n^{-\frac{1}{2}}\right) \mathbf{1}_{\{z-x \le -\eta_n\}} +$$

$$+ \left\{ O\left(n^{-\frac{1}{2\gamma_x}} (x-z)^{-\frac{3}{2}}\right) + O\left(\eta_n^{-\frac{1}{2}}\right) \right\} \mathbf{1}_{\left\{ -\eta_n \le z-x \le -n^{-\frac{1}{2\gamma_x}} \right\}}. \tag{2.2.43}$$

The proof of Lemma 2.4.6 has the following relation as an important ingredient:

$$\int_u^\infty \frac{1}{t\sqrt{t-u}}\, dt = \pi \mathbf{1}_{\{u \ge 0\}} u^{-1/2}, \tag{2.2.44}$$

which follows from: $\int_1^\infty \frac{1}{t\sqrt{t-1}} dt = \pi$. We will see that asymptotically the only term that matters is the first one in (2.2.42). It is immediate to see that $\sup_u |\int_0^u \chi_{2,n}(v-x)dv| = O(\log\left(\eta_n^{-1} n^{1/2\gamma_x}\right))$, once multiplied by $h_2/\sqrt{n \log n}$ goes to zero as $n \to \infty$. Note also that using the same computations as in equation (2.2.39) also for $\chi_{1,n}$, it follows that $D_{h_n} = 1 + o(1)$.

Denote by: $A_n^{i,x} := \left\{ n^{-\frac{1}{2\gamma_x}} \le Z_i^x \le \eta_n \right\}$, where $Z_i^x := Z_i - x$. As will be seen, the only contributing term is $\pi \mathbf{1}_{A_n^{i,x}} / \sqrt{Z_i - x}$, we denote all the remainder terms by $R_i^{n,x}$ and it is immediate to see that:

$$R_i^{n,x} = O\left(\eta_n^{-\frac{1}{2}}\right) \left\{ \mathbf{1}_{\left\{ -\eta_n \le Z_i^x \le -n^{-\frac{1}{2\gamma_x}} \right\}} + \mathbf{1}_{\{Z_i^x \le -\eta_n\}} + \mathbf{1}_{A_n^{i,x}} \right\} +$$

$$O\left(n^{-\frac{1}{2\gamma_x}} (x - Z_i)^{-\frac{3}{2}}\right) \mathbf{1}_{\left\{ -\eta_n \le Z_i^x \le -n^{-\frac{1}{2\gamma_x}} \right\}} + O\left(n^{\frac{1}{4\gamma_x}}\right) \mathbf{1}_{\left\{ -n^{-\frac{1}{2\gamma_x}} \le Z_i^x \le n^{-\frac{1}{2\gamma_x}} \right\}}$$

Analogously, we define $R_i^{n,0}$. Using the Taylor expansion of the logarithm with Peano residual, and (2.2.43):

$$\sum_{i=1}^n \log \frac{g_n(Z_i)}{g(Z_i)} = \sum_{i=1}^n \left\{ \frac{1}{m_{h_n} D_{h_n}} \Big[m_0 - m_{h_n} D_{h_n} + \frac{h_{1,n}}{2g(Z_i)} \int_{Z_i}^\infty \frac{\chi_{1,n}(v)}{\sqrt{v - Z_i}}\, dv + \right.$$

$$+ \frac{h_{2,n}}{2g(Z_i)} \int_{Z_i}^\infty \frac{\chi_{2,n}(v-x)}{\sqrt{v-Z_i}}\, dv \Big] - \frac{1}{2} \Big[\frac{m_0 - m_{h_n} D_{h_n}}{m_{h_n} D_{h_n}} + \frac{h_{1,n}}{2m_{h_n} D_{h_n} g(Z_i)} \Big(\frac{\pi \mathbf{1}_{A_n^{i,0}}}{\sqrt{Z_i}}$$

$$+ R_n^{i,0} \Big) + \frac{h_{2,n}}{2m_{h_n} D_{h_n} g(Z_i)} \Big(\frac{\pi \mathbf{1}_{A_n^{i,x}}}{\sqrt{Z_i - x}} + R_n^{i,x} \Big) \Big]^2 + o_p \Big(\Big[\frac{m_0 - m_{h_n} D_{h_n}}{m_{h_n} D_{h_n}} + \tag{2.2.45}$$

$$\left. + \frac{h_{1,n}}{2m_{h_n} D_{h_n} g(Z_i)} \Big(\frac{\pi \mathbf{1}_{A_n^{i,0}}}{\sqrt{Z_i}} + R_n^{i,0} \Big) + \frac{h_{2,n}}{2m_{h_n} D_{h_n} g(Z_i)} \Big(\frac{\pi \mathbf{1}_{A_n^{i,x}}}{\sqrt{Z_i - x}} + R_n^{i,x} \Big) \Big]^2 \Big) \right\}.$$

The Taylor expansion of the logarithm is granted by relation (2.2.46) (see Appendix 2.4 for its proof). For $\zeta_{1,n}(z) := \int_z^\infty \frac{\chi_{1,n}(v)}{\sqrt{v-z}}\,dv$, and for $\zeta_{2,n}(z-x) := \int_z^\infty \frac{\chi_{2,n}(v-x)}{\sqrt{v-z}}\,dv$ and F^* as in (2.2.36); then in probability:

$$\max_{1\le i\le n}\left|\frac{h_{1,n}\zeta_{1,n}(Z_i)}{g(Z_i)} + \frac{h_{2,n}\zeta_{2,n}(Z_i-x)}{g(Z_i)}\right| \to 0. \tag{2.2.46}$$

Now we provide a lemma on the normal behavior of the term inside the first square parenthesis in (2.2.45).

Lemma 2.2.3. *For F^* as in* (2.2.36) *and $h_{i,n}$, $i = 1, 2$ as in* (2.2.38)*:*

$$\frac{1}{m_{h_n}D_{h_n}}\sum_{i=1}^n\left\{\frac{h_{1,n}}{2g(Z_i)}\int_{Z_i}^\infty \frac{\chi_{1,n}(v)}{\sqrt{v-Z_i}}\,dv + \frac{h_{2,n}}{2g(Z_i)}\int_{Z_i}^\infty \frac{\chi_{2,n}(v-x)}{\sqrt{v-Z_i}}\,dv + \right.$$
$$\left. - m_{h_n}D_{h_n} + m_0\right\} \rightsquigarrow N\left(0, \frac{\pi^2}{8m_0^2}\left(\frac{h_1^2}{\gamma_0 g(0)} + \frac{h_2^2}{\gamma_x g(x)}\right)\right). \tag{2.2.47}$$

The proof of this lemma is provided in Appendix 2.4. Now we can partition $m_0 - m_{h_n}D_{h_n}$ into the means showed in the definition of Δ_n (use (2.4.20)). The terms appearing in Lemma 2.2.3 are a linear combination of the components of Δ_n. By Lemma 2.2.3 all such linear combinations have normal behavior. But since we can show the normal convergence for any pair of h_i, $i = 1, 2$, the Cramér-Wold device allows us to obtain the joint asymptotic normality claim for the components of Δ_n.

To determine the behavior of the leading terms inside the second square parenthesis in (2.2.45), i.e. :

$$\sum_{i=1}^n \frac{h_{1,n}^2}{4m_{h_n}^2 D_{h_n}^2 g^2(Z_i)}\frac{\pi^2\mathbf{1}_{A_n^{i,0}}}{Z_i}, \qquad \sum_{i=1}^n \frac{h_{2,n}^2}{4m_{h_n}^2 D_{h_n}^2 g^2(Z_i)}\frac{\pi^2\mathbf{1}_{A_n^{i,x}}}{Z_i - x}, \tag{2.2.48}$$

it is enough to use that (see Appendix 2.4 for its proof):

$$\frac{1}{n\log n}\sum_{i=1}^n \frac{\pi^2}{4m_{h_n}^2 g^2(Z_i)(Z_i-x)}\mathbf{1}_{\left\{\eta_n^{-1}\le \frac{1}{Z_i-x}\le n^{\frac{1}{2\gamma_x}}\right\}} \xrightarrow{\mathrm{P}} \frac{\pi^2}{8\gamma_x m_0^2 g(x)}. \tag{2.2.49}$$

The same result also applies for $A_n^{i,0}$ around 0, and this gives convergence in probability of the terms in (2.2.48). Now by grouping out properly the $h^\top = [h_1, h_2]$, we get the claim, as convergence in probability of the matrix is implied by the convergence of its components.

We now show that all the other terms converge in probability to zero. It is enough to look at multiples of the squared terms and show that these go to zero. Relation (2.2.50) gives that these remaining terms are negligible and its proof is given in Appendix 2.4. For $h_{j,n}$, $j = 1, 2$ as in (2.2.38) and accordingly $s \in \{0, x\}$:

$$\left(\frac{m_0 - m_{h_n}D_{h_n}}{m_{h_n}D_{h_n}}\right)^2 = o\left(\frac{1}{n}\right), \quad \sum_{i=1}^n\left(\frac{h_{j,n}R_n^{i,s}}{2m_{h_n}D_{h_n}g(Z_i)}\right)^2 = o_p(1). \tag{2.2.50}$$

We obtain also:

$$o_p\left(\sum_{i=1}^n\left[\frac{m_0 - m_{h_n}D_{h_n}}{m_{h_n}D_{h_n}} + \frac{h_{1,n}}{2m_{h_n}D_{h_n}g(Z_i)}\left(\frac{\pi\mathbf{1}_{A_n^{i,0}}}{\sqrt{Z_i}} + R_n^{i,0}\right)+\right.\right.$$

$$+\frac{h_{2,n}}{2m_{h_n}D_{h_n}g(Z_i)}\Big(\frac{\pi\mathbf{1}_{A_n^{i,x}}}{\sqrt{Z_i-x}}+R_n^{i,x}\Big)\Big]^2\Big)=o_p(1),$$

by combining (2.2.49) and (2.2.50). □

Remark 2.2.3. If we take $\gamma=\gamma_0=\gamma_x$, the obtained asymptotic covariance looks the same as the one obtained in [49] (cf. page 2418), but our γ_0,γ_x have a different meaning.

For the proof of the LAM (locally asymptotically minimax) theorem, the derivative of the map we are interested in estimating (cf. definition 1.10 in [7]) is needed. In particular, let $\psi : \mathcal{G}\mapsto\mathbb{R}$, where $\mathcal{G}$ is the space of the probability measures G_{F^*} associated to the random variable Z (the observations). In our case, for our fixed estimation point $x>0$, $\psi(G_{F^*})=F^*(x)\in\mathbb{R}$.

Lemma 2.2.4. *For γ_0,γ_x as in Theorem 2.2.6 greater than $\frac{1}{2}$, $h^\top=[h_1,h_2]\in\mathbb{R}^2$, $h_n=\frac{h}{\sqrt{n\log n}}$, $F^*_{h_n}$ as in* (2.2.37), *$\dot\psi=\Big[\frac{1}{2\gamma_0}(1-F^*(x))\quad -\frac{1}{2\gamma_x}\Big]^\top$, as $n\to\infty$:*

$$\sqrt{\frac{n}{\log n}}\Big(\psi(G_{F^*_{h_n}})-\psi(G_{F^*})\Big)=\sqrt{\frac{n}{\log n}}\Big(F^*_{h_n}(x)-F^*(x)\Big)\to h^\top\dot\psi.$$

The proof of this lemma is provided in Appendix 2.4. The combination of Theorem 2.2.6 and Lemma 2.2.4 yields the main result Theorem 2.2.5.

2.2.3 Conclusion

We conclude by stressing the fact that the local asymptotically minimax variance in Theorem 2.2.5 coincides with the one obtained in Theorem 2.2.1.a.. Moreover, as we naturally expect, both the estimator's behavior and the minimax theorem are obtained under some local conditions at x and 0 of the smoothness of F^*. The class of functions that are tightly Hölder continuous at these points as in Example 2.2.1 satisfy the key conditions needed for the estimator and for the minimax theorem. In general, whenever the underlying F^* satisfies both (2.2.3) and (2.2.36), then the isotonic estimator (IIE) will attain the minimax asymptotic variance in an adaptive way, and without the need of external tuning.

2.3 Efficient estimation under local constraint

In this section, we focus on the case in which the underlying cdf F^* (and therefore V) is known to be constant on an interval of positive length. We conduct the analysis on a single interval $[\underline{x},\bar{x}]$, for $\underline{x}<\bar{x}$, of constancy. The same procedures can be applied if the underlying cdf F^* is constant on multiple intervals. Note that even if F^* is constant on $[\underline{x},\bar{x}]$, we do not observe any *gap* in the data. Indeed, the density g in (2.1.1) is positive throughout the set $\{x : F^*(x)<1\}$. In this setting, we compare the Isotonic Inverse Estimator IIE that attains the $\sqrt{n}$-rate of convergence adaptively, *without* using the information that V is constant on $[\underline{x},\bar{x}]$, to three other *informed* estimators that *do use* this information and are therefore constrained to be constant on this interval.

In Section 2.3.2, we then show that all the informed estimators are asymptotically equivalent and normal, but not asymptotically equivalent to the IIE. Next in Section 2.3.3, we derive a lower bound for the local asymptotic minimax risk of any estimator sequence in

this setting, and show that this is attained by the informed estimators. In contrast, the IIE is inefficient. In fact, we show that its (non Gaussian) limit distribution is the convolution of the limit distribution of the informed estimators and a nondegenerate factor. However, the difference between the estimators is small, as we illustrate in a simulation study, in Section 2.3.4.

The scope of this section is to understand the theoretical limits of estimation when information of the constancy of F on $[\underline{x}, \bar{x}]$ is available, with the final objective of gaining a better understanding of the IIE, which does not use this information. As shown in the paper, the situation with known $[\underline{x}, \bar{x}]$ is particularly favourable, because it is characterized by normal limits, which makes the classical efficiency theory applicable (in contrast with the limit attained by the IIE which is non-normal, making even the definition of efficiency unclear). In this sense, the informed estimators serve as theoretical benchmarks. This is in the spirit of the *oracle* estimation theory. We show that they are asymptotically efficient and that the IIE is not. However, our simulation study presents a typical example in which the IIE performs close to the efficient estimators, suggesting that in practice, the loss incurred by using the IIE should be small. Therefore, except in the rare cases in which $\underline{x}$ and $\bar{x}$ are known (e.g. it is known F is the c.d.f. of a discrete random variable with known support points), we do not envision a direct practical utility for these estimators. In all other cases, precisely in view of this section, we would recommend practitioners to use the IIE, as it behaves close enough to the *oracle* informed estimators.

2.3.1 Construction of the informed estimators

In this subsection, we introduce three projection-type *informed* estimators (2.3.6)-(2.3.8). We provide explicit constructions and prove that they solve determined minimization problems. In all that follows, we assume that V is supported on $[0, M]$, so $V(y) = 0$ for all $y \in [M, \infty)$.

Using the "naive estimator" (2.2.1), we construct all the other estimators as repeated projections of V_n into designated spaces. Let $\underline{x} < \bar{x}$, $V_n(x) = 0$ for $x > M$, and a fixed partition of $[0, M]$ into intervals. Let $\mathscr{P} = \{[x_i, x_{i+1}) \,:\, i = 1, \dots, I\}$ be such that $0 = x_0 < x_1 < x_2 < \cdots < x_I = M$, and $[\underline{x}, \bar{x}] = [x_i, x_{i+1}]$ for some i. Define the spaces:

$$\mathcal{V} := \Big\{ V : [0,\infty) \mapsto [0,\infty) \;:\; V \text{ decreasing, right-cont.} \Big\}, \tag{2.3.1}$$

$$\overline{\mathcal{V}}_{\underline{x},\bar{x}} := \Big\{ V : [0,\infty) \mapsto [0,\infty) \;:\; V \text{ right-cont., piecewise-const. on } \mathscr{P} \Big\}, \tag{2.3.2}$$

$$\mathcal{V}_{\underline{x},\bar{x}} := \Big\{ V \in \mathcal{V} \;:\; V \text{ constant on } [\underline{x}, \bar{x}] \Big\}. \tag{2.3.3}$$

All the above-defined spaces are convex cones. We define projections relative to the discrepancy measure $Q^f \,:\, \mathbb{L}_2[0,\infty) \mapsto \mathbb{R}$, for a fixed function $f \in \mathbb{L}_1[0,\infty)$:

$$h \;\mapsto\; Q^f(h) := \int_0^\infty h(s)\,(h(s) - 2f(s))\,ds \tag{2.3.4}$$

Since $\|f - h\|_2^2 = Q^f(h) + \|f\|_2^2$, minimising $h \to Q^f(h)$ is equivalent to minimising the $\mathbb{L}_2$-distance to f in the case that f is square-integrable. We use the discrepancy Q^f, because in our context the naive estimator V_n is *not* square-integrable. We consider as estimators the projections of the naive estimator V_n onto the three spaces (2.3.1)-(2.3.3) and its repeated

projection onto $\mathcal{V}$ and next on $\mathcal{V}_{\underline{x},\bar{x}}$:

$$\hat{V}_n = \arg\min_{V\in\mathcal{V}} Q^{V_n}(V), \tag{2.3.5}$$

$$\hat{V}_n^{\Pi} = \arg\min_{V\in\mathcal{V}_{\underline{x},\bar{x}}} Q^{\hat{V}_n}(V). \tag{2.3.6}$$

$$V_n^{\mathscr{P}} = \arg\min_{V\in\overline{\mathcal{V}}_{\underline{x},\bar{x}}} Q^{V_n}(V), \tag{2.3.7}$$

$$V_n^{\Pi} = \arg\min_{V\in\mathcal{V}_{\underline{x},\bar{x}}} Q^{V_n}(V), \tag{2.3.8}$$

(We note that projecting on a cone and then projecting again onto a subcone does not need to give the same result as projecting directly onto the subcone). Existence and uniqueness of the solutions of these minimization problems can be shown along the lines of Theorem 1.2.1 in [35]. The first estimator $\hat{V}_n$ is the Isotonic Inverse Estimator (IIE) introduced in [52]. The other three estimators are informed estimators that take the local constancy of F^* into account.

The following proposition gives explicit constructions of the estimators (2.3.5)-(2.3.8) (the first statement of the following proposition has been proved in [52]). For a continuous function k on $[0, M]$, the Least Concave Majorant (LCM) and its right-hand side derivative, are given by:

$$k^* := \arg\min\{f \,:\, f(x) \geq k(x) \text{ for } x \in [0, M],\ f \text{ concave}\},$$
$$(k^*)'(x) := \inf_{u<x} \sup_{v\geq x} \frac{k(v) - k(u)}{v - u}.$$

By definition, $k^*(x) \geq k(x)$ for all $x \in [0, M]$, and $k^*(0) = k(0)$, $k^*(M) = k(M)$. We recall here the definitions of the functions U_n and U:

$$U_n(x) = \int_0^x V_n(y)\,dy = 2\int_0^\infty \sqrt{z}\,d\mathbb{G}_n(z) - 2\int_x^\infty \sqrt{z-x}\,d\mathbb{G}_n(z), \tag{2.3.9}$$

$$U(x) = \int_0^x V(y)\,dy = \frac{\pi}{2m_0}\int_0^x (1 - F(y))\,dy. \tag{2.3.10}$$

The last equality follows from the first equation in (2.1.2).

Proposition 2.3.1 (Construction of the estimators)**.** *Let U_n be as in* (2.3.9)*, $\underline{x} < \bar{x}$ with $[\underline{x}, \bar{x}) \in \mathscr{P}$ and $x \in [0, \infty)$. The solutions of the minimizations* (2.3.5)-(2.3.8) *are:*

$$\hat{V}_n(x) = (U_n^*)'(x). \tag{2.3.11}$$

$$\hat{V}_n^{\Pi}(x) = \begin{cases} \frac{U_n^*(\bar{x}) - U_n^*(\underline{x})}{\bar{x}-\underline{x}}, & x \in [\underline{x}, \bar{x}], \\ \hat{V}_n(x), & x \notin [\underline{x}, \bar{x}]. \end{cases} \tag{2.3.12}$$

$$V_n^{\mathscr{P}}(x) = V_n^{(x_i, x_{i+1})}(x), \quad x \in [x_i, x_{i+1}), \quad i = 1, 2, \ldots, I, \tag{2.3.13}$$

where for $x \in [x_i, x_{i+1}) \in \mathscr{P}$:

$$V_n^{(x_i,x_{i+1})}(x) = \frac{U_n(x_{i+1}) - U_n(x_i)}{x_{i+1} - x_i}, \tag{2.3.14}$$

$$V_n^{\Pi}(x) = (\widetilde{U}_n^*)'(x), \tag{2.3.15}$$

where:

$$\widetilde{U}_n(t) := \begin{cases} \frac{U_n(\bar{x})-U_n(\underline{x})}{\bar{x}-\underline{x}}(t-\underline{x}) + U_n(\underline{x}), & t \in [\underline{x}, \bar{x}], \\ U_n(t), & t \notin [\underline{x}, \bar{x}]. \end{cases} \tag{2.3.16}$$

The proof of Proposition 2.3.1 is given in Appendix 2.4 and the asymptotic properties of the constructed estimators are given in Theorem 2.3.1 below.

Remark 2.3.1. For any $v \in [\underline{x}, \bar{x}]$ and $x \in [\underline{x}, \bar{x}]$ with $v \neq x$, we have:

$$U(v) - U(x) - V(x)(v - x) = 0,$$

and thus by evaluating the above relation at $v = \bar{x}$ and $x = \underline{x}$ we get:

$$U(\bar{x}) - U(\underline{x}) - V(x)(\bar{x} - \underline{x}) = 0. \tag{2.3.17}$$

This gives an intuition for (2.3.14) and explains why $V_n^{(\underline{x},\bar{x})}(x)$, i.e. $V_n^{\mathscr{P}}(x)$ for $x \in [\underline{x}, \bar{x})$, can be called "empirical slope" on $[\underline{x}, \bar{x})$. It what follows we use the notation $V_n^{(\underline{x},\bar{x})}(x)$ to indicate $V_n^{\mathscr{P}}(x)$ for $x \in [\underline{x}, \bar{x})$.

The isotonic estimator in (2.3.5) can be implemented using (2.3.11), where the least concave majorant of a function can be computed by classical algorithms like the PAVA algorithm (cf. [53]). Similarly, the solutions to (2.3.6)-(2.3.8) can be easily implemented using (2.3.12), (2.3.14) and (2.3.15). Alternatively, the solution to the minimization problem (2.3.8) can be computed algorithmically using a *profile procedure* as follows. First, fix $a > 0$ and compute the minimizer of $Q^{V_n}(V)$ over the space of decreasing, right-continuous functions V that have value a on $[\underline{x}, \bar{x}]$, and call such minimizer V_n^a. Next we optimize the mapping $a \mapsto Q^{V_n}(V_n^a)$ over $a > 0$ and obtain the desired projection. See [44] for a detailed version of this algorithm.

2.3.2 Asymptotic distributions of the informed estimators

In Theorem 2.2.4, it was shown that if $V \in \mathcal{V}_{\underline{x},\bar{x}}$ and $x \in (\underline{x}, \bar{x})$, then as $n \to \infty$:

$$\sqrt{n}\left(\hat{V}_n(x) - V(x)\right) \rightsquigarrow L_x, \tag{2.3.18}$$

where, for any $a \in \mathbb{R}$: $\mathrm{P}(L_x \leq a) = \mathrm{P}(\arg\max_{s \in [\underline{x},\bar{x}]} \{\mathbb{Z}_x(s) - as\} \leq x)$, for $(x, t) \mapsto \mathbb{Z}_x(t)$ a centered continuous Gaussian Process with covariance structure given by:

$$\mathrm{Cov}\left(\mathbb{Z}_y(t), \mathbb{Z}_x(s)\right) = 4\,\mathrm{Cov}\left(\sqrt{(Z_y)_+} - \sqrt{(Z_t)_+},\, \sqrt{(Z_x)_+} - \sqrt{(Z_s)_+}\right). \tag{2.3.19}$$

where $(\cdot)_+ = \max\{0, \cdot\}$ and $Z_u := Z - u$, $Z \sim g$, $u \in \mathbb{R}$. In the present section, we use the characterizations of the *informed* estimators to obtain the first main result of this section that puts the different estimators in perspective.

Theorem 2.3.1 (Asymptotics)**.** *Let F^* be the distribution function of the squared sphere radii and g the corresponding density of the squared circle radii Z according to* (2.1.1)*. Let $x \geq 0$, and $K := [\underline{x}, \bar{x}]$, for $\underline{x} < \bar{x}$, the biggest interval that contains x on which F^* is constant and let* (2.2.6) *hold true. Then, as $n \to \infty$:*

a. The estimators $V_n^{(\underline{x},\bar{x})}$, V_n^{Π} *and* $\hat{V}_n^{\Pi}$ *are all asymptotically equivalent:*

$$\sqrt{n}\left(V_n^{\Pi}(x) - V_n^{(\underline{x},\bar{x})}(x)\right) = o_p(1), \qquad \sqrt{n}\left(\hat{V}_n^{\Pi}(x) - V_n^{(\underline{x},\bar{x})}(x)\right) = o_p(1).$$

Furthermore, they all attain the same limiting distribution, given by:

$$\sqrt{n}\left(V_n^{(\underline{x},\bar{x})}(x) - V(x)\right) \rightsquigarrow N(0, \sigma^2_{\underline{x},\bar{x}})$$

where:

$$\sigma^2_{\underline{x},\bar{x}} = \mathrm{Var}\left(2\,(\bar{x} - \underline{x})^{-1}\{\sqrt{(Z_{\underline{x}})_+} - \sqrt{(Z_{\bar{x}})_+}\}\right). \tag{2.3.20}$$

b. The isotonic estimator $\hat{V}_n$ *is* ***not*** *asymptotically equivalent to* $V_n^{(\underline{x},\bar{x})}$*, as:*

$$\sqrt{n}\left(\hat{V}_n(x) - V_n^{(\underline{x},\bar{x})}(x)\right) \rightsquigarrow W, \tag{2.3.21}$$

where, for any $a \in \mathbb{R}$, $\mathrm{P}(W \le a) = \mathrm{P}\left(\arg\max_{s\in K}\{\mathbb{Z}(s) - as\} \le x\right)$*, for*

$$\mathbb{Z}(t) = \frac{\bar{x} - t}{\bar{x} - \underline{x}}\mathbb{Z}_{\underline{x}}(t) + \frac{t - \underline{x}}{\bar{x} - \underline{x}}\mathbb{Z}_{\bar{x}}(t), \tag{2.3.22}$$

and $(t, x) \mapsto \mathbb{Z}_x(t)$ *the centered Gaussian Process with covariance structure* (2.3.19)*.*

c. Under the assumptions of Proposition 2.3.2, the sequences

$$\sqrt{n}\left(V_n^{(\underline{x},\bar{x})}(x) - V(x)\right) \text{ and } \sqrt{n}\left(\hat{V}_n(x) - V_n^{(\underline{x},\bar{x})}(x)\right),$$

are asymptotically independent.

The proofs of Theorem 2.3.1.a., 2.3.1.b. and 2.3.1.c. are given in Appendix 2.4. Combining 2.3.1.a., 2.3.1.b. and 2.3.1.c. of the Theorem, we see that the limit L_x of the isotonic estimator in (2.3.18) is the convolution of the normal limit $N(0, \sigma^2_{\underline{x},\bar{x}})$ of the informed estimators and the distribution of W in (2.3.21). Thus the limit distribution of the isotonic estimator is less concentrated than the limit distribution of the informed estimators.

2.3.3 Local asymptotic minimax risk under constraint

In this subsection, we show that the estimator (2.3.7) is asymptotically efficient, in the sense that its asymptotic variance is the smallest attainable. This proves that also (2.3.8) and (2.3.6), which are asymptotically equivalent to (2.3.7), are efficient. The claim is valid in the sense of both the local asymptotic minimax theorem and the convolution theorem (c.f. Chapter 25 of [102]). The key is a LAN expansion for a submodel constructed by perturbing the true function V in the least favorable direction.

We consider the estimation of V at a point x in $(\underline{x}, \bar{x})$, knowing that F^*, and thus V, are constant on this interval (cf. (2.4.58)). The associated version of the density of the observations in Wicksell's problem is expressed in V as:

$$g_V(z) = -\int_z^M \frac{dV(s)}{\pi\sqrt{s - z}} \quad \left(= \frac{1}{2m_0}\int_z^M \frac{dF^*(s)}{\sqrt{s - z}}\right). \tag{2.3.23}$$

We denote the corresponding probability measure by G_V. We dexcribe a so called "least favorable" submodel in terms of V. We assume that the true parameter V satisfies the following assumption.

Assumption. Let $V \in \mathcal{V}$ be supported on $[0, M]$, constant on $[\underline{x}, \bar{x}]$, for $\underline{x} < \bar{x}$ and possess Lipschitz continuous density v on $[\underline{x}, \bar{x}]^c$. Moreover $\exists\ \eta > 0$ such that:

$$\int_{[\underline{x},\bar{x}]^c} \frac{1}{|v(s)|}\, ds + \int_{\underline{x}-\eta}^{\underline{x}} \frac{\log^2(\underline{x}-s)}{|v(s)|}\, ds + \int_{\bar{x}}^{\bar{x}+\eta} \frac{\log^2(s-\bar{x})}{|v(s)|}\, ds < \infty. \tag{2.3.24}$$

Define functions $k^{(\underline{x},\bar{x})}$, h and h_0 by, for $\underline{x} < \bar{x}$:

$$k^{(\underline{x},\bar{x})}(z) := 2\frac{\sqrt{(z-\underline{x})_+} - \sqrt{(z-\bar{x})_+}}{\bar{x}-\underline{x}},$$

$$h(z) := \begin{cases} -\frac{1}{\sqrt{\pi} v(z)} \int_z^M \frac{(k^{(\underline{x},\bar{x})} g_V)'(s)}{\sqrt{s-z}}\, ds, & z \notin [\underline{x}, \bar{x}], \\ 0, & z \in [\underline{x}, \bar{x}], \end{cases} \tag{2.3.25}$$

$$h_0(z) := h(z) + \frac{2}{\pi} \int_0^M \sqrt{s} h(s)\, dV(s). \tag{2.3.26}$$

The *least favorable submodel* is defined by:

$$dV_t(s) = \frac{1}{c_t}(1 + t h_0(s))_+\, dV(s), \quad c_t := -\frac{2}{\pi} \int_0^M \sqrt{s}(1 + t h_0(s))_+\, dV(s). \tag{2.3.27}$$

Note that $\int_0^\infty \sqrt{s}\, dV_t(s) = -\frac{\pi}{2}$, that $c_t \geq 0$ and that the obtained V_t is decreasing. In Appendix 2.4 we show that the score function of the model $t \mapsto G_{V_t}$ at $t = 0$ is the centered function $k^{(\underline{x},\bar{x})}$, the influence function of the informed estimators. Assumption 2.3.3 ensures that indeed h_0 is mapped to $k^{(\underline{x},\bar{x})}$ via the score operator and that $-\frac{2}{\pi} \int h_0^2(s) \sqrt{s}\, dV(s) < \infty$. A detailed explanation is contained in the study of the tangent spaces in the context of our problem in Appendix 2.4.

The next result shows the model $t \mapsto G_{V_t}$ is locally asymptotically normal (LAN, see [102], Chapter 7). This result is fundamental because both Theorem 2.3.1.c. and Theorem 2.3.2 rely on it. The proof of Proposition 2.3.2 is given in Appendix 2.4.

Proposition 2.3.2 (LAN expansion)**.** *Let V_t be as in* (2.3.27) *and $t_n = t/\sqrt{n}$ with $t \in \mathbb{R}$. If* (2.2.6) *and Assumption 2.3.3 hold true, then:*

$$\sum_{i=1}^n \log \frac{g_{V_{t_n}}(Z_i)}{g_V(Z_i)} = t\,\Delta_n - \frac{t^2}{2}\,\sigma^2_{\underline{x},\bar{x}} + o_p(1) \tag{2.3.28}$$

where: $\Delta_n := \frac{1}{\sqrt{n}} \sum_{i=1}^n \left(k^{(\underline{x},\bar{x})}(Z_i) - \mathrm{E}\, k^{(\underline{x},\bar{x})}\right) \rightsquigarrow N(0, \sigma^2_{\underline{x},\bar{x}})$ *and $\sigma^2_{\underline{x},\bar{x}}$ is as in* (2.3.20).

The second fundamental result of this section gives a lower bound for the local asymptotic minimimax (LAM) risk of an arbitrary estimator, and the related convolution theorem.

Theorem 2.3.2 (LAM & Convolution)**.** *Let* (2.2.6) *and Assumption 2.3.3 hold true and let $\sigma^2_{\underline{x},\bar{x}}$ be as in* (2.3.20)*. Let then $\ell : \mathbb{R} \mapsto [0, \infty)$ be symmetric and subconvex. Then for any $x \in (\underline{x}, \bar{x})$, and every estimator sequence $(V_n(x))_{n\in\mathbb{N}}$:*

$$\sup_{\substack{I \subset \mathbb{R} \\ I \text{ finite}}} \liminf_{n\to\infty} \sup_{t\in I} \mathrm{E}_t\, \ell\left(\sqrt{n}\left(V_n(x) - V_{t/\sqrt{n}}(x)\right)\right) \geq \int \ell\, dN(0, \sigma^2_{\underline{x},\bar{x}}), \tag{2.3.29}$$

where E_t *denotes the expectation under $G_{t/\sqrt{n}}$. Moreover, for any estimator sequence such that $\sqrt{n}\left(V_n(x) - V_{t/\sqrt{n}}(x)\right) \rightsquigarrow K$ under $G_{t/\sqrt{n}}$, for every $t \in \mathbb{R}$, and some given distribution K, this distribution is the convolution of $N(0, \sigma^2_{\underline{x},\bar{x}})$ and some other distribution.*

The proof of Theorem 2.3.2 is given in Appendix 2.4. By Theorem 2.3.1 the informed estimators attain equality in the LAM risk (for bounded continuous loss functions), and their limit distribution is $N(0, \sigma^2_{\underline{x},\bar{x}})$ without extra convolution factor. On the other hand, the limit distribution of the isotonic estimator contains the distribution of W as in (2.3.21) as extra factor. Hence the isotonic estimator is not LAM.

2.3.4 Simulation study

In this subsection, we present a simulation study that shows that the isotonic inverse estimator, even if not efficient, behaves very closely to the efficient estimator. In practice, the information whether or not the cdf F^* is constant on a specific interval is generally unavailable. In that situation the best choice would be to use the isotonic estimator, given the efficiency theory and the adaptivity results developed in the previous chapter. However, if the cdf is known to be constant on some known interval, then the isotonic estimator will incur a loss in terms of efficiency. In this section, we perform a simulation study to illustrate the difference in terms of performance between the isotonic estimator $\hat{V}_n$ and the efficient informed estimator $V^{(\underline{x},\bar{x})}$. We consider an underlying distribution F^* of the squared sphere radii, which is constant on $[2, 3]$ (see Figure 2.7).

Figure 2.8 shows the true underlying V with all estimators defined in this section within the specified interval. As $\hat{V}_n^{\Pi}$, V_n^{Π}, and $V_n^{(\underline{x},\bar{x})}$ are all asymptotically equivalent by Theorem 2.3.1, we focus on the comparison between $V_n^{(\underline{x},\bar{x})}$ and $\hat{V}_n$.

In Figure 2.9, the scatter plot reveals a strong positive correlation between the two estimators, while the Tukey mean-difference plot indicates a lack of inherent bias. Taken together, the two plots in Figure 2.9 demonstrate a substantial agreement between the two estimation methods.



$$F(x) = \begin{cases} 1 - e^{-x^2/20}, & 0 \le x < 2 \\ 1 - e^{-1/5}, & x \in [2, 3] \\ 1 - e^{-(x-1)^2/20}, & x > 3 \end{cases}$$

$$g_V(x) = \frac{1}{2m_0} \int_x^M \frac{s e^{-\frac{s^2}{10}} \mathbf{1}_{[0,2]}(s)}{10\sqrt{s-x}}\, ds + \frac{1}{2m_0} \int_x^M \frac{(s-1) e^{-\frac{(s-1)^2}{20}} \mathbf{1}_{[3,M]}(s)}{10\sqrt{s-x}}\, ds$$

Figure 2.7: Underlying cdf F^* and in light blue a histogram of a sample of size 1000 from g.

In Figure 2.10, we observe that the distributions of $\sqrt{n}(V_n^{(\underline{x},\bar{x})}(x) - V(x))$ and $\sqrt{n}(\hat{V}_n(x) - V(x))$ closely resemble each other. Moreover, we observe a strong resemblance not only between the estimators but also between their respective limiting distributions, L_x and $N(0, \sigma^2_{\underline{x},\bar{x}})$. However, the violin plot with the standard percentiles, the second histogram with $\sqrt{n}(\hat{V}_n(x) - V_n^{(\underline{x},\bar{x})}(x))$ and W and Table 2.1 indicate that the variance of $\hat{V}_n(x)$ is slightly bigger than the one of $V_n^{(\underline{x},\bar{x})}(x)$.

The values in Table 2.1 were computed using varying sample size n from g as indicated, with 2000 samples from $\hat{V}_n(x)$ and $V_n^{(\underline{x},\bar{x})}(x)$. All the Kernel Density Estimator plots were computed using Scott's rule (where the bin size $h \sim 3.5\hat{\sigma}n^{-1/3}$) for the chosen bandwidth.



Figure 2.8: All estimators and the true function V, which is constant on $[2, 3]$, for a sample of size $n = 300$.

For W and L_x, 20000 samples were used. Taken together, the simulation study confirms the theoretical findings of Theorem 2.3.1 and 2.3.2: the informed estimators are efficient and not asymptotically equivalent to the IIE. However, the empirical performances of $V_n^{(\underline{x},\bar{x})}$ and $\hat{V}_n$ show a high degree of proximity in the estimation scenario where the underlying function V is constant on an interval. Even if $\hat{V}_n$ is used in cases where it is indeed true that V is constant within an interval, the incurred loss appears to be small.



Figure 2.9: Scatter and Tukey mean-difference plots for $n = 1000$ and 300 repetitions of $\hat{V}_n$ and $V_n^{(\underline{x},\bar{x})}$.





Figure 2.10: Based on 1000 samples from g, centered and rescaled histograms based on 300 repetitions, $V_n^{(\underline{x},\bar{x})}$ and their difference compared with respective limiting distributions L_x, $N(0, \sigma^2_{\underline{x},\bar{x}})$ and W.

n	100	200	400	1000	2000		limit
$\sqrt{n}(V_n^{(\underline{x},\bar{x})} - V)(x)$	0.5441	0.5439	0.5465	0.5467	0.5464	$N(0, \sigma^2_{\underline{x},\bar{x}})$	0.5466
$\sqrt{n}(\hat{V}_n - V)(x)$	0.5537	0.5592	0.5627	0.5664	0.5697	L_x	0.5703

Table 2.1: Standard deviations

2.3.5 Conclusion

We conclude by stressing the fact that the local asymptotically minimax variance in Theorem 2.3.2 coincides with the one obtained in Theorem 2.3.1.a.. This shows that in the setting in which V is constant on $[\underline{x}, \bar{x}]$, for $\underline{x} < \bar{x}$, the informed estimators are asymptotically efficient, whereas the isotonic inverse estimator is not, as it is clearly shown by Theorem 2.3.2. However, we conjecture that the isotonic estimator remains asymptotically efficient in the situations in which the boundary points $\underline{x}, \bar{x}$ are allowed to converge to x at an arbitrarily polynomial rate (or slower). In this case, the rate of convergence drops to $\sqrt{n/\log(\bar{x} - \underline{x})^{-1}}$.

From a practical point of view, both the results of the current section and Theorems 2.2.4-2.2.3 are difficult to use to obtain confidence bands, as they require estimating the density g, either at x or globally to compute $\sigma^2_{\underline{x},\bar{x}}$ or to compute the covariance structure of the limiting Gaussian process. We envision Theorems 2.2.4-2.2.3 and the asymptotic results of the current section as the theoretical foundation, while the practical computation of confidence bands may be better addressed through bootstrap methods. A potential direction for further research would be proving formally why the bootstrap works. We believe that a formal justification would combine our theoretical results with the techniques used in [92] (which already shows consistency of the bootstrap for $\gamma = 1$ in the limiting variance of Theorem 2.2.4).

Another potential direction for further investigation is whether the IIE is efficient in the setting in which there exists an interval of constancy for V but the boundary points $\underline{x}, \bar{x}$ are unknown. However, the definition and assessment of *efficiency* in this context are unclear. For practical purposes, if the boundary points $\underline{x}, \bar{x}$ are unknown, the IIE remains the state-of-art as

it behaves very closely to the efficient estimator and in the $\sqrt{n/\log n}$ rate setting is efficient and adaptive to the level of smoothness of V.

2.4 Appendix

2.4.1 Proofs of complemental statements for Theorem 2.2.3.

First we provide some proofs about the fundamental behavior of the covariance structure of the process $Z_n(t)$. We start by showing the case of the variance and we will proceed by giving a proof along the same lines of (2.2.13). We will use several times how samples of Z with density g (see (2.1.1)) are obtained in Wicksell's problem. First, note that the observable radii in the cross sections cannot exceed the radii of the associated sphere; the spheres with bigger radius are more likely to be sampled. Thus the distribution of the squared radii of the spheres actually cut is not F^* but a biased version of it; thus we write X with distribution function F for the squared radius of the spheres actually cut. Since we assume the samples are gathered uniformly, the observed squared circle radius Z is related to X via:

$$Z = (1-U^2)X = VX, \quad U \sim U[0,1], \quad V \sim \mathrm{Be}(1,1/2) \tag{2.4.1}$$

Lemma 2.4.1. *Fix $x > 0$. If $g(x) < \infty$, then:*

$$\mathrm{E}\Big(\sqrt{Z-x}\mathbf{1}_{\{Z>x\}} - \sqrt{Z-x-\varepsilon}\mathbf{1}_{\{Z>x+\varepsilon\}}\Big)^2 \sim \varepsilon^2 \log\left(\frac{1}{\varepsilon}\right) g(x), \quad \textit{as} \quad \varepsilon \downarrow 0.$$

Proof. Define $Z_x := Z - x$. The left side is:

$$\underbrace{\mathrm{E}Z_x\mathbf{1}_{\{Z_x<\varepsilon\}}}_{(1)} + \underbrace{\mathrm{E}\Big(\sqrt{Z_x} - \sqrt{Z_x-\varepsilon}\Big)^2\mathbf{1}_{\{Z_x>\varepsilon\}}}_{(2)}.$$

We start with (1). By making a substitution below with $vX - x = y(X-x)$, which means $1 - v = (1-y)\frac{X-x}{X}$, we obtain:

$$\begin{aligned}\mathrm{E}\,Z_x\mathbf{1}_{Z_x<\varepsilon} &\overset{(2.4.1)}{=} \mathrm{E}\left\{\mathbf{1}_{X>x}\int_{0<vX-x<\varepsilon}(vX-x)\frac{\mathbf{1}_{0<v<1}}{2\sqrt{1-v}}\,dv\right\}\\ &= \mathrm{E}\left\{\mathbf{1}_{X>x}\int_0^{\frac{\varepsilon}{X-x}\wedge 1}\frac{y}{2\sqrt{1-y}}\frac{(X-x)^{3/2}}{\sqrt{X}}\,dy\right\}.\end{aligned}$$

If we define $y(z) = \int_0^z \frac{y}{2\sqrt{1-y}}\,dy$ then the above equals to:

$$\mathrm{E}\left\{y\left(\frac{\varepsilon}{X-x}\wedge 1\right)\mathbf{1}_{X>x}\frac{(X-x)^{3/2}}{\sqrt{X}}\right\}.$$

Using: $\int \frac{y}{2\sqrt{1-y}}\,dy = \frac{(1-y)^{3/2}}{3} - \sqrt{1-y} + c$, we can see that:

$$y\left(\frac{\varepsilon}{X-x}\wedge 1\right) = \int_0^{\frac{\varepsilon}{X-x}\wedge 1}\frac{y}{2\sqrt{1-y}}\,dy = \frac{1}{4}\left(\frac{\varepsilon}{X-x}\right)^2 + O\left(\left(\frac{\varepsilon}{X-x}\right)^3\right).$$

Moreover it is also true $y(z) \le z^2$, $\forall\, z \in (0,1)$. The above proves:

$$\mathrm{E}\, Z_x \mathbf{1}_{Z_x<\varepsilon} = O(\varepsilon^2). \tag{2.4.2}$$

Therefore (1) is negligible. Now we proceed with (2). Using Taylor's theorem with integral form of the remainder we obtain:

$$\begin{aligned}
&\mathrm{E}\Big(\sqrt{Z_x} - \sqrt{Z_x-\varepsilon}\Big)^2 \mathbf{1}_{\{Z_x>\varepsilon\}} = \mathrm{E}\int_0^1\int_0^1 \frac{\varepsilon}{2\sqrt{Z_x-\varepsilon s}}\frac{\varepsilon}{2\sqrt{Z_x-\varepsilon t}}\,ds\,dt\,\mathbf{1}_{\{Z_x>\varepsilon\}} \\
&\overset{(2.4.1)}{=} \mathrm{E}\int_0^1\int_0^1\int_0^1 \frac{\varepsilon}{2\sqrt{vX-x-\varepsilon s}}\frac{\varepsilon}{2\sqrt{vX-x-\varepsilon t}}\frac{1}{2\sqrt{1-v}}\mathbf{1}_{\{vX-x>\varepsilon\}}\,dv\,ds\,dt.
\end{aligned}$$

Now we use again the substitution $vX - x = y(X-x)$ which implies $1 - v = (1-y)\frac{X-x}{X}$ and $y(\varepsilon, s, t) = \int_\varepsilon^1 \frac{1}{\sqrt{y-\varepsilon s}}\frac{1}{\sqrt{y-\varepsilon t}}\frac{1}{\sqrt{1-y}}\,dy$ to obtain:

$$\begin{aligned}
&= \frac{\varepsilon^2}{8}\mathrm{E}\int_0^1\int_0^1\int_0^1 \frac{1}{\sqrt{y-\frac{\varepsilon s}{X-x}}}\frac{1}{\sqrt{y-\frac{\varepsilon t}{X-x}}}\frac{1}{\sqrt{1-y}}\frac{\mathbf{1}_{\{y(X-x)>\varepsilon\}}}{\sqrt{X-x}\sqrt{X}}\,dy\,ds\,dt \\
&= \frac{\varepsilon^2}{8}\mathrm{E}\int_0^1\int_0^1 y\Big(\frac{\varepsilon}{X-x}, s, t\Big)\frac{\mathbf{1}_{\{X-x>\varepsilon\}}}{\sqrt{X-x}\sqrt{X}}\,ds\,dt.
\end{aligned}$$

Now we show the two following relations $\forall\ 0 < \varepsilon, s, t < 1$:

$$y(\varepsilon, s, t) \sim \log\frac{1}{\varepsilon} - \frac{1}{2}\log\log\frac{1}{\varepsilon} + O(1) \quad \text{as } \varepsilon \downarrow 0 \tag{2.4.3}$$

$$y(\varepsilon, s, t) \le \left(\log\frac{1}{\varepsilon} + \log\frac{e^2}{1 - s\vee t}\right)\frac{1}{\sqrt{1-s\vee t}} + 1. \tag{2.4.4}$$

From this it will follow that:

$$\mathrm{E}\Big(\sqrt{Z_x} - \sqrt{Z_x-\varepsilon}\Big)^2 \mathbf{1}_{\{Z_x>\varepsilon\}} \sim \frac{\varepsilon^2}{4} g(x)\log\frac{1}{\varepsilon}. \tag{2.4.5}$$

Fix $\varepsilon < b < 1$, then we can split the integral in the definition of $y(\varepsilon, s, t)$ into two integrals: one from ε to b and another one from b to 1. Let us look at the second one:

$$\int_b^1 \frac{(1-y)^{-\frac{1}{2}}}{y\sqrt{y-\varepsilon s}}\frac{y}{\sqrt{y-\varepsilon t}}\,dy \le \frac{b}{b-\varepsilon}\int_b^1 \frac{1}{y\sqrt{1-y}}\,dy = \frac{b}{b-\varepsilon} 2\tanh^{-1}(\sqrt{1-b}).$$

Here we used the fact that the mapping $y \mapsto \frac{y}{y-\varepsilon s}$ is decreasing on $[\varepsilon s, 1]$, that $\frac{b}{b-\varepsilon s} \le \frac{b}{b-\varepsilon}$ and $\int \frac{1}{y}\frac{1}{\sqrt{1-y}}\,dy = -2\tanh^{-1}(\sqrt{1-y}) + c$. This shows that the asymptotic behavior will be determined by the integral between ε and b;

$$\begin{aligned}
&\left|\int_\varepsilon^b \frac{1}{\sqrt{y-\varepsilon s}}\frac{1}{\sqrt{y-\varepsilon t}}\frac{1}{\sqrt{1-y}}\,dy - \int_\varepsilon^b \frac{1}{\sqrt{y-\varepsilon s}}\frac{1}{\sqrt{y-\varepsilon t}}\,dy\right| \\
&\qquad \le \int_\varepsilon^b \frac{1}{\sqrt{y-\varepsilon s}}\frac{1}{\sqrt{y-\varepsilon t}}\,dy\left(\frac{1}{\sqrt{1-b}} - 1\right).
\end{aligned}$$

On the other hand, since:

$$\int_\varepsilon^b \frac{1}{y-\varepsilon s\wedge t}\,dy \leqslant \int_\varepsilon^b \frac{1}{\sqrt{y-\varepsilon s}}\frac{1}{\sqrt{y-\varepsilon t}}\,dy \leqslant \int_\varepsilon^b \frac{1}{y-\varepsilon s\vee t}\,dy.$$

Therefore we have:

$$\int_\varepsilon^b \frac{1}{\sqrt{y-\varepsilon s}}\frac{1}{\sqrt{y-\varepsilon t}}\frac{1}{\sqrt{1-y}}\,dy \begin{cases} \leq \log\left\{\frac{b-\varepsilon s\vee t}{\varepsilon(1-s\vee t)}\right\}\frac{1}{\sqrt{1-b}}, \\ \geq \log\left\{\frac{b-\varepsilon s\wedge t}{\varepsilon(1-s\wedge t)}\right\}\left(2-\frac{1}{\sqrt{1-b}}\right). \end{cases}$$

Using $b=\frac{1}{\sqrt{\log\frac{1}{\varepsilon}}}$ we obtain for fixed c:

$$\log\left\{\frac{1/\sqrt{\log\frac{1}{\varepsilon}}-\varepsilon c}{\varepsilon(1-c)}\right\} = \log\left\{1-\varepsilon\sqrt{\log\frac{1}{\varepsilon}}c\right\} + \log\left\{\frac{1}{\sqrt{\log\frac{1}{\varepsilon}}}\right\} + \log\frac{1}{\varepsilon} + \\ + \log\frac{1}{1-c} \sim \log\frac{1}{\varepsilon} - \frac{1}{2}\log\log\frac{1}{\varepsilon} + O(1), \qquad \text{as } \varepsilon\downarrow 0.$$

This proves the first relation (2.4.3). Now we proceed with relation (2.4.4). Assume $s\leq t$. By performing a substitution $y-\varepsilon t = z(1-\varepsilon t)$ which means $1-y=(1-\varepsilon t)(1-z)$ we obtain:

$$y(\varepsilon,s,t) \leq \int_\varepsilon^1 \frac{1}{y-\varepsilon t}\frac{1}{\sqrt{1-y}}\,dy = \frac{1}{\sqrt{1-\varepsilon t}}\int_{\frac{\varepsilon(1-t)}{1-\varepsilon t}}^1 \frac{1}{z}\frac{1}{\sqrt{1-z}}\,dz.$$

We used that we can compute the primitive: $\int \frac{1}{y}\frac{1}{\sqrt{1-y}}\,dy = -2\tanh^{-1}(\sqrt{1-y})+c$. Therefore:

$$\frac{1}{\sqrt{1-\varepsilon t}}\int_{\frac{\varepsilon(1-t)}{1-\varepsilon t}}^1 \frac{1}{z}\frac{1}{\sqrt{1-z}}\,dz = 2\tanh^{-1}\left(\sqrt{1-\frac{\varepsilon(1-t)}{1-\varepsilon t}}\right)\frac{1}{\sqrt{1-\varepsilon t}}$$
$$= \frac{1}{\sqrt{1-\varepsilon t}}\log\left\{\frac{1+\sqrt{\frac{1-\varepsilon}{1-\varepsilon t}}}{1-\sqrt{\frac{1-\varepsilon}{1-\varepsilon t}}}\right\} = \log\left\{\frac{\left(\sqrt{1-\varepsilon t}+\sqrt{1-\varepsilon}\right)^2}{\varepsilon-\varepsilon t}\right\}\frac{1}{\sqrt{1-\varepsilon t}}$$
$$\leq \left(\log\frac{1}{\varepsilon}+\log\frac{1-\varepsilon t}{1-t}+2\right)\frac{1}{\sqrt{1-\varepsilon t}} \leq \left(\log\frac{1}{\varepsilon}+\log\frac{e^2}{1-t}\right)\frac{1}{\sqrt{1-t}}+1.$$

And this proves (2.4.4). Now we proceed by using the two relations we just proved. First of all, (2.4.3) as $\varepsilon\downarrow 0$:

$$\frac{y\left(\frac{\varepsilon}{X_x},s,t\right)\mathbf{1}_{\{X_x>\varepsilon\}}}{\log\frac{1}{\varepsilon}} = \frac{y\left(\frac{\varepsilon}{X_x},s,t\right)\mathbf{1}_{\{X_x>\varepsilon\}}}{\log\frac{X_x}{\varepsilon}}\cdot\frac{\log X_x+\log\frac{1}{\varepsilon}}{\log\frac{1}{\varepsilon}} \to \mathbf{1}_{\{X>x\}}$$

Now using $\log xy \leq \log x\mathbf{1}_{x>1}+\log y$ and (2.4.4) we obtain:

$$\frac{y\left(\frac{\varepsilon}{X-x},s,t\right)\mathbf{1}_{\{X-x>\varepsilon\}}}{\log\frac{1}{\varepsilon}} \leq \frac{\left(\log\frac{X_x}{\varepsilon}+\log\frac{e^2}{1-s\vee t}\right)\frac{1}{\sqrt{1-s\vee t}}+1}{\log\frac{1}{\varepsilon}}$$
$$\lesssim \left(\log X_x\mathbf{1}_{X_x>1}+\log\frac{e^2}{\sqrt{1-s\vee t}}\right)\frac{1}{\sqrt{1-s\vee t}}+1.$$

Since the idea is to apply the Dominated Convergence Theorem to:

$$\frac{1}{\log\frac{1}{\varepsilon}}\mathrm{E}\int_0^1\int_0^1 y\Big(\frac{\varepsilon}{X-x},s,t\Big)\frac{1}{2\sqrt{X-x}\sqrt{X}}\mathbf{1}_{\{X-x>\varepsilon\}}\,ds\,dt,$$

we need to verify that all the upper bounds have finite expectations. First $\frac{\log X_x\mathbf{1}_{\{X_x>1\}}}{\sqrt{X}}\lesssim 1$ which means:

$$\int_0^1\int_0^1\frac{\log X_x\mathbf{1}_{\{X_x>1\}}}{\sqrt{X}}\frac{1}{\sqrt{1-s\vee t}}\,ds\,dt\lesssim 2\int_0^1\int_0^t\frac{1}{\sqrt{1-t}}\,ds\,dt<\infty.$$

On the other hand:

$$\begin{aligned}\int_0^1\int_0^1\Big(\log\frac{1}{1-s\vee t}\Big)\frac{1}{\sqrt{1-s\vee t}}\,ds\,dt&\lesssim 2\int_0^1\int_0^t\Big(\log\frac{1}{1-t}\Big)\frac{1}{\sqrt{1-t}}\,ds\,dt\\&=2\int_0^1 t\Big(\log\frac{1}{1-t}\Big)\frac{1}{\sqrt{1-t}}\,dt<\infty.\end{aligned}$$

So by the Dominated Convergence Theorem, if $g(x)=\mathrm{E}\frac{1}{2\sqrt{X-x}}\frac{1}{\sqrt{X}}\mathbf{1}_{\{X>x\}}<\infty$ we obtain the claim. □

Proof of (2.2.13). Fix $s,t\in I_x$. Take the case $0\le s\le t$. Fix μ; we define the function $\phi(\cdot,\mu)$ as follows:

$$\phi(z,\mu)=\sqrt{z-x}\,\mathbf{1}_{[x,\infty)}(z)-\sqrt{z-x-\mu}\,\mathbf{1}_{[x+\mu,\infty)}(z).$$

Using the i.i.d. property, we can write:

$$\mathrm{Cov}\left(Z_n(s),Z_n(t)\right)=4\delta_n^{-2}(\delta_n^*)^{-2}n^{-1}\,\mathrm{Cov}\left(\phi\left(Z,\delta_n^*t\right),\phi\left(Z,\delta_n^*s\right)\right),\tag{2.4.6}$$

where Z has density g. Since $g(x)<\infty$, we will prove:

$$\mathrm{E}\phi\left(Z,\delta_n^*s\right)\phi\left(Z,\delta_n^*t\right)=-\frac{1}{4}stg(x)(\delta_n^*)^2\log\delta_n^*+O\Big((\delta_n^*)^2\Big),\tag{2.4.7}$$

which implies (2.2.13) for $\delta_n=n^{-1/2}\sqrt{\log n}$ and $\delta_n^*=(\delta_n)^{1/\gamma_x}$. Define $Z_x:=Z-x$:

$$\begin{aligned}&\mathrm{E}\Big[\sqrt{Z_x}\mathbf{1}_{\{Z_x>0\}}-\sqrt{Z_x-\delta_n^*t}\mathbf{1}_{\{Z_x>\delta_n^*t\}}\Big]\Big[\sqrt{Z_x}\mathbf{1}_{\{Z_x>0\}}-\sqrt{Z_x-\delta_n^*s}\mathbf{1}_{\{Z_x>\delta_n^*s\}}\Big]\\&=\underbrace{\mathrm{E}Z_x\mathbf{1}_{\{0<Z_x<\delta_n^*s\}}}_{(1)}+\underbrace{\mathrm{E}\Big\{\sqrt{Z_x}\Big(\sqrt{Z_x}-\sqrt{Z_x-\delta_n^*s}\Big)\mathbf{1}_{\{\delta_n^*s\le Z_x\le\delta_n^*t\}}\Big\}}_{(2)}+\\&+\underbrace{\mathrm{E}\Big\{\Big(\sqrt{Z_x}-\sqrt{Z_x-\delta_n^*t}\Big)\Big(\sqrt{Z_x}-\sqrt{Z_x-\delta_n^*s}\Big)\mathbf{1}_{\{Z_x>\delta_n^*t\}}\Big\}}_{(3)}.\end{aligned}$$

Let us start with (1). We can use (2.4.2) to see that for δ_n^* going to zero:

$$\mathrm{E}Z_x\mathbf{1}_{\{0<Z_x<\delta_n^*s\}}\le(\delta_n^*s)^2\mathrm{E}\frac{1}{\sqrt{X-x}}\frac{1}{\sqrt{X}}\mathbf{1}_{\{X>x\}}=O((\delta_n^*s)^2).$$

This shows that this term is negligible as for the covariance it is multiplied by $4\delta_n^{-2}(\delta_n^*)^{-2}n^{-1}=4\frac{1}{\log n}(\delta_n^*)^{-2}$, will be of order $\frac{1}{\log n}$ and it will go to zero.

Let us continue with (2). First note one can upper bound (2) using Hölder's inequality and notice that by (2.4.5) we have:

$$\mathrm{E}\Big(\sqrt{Z_x}-\sqrt{Z_x-\delta_n^* s}\Big)^2\,\mathbf{1}_{\{Z_x>\delta_n^* s\}}\sim\frac{(\delta_n^* s)^2}{4}\log\frac{1}{\delta_n^* s}g(x). \tag{2.4.8}$$

Which implies that the square root of this term is of the order $O(\delta_n^*\sqrt{\log\frac{1}{\delta_n^*}})$. Moreover, notice that by (2.4.2):

$$\Big\{\mathrm{E}Z_x\mathbf{1}_{\{\delta_n^* s<Z_x<\delta_n^* t\}}\Big\}^{1/2}\le\Big\{\mathrm{E}Z_x\mathbf{1}_{\{Z_x>\delta_n^* s\}}\Big\}^{1/2}=O(\delta_n^*).$$

All together we obtain that (2) is of the order $O((\delta_n^*)^2\sqrt{\log(\delta_n^*)^{-1}})$. This quantity for the covariance structure will be multiplied by $4\delta_n^{-2}(\delta_n^*)^{-2}n^{-1}$, and will be of order $O(\frac{1}{\sqrt{\log n}})$ thus it will go to zero because $\sqrt{\log\frac{1}{\delta_n^*}}\sim\sqrt{\log n}$.
Now we argue the order of (3). We argue like in the proof of (2.4.5):

$$\begin{aligned}(3)&=(\delta_n^*)^2 st\mathrm{E}\int_0^1\int_0^1\frac{1}{2\sqrt{Z_x-\delta_n^* sp}}\frac{1}{2\sqrt{Z_x-\delta_n^* tq}}\,dp\,dq\,\mathbf{1}_{\{Z_x>\delta_n^* t\}}\\&=\mathrm{E}\int_0^1\int_0^1\int_0^1\frac{(\delta_n^*)^2 st}{2\sqrt{vX-x-\delta_n^* sp}}\frac{\mathbf{1}_{\{vX-x>\delta_n^* t\}}}{2\sqrt{vX-x-\delta_n^* tq}}\frac{1}{2\sqrt{1-v}}\,dv\,dp\,dq.\end{aligned}$$

Now we use again the substitution $vX-x=y(X-x)$ which implies $1-v=(1-y)\frac{X-x}{X}$ and $y(\delta_n^*,s,t)=\int_{\delta_n^*}^1\frac{1}{\sqrt{y-\delta_n^* s}}\frac{1}{\sqrt{y-\delta_n^* t}}\frac{1}{\sqrt{1-y}}dy$ to obtain:

$$\begin{aligned}&=\mathrm{E}\int_0^1\int_0^1\int_0^1\frac{(\delta_n^*)^2 st}{8\sqrt{y-\frac{\delta_n^* t}{X-x}q}}\frac{\mathbf{1}_{\{y(X-x)>\delta_n^* t\}}}{\sqrt{y-\frac{\delta_n^* s}{X-x}p}}\frac{1}{\sqrt{1-y}}\frac{1}{\sqrt{X-x}\sqrt{X}}\,dy\,dp\,dq\\&=\frac{(\delta_n^*)^2}{8}st\mathrm{E}\int_0^1\int_0^1 y\left(\frac{\delta_n^*}{X-x},sp,tq\right)\frac{1}{\sqrt{X-x}\sqrt{X}}\mathbf{1}_{\{X-x>\delta_n^* t\}}\,dp\,dq.\end{aligned}$$

Then one can obtain the claim by using (2.4.3) and (2.4.4). The reason why we can apply the Dominated Convergence theorem is as in the proof of Lemma 2.4.1. The cases $s\le 0\le t$ and $s\le t\le 0$ can be argued in a similar way. □

Proof of (2.2.14). We verify the Lindeberg condition for $Z_n(1)$. Recall from (2.2.12) the definition of $Z_n(1)$. Then set $Z_n(1)=\sum_{k=1}^n(X_k^n-\mathrm{E}X_k^n)$ and thus $s_n^2=\sum_{k=1}^n\mathrm{Var}(X_k^n)$, which on the other hand means:

$$X_k^n=\frac{2\delta_n^{-1}}{n}\frac{\sqrt{Z_k-x}\mathbf{1}_{[x,\infty)}(Z_k)-\sqrt{Z_k-x-\delta_n^*}\mathbf{1}_{[x+\delta_n^*,\infty)}(Z_k)}{\delta_n^*}.$$

Therefore by (2.2.13): $s_n^2=\mathrm{Var}(Z_n(1))=\frac{1}{2\gamma_x}g(x)\Big(1-\frac{\log\log n}{\log n}\Big)+O\Big(\frac{1}{\log n}\Big)$. We show $\mathrm{E}X_k^n=\frac{(\delta_n^*)^{-1}\delta_n^{-1}}{n}O(\delta_n^*)=O(\frac{1}{\sqrt{n\log n}})$. It is enough to prove for $\delta_n^*\downarrow 0$:

$$\int\Big\{\sqrt{z-x}\mathbf{1}_{[x,\infty)}(z)-\sqrt{z-x-\delta_n^*}\mathbf{1}_{[x+\delta_n^*,\infty)}(z)\Big\}g(z)dz=O(\delta_n^*). \tag{2.4.9}$$

Let $Z_x := Z - x$ therefore the right-hand side of the above can be written as:

$$\underbrace{\mathrm{E}\left[\sqrt{Z_x}\mathbf{1}_{\{0<Z_x<\delta_n^*\}}\right]}_{=(a)} + \underbrace{\mathrm{E}\left(\sqrt{Z_x} - \sqrt{Z_x - \delta_n^*}\right)\mathbf{1}_{\{Z_x>\delta_n^*\}}}_{=(b)}.$$

We show $(a) = O(\delta_n^*)$. By using (2.4.1) (we denote X^b by X for brevity) and making a substitution below with $vX - x = y(X - x)$, which means $1 - v = (1 - y)\frac{X-x}{X}$, we obtain:

$$\begin{aligned}\mathrm{E}\sqrt{Z_x}\mathbf{1}_{Z_x<\delta_n^*} &= \mathrm{E}\left\{\mathbf{1}_{X>x}\int_{0<vX-x<\delta_n^*}\sqrt{vX-x}\frac{\mathbf{1}_{0<v<1}}{2\sqrt{1-v}}\,dv\right\}\\ &= \mathrm{E}\left\{\mathbf{1}_{X>x}\int_0^{\frac{\delta_n^*}{X-x}\wedge 1}\frac{\sqrt{y}}{2\sqrt{1-y}}\frac{(X-x)}{\sqrt{X}}\,dy\right\}.\end{aligned}$$

For $y(z) = \int_0^z \frac{\sqrt{y}}{2\sqrt{1-y}}dy$, the above can be upper bounded for all δ_n^* small:

$$\mathrm{E}\left\{y\left(\frac{\delta_n^*}{X-x}\wedge 1\right)\mathbf{1}_{X>x}\frac{X-x}{\sqrt{X}}\right\} \lesssim \mathrm{E}\delta_n^*\frac{1}{\sqrt{X}}\mathbf{1}_{\{X>x\}} = O(\delta_n^*). \tag{2.4.10}$$

That is because: $\int_0^z \frac{\sqrt{y}}{2\sqrt{1-y}}dy = \frac{1}{2}\sin^{-1}(\sqrt{z}) - \frac{1}{2}\sqrt{z(1-z)} \le z$ for $z \in (0,1)$. Now we proceed with (b). By using again (2.4.1) and by making a substitution below with $vX - x = y(X - x)$, which means $1 - v = (1 - y)\frac{X-x}{X}$, we obtain:

$$\begin{aligned}\mathrm{E}\left(\sqrt{Z_x} - \sqrt{Z_x - \delta_n^*}\right)\mathbf{1}_{\{Z_x>\delta_n^*\}} &= \mathrm{E}\int_0^1 \frac{\delta_n^*}{2\sqrt{Z_x - \delta_n^* s}}\,ds\,\mathbf{1}_{\{Z_x>\delta_n^*\}}\\ &= \mathrm{E}\int_0^1\int_0^1 \frac{\delta_n^*}{2\sqrt{vX - x - \delta_n^* s}}\frac{1}{2\sqrt{1-v}}\mathbf{1}_{\{Z_x>\delta_n^*\}}\,dv\,ds\\ &= \frac{1}{4}\mathrm{E}\int_0^1\int_0^1 \frac{\delta_n^*}{\sqrt{y(X-x) - \delta_n^* s}}\frac{1}{\sqrt{1-y}}\frac{\sqrt{X}}{\sqrt{X-x}}\frac{X-x}{X}\,dy\,ds\\ &= \frac{\delta_n^*}{4}\mathrm{E}\int_0^1 y\left(\frac{\delta_n^*}{X-x}, s\right)\frac{1}{\sqrt{X}}\mathbf{1}_{\{X>x+\delta_n^*\}}\,ds,\end{aligned}$$

where $y(\varepsilon, s) = \int_\varepsilon^1 \frac{1}{\sqrt{y-\varepsilon s}}\frac{1}{\sqrt{1-y}}dy$. For $b \in (\varepsilon, 1)$, as done previously, we split the integral between ε and b and between b and 1. For the first integral we have:

$$\left|\int_\varepsilon^b \frac{1}{\sqrt{y-\varepsilon s}}\frac{1}{\sqrt{1-y}}\,dy - \int_\varepsilon^b \frac{1}{\sqrt{y-\varepsilon s}}\,dy\right| \le \int_\varepsilon^b \frac{1}{\sqrt{y-\varepsilon s}}\,dy\left(\frac{1}{\sqrt{1-b}} - 1\right).$$

From which we conclude:

$$\int_\varepsilon^b \frac{1}{\sqrt{y-\varepsilon s}}\frac{1}{\sqrt{1-y}}\,dy = \begin{cases} \le (2\sqrt{b-\varepsilon s} - 2\sqrt{\varepsilon(1-s)})\left(\frac{1}{\sqrt{1-b}} - 1\right),\\ \ge (2\sqrt{b-\varepsilon s} - 2\sqrt{\varepsilon(1-s)})\left(2 - \frac{1}{\sqrt{1-b}}\right).\end{cases}$$

For the second integral we have:

$$\int_b^1 \frac{1}{\sqrt{y-\varepsilon s}}\frac{1}{\sqrt{1-y}}\,dy \le \frac{1}{\sqrt{b^2 - b\varepsilon}}\left\{\sqrt{b(1-b)} - \sin^{-1}(\sqrt{1-b})\right\}.$$

Putting together the above computations, for all δ_n^* small and $b \sim \sqrt{\frac{1}{\log \frac{1}{\delta_n^*}}}$:

$$\mathrm{E}\Big(\sqrt{Z_x} - \sqrt{Z_x - \delta_n^*}\Big)\mathbf{1}_{\{Z_x > \delta_n^*\}} \lesssim \delta_n^* \mathrm{E}\frac{\mathbf{1}_{\{X > x+\delta_n^*\}}}{\sqrt{X}} = O(\delta_n^*).$$

This proves that (2.4.9) is true.

Let us now look at the indicator $\mathbf{1}_{\{|X_k^n - \mathrm{E}X_k^n| > \varepsilon s_n\}}$. We look separately at the cases: $X_k^n - \mathrm{E}X_k^n > \varepsilon s_n$ and $X_k^n - \mathrm{E}X_k^n < -\varepsilon s_n$. Using the following facts:

$$\sqrt{Z_x}\mathbf{1}_{\{Z_x>0\}} - \sqrt{Z_x - \delta_n^*}\mathbf{1}_{\{Z_x > \delta_n^*\}} = \sqrt{Z_x}\mathbf{1}_{\{0<Z_x<\delta_n^*\}} + \Big(\sqrt{Z_x} +$$
$$- \sqrt{Z_x - \delta_n^*}\Big)\mathbf{1}_{\{Z_x>\delta_n^*\}} \leq \sqrt{\delta_n^*} + \int_0^1 \frac{\delta_n^* \mathbf{1}_{\{Z_x>\delta_n^*\}}}{2\sqrt{Z_x - \delta_n^* s}}\, ds \leq \sqrt{\delta_n^*} + \delta_n^* \frac{\mathbf{1}_{\{Z_x>\delta_n^*\}}}{2\sqrt{Z_x - \delta_n^*}}.$$

So we can rewrite the set $\{X_k^n - \mathrm{E}X_k^n > \varepsilon s_n\}$, using the fact that $\mathrm{E}X_k^n = O\left(\frac{1}{\sqrt{n \log n}}\right)$, and noticing that the last inclusion is valid for all n big enough:

$$\left\{\frac{\sqrt{Z_x}\mathbf{1}_{\{Z_x>0\}} - \sqrt{Z_x - \delta_n^*}\mathbf{1}_{\{Z_x>\delta_n^*\}}}{\delta_n^*} > \varepsilon s_n \sqrt{n \log n} + O(1)\right\}$$
$$\subseteq \left\{\frac{\mathbf{1}_{\{Z_x>\delta_n^*\}}}{2\sqrt{Z_x - \delta_n^*}} > \varepsilon s_n \sqrt{n \log n} - \frac{1}{\sqrt{\delta_n^*}} + O(1)\right\}$$
$$\subseteq \left\{Z_x < \frac{\delta_n^*}{4\Big(\sqrt{\delta_n^* n \log n}\varepsilon s_n - 1 + O(\sqrt{\delta_n^*})\Big)^2} + \delta_n^*\right\}.$$

Since $\frac{\delta_n^*}{4\left(\sqrt{\delta_n^* n \log n}\varepsilon s_n - 1 + O(\sqrt{\delta_n^*})\right)^2} \sim \frac{1}{n \log n}$, for all n big enough we conclude:

$$\{X_k^n - \mathrm{E}X_k^n > \varepsilon s_n\} \subseteq \left\{Z_x < \frac{1}{n \log n} + \delta_n^*\right\} =: A_n.$$

Now we see that the set $\{X_k^n - \mathrm{E}X_k^n < -\varepsilon s_n\}$ is negligible, because the object on the left-hand side in the following expression is nonnegative:

$$\left\{\frac{\sqrt{Z_x}\mathbf{1}_{\{Z_x>0\}} - \sqrt{Z_x - \delta_n^*}\mathbf{1}_{\{Z_x>\delta_n^*\}}}{\delta_n^*} < -\varepsilon s_n \sqrt{n \log n} + O(1)\right\}.$$

Therefore we see that the integral over the set $\left|X_k^n - \mathrm{E}X_k^n\right| > \varepsilon s_n$ for all n big enough has the same behavior as integrating over the set: A_n. Using i.i.d.:

$$\sum_{k=1}^{n} \mathrm{E}[(X_k^n - \mathrm{E}X_k^n)^2 \mathbf{1}_{A_n}] \leq \underbrace{2n\mathrm{E}[(X_k^n)^2 \mathbf{1}_{A_n}]}_{(1)} + \underbrace{2n(\mathrm{E}X_k^n)^2 \mathrm{P}(A_n)}_{(2)} \to 0.$$

Since $\mathrm{E}X_k^n = O(\frac{1}{\sqrt{n \log n}})$ we see immediately that (2) goes to 0. Now we prove that (1) also does. We need to look at the following upper bound:

$$\frac{(\delta_n^*)^{-2}}{\log n}\mathrm{E}\left[\left\{Z_x \mathbf{1}_{\{Z_x>0\}} + (Z_x - \delta_n^*)\mathbf{1}_{\{Z_x>\delta_n^*\}}\right\}\mathbf{1}_{A_n}\right] \leq \frac{2(\delta_n^*)^{-2}}{\log n}\mathrm{E}\left[Z_x \mathbf{1}_{\{Z_x>0\}\cap A_n}\right]$$

From (2.4.2), we recall that: $\mathrm{E}Z_x\mathbf{1}_{0<Z_x<\varepsilon} = O(\varepsilon^2)$ for $\varepsilon \downarrow 0$. We conclude that:

$$\frac{(\delta_n^*)^{-2}}{\log n}\mathrm{E}\left[Z_x\mathbf{1}_{\{0<Z_x<\delta_n^*+\frac{1}{n\log n}\}}\right] \sim \frac{1}{\log n}\left(\frac{n}{\log n}\right)^{\frac{1}{\gamma_x}}\left(\left(\frac{\log n}{n}\right)^{\frac{1}{2\gamma_x}} + \frac{1}{n\log n}\right)^2 \sim$$

$$\sim \frac{1}{\log n} + \frac{1}{\log n}\left(\frac{n}{\log n}\right)^{1/2\gamma_x}\frac{1}{n\log n} + \frac{1}{\log n}\left(\frac{n}{\log n}\right)^{1/\gamma_x}\frac{1}{n^2\log^2 n} \to 0.$$

For $\gamma_x > \frac{1}{2}$. Therefore the Lindeberg condition is satisfied for $Z_n(1)$. □

Proof of (2.2.16). We start with the class of functions $h_\theta : [0,\infty) \to \mathbb{R}$:

$$\mathscr{H} = \left\{h_\theta \ : \ h_\theta(z) = 2\sqrt{z-x}\mathbf{1}_{[x,\infty)}(z) - 2\sqrt{z-\theta}\mathbf{1}_{[\theta,\infty)}(z) - a_0(x-\theta), \theta \geq 0\right\}$$

and, for $\eta > 0$, its subclass $\mathscr{H}_\eta = \{h_\theta \in \mathscr{H} : |\theta - x| \leq \eta\}$. In [52], they show that the envelope of $\mathscr{H}_\eta$, $H_\eta(z) = 2\left(\sqrt{z-x}\mathbf{1}_{[x,\infty)}(z) - \sqrt{z-x-\eta}\mathbf{1}_{[x-\eta,\infty)}(z)\right) + a_0\eta$, satisfies for all η small (cf. (11) in [52]):

$$\int H_\eta^2(z)g(z)\,dz \leq -2g(x)\eta^2\log\eta. \tag{2.4.11}$$

The first thing that needs to be proved is that for $|\theta - x|$ small, we have:

$$\int h_\theta(z) - h_x(z)\,dG(z) \lesssim -|\theta - x|^{\gamma_x+1}.$$

Using a Taylor expansion for $\int h_\theta(z)dG(z)$ around x (note $h_x(z) = 0$), we get, using the definition of U: $U(x) = \frac{\pi}{2m_0}\int_0^x(1-F(y))\,dy = \int_0^x V(y)\,dy$, that there exists $\eta_0 > 0$ such that:

$$\int h_\theta(z)\,dG(z) = 2\int \sqrt{z-x}\mathbf{1}_{[x,\infty)}(z) - \sqrt{z-\theta}\mathbf{1}_{[\theta,\infty)}(z)\,dG(z) - V(x)(\theta-x)$$

$$= U(x) - U(\theta) - V(x)(x-\theta) \overset{(T)}{=} \frac{(\theta-x)\pi}{2m_0}\int_0^1 F(x+u(\theta-x)) - F(x)\,du$$

$$\overset{(2.2.3)}{=} -\frac{\pi}{2m_0}|\theta-x|^{\gamma_x+1}(K_x + o(1)),$$

for all θ with $|\theta - x| < \eta_0$ and for some constant $K > 0$. Note that here we used the same reasoning as in (2.2.11) and we also used assumption (2.2.3).

Now we show the weak consistency of $T_n(a_0 + \delta_n a)$ for $x = V^{-1}(a_0)$. This will follow from Theorem 3, Theorem 4 in [52] and the switch relation (2.2.8). In other words we prove: $\forall\ \varepsilon > 0$ then $\mathrm{P}(|T_n(a_0+\delta_n a) - x| > \varepsilon) \to 0$. Indeed using the switch relation and consistency of $\hat{V}_n$ we get:

$$\begin{aligned}
&\mathrm{P}(|T_n(a_0+\delta_n a) - x| > \varepsilon)\\
&\quad \leq \mathrm{P}(T_n(V(x)+\delta_n a) > \varepsilon + x) + \mathrm{P}(T_n(V(x)+\delta_n a) < x - \varepsilon)\\
&\quad = \mathrm{P}\left(\hat{V}_n(x+\varepsilon) > V(x) + \delta_n a\right) + \mathrm{P}\left(\hat{V}_n(x-\varepsilon) < V(x) + \delta_n a\right) \longrightarrow 0.
\end{aligned}$$

As by consistency $\hat{V}_n(x+\varepsilon) \to V(x+\varepsilon)$, $\delta_n \downarrow 0$ and V is strictly decreasing at x, it will never be the case that $V(x+\varepsilon) \geq V(x)$, nor $V(x-\varepsilon) \leq V(x)$. Note that V is already decreasing by

definition, and it cannot be constant in a neighborhood of x, as we would get a contradiction with the fact that F is not constant in a neighborhood of x.
Now we will proceed by deriving the rate of convergence of $T_n(a_0 + \delta_n a)$ to $x = V^{-1}(a_0)$. We will do it by a standard technique using a peeling argument, combined later with what we showed above.

First it needs to be proved that $\forall\, \varepsilon > 0$ there exists $\{M_n\}$ sequence of stochastically bounded random variables and a positive constant η_1 such that:

$$2\left|\int \left\{\sqrt{z-x}\mathbf{1}_{[x,\infty)}(z) - \sqrt{z-\theta}\mathbf{1}_{[\theta,\infty)}(z)\right\} d\left(\mathbb{G}_n - G\right)(z)\right| \tag{2.4.12}$$

$$\leq \varepsilon|x-\theta|^{\gamma_x+1} + (\delta_n^*)^{\gamma_x+1} M_n, \tag{2.4.13}$$

for all θ with $|\theta - x| < \eta_1$, and γ as in assumption (2.2.3). Remember that x has been fixed and that $\delta_n = n^{-1/2}\sqrt{\log n}$, whereas $\delta_n^* = (\delta_n)^{1/\gamma_x}$. Now we proceed with the proof of (2.4.12). Fix $\varepsilon > 0$, and let $\{M_n(\omega)\}$ be the random variable defined omega-wise as the infimum of all positive numbers η so that:

$$\left|\int h_\theta(z)\, d\left(\mathbb{G}_n - G\right)(z;\omega)\right| \leq \varepsilon|x-\theta|^{\gamma_x+1} + (\delta_n^*)^{\gamma_x+1}\eta, \tag{2.4.14}$$

holds for all $\theta \in [x - \eta_1, x + \eta_1]$. If we define, for $n, j \geq 1$,

$$A(n,j) = \{\theta \geq 0 : (j-1)\delta_n^* \leq |\theta - x| \leq j\delta_n^*\},$$

we can write, for all $n \geq 1$ and ν constant:

$$\begin{aligned}
&\mathrm{P}\{M_n > \nu\} \leq \\
&\leq \sum_{j=1}^{\eta_1/\delta_n^*} \mathrm{P}\left\{\sup_{\theta\in A(n,j)} (\delta_n^*)^{-(\gamma_x+1)}\left|\int h_\theta(z)\, d\left(\mathbb{G}_n - G\right)(z)\right| > \varepsilon(j-1)^{\gamma_x+1} + \nu\right\} \\
&\leq \sum_{j=1}^{\eta_1/\delta_n^*} \frac{\mathrm{E}\left\{\sup_{|\theta-x|\leq j\delta_n^*} (\delta_n^*)^{-2(\gamma_x+1)}\left[\int h_\theta(z)\, d\left(\mathbb{G}_n - G\right)(z)\right]^2\right\}}{\left[\varepsilon(j-1)^{\gamma_x+1} + \nu\right]^2}.
\end{aligned} \tag{2.4.15}$$

By maximal inequality 3.1 (ii) in [65] and relation (2.4.11) (with $\eta = \delta_n^* j$), $\exists$ a $C > 0$ such that, for each $j \leq \eta_1/\delta_n^*$ and using that $(\delta_n^*)^{-2(\gamma_x+1)} = (\delta_n^*)^{-2}\delta_n^{-2}$:

$$\begin{aligned}
&\mathrm{E}\left\{\sup_{|\theta-x|\leq j\delta_n^*} (\delta_n^*)^{-2(\gamma_x+1)}\left[\int h_\theta(z)\, d\left(\mathbb{G}_n - G\right)(z)\right]^2\right\} \\
&= (\delta_n^*)^{-2}\frac{n}{\log n}\mathrm{E}\left\{\sup_{|\theta-x|\leq j\delta_n^*}\left[\int h_\theta(z)\, d\left(\mathbb{G}_n - G\right)(z)\right]^2\right\} \\
&\leq (\delta_n^*)^{-2}\frac{J(1)^2}{\log n}\left(-2g(x)j^2(\delta_n^*)^2\log\left(\delta_n^*\cdot j\right)\right) = Cg(x)j^2 + O\left((\log n)^{-1}\right).
\end{aligned}$$

where $J(1)$ is the uniform entropy integral appearing from the application of the maximal inequality 3.1 (ii) in [65]. We argue that $J(1)$ is finite in the proof of Lemma 2.2.1. Applying this inequality to (2.4.15), it follows that we can upper bound it by: $\sum_{j=1}^{\eta_1/\delta_n^*} \frac{Cg(x)j^2 + o(1)}{\left[\varepsilon(j-1)^{\gamma_x+1}+\nu\right]^2}$, by

choosing ν sufficiently large, $\mathrm{P}\{M_n > \nu\}$ can be made arbitrarily small uniformly in n, which proves (2.4.12).

We are now ready to conclude that $T_n(a_0 + \delta_n a)$ is a δ_n^*-consistent estimator for $x = V^{-1}(a_0)$. From the above we know with probability tending to 1, $|T_n(a_0 + \delta_n a) - x| < \eta_0 \wedge \eta_1$, for $n \to \infty$. If $|T_n(a_0 + \delta_n a) - x| < \eta_0 \wedge \eta_1$, take ε in relation (2.4.14) equal to $\frac{\pi K}{4m_0}$ and obtain that:

$$\begin{aligned} 0 &= \int h_x(z)\, d\mathbb{G}_n(z) \leq \int h_{T_n(a_0+\delta_n a)}(z)\, d\mathbb{G}_n(z) \\ &\leq \int h_{T_n(a_0+\delta_n a)}(z)\, dG(z) + \frac{\pi K_x}{4m_0} |x - T_n(a_0 + \delta_n a)|^{\gamma_x+1} + (\delta_n^*)^{\gamma_x+1} M_n \\ &= \left(\frac{\pi K_x}{4m_0} - \frac{\pi K_x}{2m_0} + o(1)\right) |x - T_n(a_0 + \delta_n a)|^{\gamma_x+1} + (\delta_n^*)^{\gamma_x+1} M_n, \end{aligned}$$

where the sequence $\{M_n\}$ is stochastically bounded by the above part. This proves that $T_n(a_0 + \delta_n a) - x = O_p(\delta_n^*)$ for $n \to \infty$. □

Lemma 2.4.2. *Condition* (2.2.20) *implies that F^* is not constant in a neighborhood of x and that: $\int_x^\infty \frac{dF^*(s)}{\sqrt{s-x}} < \infty$.*

Proof. Assume (2.2.20). Suppose F^* is constant in a neighborhood of x then the limit in (2.2.20) is zero, a contradiction. From condition (2.2.20), it follows that we can fix $K > \epsilon > 0$ knowing that there exists $\delta > 0$ such that $\forall\, t \in \mathbb{R}^+ : t \leq \delta$ then: $(K-\epsilon)t^{\gamma_x} \leq F^*(x+t) - F^*(x) \leq (K+\epsilon)t^{\gamma_x}$. Let us set a grid that goes from x to $x+\delta$ as: $x_0 = x+\delta \geq x_1 \geq ... \geq x$, $x_j = x + 2^{-j}\delta$. It is enough to look at:

$$\int_x^{x+\delta} \frac{dF^*(s)}{\sqrt{s-x}} \leq \sum_{j=0}^{\infty} \int_{x_{j+1}}^{x_j} \frac{dF^*(s)}{\sqrt{s-x}} \leq \sum_{j=0}^{\infty} \frac{(K+\epsilon)2^{-\gamma_x(j+1)}}{\sqrt{2^{-(j+1)}\delta}} - \sum_{j=0}^{\infty} \frac{(K-\epsilon)2^{-\gamma_x j}}{\sqrt{2^{-(j+1)}\delta}},$$

where the last expression is finite as $\gamma_x > \frac{1}{2}$. □

Lemma 2.4.3. *If F^* is locally Hölder continuous of degree $\gamma_x > 1/2$ around x_0, then the density of the observations g is bounded and continuous around x_0.*

Proof. Fix any z in a neighborhood of x_0, so that $\exists\, \eta > 0 : F^*$ is Hölder continuous on $(z-\eta, z+\eta)$. Set $z_j = z + \eta 2^{-j}$ for $j = 0, 1, 2...$ so that $z_0 = z + \eta \geq z_1 \geq ... \geq z$ and therefore $z_j - z_{j+1} = \eta 2^{-j-1}$. We have (note $m_0 < \infty$ by (2.2.6)):

$$\begin{aligned} g(z)m_0 &= \int_z^\infty \frac{1}{\sqrt{x-z}} dF^*(x) = \sum_{j=0}^{\infty} \int_{z_{j+1}}^{z_j} \frac{1}{\sqrt{x-z}} dF^*(x) + \int_{z+\eta}^{\infty} \frac{1}{\sqrt{x-z}} dF^*(x) \\ &\leq \sum_{j=0}^{\infty} \frac{(F^*(z_j) - F^*(z_{j+1}))}{\sqrt{z_{j+1} - z}} + \frac{1}{\sqrt{\eta}} \leq \frac{1}{\sqrt{\eta}} \left(\sum_{j=0}^{\infty} \frac{2^{-\gamma(j+1)}}{2^{-\frac{1}{2}(j+1)}} + 1 \right), \end{aligned}$$

where the last expression is $< \infty$ if and only if $\gamma_x > \frac{1}{2}$. Now we know that the function g is bounded around x_0. We have to prove that it is continuous. Using the same reasoning as in the previous part we have that:

$$g(z) = \frac{1}{m_0} \underbrace{\sum_{j=0}^{\infty} \int_{z_{j+1}}^{z_j} \frac{1}{\sqrt{x-z}} dF^*(x)}_{(1)} + \frac{1}{m_0} \underbrace{\int_{z+\eta}^{\infty} \frac{1}{\sqrt{x-z}} dF^*(x)}_{(2)}.$$

Since in the previous part of the proof we have been able to bound (1) with the convergent series independent of z: $\sum_{j=0}^{\infty} \frac{2^{-\gamma(j+1)}}{2^{-\frac{1}{2}(j+1)}}$, then the Weierstrass criterion for uniform convergence of series of functions (M-test) yields that the series (1) converges uniformly. Now we need to show that $z \mapsto \int_{z_{j+1}}^{z_j} \frac{1}{\sqrt{x-z}} dF^*(x)$ are continuous $\forall\ j$. We also need to show that (2) is continuous. First of all, notice that the following function: $f(s) = \frac{\mathbf{1}_{[\eta 2^{-j-1}, \eta 2^{-j}]}(s)}{\sqrt{s}}$, is continuous on the compact set $[\eta 2^{-j-1}, \eta 2^{-j}]$, thus uniformly continuous. By taking $s = x - z$, one can retrieve $\frac{\mathbf{1}_{[z+\eta 2^{-j-1}, z+\eta 2^{-j}]}(x)}{\sqrt{x-z}}$. Now take δ so that whenever $|z_1 - z_2| < \delta$, by uniform continuity we have:

$$\left| \frac{\mathbf{1}_{[z_1+\eta 2^{-j-1}, z_1+\eta 2^{-j}]}(x)}{\sqrt{x-z_1}} - \frac{\mathbf{1}_{[z_2+\eta 2^{-j-1}, z_2+\eta 2^{-j}]}(x)}{\sqrt{x-z_2}} \right| < \varepsilon.$$

Then we have, using the triangle inequality for integrals:

$$\left| \int \frac{\mathbf{1}_{[z_1+\eta 2^{-j-1}, z_1+\eta 2^{-j}]}(x)}{\sqrt{x-z_1}} - \frac{\mathbf{1}_{[z_2+\eta 2^{-j-1}, z_2+\eta 2^{-j}]}(x)}{\sqrt{x-z_2}} dF^*(x) \right| \leq \varepsilon \int dF^*(x) = \varepsilon.$$

Finally, (2) is continuous because $\frac{\mathbf{1}_{[z+\eta,\infty)}(x)}{\sqrt{x-z}} = \frac{\mathbf{1}_{[\eta,\infty)}(x-z)}{\sqrt{x-z}}$ is uniformly continuous on $[\eta, \infty)$. The Weierstrass M-test together with the uniform limit theorem yield continuity of g. □

Lemma 2.4.4. *For every $a \in \mathbb{R}$, $\mathrm{P}(\hat{s}(a) = 0) = 0$, where $\hat{s}(a) := \arg\max_{t \in I_x} \{tX - \frac{\pi K_x}{2m_0}|t|^{1+\gamma_x} - at\}$ and $X \sim N(0,\ g(x)/(2\gamma_x))$, $x > 0$.*

Proof. Set $c := X - a$ and $\varphi(s) := cs - \frac{\pi K_x}{2m_0}|s|^{1+\gamma_x}$. Since φ is strictly concave on each half-line and $\varphi(s) \to -\infty$ as $|s| \to \infty$, the unique maximiser is

$$\hat{s}(a) = \mathrm{sign}(c)\Big(\frac{2m_0|c|}{\pi K_x(1+\gamma_x)}\Big)^{1/\gamma_x},$$

a strictly increasing function of $c = X - a \sim N(-a,\ g(x)/(2\gamma_x))$. Since this normal has no atoms, neither does the law of $\hat{s}(a)$; in particular $\mathrm{P}(\hat{s}(a) = 0) = 0$. The cdf of the limit is continuous, with explicit form $\mathrm{P}(\hat{s}(a) \leq x) = \mathrm{P}(c \leq \frac{\pi K_x}{2m_0}(1+\gamma_x)\,\mathrm{sign}(x)|x|^{\gamma_x})$. □

***Proof of Lemma** 2.2.1.* Because

$$U_n(s) := \frac{2}{n} \sum_{i=1}^{n} \left\{ \sqrt{(Z_i)_+} - \sqrt{(Z_i - s)_+} \right\},$$

the stated convergence is equivalent to saying that the class of functions $z \to \sqrt{(z)_+} - \sqrt{(z-s)_+}$, $s \geq 0$ is Donsker. From Lemma 2.6.16 in [103], it follows that the class of functions $\{z - s : s \geq 0\}$ is VC, next by Lemma 2.6.18 (ii) in [103] we see that the class $\{(z-s)_+ : s \geq 0\}$ is VC. Again by Lemma 2.6.18 (vii), (iv) and (v) in [103] we conclude that the class of functions $\{\sqrt{(z)_+} - \sqrt{(z-s)_+} : s \geq 0\}$ is VC. Such class has envelope: $0 \leq \sqrt{(z)_+} - \sqrt{(z-s)_+} \leq \sqrt{(z)_+}$. Therefore the class is P-Donsker for any P with $\int_0^\infty z dP(z) < \infty$, which is the case here by assumption (2.2.6) with $k = \frac{3}{2}$. Now we prove that $\mathbb{Z}$ is a Gaussian process with a.s. continuous sample paths. Consider:

$$\mathrm{E}\left(\mathbb{Z}(t_j) - \mathbb{Z}(t_i)\right)^2 = \mathrm{E}\left| \sqrt{(Z - t_j)_+} - \sqrt{(Z - t_i)_+} \right|^2 \leq |t_j - t_i|.$$

Where we use the fact that the square root is Hölder continuous of degree $1/2$.

Since the intrinsic semimetric $d(s,t) := (\mathrm{E}\,(\mathbb{Z}(s) - \mathbb{Z}(t))^2)^{1/2} \leq \sqrt{|s-t|}$, we obtain sample path continuity of $\mathbb{Z}$.

By the very same computation as above, we deduce that $\mathrm{Var}(\mathbb{Z}(t) - \mathbb{Z}(s)) \neq 0$ for $s \neq t$ and since $\mathbb{Z}$ is indexed by a σ-compact metric space, we obtain from Lemma 2.6 in [65] that the maxima of the sample paths of $\mathbb{Z}$ are a.s. unique. Finally the covariance structure of $\mathbb{Z}$ is given by direct computation:

$$\mathrm{Cov}\,(\mathbb{Z}(t), \mathbb{Z}(s)) = \mathrm{Cov}\Big(\sqrt{(Z)_+} - \sqrt{(Z_t)_+},\ \sqrt{(Z)_+} - \sqrt{(Z_s)_+}\Big).$$

□

Proof of (2.2.31). Note that:

$$\sup_{s \in K_n^c} \left\{ Z_n(s) - \sqrt{n} h_x(s) - as \right\} \leq \sup_{s \geq 0} Z_n(s) - \inf_{\substack{s < \underline{x} - \varepsilon_n \\ s > \bar{x} + \varepsilon_n}} \left\{ \sqrt{n} h_x(s) + as \right\},$$

where the first term on the right-hand side is $O_p(1)$ by the convergence given in Lemma 2.2.1. The second infimum is the minimum over the infima over the intervals $[0, \underline{x} - c]$, $[\underline{x} - c, \underline{x} - \varepsilon_n]$, $[\bar{x} + \varepsilon_n, \bar{x} + 1]$ and $[\bar{x} + 1, \infty)$ for a constant $\varepsilon_1 < c < \underline{x}$. Because F^* is constant in $[\underline{x}, \bar{x}]$, we have $U(s) - U(x) - V(x)(s - x) = 0$ for $s \in [\underline{x}, \bar{x}]$ and from the concavity of U:

$$U(s) - U(x) - V(x)(s - x) < 0, \quad \text{for} \quad s \notin [\underline{x}, \bar{x}].$$

Again from the fact that U is concave and because $[\underline{x}, \bar{x}]$ is maximal:

$$\begin{aligned} &\forall\, \delta > 0,\ \exists\, \eta > 0\ :\ U(s) - U(x) - V(x)(s - x) < -\eta |s - \underline{x}| \quad \text{for} \quad s < \underline{x} - \delta, \\ &\forall\, \delta > 0,\ \exists\, \eta > 0\ :\ U(s) - U(x) - V(x)(s - x) < -\eta |s - \bar{x}| \quad \text{for} \quad s > \bar{x} + \delta. \end{aligned}$$

Therefore using these three facts stated above we have: on $[\bar{x} + 1, \infty)$,

$$\sqrt{n} h_x(s) + as \geq \sqrt{n}\eta |s - \bar{x}| - |a| s \gtrsim \sqrt{n} \to \infty$$

and on $[\bar{x} + \varepsilon_n, \bar{x} + 1]$,

$$\sqrt{n} h_x(s) + as \geq \sqrt{n} h_x(\bar{x} + \varepsilon_n) - |a| \to \infty$$

by the construction of ε_n. The other two intervals can be handled similarly. □

Proof of (2.2.33). First note:

$$\begin{aligned} 0 &\leq \sup_{s \in F \cap K_n} \left\{ Z_n(s) - \sqrt{n} h_x(s) - as \right\} - \sup_{s \in F \cap K} \{ Z_n(s) - as \} \\ &\leq \sup_{s \in F \cap K_n} \{ Z_n(s) - as \} - \sup_{s \in F \cap K} \{ Z_n(s) - as \} \end{aligned}$$

Define the process:

$$\widetilde{Z}_n(s) := \begin{cases} Z_n(s) - as, & \text{if } s \in [\underline{x}, \bar{x}] \\ Z_n(\underline{x}) - a\underline{x}, & \text{if } s \in [0, \underline{x}] \\ Z_n(\bar{x}) - a\bar{x}, & \text{if } s \in [\bar{x}, \infty) \end{cases}$$

because: $\sup_{s\in F\cap K_n}\widetilde{Z}_n(s) = \sup_{s\in F\cap K}\widetilde{Z}_n(s) = \sup_{s\in F\cap K}\{Z_n(s)-as\}$, then

$$\begin{aligned}
&\left|\sup_{s\in F\cap K_n}\{Z_n(s)-as\} - \sup_{s\in F\cap K}\{Z_n(s)-as\}\right| \\
&= \left|\sup_{s\in F\cap K_n}\{Z_n(s)-as\} - \sup_{s\in F\cap K_n}\widetilde{Z}_n(s)\right| \le \sup_{s\in F\cap K_n}\left|Z_n(s)-as-\widetilde{Z}_n(s)\right| \\
&\le \sup_{s\in F\cap K_n\cap[0,\underline{x}]}\left|Z_n(s)-as-\widetilde{Z}_n(\underline{x})\right| \vee \sup_{s\in F\cap K_n\cap[\bar{x},\infty)}\left|Z_n(s)-as-\widetilde{Z}_n(\bar{x})\right| \to 0
\end{aligned}$$

where we used the asymptotic equicontinuity of the Z_n. □

***Proof of Theorem** 2.2.1.* The claim is a consequence of $1-F(x) = V(x)/V(0)$ and by analogy, we construct the estimator using the adaptive isotonic estimator. Apply Slutsky's Lemma together with Theorem 2.2.3 (cf. Corollary 1 in [52]) to get Theorem 2.2.1.a.. To prove Theorem 2.2.1.b., it is possible to follow the same steps as in the proof of Theorem 2.2.4 and show the joint convergence, for all $a, b \in \mathbb{R}$:

$$\begin{aligned}
&\mathrm{P}\Big(\sqrt{n}\left(\hat{V}_n(x)-V(x)\right)\le a,\ \sqrt{n}\left(\hat{V}_n(0)-V(0)\right)\le b\Big)\to \\
&\to \mathrm{P}\Big(\arg\max_{s\in K_x}\{\mathbb{Z}_x(s)-as\}\le x, \arg\max_{s\in K_0}\{\mathbb{Z}_0(s)-bs\}\le 0\Big) \\
&= \mathrm{P}\left(L_x\le a, L_0\le b\right),
\end{aligned}$$

where in the last step we used the switch relation. Using again Slutsky's Lemma with the above convergence and the asymptotic independence of $\hat{V}_n(x)$ and $\hat{V}_n(0)$ (which follows from the fact that $K_x\cap K_0=\emptyset$), we get claim. □

Lemma 2.4.5. *x is a continuity point of the distribution of $\hat{s}(a) := \arg\max_{s\in K_x}\{\mathbb{Z}_x(s)-as\}$, i.e.*

$$\mathrm{P}(\hat{s}(a)=x)=0.$$

Proof. It suffices to prove

$$\sup_{s>x}\{\mathbb{Z}_x(s)-a(s-x)\}>0 \quad \text{a.s.}, \tag{2.4.16}$$

since $\mathbb{Z}_x(x)=0$ and the sup over $s<x$ is handled symmetrically.

Set $\psi_s(z) := 2\sqrt{(z-x)_+} - 2\sqrt{(z-s)_+}$ so that $\mathbb{Z}_x(s) = \mathbb{G}\psi_s$ where $\mathbb{G}$ is the G-Brownian bridge. Define

$$\sigma(s) := \int_s^\infty \frac{dG(z)}{z-x}, \qquad s>x,$$

which decreases continuously from $+\infty$ (as $s\downarrow x$) to 0 (as $s\to\infty$); write $s_K := \sigma^{-1}(K)$ for $K>0$. Under the standing assumptions, we prove

$$\limsup_{K\to\infty}\frac{\mathbb{G}\psi_{s_K}}{(s_K-x)\sqrt{2K\log\log K}} = 1 \quad \text{a.s.}$$

Write $f(z) := \dfrac{1}{\sqrt{z-x}}\,\mathbf{1}_{z>x}$ and decompose

$$\psi_s(z) = (s-x)\,f(z)\,\mathbf{1}_{[s,\infty)}(z) + r_s(z),$$

where

$$r_s(z) := \begin{cases} 0 & z < x, \\ 2\sqrt{z-x} & x < z < s, \\ \dfrac{s-x}{\sqrt{z-x}}\,\phi\Big(\dfrac{z-s}{z-x}\Big) & z > s, \end{cases} \qquad \phi(t) := \frac{1-\sqrt{t}}{1+\sqrt{t}}.$$

Represent $\mathbb{G}$ via the G-Brownian motion $\mathbb{W}$: $\mathbb{G}h = \mathbb{W}h - (\mathbb{W}\mathbf{1})\int h\,dG$. Then

$$\mathbb{G}\psi_s = (s-x)\,\mathbb{W}(f\mathbf{1}_{[s,\infty)}) - (s-x)\,(\mathbb{W}\mathbf{1})\int_{[s,\infty)} f\,dG + \mathbb{G}r_s. \tag{2.4.17}$$

Leading term. Since $\mathbb{W}(f\mathbf{1}_{[\sigma^{-1}(\cdot),\infty)})$ is a standard Brownian motion in the parameter (variance = $\sigma(s)$), the LIL gives

$$\limsup_{K\to\infty} \frac{\mathbb{W}(f\mathbf{1}_{[s_K,\infty)})}{\sqrt{2K\log\log K}} = 1 \quad \text{a.s.}$$

Second term. $\mathbb{W}\mathbf{1} = O(1)$ a.s. and $\int_{[s,\infty)} f\,dG \le \int f\,dG < \infty$, so the second term in (2.4.17) is $O(s_K - x) = o((s_K - x)\sqrt{2K\log\log K})$.

Remainder. Since $1 - t \lesssim \phi(t) \lesssim 1 - t$ for $0 \le t \le 1$,

$$\int_x^s r_s^2\,dG \lesssim \|g\|_\infty \int_x^s (z-x)\,dz \lesssim (s-x)^2,$$
$$\int_s^\infty r_s^2\,dG \lesssim (s-x)^2 \int_s^\infty \frac{(s-x)^2}{(z-x)^3}\,dG(z) \lesssim (s-x)^2.$$

Hence $\mathrm{Var}(\mathbb{G}r_{s_K}) \le c(s_K - x)^2$, and for every $\varepsilon > 0$,

$$\sum_{K=1}^\infty \mathrm{P}\left(\frac{|\mathbb{G}r_{s_K}|}{(s_K-x)\sqrt{2K\log\log K}} \ge \varepsilon\right) \le \sum_{K=1}^\infty 2\exp\Big(-\tfrac{\varepsilon^2}{c^2}K\log\log K\Big) < \infty.$$

Borel–Cantelli gives $|\mathbb{G}r_{s_K}| = o((s_K - x)\sqrt{2K\log\log K})$ a.s.

Combining all three parts yields the lemma, since

$$\limsup_{K\to\infty} \frac{\mathbb{G}\psi_{s_K} - a(s_K - x)}{(s_K - x)\sqrt{2K\log\log K}} = 1 > 0 \quad \text{a.s.},$$

so $\mathbb{G}\psi_{s_K} - a(s_K - x) > 0$ for all sufficiently large K, and hence

$$\sup_{s>x}\{\mathbb{Z}_x(s) - a(s-x)\} > 0 \quad \text{a.s.} \qquad \square$$

□

2.4.2 Proofs complemental statements for the LAN expansion.

***Proof of Lemma** 2.2.2.* It is enough to consider the case $h_1 > 0$ and $h_2 > 0$, (depicted in Figure 2.6 around x) and we prove in this case that $F^*_{h_n}$ is non-decreasing for all n big enough. We check only around x on ${}^{l}\mathrm{K}_n^x$, as around 0 is the same. From (2.2.36), for any $h_2 \ge 0$,

$\exists\, N^*$ such that $\forall\, n \geq N^*$, all $x_1, x_2 \in K_n^x$, $x_1 < x_2$: $F^*(x_2) - F^*(x_1) \geq \frac{h_2}{\sqrt{n \log n}} \log\left\{\frac{x-x_2}{x-x_1}\right\}$, and therefore eventually:

$$D_{h_n}\left(F^*_{h_n}(x_2) - F^*_{h_n}(x_1)\right) = F^*(x_2) - F^*(x_1) - \frac{h_2}{\sqrt{n \log n}} \log\left\{\frac{x-x_2}{x-x_1}\right\} \geq 0.$$

□

Lemma 2.4.6. *Let* $\zeta_n(x) := \int_x^\infty \frac{\mathbf{1}_{[\delta_n \leq |v| \leq \eta_n]}}{v\sqrt{v-x}}\, dv$, *then:*

$$\zeta_n(x) = 0 \quad \text{if } x > \eta_n, \qquad \left|\zeta_n(x)\right| \lesssim \frac{\delta_n}{(-x)^{3/2}} + \frac{1}{\sqrt{\eta_n}} \quad \text{if } -\eta_n \leq x \leq -\delta_n,$$
$$\sup_{[\delta_n, \eta_n]} \left|\zeta_n(x) - \frac{\pi}{\sqrt{x}}\right| \leq \frac{\pi}{\sqrt{\eta_n}}, \quad \sup_{[-\delta_n, \delta_n]} \left|\zeta_n(x)\right| \leq \frac{\pi}{\sqrt{\delta_n}}, \quad \sup_{x \leq -\eta_n} \left|\zeta_n(x)\right| \lesssim \frac{1}{\sqrt{\eta_n}}.$$

Remark 2.4.1. Another more compact way of stating the above lemma is (cf. [49]):

$$\zeta_n(x) = \begin{cases} 0, & \text{if } x > \eta_n, \\ \pi x^{-1/2} + O\left(\eta_n^{-1/2}\right), & \text{if } \delta_n \leq x \leq \eta_n, \\ O\left(\delta_n^{-1/2}\right), & \text{if } -\delta_n \leq x \leq \delta_n, \\ O\left(\delta_n(-x)^{-3/2}\right) + O\left(\eta_n^{-1/2}\right), & \text{if } -\eta_n \leq x \leq -\delta_n, \\ O\left(\eta_n^{-1/2}\right), & \text{if } x \leq -\eta_n. \end{cases} \tag{2.4.18}$$

Proof. 1. If $x > \eta_n$, then clearly $\zeta_n(x) = 0$, due to the indicator function.

2. If $\delta_n \leq x \leq \eta_n$, using (2.2.44) and the triangle inequality:

$$\begin{aligned} \left|\zeta_n(x) - \frac{\pi}{\sqrt{x}}\mathbf{1}_{x\geq 0}\right| &= \left|\int_x^{\eta_n} \frac{1}{v\sqrt{v-x}}\, dv - \int_x^\infty \frac{1}{v\sqrt{v-x}}\, dv\right| \\ &= \left|\int_{\eta_n}^\infty \frac{1}{v\sqrt{v-x}}\, dv\right| \leq \int_{\eta_n}^\infty \frac{1}{v\sqrt{v-\eta_n}}\, dv = \frac{\pi}{\sqrt{\eta_n}}. \end{aligned}$$

3. If $-\delta_n \leq x \leq \delta_n$, using (2.2.44):

$$\left|\zeta_n(x)\right| = \int_{\delta_n}^{\eta_n} \frac{1}{v\sqrt{v-x}}\, dv \leq \int_{\delta_n}^\infty \frac{1}{v\sqrt{v-x}}\, dv \leq \int_{\delta_n}^\infty \frac{1}{v\sqrt{v-\delta_n}}\, dv = \frac{\pi}{\sqrt{\delta_n}}.$$

4. If $-\eta_n \leq x \leq -\delta_n$, $J(y) := \int_y^1 \frac{1}{v}\left(\frac{1}{\sqrt{v+1}} - \frac{1}{\sqrt{1-v}}\right) dv$, $H(y) := \int_1^y \frac{1}{v\sqrt{v+1}} dv$:

$$\begin{aligned} \zeta_n(x) &= \int_x^{-\delta_n} \frac{1}{v\sqrt{v-x}}\, dv + \int_{\delta_n}^{\eta_n} \frac{1}{v\sqrt{v-x}}\, dv \\ &= \int_{\delta_n}^{-x} \frac{1}{v}\left(\frac{1}{\sqrt{v-x}} - \frac{1}{\sqrt{-v-x}}\right) dv + \int_{-x}^{\eta_n} \frac{1}{v\sqrt{v-x}}\, dv \\ &= \frac{1}{\sqrt{-x}}\left(J\left(\frac{\delta_n}{-x}\right) + H\left(\frac{\eta_n}{-x}\right)\right). \end{aligned}$$

Because $H(\infty) = -J(0) = \log(3 + 2\sqrt{2})$ we obtain:

$$\begin{aligned}
\left|\zeta_n(x)\right| &= \left|\frac{1}{\sqrt{-x}}\left(J\left(\frac{\delta_n}{-x}\right) - J(0) - H(\infty) + H\left(\frac{\eta_n}{-x}\right)\right)\right| = \\
&= \left|\frac{1}{\sqrt{-x}}\left(\int_0^{\delta_n/-x} \frac{1}{v}\left(\frac{1}{\sqrt{v+1}} - \frac{1}{\sqrt{1-v}}\right) dv + \int_{\eta_n/-x}^{\infty} \frac{1}{v\sqrt{v+1}}\, dv\right)\right| \\
&\lesssim \frac{\delta_n}{(-x)^{3/2}} + \frac{1}{\sqrt{\eta_n}}
\end{aligned}$$

5. If $x \leq -\eta_n$, using the same reasoning as above:

$$\left|\zeta_n(x)\right| = \frac{1}{\sqrt{-x}}\left|\int_{\delta_n/-x}^{\eta_n/-x} \frac{1}{v}\left(\frac{1}{\sqrt{v+1}} - \frac{1}{\sqrt{1-v}}\right)\right| dv \leq \frac{\log(3+2\sqrt{2})}{\sqrt{\eta_n}}.$$

□

Proof of (2.2.46). Fix $\epsilon > 0$, using the triangle inequality, we can just look separately at $p \in \{0, x\}$ (where for $j = 1$, $p = 0$; for $j = 2$, $p = x$ and recall $h_{j,n}$, $j = 1, 2$ as in (2.2.38)):

$$\begin{aligned}
&\mathrm{P}\left(\max_{1\leq i\leq n}\left|\frac{h_{j,n}\zeta_{j,n}(Z_i - p)}{g(Z_i)}\right| > \epsilon\right) \leq \sum_{i=1}^{n} \mathrm{P}\left(\left|\frac{h_{j,n}\zeta_{j,n}(Z_i - p)}{g(Z_i)}\right| > \epsilon\right) \\
&\leq \sum_{i=1}^{n} \mathrm{P}\left(\left|\frac{\zeta_{j,n}(Z_i - p)}{g(Z_i)}\right| > \epsilon\frac{\sqrt{n\log n}}{h_j}\right), \qquad j = 1, 2.
\end{aligned}$$

Consider now the expansion in (2.2.43). Define:

$$A_n^{i,x} := \{n^{-\frac{1}{2\gamma_x}} \leq Z_i - x \leq \eta_n\}, \tag{2.4.19}$$

and analogously $A_n^{i,0}$. Now we can use (2.2.43) and, in particular, notice that the term in (2.2.42) is the one of greatest order. So if we show that for $j = 1, 2$ and accordingly $p \in \{0, x\}$:

$$\sum_{i=1}^{n} \mathrm{P}\left(\frac{\pi}{g(Z_i)\sqrt{Z_i - p}}\mathbf{1}_{A_n^{i,p}} > \epsilon\frac{\sqrt{n\log n}}{h_j}\right) \to 0,$$

where $A_n^{i,p}$ is as in (2.4.19), we obtain the claim. The above summation is zero for all n big enough because the indicator on $A_n^{i,p}$ requires $1/\sqrt{Z_i - p}$ to be bounded by $n^{\frac{1}{4\gamma_p}} \leq \sqrt{n\log n}(C + o(1))$, for $\gamma_p > \frac{1}{2}$, $C > 0$ and n big enough. Moreover the density of the observations on $A_n^{i,p}$ is bounded away from zero. □

Proof of Lemma *2.2.3.* We will structure the proof as follows: first, we show that the expression inside the parenthesis in (2.2.47) is mean zero, secondly, we show that by using the expansion (2.2.43) the term in line (2.2.42) is the one that contributes to the normal behavior (both around x and around 0) and lastly we show that all the other terms converge to zero in probability. First note:

$$\int_0^{\infty}\int_z^{\infty} \frac{\chi_{2,n}(v-x)dv}{2g(z)\sqrt{v-z}} g(z)dz = \int_0^{\infty}\int_0^{v} \frac{\chi_{2,n}(v-x)}{2\sqrt{v-z}}\, dzdv = \int_0^{\infty} \frac{\chi_{2,n}(v-x)}{v^{-\frac{1}{2}}}\, dv$$

and it is immediate to see, using the definition of m_{h_n} and m_0, that:

$$D_{h_n} m_{h_n} = m_0 + h_{1,n} \int_0^\infty \sqrt{v} \chi_{1,n}(v)\, dv + h_{2,n} \int_0^\infty \sqrt{v} \chi_{2,n}(v - x)\, dv. \tag{2.4.20}$$

This proves that the expression inside the parenthesis in the statement of Lemma 2.2.3 is mean zero. Let $\overline{k}_n^x$ be the mean of the term first term on the right hand side of the equality in (2.2.42) multiplied $1/(2m_{h_n}\sqrt{n \log n})$ and analogously $\overline{k}_n^0$ the same mean with 0 instead of x. Define (independently of i and recall (2.2.38)):

$$\overline{k}_n := \overline{k}_n^x + \overline{k}_n^0 = \mathrm{E}\left\{ \frac{\pi}{2m_{h_n}} \frac{h_{2,n}}{g(Z_i)\sqrt{Z_i - x}} \mathbf{1}_{A_n^{i,x}} \right\} + \mathrm{E}\left\{ \frac{\pi}{2m_{h_n}} \frac{h_{1,n}}{g(Z_i)\sqrt{Z_i}} \mathbf{1}_{A_n^{i,0}} \right\},$$

where we used again the definition of $A_n^{i,p}$ in (2.4.19). Note that we just isolate the term $\overline{k}_n$ from the centering $(m_{h_n} D_{h_n} - m_0)/(m_{h_n})$ which already contained it. We need to isolate $\overline{k}_n$ for the next part (note $\overline{k}_n = o(1)$). Finally, denote by Σ_h the asymptotic variance in (2.2.47). Using the fact that $D_{h_n} = 1 + o(1)$, we can prove that for $h_{i,n},\ i = 1, 2$ as in (2.2.38):

$$\frac{\pi}{2m_{h_n}} \sum_{i=1}^n \left\{ \frac{h_{1,n}}{g(Z_i)\sqrt{Z_i}} \mathbf{1}_{A_n^{i,0}} + \frac{h_{2,n}}{g(Z_i)\sqrt{Z_i - x}} \mathbf{1}_{A_n^{i,x}} - \overline{k}_n \right\} \rightsquigarrow N(0, \Sigma_h).$$

We apply the Lindeberg Central Limit Theorem. If we look at:

$$s_n^2 := \frac{\pi^2}{4m_{h_n}^2} \sum_{i=1}^n \mathrm{Var}\left\{ \frac{h_{1,n}}{g(Z_i)\sqrt{Z_i}} \mathbf{1}_{A_n^{i,0}} + \frac{h_{2,n}}{g(Z_i)\sqrt{Z_i - x}} \mathbf{1}_{A_n^{i,x}} \right\},$$

the covariance of the two terms above inside the parenthesis goes to zero because for all n sufficiently large $\mathbf{1}_{A_i^0} \mathbf{1}_{A_i^x} = 0$. Now we focus on the variance:

$$\sum_{i=1}^n \mathrm{Var}\left\{ \frac{h_2}{\sqrt{n \log n}} \frac{\pi}{2m_{h_n} g(Z_i)\sqrt{Z_i - x}} \mathbf{1}_{A_n^{i,x}} \right\} = \frac{h_2^2 \pi^2}{8\gamma_x m_0^2 g(x)} + o(1). \tag{2.4.21}$$

Indeed using the i.i.d. assumption and leaving out the constants $(h_i^2\pi^2)/(4m_0^2)$ (recall $m_{h_n}/m_0 \to 1$), we obtain that the left hand side of (2.4.21) is equivalent to:

$$\frac{1}{\log n} \mathrm{E}\left[\frac{1}{g^2(Z)(Z - x)} \mathbf{1}_{A_n^x} \right] - \left(\frac{1}{\sqrt{\log n}} \mathrm{E}\left[\frac{1}{g(Z)\sqrt{Z - x}} \mathbf{1}_{A_n^x} \right] \right)^2.$$

Now analyze the first term by taking limit in n. Using the continuity of g at x and that $g(x) > 0$, together with $0 < \eta_n < 1$, and $\eta_n \gtrsim \frac{1}{\log^\alpha(n)}$:

$$\lim_{n\to\infty} \frac{1}{\log n} \int_{x + \frac{1}{n^{1/2\gamma_x}}}^{x+\eta_n} \frac{1}{z - x} \frac{1}{g(z)}\, dz = \frac{1}{2\gamma_x g(x)}. \tag{2.4.22}$$

The second term converges to zero because: $\lim_{n\to\infty} \frac{1}{\sqrt{\log n}} \int_{x + \frac{1}{n^{1/2\gamma_x}}}^{x+\eta_n} \frac{1}{\sqrt{z-x}}\, dz = 0$ just by integrating and taking the limit. This proves the claim about the variance. Using the same relation (2.4.21) with 0 instead of x, we obtain:

$$s_n^2 = \frac{h_1^2 \pi^2}{8\gamma_0 m_0^2 g(0)} + \frac{h_2^2 \pi^2}{8\gamma_x m_0^2 g(x)} + o(1). \tag{2.4.23}$$

Now we look at Lindeberg condition; define:

$$B_n^\epsilon := \left\{ \left| \frac{\pi}{2m_{h_n}\sqrt{n\log n}} \left(\frac{\mathbf{1}_{A_n^{i,x}}}{g(Z_i)\sqrt{Z_i - x}} + \frac{\mathbf{1}_{A_n^{i,0}}}{g(Z_i)\sqrt{Z_i}} \right) - \bar{k}_n \right| > \epsilon s_n \right\}.$$

Then $\forall\, \epsilon > 0$, as $n \to \infty$, we show:

$$\frac{1}{s_n^2} \sum_{i=1}^{n} \mathrm{E}\left[\left(\frac{\pi h_{1,n} \mathbf{1}_{A_n^{i,0}}}{2g(Z_i) m_{h_n} \sqrt{Z_i}} + \frac{\pi h_{2,n} \mathbf{1}_{A_n^{i,x}}}{2g(Z_i) m_{h_n} \sqrt{Z_i - x}} - \bar{k}_n \right)^2 \mathbf{1}_{B_n^\epsilon} \right] \to 0. \tag{2.4.24}$$

By using the triangle inequality, the fact that $\bar{k}_n = O(1/\sqrt{n\log n})$, that from (2.4.23) s_n converges to a constant, and that $\mathbf{1}_{A_n^{i,p}}/g(Z_i) = \mathbf{1}_{A_n^{i,p}}/g(p) + o(1)$, for $p \in \{0, x\}$ and all n big enough, we can work with the superset of B_n^ϵ:

$$C_{n,i}^\epsilon := \left\{ \frac{\mathbf{1}_{A_n^{i,x}}}{(g(x) + o(1))\sqrt{Z_i - x}} + \frac{\mathbf{1}_{A_n^{i,0}}}{(g(0) + o(1))\sqrt{Z_i}} > \frac{\epsilon 2 m_0 s_n}{\pi (n\log n)^{-1/2}} + O(1) \right\}.$$

Since $(\bar{k}_n)^2 = o(1)$, the term in (2.4.24) involving $(\bar{k}_n)^2$ goes to zero. Alike, the cross-product terms go to zero. So the only terms that matter are the squared terms. Thus we look at:

$$\frac{\pi^2}{4m_{h_n}^2 s_n^2 n\log n} \sum_{i=1}^{n} \mathrm{E}\left[\frac{1}{g^2(Z_i)(Z_i - p)} \mathbf{1}_{A_n^{i,p}} \mathbf{1}_{C_{n,i}^\epsilon} \right], \qquad \text{for } p \in \{0, x\},$$

which are zero for all n sufficiently large as for all such n we have $\mathbf{1}_{A_n^{i,p}} \mathbf{1}_{C_{n,i}^\epsilon} = 0$.

Now we proceed with the last part of the proof and we show that all the terms that we did not take into account in the computations above converge to zero by using the Bernstein inequality (see for instance [83] page 193). We will do it only for the terms deriving from (2.2.43) involving the perturbation around x. The analogous terms deriving from the perturbation around 0 can be treated in the same way.

To apply Bernstein's inequality we need to center the random variables of interest. The means of all the terms below are already contained in $(m_{h_n} D_{h_n} - m_0)/(m_{h_n})$ (as it was for $\bar{k}_n$), thus we only need to isolate them for the next part. Recall also the assumptions on F^* and g (continuous at x and 0 and $g(x) > 0, g(0) > 0$).

1. Let $A_n^i := \{\frac{1}{n^{1/(2\gamma_x)}} \le Z_i - x \le \eta_n\}$ and $Y_n^i = O(\eta_n^{-1/2})(\mathbf{1}_{A_n^i}/g(Z_i) - \mathrm{E}\mathbf{1}_{A_n^i}/g(Z_i))$ then $\mathrm{Var}(Y_n^i) = O(1)$. Define:

$$\widetilde{Y}_n^i = \eta_n^{1/2} Y_n^i.$$

Then $\sqrt{\mathrm{Var}(Y_n^i)} = \eta_n^{-1/2} \sqrt{\mathrm{Var}(\widetilde{Y}_n^i)}$ and by the assumptions on g we have $|\widetilde{Y}_n^i| \le L$ bounded for all n big enough, therefore $\forall\, \varepsilon > 0$:

$$\mathrm{P}\left(\frac{1}{\sqrt{n\log n}} \left| \sum_{i=1}^{n} Y_n^i \right| > \varepsilon \right) = \mathrm{P}\left(\frac{\sqrt{\mathrm{Var}(Y_n^i)}}{\sqrt{\mathrm{Var}(Y_n^i)}\sqrt{n\log n}} \left| \sum_{i=1}^{n} Y_n^i \right| > \varepsilon \right) \tag{2.4.25}$$

$$\le \mathrm{P}\left(\frac{O(1)}{\sqrt{\mathrm{Var}(\widetilde{Y}_n^i)}\sqrt{n\log n}} \left| \sum_{i=1}^{n} \widetilde{Y}_n^i \right| > \varepsilon \right) \lesssim 2e^{-\frac{\varepsilon^2 \log n}{2}\left(1 + \frac{\varepsilon L \sqrt{\log n}}{O(\sqrt{n_n})\sqrt{n}}\right)^{-1}} \to 0.$$

2. Let $A_n^i := \left\{-\frac{1}{n^{1/(2\gamma_x)}} \le Z_i - x \le \frac{1}{n^{1/(2\gamma_x)}}\right\}$, $Y_n^i = O(n^{\frac{1}{4\gamma_x}})(\mathbf{1}_{A_n^i}/g(Z_i) - \mathrm{E}\mathbf{1}_{A_n^i}/g(Z_i))$ then $\mathrm{Var}(Y_n^i) = O(1)$. Define:

$$\widetilde{Y}_n^i = n^{-\frac{1}{4\gamma_x}} Y_n^i.$$

Then $\sqrt{\mathrm{Var}(Y_n^i)} = n^{\frac{1}{4\gamma_x}} \sqrt{\mathrm{Var}(\widetilde{Y}_n^i)}$ and by the assumptions on g we have $|\widetilde{Y}_n^i| \le L$ bounded for all n big enough, using the same steps as for the previous term in (2.4.25) and because $\gamma_x > \frac{1}{2}$, $\forall\, \varepsilon > 0$:

$$\mathrm{P}\left(\frac{1}{\sqrt{n\log n}}\left|\sum_{i=1}^n Y_n^i\right| > \varepsilon\right) \lesssim 2e^{-\frac{\varepsilon^2\log n}{2}\left(1+\frac{\varepsilon L\sqrt{\log n}}{O(n^{-1/4\gamma_x})\sqrt{n}}\right)^{-1}} \to 0.$$

3. Let $A_n^i := \{-\eta_n \le Z_i - x \le -\frac{1}{n^{1/(2\gamma_x)}}\}$, $Y_n^i = \frac{O(n^{-1/2\gamma_x})}{g(Z_i)(x-Z_i)^{3/2}}\mathbf{1}_{A_n^i} - \mathrm{E}\frac{O(n^{-1/2\gamma_x})}{g(Z_i)(x-Z_i)^{3/2}}\mathbf{1}_{A_n^i}$, then $\mathrm{Var}(Y_n^i) = O(1)$. Define:

$$\widetilde{Y}_n^i = n^{-\frac{1}{4\gamma_x}} Y_n^i.$$

Then $\sqrt{\mathrm{Var}(Y_n^i)} = n^{\frac{1}{4\gamma_x}} \sqrt{\mathrm{Var}(\widetilde{Y}_n^i)}$ and by the assumptions on g we have $|\widetilde{Y}_n^i| \le L$ bounded for all n big enough, using the same steps as for the previous term in (2.4.25) and because $\gamma_x > \frac{1}{2}$, $\forall\, \varepsilon > 0$:

$$\mathrm{P}\left(\frac{1}{\sqrt{n\log n}}\left|\sum_{i=1}^n Y_n^i\right| > \varepsilon\right) \lesssim 2e^{-\frac{\varepsilon^2\log n}{2}\left(1+\frac{\varepsilon L\sqrt{\log n}}{O(n^{-1/4\gamma_x})\sqrt{n}}\right)^{-1}} \to 0.$$

4. Let $A_n^i := \{Z_i - x \le -\eta_n\}$, and $Y_n^i = O(\eta_n^{-\frac{1}{2}})(\mathbf{1}_{A_n^i}/g(Z_i) - \mathrm{E}\mathbf{1}_{A_n^i}/g(Z_i))$ then $\mathrm{Var}(Y_n^i) = O(\eta_n^{-1})\left(\int_0^x 1/g(z)\,dz - x^2\right)$ (using the assumptions on g we have $\int_0^x 1/g(z)\,dz > 0$). Define:

$$\widetilde{Y}_n^i = \eta_n^{1/2} Y_n^i.$$

Then $\sqrt{\mathrm{Var}(Y_n^i)} = \eta_n^{-1/2}\sqrt{\mathrm{Var}(\widetilde{Y}_n^i)}$ and by the assumptions on g we have $|\widetilde{Y}_n^i| \le L$ bounded for all n big enough, therefore $\forall\, \varepsilon > 0$:

$$\mathrm{P}\left(\frac{1}{\sqrt{n\log n}}\left|\sum_{i=1}^n Y_n^i\right| > \varepsilon\right) \le \mathrm{P}\left(\frac{O(\eta_n^{-\frac{1}{2}})}{\sqrt{\mathrm{Var}(Y_n^i)}\sqrt{n\log n}}\left|\sum_{i=1}^n Y_n^i\right| > \varepsilon\right) =$$
$$\mathrm{P}\left(\frac{O(1)}{\sqrt{n}\sqrt{\mathrm{Var}(\widetilde{Y}_n^i)}}\left|\sum_{i=1}^n \widetilde{Y}_n^i\right| > \varepsilon\sqrt{\eta_n\log n}\right) \lesssim 2e^{-\frac{\varepsilon^2\log n}{2(\log n)^\alpha}\left(1+\frac{\varepsilon L\sqrt{\log n}}{\sigma\eta_n^{-1/2}\sqrt{n}}\right)^{-1}} \to 0.$$

where $\alpha \in (0,1)$ and $\sigma^2 := \int_0^x 1/g(z)\,dz - x^2$.

5. The last term can be treated identically to 1.

□

Proof of (2.2.49). Let A_n^i as in (2.4.19). We are going to verify the conditions of Theorem 2.2.6. from [30] to prove:

$$\frac{1}{n\log n}\left\{\sum_{i=1}^{n}\frac{1}{g^2(Z_i)(Z_i-x)}\mathbf{1}_{A_n^i}-n\int_{x+\frac{1}{n^{1/2\gamma_x}}}^{x+\eta_n}\frac{1}{z-x}\frac{1}{g(z)}\,dz\right\}\to 0. \tag{2.4.26}$$

By (2.4.22) we know how $\frac{1}{\log n}\int_{x+\frac{1}{n^{1/2\gamma_x}}}^{x+\eta_n}\frac{1}{z-x}\frac{1}{g(z)}dz$ behaves; for this reason, if we show (2.4.26), we get the claim. We verify condition (i) of Theorem 2.2.6. in [30]:

$$n\int_{\left\{\frac{1}{g^2(z)(z-x)}\mathbf{1}_{\left\{\eta_n^{-1}\le\frac{1}{z-x}\le n^{1/2\gamma_x}\right\}}>n\log n\right\}}g(z)\,dz=0,$$

because the set over which we are integrating is empty for all n big enough. Now we check condition (ii) of Theorem 2.2.6. from [30]; we have, using Leibniz integral rule together with de L'Hôpital rule and the fact that $\frac{1}{2\gamma_x}+1>\frac{1}{\gamma_x}$:

$$\lim_{n\to\infty}\frac{1}{n^2\log^2 n}n\int_{x+\frac{1}{n^{1/2\gamma_x}}}^{x+\eta_n}\frac{1}{(z-x)^2}\frac{1}{g^3(z)}\,dz=0.$$

The above computations now yield (2.4.26). By combining (2.4.26) and (2.4.22), together with: $\frac{\pi^2}{4m_{h_n}^2}\to\frac{\pi^2}{4m_0^2}$ we obtain the claim. □

Proof of (2.2.50). Using (2.4.20), the definition of $h_{i,n}$, $i=1,2$ in (2.2.38), the fact that $\int_0^\infty\sqrt{v}\chi_n(v)\,dv=O(\sqrt{\eta_n})$ and that $\int_0^\infty\sqrt{v}\chi_n(v-x)\,dv=O(\eta_n)$:

$$\left(\frac{m_0-m_{h_n}D_{h_n}}{m_{h_n}D_{h_n}}\right)^2=O\left(\frac{\eta_n}{n\log n}\right)=O\Big(\frac{1}{n\log^{1+\alpha}(n)}\Big).$$

Now we look only at the terms derived from the perturbation around x (i.e. the ones for $p=x$), as the one derived from the perturbation around 0 behave alike. It is enough to consider all the squared terms in $(R_n^{i,x})^2$. The ones that are at most $O(\eta_n^{-1})$, go to zero as $\frac{nO(\log^\alpha n)}{n\log n}\to 0$, $\alpha\in(0,1)$. The remaining terms converge in probability to 0 by Bernstein's inequality as in the proof of Lemma 2.2.3. □

Proof of Lemma *2.2.4.*

$$\sqrt{\frac{n}{\log n}}\left(F^*_{h_n}(x)-F^*(x)\right)=\frac{\sqrt{n}}{D_{h_n}\sqrt{\log n}}\left((1-D_{h_n})F^*(x)\right)+\frac{h_1\int_0^x\chi_{1,n}(v)\,dv}{D_{h_n}\log n}$$
$$+\frac{h_2\int_0^x\chi_{2,n}(v-x)\,dv}{D_{h_n}\log n}=\left(-\frac{h_1\int_0^\infty\chi_{1,n}(v)\,dv}{D_{h_n}\log n}-\frac{h_2\int_0^\infty\chi_{2,n}(v-x)\,dv}{D_{h_n}\log n}\right)F^*(x)$$
$$+\frac{h_1\int_0^x\chi_{1,n}(v)\,dv}{D_{h_n}\log n}+\frac{h_2\int_0^x\chi_{2,n}(v-x)\,dv}{D_{h_n}\log n}.$$

By using: $\int_0^\infty\chi_{1,n}(v)\,dv=-(1+o(1))\log n^{-1/(2\gamma_0)}$, $\int_0^\infty\chi_{2,n}(v-x)\,dv=o(1)$, $\int_0^x\chi_{1,n}(v)\,dv=-(1+o(1))\log n^{-1/(2\gamma_0)}$, and that $\int_0^x\chi_{2,n}(v-x)\,dv=(1+o(1))\log n^{-1/(2\gamma_x)}$, the above for all n big is equal to (using also $1/D_{h_n}\to 1$):

$$=\frac{h_1}{D_{h_n}\log n}(1+o(1))(F^*(x)-1)\log n^{-\frac{1}{2\gamma_0}}-\frac{h_2}{D_{h_n}\log n}(o(1))F^*(x)+$$

$$\frac{h_2}{D_{h_n}\log n}(1+o(1))\log n^{-\frac{1}{2\gamma_x}} \to \frac{h_1}{2\gamma_0}(1-F^*(x)) - \frac{h_2}{2\gamma_x} = h^\top \begin{bmatrix} \frac{1}{2\gamma_0}(1-F^*(x)) \\ -\frac{1}{2\gamma_x} \end{bmatrix}.$$

□

***Proof of Theorem** 2.2.2.* Because for any neighborhood in supremum norm of F^* there exists $n \in \mathbb{N}$ such that the "least favorable" perturbation $F^*_{h_n}$ (cf. (2.2.37)) is in such neighbourhood, the expression in (2.2.5) is lower-bounded by the expression in (2.2.40) (cf. pag. 117 in [102]). □

2.4.3 Proofs of Section 2.3.

To prove Proposition 2.3.1, we need an additional lemma that will be essential to show that the estimators (2.3.12)-(2.3.15) are indeed the projections. The proof of this lemma uses similar reasoning as contained in [40] and [41]. Below for a function f, $f(t^-)$ indicates the left limit of f at t.

Lemma 2.4.7. *Let $\overline{V} \in \mathbb{L}_1$ be given, define $V \mapsto Q^{\overline{V}}(V)$ by* (2.3.4) *and $\hat{V} \in \mathcal{V}_{\underline{x},\bar{x}}$. Suppose the following conditions hold true, for $\underline{x} < \bar{x}$:*

1. $\int_0^t (\hat{V}(x) - \overline{V}(x))\,dx \geq 0, \quad \forall\, t \in [0,\infty) \cap [\underline{x},\bar{x}]^c,$
2. $\int_0^t (\hat{V}(x) - \overline{V}(x))\,dx = 0, \quad \forall\, t \in [0,\infty) \cap [\underline{x},\bar{x}]^c$ *with* $\hat{V}(t) < \hat{V}(t^-)$.
3. *The measure defined via $\hat{V}$ and the one defined via $\overline{V}$ are compactly supported on the positive real line.*

Then $\hat{V} = \underset{V \in \mathcal{V}_{\underline{x},\bar{x}}}{\arg\min}\, Q^{\overline{V}}(V)$.

Proof. Let us recall the concept of Gateaux differentiability at a point $V \in \mathcal{V}$, for some function space $\mathcal{V}$. Let $Q : \mathcal{V} \mapsto \mathbb{R}$ be an arbitrary functional. It is called Gateaux differentiable at the point $V \in \mathcal{V}$ if the limit:

$$\partial Q(V,h) = \lim_{\varepsilon \to 0} \frac{Q(V+\varepsilon h) - Q(V)}{\varepsilon}$$

exists for every h such that $V + \varepsilon h \in \mathcal{V}$ for small enough ε. By straightforward computations, one can verify that:

$$Q^{\overline{V}}(\hat{V} + \varepsilon \mathbf{1}_{[0,t]}) - Q^{\overline{V}}(\hat{V}) = 2\varepsilon \int_0^t \hat{V}(x)\,dx + \varepsilon^2 t - 2\varepsilon \int_0^t \overline{V}(x)\,dx$$

Therefore the Gateaux derivative of $Q^{\overline{V}}$ is given by:

$$\partial Q^{\overline{V}}(\hat{V},h) = 2\int_0^\infty h(x)\left(\hat{V}(x) - \overline{V}(x)\right)dx \tag{2.4.27}$$

Note that, since $\partial Q^{\overline{V}}(\hat{V}, \mathbf{1}_{[0,t]}) = \int_0^t \hat{V}(x) - \overline{V}(x)\,dx$, the first two conditions of Lemma 2.4.7 entail the Gateaux derivative of the functional $Q^{\overline{V}}$ along the test functions $\mathbf{1}_{[0,t]}$, $\forall\, t \geq 0$ but $t \notin [\underline{x},\bar{x}]$.

Now we show that for any $V \in \mathcal{V}_{\underline{x},\bar{x}}$, $\partial Q^{\overline{V}}(\hat{V}, V - \hat{V}) \geq 0$. First notice:

$$\hat{V}(x) - \overline{V}(x) = \frac{d(\hat{U} - \overline{U})}{dx}(x)$$

Without loss of generality, we assume that the two supports of the measures defined via $\hat{V}$ and $\overline{V}$ are contained in $[0, M_1]$ and $[0, M_2]$. Defining $M := \max\{M_1, M_2\}$. Using integration by parts and assumption 3, $\hat{U}(0) - \overline{U}(0) = \hat{U}(M) - \overline{U}(M) = 0$:

$$\begin{aligned}
\partial Q^{\overline{V}}(\hat{V}, V - \hat{V}) &= 2 \int_0^{\infty} \Big(V(x) - \hat{V}(x)\Big)\Big(\hat{V}(x) - \overline{V}(x)\Big)\, dx \\
&= 2 \int_0^{M} \Big(V(x) - \hat{V}(x)\Big) d(\hat{U} - \overline{U})(x) \\
&= 2 \underbrace{\Big(V(x) - \hat{V}(x)\Big)\Big(\hat{U}(x) - \overline{U}(x)\Big)\Big|_0^M}_{=0} + \\
&\qquad - 2 \int_0^{M} \Big(\hat{U}(x) - \overline{U}(x)\Big) d\Big(V(x) - \hat{V}(x)\Big)
\end{aligned}$$

By assumption 2, the support of the measure $-d\hat{V}$ is included in the set $\{x : \overline{U}(x) = \hat{U}(x)\}$ (cf. Lemma 1 in [40]), therefore: $\int_0^{\infty} \Big(\hat{U}(x) - \overline{U}(x)\Big) d\hat{V}(x) = 0$. On the other hand, because $V \in \mathcal{V}_{\underline{x},\bar{x}}$ the negative measure defined based on V does not have mass in $[\underline{x}, \bar{x}]$ and then, by using assumption 1, we obtain:

$$\int_0^{\infty} \Big(\overline{U}(x) - \hat{U}(x)\Big) dV(x) \geq 0$$

which proves indeed $\partial Q^{\overline{V}}(\hat{V}, V - \hat{V}) \geq 0$. Now define the convex function $t \mapsto u(t) = Q^{\overline{V}}(\hat{V} + t(V - \hat{V}))$ where $t \in [0, 1]$. Using (2.4.27) and what we just proved, we have that $u'(0) = \partial Q^{\overline{V}}(\hat{V}, V - \hat{V}) \geq 0$. Since u is convex, the fact $u'(0) \geq 0$ implies $u(1) \geq u(0)$, which indeed is $Q^{\overline{V}}(V) \geq Q^{\overline{V}}(\hat{V})$. □

***Proof of Proposition** 2.3.1.* The proof of (2.3.5) is provided in [52] (c.f. Lemma 2 in [52]).

Proof of (2.3.6). First, we show that $\hat{V}_n^{\Pi} \in \mathcal{V}_{\underline{x},\bar{x}}$. By definition, for $x \in [\underline{x}, \bar{x}]$, for $\underline{x} < \bar{x}$:

$$\hat{V}_n^{\Pi}(x) = \frac{U_n^*(\bar{x}) - U_n^*(\underline{x})}{\bar{x} - \underline{x}}$$

Let Z_k^* for $k \in \{1, ..., m-1\}$ be the points where U_n^* changes slope in $[\underline{x}, \bar{x}]$. Denote by Z_0^* the closest point where U_n^* changes slope to the left of $\underline{x}$. Similarly, denote by Z_m^* the closest point where U_n^* changes slope to the right of $\bar{x}$. Now consider the fact that:

$$\begin{aligned}
U_n^*(\underline{x}) &= \frac{U_n(Z_1^*) - U_n(Z_0^*)}{Z_1^* - Z_0^*}(\underline{x} - Z_1^*) + U_n(Z_1^*) \\
U_n^*(\bar{x}) &= \frac{U_n(Z_m^*) - U_n(Z_{m-1}^*)}{Z_m^* - Z_{m-1}^*}(\bar{x} - Z_{m-1}^*) + U_n(Z_{m-1}^*)
\end{aligned}$$

By telescoping we have:

$$U_n(Z_1^*) - U_n(Z_{m-1}^*) = -\sum_{k=2}^{m-1} U_n(Z_k^*) - U_n(Z_{k-1}^*)$$

And therefore all together we have on $[\underline{x}, \bar{x}]$, for $\underline{x} < \bar{x}$:

$$\begin{aligned}\hat{V}_n^{\Pi}(x) := &\sum_{k=2}^{m-1} \frac{Z_k^* - Z_{k-1}^*}{\bar{x} - \underline{x}} \frac{U_n(Z_k^*) - U_n(Z_{k-1}^*)}{Z_k^* - Z_{k-1}^*} + \frac{Z_1^* - \underline{x}}{\bar{x} - \underline{x}} \frac{U_n(Z_1^*) - U_n(Z_0^*)}{Z_1^* - Z_0^*} \\ &+ \frac{\bar{x} - Z_{m-1}^*}{\bar{x} - \underline{x}} \frac{U_n(Z_m^*) - U_n(Z_{m-1}^*)}{Z_m^* - Z_{m-1}^*}\end{aligned} \tag{2.4.28}$$

The fact that $\hat{V}_n^{\Pi} \in \mathcal{V}_{\underline{x},\bar{x}}$ is now clear, because the computations given above prove that $\hat{V}_n^{\Pi}$ on $[\underline{x}, \bar{x}]$ is just a weighted average of the values chosen by the isotonic estimator over $[\underline{x}, \bar{x}]$. This ensures both the monotonicity constraint and right-continuity, other than the fact that $\hat{V}_n^{\Pi}$ is constant on $[\underline{x}, \bar{x}]$. Let us now verify condition 1 of Lemma 2.4.7. Clearly, for $t \leq \underline{x}$, the condition is satisfied as it is zero. Take $t \geq \bar{x}$. The primitive of $\hat{V}$ is U_n^*. We verify the main condition. We split the integral into three bits. Clearly $\int_0^{\underline{x}} \hat{V}_n^{\Pi}(x) - \hat{V}_n(x)\, dx = \int_{\bar{x}}^{t} \hat{V}_n^{\Pi}(x) - \hat{V}_n(x)\, dx = 0$. Now for $[\underline{x}, \bar{x}]$: $\int_{\underline{x}}^{\bar{x}} \hat{V}_n^{\Pi}(x) - \hat{V}_n(x)\, dx = 0$ by (2.3.12). Condition 2 can be verified immediately using the same strategy and condition 3 is verified as well by the definition of the isotonic estimator.

Proof of (2.3.7). If we minimize Q^{V_n} over a space where we do not require the solution to be decreasing (but still constant on each $[x_i, x_{i+1})$), then we can split the minimization problem into different integrals over each $[x_i, x_{i+1})$. When we minimize over $[x_i, x_{i+1})$ we obtain, setting $V(x) = c \in \mathbb{R}$ for $x \in [x_i, x_{i+1})$:

$$\begin{aligned}&\int_{x_i}^{x_{i+1}} V(x)\,(V(x) - 2V_n(x))\, dx = \int_{x_i}^{x_{i+1}} c^2 - 2cV_n(x)dx = \\ &= (x_{i+1} - x_i)c^2 - 2c\frac{1}{n}\sum_{i=1}^{n} \int_{x_i}^{x_{i+1}} (Z_i - x)^{-1/2}\, dx \\ &= (x_{i+1} - x_i)c^2 - 2c\,(U_n(x_{i+1}) - U_n(x_i))\,.\end{aligned}$$

By optimizing, we obtain $c = \frac{U_n(x_{i+1}) - U_n(x_i)}{x_{i+1} - x_i}$.

Proof of (2.3.8). We use Lemma 2.4.7. By (2.3.15), $V_n^{\Pi} \in \mathcal{V}_{\underline{x},\bar{x}}$. Moreover, we have that condition 1 is obtained because it requires that $\forall\, t \in [\underline{x}, \bar{x}]^c \cap [0, \infty)$:

$$\int_0^t V_n^{\Pi}(x) - V_n(x)\, dx = \widetilde{U}_n^*(t) - U_n(t) \geq 0$$

but by construction $\widetilde{U}_n^*(t) \geq \widetilde{U}_n(t) = U_n(t)$ for $t \notin [\underline{x}, \bar{x}]$. For condition 2, if $V_n^{\Pi}(t) < V_n^{\Pi}(t^-)$ then t is a support point of the $\widetilde{U}_n^*$ and we also know by construction that all the support points of $\widetilde{U}_n^*$ satisfy $\widetilde{U}_n^*(t) = U_n(t)$. Condition 3 is trivially satisfied. □

All the proofs that follow make use of the explicit constructions of the estimators (2.3.5)-(2.3.6) given in Proposition 2.3.1. We present the proof of Theorem 2.3.1.b. first, as part of the proof of Theorem 2.3.1.a. can be viewed as a special case of it.

To prove Theorem 2.3.1.a. and Theorem 2.3.1.b. we need the following additional lemma.

Lemma 2.4.8. *Let $\ell^\infty(K)$ be endowed with the uniform norm and $\forall\, s \in K$:*

$$Z_n(s) := \frac{\sqrt{n}}{\bar{x}-\underline{x}}\Big((U_n(s)-U_n(\underline{x}))(\bar{x}-s)-(U_n(\bar{x})-U_n(s))(s-\underline{x}) \\ -(U(s)-U(\underline{x}))(\bar{x}-s)-(U(\bar{x})-U(s))(s-\underline{x})\Big). \tag{2.4.29}$$

Then

$$Z_n \rightsquigarrow \mathbb{Z} \quad \text{in } \ell^\infty(K), \tag{2.4.30}$$

where $\mathbb{Z}$ is the zero mean Gaussian Process given in (2.3.22) *and has a version with continuous sample paths, with unique point of maximum.*

Proof. Because

$$(U_n(s)-U_n(\underline{x}))(\bar{x}-s)-(U_n(\bar{x})-U_n(s))(s-\underline{x}) \\ = \frac{2}{n}\sum_{i=1}^{n}\left\{\left(\sqrt{(Z_i-\underline{x})_+}-\sqrt{(Z_i-s)_+}\right)(\bar{x}-s)-\left(\sqrt{(Z_i-s)_+}-\sqrt{(Z_i-\bar{x})_+}\right)(s-\underline{x})\right\},$$

the stated convergence is equivalent to saying that the class of functions $z \to (\sqrt{(z-\underline{x})_+}-\sqrt{(z-s)_+})(\bar{x}-s)-(\sqrt{(z-s)_+}-\sqrt{(z-\bar{x})_+})(s-\underline{x})$, $s \in K$ is Donsker. From Lemma 2.6.16 in [103], it follows that the class of functions $\{z-s\,:\,s\geq 0\}$ is VC, next by Lemma 2.6.18 (ii) in [103] we see that the class $\{(z-s)_+\,:\,s\geq 0\}$ is VC. Again by Lemma 2.6.18 (vii), (iv) and (v) in [103] we conclude that the class of functions $\{\sqrt{(z-x)_+}-\sqrt{(z-s)_+}\,:\,s\geq 0\}$ is VC. Similarly, we conclude that the class of functions $z\mapsto\sqrt{(z-s)_+}-\sqrt{(z-\bar{x})_+}$ is VC. The two above-mentioned classes of functions have respective envelopes: $0\leq\sqrt{(z-x)_+}-\sqrt{(z-s)_+}\leq\sqrt{(z-x)_+}$ and $0\leq\sqrt{(z-s)_+}-\sqrt{(z-\bar{x})_+}\leq\sqrt{z}$. Therefore these classes are P-Donsker for any P with $\int_0^\infty z dP(z)<\infty$. This is the case by assumption (2.2.6). Again by Lemma 2.6.16 in [103] we have that the classes of functions: $\{(\bar{x}-s)\,:\,s\in K\}$ and $\{(\underline{x}-s)\,:\,s\in K\}$ are VC. They have clearly square integrable envelope as $K=[\underline{x},\bar{x}]$ is compact. We therefore conclude that the uniform entropy integral for the class of functions of our interest is finite, using Lemma 7.21 (i) and (ii) from [91].

To prove that $\mathbb{Z}$ possesses a version with continuous sample paths, it suffices to show:

$$\mathrm{E}\,(\mathbb{Z}(s)-\mathbb{Z}(t))^2 \lesssim |s-t|+|s-t|^2. \tag{2.4.31}$$

This follows from the fact that the square root is Hölder continuous of degree $1/2$ and the fact that:

$$\frac{1}{\bar{x}-\underline{x}}\left(\sqrt{(z-\underline{x})_+}-\sqrt{(z-s)_+})(\bar{x}-s)-(\sqrt{(z-s)_+}-\sqrt{(z-\bar{x})_+})(s-\underline{x})\right) \\ =\sqrt{(z-\underline{x})_+}-\sqrt{(z-s)_+}+\frac{1}{\bar{x}-\underline{x}}(\sqrt{(z-\underline{x})_+}-\sqrt{(z-\bar{x})_+})(\underline{x}-s)$$

The covariance function of $\mathbb{Z}$ is given by, for $Z_x := Z-x$:

$$\mathrm{Cov}\,(\mathbb{Z}(t),\mathbb{Z}(s)) \\ =4\mathrm{Cov}\left(\frac{(\sqrt{(Z_{\underline{x}})_+}-\sqrt{(Z_t)_+})(\bar{x}-t)-(\sqrt{(Z_t)_+}-\sqrt{(Z_{\bar{x}})_+})(t-\underline{x})}{\bar{x}-\underline{x}},\right.$$

$$\frac{(\sqrt{(Z_{\underline{x}})_+} - \sqrt{(Z_s)_+})(\bar{x} - s) - (\sqrt{(Z_s)_+} - \sqrt{(Z_{\bar{x}})_+})(s - \underline{x})}{\bar{x} - \underline{x}}\Big).$$

This coincides with the covariance function of the process in (2.3.22).

By (2.4.31), we deduce that $\mathrm{Var}(\mathbb{Z}(t) - \mathbb{Z}(s)) \neq 0$ for $s \neq t$. Since $\mathbb{Z}$ is indexed by a σ-compact metric space, we obtain from Lemma 2.6 in [65] that the location of the maximum of the sample paths of $\mathbb{Z}$ is a.s. unique. □

***Proof of Theorem** 2.3.1.b.* For $a > 0$, the "switch relation" (see [53]) gives:

$$\hat{V}_n(x) \leq a \quad \Leftrightarrow \quad \arg\max_{s\geq 0}{}^{-} \{U_n(s) - as\} \leq x. \tag{2.4.32}$$

Here the argmax is defined as:

$$\arg\max_{s\geq 0}{}^{-} \{U_n(s) - as\} := \inf\{s \geq 0 : U_n(s) - as \text{ is maximal}\}.$$

Therefore for each fixed $a \in \mathbb{R}$ we can, for all n sufficiently large, write:

$$\begin{aligned}
&\sqrt{n}\left(\hat{V}_n(x) - V_n^{(\underline{x},\bar{x})}(x)\right) \leq a \\
\Leftrightarrow \quad &\arg\max_{s\geq 0}{}^{-} \left\{U_n(s) - \left(V_n^{(\underline{x},\bar{x})}(x) + a/\sqrt{n}\right)s\right\} \leq x \\
\Leftrightarrow \quad &\arg\max_{s\geq 0}{}^{-} \{ \sqrt{n}(U_n(s) - U_n(\underline{x}) - U(s) + U(\underline{x})) + \\
&\qquad - \sqrt{n}((U_n(\bar{x}) - U_n(\underline{x}))/(\bar{x} - \underline{x}) - V(x))(s - \underline{x}) + \\
&\qquad + \sqrt{n}(U(s) - U(\underline{x}) - V(x)(s - \underline{x})) - as\} \leq x.
\end{aligned}$$

Here we used that the location of a maximum of a function is equivariant under translations and invariant under multiplication by a positive number and addition of a constant. We obtain that the above is equivalent to:

$$\arg\max_{s\geq 0}{}^{-} \left\{Z_n(s) - as - \sqrt{n}h_x(s)\right\} \leq x,$$

where:

$$\begin{aligned}
Z_n(s) :=\ & \sqrt{n}(U_n(s) - U_n(\underline{x}) - U(s) + U(\underline{x})) \\
& - \sqrt{n}((U_n(\bar{x}) - U_n(\underline{x}))/(\bar{x} - \underline{x}) - V(x))(s - \underline{x}), \\
h_x(s) :=\ & U(\underline{x}) - U(s) - V(x)(\underline{x} - s).
\end{aligned}$$

One can easily show that Z_n can be written as in (2.4.29). Therefore by Lemma 2.4.8 and the above computations, we conclude that the limiting behavior of:

$$\widetilde{Z}_n(s) := Z_n(s) - as - \sqrt{n}h_x(s),$$

is determined by the process $\mathbb{W}$ defined by, for $\underline{x} < \bar{x}$:

$$\mathbb{W}(s) = \begin{cases} \mathbb{Z}(s) - as, & \text{for } s \in [\underline{x}, \bar{x}] \\ -\infty, & \text{for } s \notin [\underline{x}, \bar{x}]. \end{cases}$$

This is because $h_x(s) > 0$ for all $s \in (0, \underline{x}) \cup (\bar{x}, \infty)$, and therefore multiplied by $-\sqrt{n}$ will go to $-\infty$, while $h_x(s) = 0,\ \forall\ s \in [\underline{x}, \bar{x}]$, and thus on that interval the limiting behavior is completely determined by $\mathbb{Z}(s) - as$.

Let:

$$\hat{s}_n := \arg\max_{s\geq 0}{}^{-} \left\{ Z_n(s) - as - \sqrt{n} h_x(s) \right\}, \tag{2.4.33}$$

$$\hat{s} := \arg\max_{s\geq 0} \{\mathbb{W}(s)\} = \arg\max_{s\in[\underline{x},\bar{x}]} \{\mathbb{Z}(s) - as\}. \tag{2.4.34}$$

To prove convergence in distribution of $\hat{s}_n$ to $\hat{s}$, we use the Portmanteau Lemma and show for every closed subset F:
$\limsup_{n\to\infty} \mathrm{P}(\hat{s}_n \in F) \leq \mathrm{P}(\hat{s} \in F)$. We choose a sequence $\varepsilon_n \downarrow 0$ such that $\sqrt{n} h_x(\underline{x} - \varepsilon_n) \to \infty$ and $\sqrt{n} h_x(\bar{x} + \varepsilon_n) \to \infty$. For $K_n := [\underline{x} - \varepsilon_n, \bar{x} + \varepsilon_n]$ and for $K := [\underline{x}, \bar{x}]$:

$$\begin{aligned}
&\limsup_{n\to\infty} \mathrm{P}(\hat{s}_n \in F) \leq \limsup_{n\to\infty} \mathrm{P}(\hat{s}_n \in F \cap K_n) + \limsup_{n\to\infty} \mathrm{P}(\hat{s}_n \in K_n^c) \\
&\leq \underbrace{\limsup_{n\to\infty} \mathrm{P}\left(\sup_{s\in F\cap K_n} \widetilde{Z}_n(s) \geq \sup_{s\in K_n} \widetilde{Z}_n(s) \right)}_{(1)} + \underbrace{\limsup_{n\to\infty} \mathrm{P}(\hat{s}_n \in K_n^c)}_{(2)}.
\end{aligned} \tag{2.4.35}$$

We show that (2) in (2.4.35) goes to zero. If $\forall\ s \in K^c$ we have Z_n as in (2.4.29), then:

$$\sup_{s\in K_n^c} \widetilde{Z}_n(s) = \sup_{s\in K_n^c} \{Z_n(s) - as - \sqrt{n} h_x(s)\} \xrightarrow{\mathrm{P}} -\infty \tag{2.4.36}$$

Note that:

$$\begin{aligned}
&\sup_{s\in K_n^c} \left\{ Z_n(s) - \sqrt{n} h_x(s) - as \right\} \leq \sup_{s\in K^c} Z_n(s) - \inf_{\substack{s<\underline{x}-\varepsilon_n \\ s>\bar{x}+\varepsilon_n}} \left\{ \sqrt{n} h_x(s) + as \right\} \\
&\leq \sup_{s\in K^c} \{ \sqrt{n}(U_n(s) - U_n(\underline{x}) - U(s) + U(\underline{x})) \} - \inf_{\substack{s<\underline{x}-\varepsilon_n \\ s>\bar{x}+\varepsilon_n}} \left\{ \sqrt{n} h_x(s) + as \right\} \\
&\qquad + \sup_{s\geq 0} \left\{ -\sqrt{n} \left(U_n(\bar{x}) - U_n(\underline{x}) - U(\bar{x}) + U(\underline{x}) \right) \frac{s - \underline{x}}{\bar{x} - \underline{x}} \right\} \\
&= \sup_{s\in K^c} \{ \sqrt{n}(U_n(s) - U_n(\underline{x}) - U(s) + U(\underline{x})) \} - \inf_{\substack{s<\underline{x}-\varepsilon_n \\ s>\bar{x}+\varepsilon_n}} \left\{ \sqrt{n} h_x(s) + as \right\} \\
&\qquad + \sqrt{n} \frac{U_n(\bar{x}) - U_n(\underline{x}) - U(\bar{x}) + U(\underline{x})}{\bar{x} - \underline{x}} \underline{x}
\end{aligned}$$

where the first and the last term on the right-hand side are $O_p(1)$ by the convergence given in Lemma 2.2.1. The rest of the proof is the same as in the proof of Theorem 2.2.4, and it shows that: $-\inf_{\substack{s<\underline{x}-\varepsilon_n \\ s>\bar{x}+\varepsilon_n}} \left\{ \sqrt{n} h_x(s) + as \right\} \to -\infty$. Using (2.4.36):

$$\begin{aligned}
\mathrm{P}(\hat{s}_n \in K_n^c) &\leq \mathrm{P}\left(\sup_{s\in K_n^c} \widetilde{Z}_n(s) \geq \sup_{s\in K_n} \widetilde{Z}_n(s) \right) \\
&\leq \mathrm{P}\left(\sup_{s\in K_n^c} \widetilde{Z}_n(s) \geq \sup_{s\in K} \{Z_n(s) - as\} \right) \to 0.
\end{aligned} \tag{2.4.37}$$

The last convergence is a consequence of (2.4.36) and the convergence $Z_n \rightsquigarrow \mathbb{Z}$ on $\ell^\infty(K)$, combined with the continuous mapping theorem (which implies that the term on the right-hand side of the last inequality is $O_p(1)$).

Now we argue the behavior of (1) in (2.4.35). For that, we need an additional convergence in probability given in (2.4.38):

$$\sup_{s\in F\cap K_n}\left\{Z_n(s) - \sqrt{n}h_x(s) - as\right\} - \sup_{s\in F\cap K}\{Z_n(s) - as\} \xrightarrow{\mathrm{P}} 0. \tag{2.4.38}$$

First note:

$$\begin{aligned} 0 &\leq \sup_{s\in F\cap K_n}\left\{Z_n(s) - \sqrt{n}h_x(s) - as\right\} - \sup_{s\in F\cap K}\{Z_n(s) - as\} \\ &\leq \sup_{s\in F\cap K_n}\{Z_n(s) - as\} - \sup_{s\in F\cap K}\{Z_n(s) - as\}. \end{aligned}$$

Define the process, for $\underline{x} < \bar{x}$:

$$\widetilde{Z}_n(s) := \begin{cases} Z_n(s) - as, & \text{if } s \in [\underline{x}, \bar{x}], \\ Z_n(\underline{x}) - a\underline{x}, & \text{if } s \in [0, \underline{x}], \\ Z_n(\bar{x}) - a\bar{x}, & \text{if } s \in [\bar{x}, \infty). \end{cases}$$

Because: $\sup_{s\in F\cap K_n}\widetilde{Z}_n(s) = \sup_{s\in F\cap K}\widetilde{Z}_n(s) = \sup_{s\in F\cap K}\{Z_n(s) - as\}$, it follows that

$$\begin{aligned} &\left|\sup_{s\in F\cap K_n}\{Z_n(s) - as\} - \sup_{s\in F\cap K}\{Z_n(s) - as\}\right| \\ &= \left|\sup_{s\in F\cap K_n}\{Z_n(s) - as\} - \sup_{s\in F\cap K_n}\widetilde{Z}_n(s)\right| \leq \sup_{s\in F\cap K_n}\left|Z_n(s) - as - \widetilde{Z}_n(s)\right| \\ &\leq \sup_{s\in F\cap K_n\cap[0,\underline{x}]}\left|Z_n(s) - as - \widetilde{Z}_n(\underline{x})\right| \vee \sup_{s\in F\cap K_n\cap[\bar{x},\infty)}\left|Z_n(s) - as - \widetilde{Z}_n(\bar{x})\right| \to 0, \end{aligned}$$

where we used the asymptotic equicontinuity of the Z_n. Using that:

$$\sup_{s\in F\cap K}\{Z_n(s) - as\} \rightsquigarrow \sup_{s\in F\cap K}\{\mathbb{Z}_x(s) - as\},$$

we conclude using Theorem 2.7 (iv) from [102] and (2.4.38):

$$\sup_{s\in F\cap K_n}\left\{Z_n(s) - \sqrt{n}h_x(s) - as\right\} \rightsquigarrow \sup_{s\in F\cap K}\{\mathbb{Z}_x(s) - as\}.$$

Term (1) in (2.4.35) can be upper bounded by:

$$\begin{aligned} &\limsup \mathrm{P}\left(\sup_{s\in F\cap K_n}\left\{Z_n(s) - \sqrt{n}h_x(s) - as\right\} \geq \sup_{s\in K}\{Z_n(s) - as\}\right) \\ &\leq \mathrm{P}\left(\sup_{F\cap K}\{\mathbb{Z}_x(s) - as\} \geq \sup_{s\in K}\{\mathbb{Z}_x(s) - as\}\right) \leq \underbrace{\mathrm{P}(\hat{s} \in K^c)}_{=0} + \mathrm{P}(\hat{s} \in F), \end{aligned} \tag{2.4.39}$$

where we used the above derivations, together with Lemma 2.4.8 and Theorem 2.7 (v) from [102]. Therefore by (2.4.35) combined with (2.4.37), (2.4.39) and the Portmanteau Lemma we obtain the desired convergence in distribution. □

To prove Theorem 2.3.1.a. we need the following additional lemma, which uses similar arguments as in [39].

Lemma 2.4.9. *If U is linear with positive slope on $[\underline{x}, \bar{x}]$, for $\underline{x} < \bar{x}$, and there does not exist a larger interval containing $[\underline{x}, \bar{x}]$ on which U is linear, then:*

$$U_n^*(\bar{x}) - U_n(\bar{x}) = o_p(1/\sqrt{n}) \quad \textit{and} \quad U_n^*(\underline{x}) - U_n(\underline{x}) = o_p(1/\sqrt{n})$$

Proof. Because $U_n(x) = \frac{2}{n}\sum_{i=1}^n \left\{\sqrt{(Z_i)_+} - \sqrt{(Z_i - x)_+}\right\}$, the same arguments as in the proof of Lemma 2.4.8 give in $\ell^\infty(0,\infty)$:

$$\sqrt{n}\,(U_n - U) \rightsquigarrow \widetilde{\mathbb{Z}} \tag{2.4.40}$$

for a Gaussian Process $\widetilde{\mathbb{Z}}$. By assumption U is linear on $[\underline{x}, \bar{x}]$. Let $l(t) = \frac{U(\bar{x})-U(\underline{x})}{\bar{x}-\underline{x}}(t-\underline{x}) + U(\underline{x})$ be the line which is equal to $U(t)$ on $[\underline{x}, \bar{x}]$. Recall that the upper script $*$ denotes the operation of taking the least concave majorant, for instance U^* is the least concave majorant of U. Then consider:

$$\begin{aligned}\sqrt{n}\,(U_n^*(\underline{x}) - U_n(\underline{x})) &= \sqrt{n}\,[\{U_n - U + U - l + l\}^*(\underline{x}) - U_n(\underline{x})] \\ &= \left[\sqrt{n}\,\{U_n - U\} + \sqrt{n}\,\{U - l\}\right]^*(\underline{x}) - \sqrt{n}\,(U_n(\underline{x}) - U(\underline{x}))\,,\end{aligned} \tag{2.4.41}$$

where we use that the least concave majorant of the sum of any function h and a linear function l is equal to the sum of the linear function l and the least concave majorant of h.

By Skorokhod's representation theorem there exists a sequence of random variables $\widetilde{Z}_n$, with same distribution as $\sqrt{n}\,\{U_n - U\}$, which converges almost surely to $\widetilde{\mathbb{Z}}$ (or see [72] for strong approximation type of result). Furthermore taking the Least Concave Majorant is a continuous function (see [15]). Thus the expression in (2.4.41) has the same limiting distribution as:

$$\left[\widetilde{\mathbb{Z}} + \sqrt{n}\,\{U - l\}\right]^*(\underline{x}) - \widetilde{\mathbb{Z}}(\underline{x}).$$

We proceed by an analytic argument to show that the above tends to zero in probability. Fix a sample path k of $\widetilde{Z}$, which is a bounded and continuous function on $[0, M]$. Because $0 \le \sqrt{n}(U_n^*(\underline{x}) - U_n(\underline{x}))$, for $\varepsilon > 0$ we need to show $\forall\, n$ large enough:

$$\{k + \sqrt{n}[U - l]\}^*(\underline{x}) < k(\underline{x}) + \varepsilon \tag{2.4.42}$$

To do so, we construct a concave majorant of $k + \sqrt{n}[U - l]$ on $(\underline{x}, M]$ passing through the point $(\underline{x}, k(\underline{x}) + \varepsilon)$ and we show that for all n big enough it constitutes a concave majorant of $k + \sqrt{n}[U - l]$ also on $[0, \underline{x}]$, thus eventually on the whole $[0, M]$.

Let S_n denote the largest slope of lines connecting the point $(\underline{x}, k(\underline{x}) + \varepsilon)$ with $(t, k(t) + \sqrt{n}[U - l](t))$ for $t > \underline{x}$. Note that $S_n < \infty$ for all n. Since U is concave, increasing and linear on $[\underline{x}, \bar{x}]$, $U - l \le 0$ and thus $\sqrt{n}[U - l]$ becomes progressively smaller on $(\bar{x}, M]$, implying $S_1 \ge S_2 \ge \cdots$.

Let l_n be the line with slope S_n that passes through the point $(\underline{x}, k(\underline{x}) + \varepsilon)$. (2.4.42) holds true if and only if $k(t) + \sqrt{n}[U(t) - l(t)] < l_n(t)$ for all $t \in [0, \underline{x}]$. Since $S_1 \ge S_n$, it suffices to show that for all $t \in [0, \underline{x}]$ for all n large enough

$$k(t) + \sqrt{n}[U(t) - l(t)] < l_1(t), \tag{2.4.43}$$

as for $t > \underline{x}$, $k + \sqrt{n}[U - l] < l_1$ by construction and $S_1 \geq S_n$. These imply:

$$\{k + \sqrt{n}[U - l]\}^* \leq l_1.$$

We prove (2.4.43). Because $[U - l] \leq 0$, $\forall\, \delta > 0$, as $n \to \infty$

$$\sup_{0 \leq t < \underline{x} - \delta} (k + \sqrt{n}[U - l])(t) \to -\infty$$

this implies that for any $\delta > 0$, on $[0, \underline{x} - \delta)$ for sufficiently large n: $k + \sqrt{n}[U - l] < l_1$. For $\underline{x} - \delta \leq t < \underline{x}$ and $\delta < \underline{x} - \varepsilon/(2S_1)$,

$$l_1(t) = k(\underline{x}) + \varepsilon + (t - \underline{x})S_1 \geq k(\underline{x}) + \varepsilon/2.$$

For δ small enough on $[\underline{x} - \delta, \underline{x})$ and sufficiently large n, by the continuity of k:

$$\sup_{\underline{x} - \delta \leq t < \underline{x}} (k + \sqrt{n}[U - l])(t) \leq \sup_{\underline{x} - \delta \leq t < \underline{x}} k(t) \leq k(\underline{x}) + \varepsilon/2.$$

This implies that for δ small enough, on $[\underline{x} - \delta, \underline{x})$ for sufficiently large n: $k + \sqrt{n}[U - l] < l_1$.

The result for $\bar{x}$ follows analogously. □

***Proof of Theorem** 2.3.1.a.* First, we show:

$$\sqrt{n}\left(V_n^{\Pi}(x) - V_n^{(\underline{x},\bar{x})}(x)\right) = o_p(1), \tag{2.4.44}$$

Because V_n^{Π} is the left derivative of the least concave majorant of $\widetilde{U}_n$, the switch relation gives, for $a \in \mathbb{R}$ and sufficiently large n,

$$\begin{aligned}
&\sqrt{n}\left(V_n^{\Pi}(x) - V_n^{(\underline{x},\bar{x})}(x)\right) \leq a \\
\Leftrightarrow \quad &\arg\max_{s \geq 0}{}^{-}\{ \sqrt{n}(\widetilde{U}_n(s) - \widetilde{U}_n(\underline{x}) - U(s) + U(\underline{x})) + \\
&\qquad - \sqrt{n}(V_n^{(\underline{x},\bar{x})}(x) - V(x))(s - \underline{x}) + \\
&\qquad + \sqrt{n}(U(s) - U(\underline{x}) - V(x)(s - \underline{x})) - as\} \leq x,
\end{aligned}$$

where we used the usual properties of the $\arg\max^{-}$ as in the proof of Theorem 2.3.1.b.. By the definition of $\widetilde{U}_n$ in (2.3.16) and $V_n^{(\underline{x},\bar{x})}(x)$ in (2.3.14), the above can be rewritten as:

$$\arg\max_{s \geq 0}{}^{-}\left\{Z_n(s) - as - \sqrt{n}h_x(s)\right\} \leq x,$$

where:

$$Z_n(s) := \begin{cases} \frac{\sqrt{n}}{\bar{x} - \underline{x}}\Big((U_n(s) - U_n(\underline{x}))(\bar{x} - s) - (U_n(\bar{x}) - U_n(s))(s - \underline{x}) \\ \qquad -(U(s) - U(\underline{x}))(\bar{x} - s) - (U(\bar{x}) - U(s))(s - \underline{x})\Big), & s \notin [\underline{x}, \bar{x}], \\ 0, & s \in [\underline{x}, \bar{x}], \end{cases}$$

$$h_x(s) := U(\underline{x}) - U(s) - V(x)(\underline{x} - s).$$

From here on, we apply the argument of Theorem 2.3.1.b., using also for the current case the convergence in equation 2.4.36. Presently

$$\arg\max_{s\geq 0}{}^{-}\left\{Z_n(s) - as - \sqrt{n}h_x(s)\right\} \rightsquigarrow \arg\max_{s\in K}\{-as\},$$

as in this case the limiting process $\mathbb{Z}$ is constantly 0 on K. The argmax on the right is $\bar{x}$ if $a < 0$ and $\underline{x}$ if $a > 0$. Since $x \in (\underline{x}, \bar{x})$ it follows that

$$\begin{aligned}&\mathrm{P}\left(\arg\max_{s\geq 0}{}^{-}\left\{Z_n(s) - as - \sqrt{n}h_x(s)\right\} \leq x\right)\\ &\overset{n\to\infty}{\longrightarrow} \mathrm{P}\left(\arg\max_{s\in K}\{-as\} \leq x\right) = \begin{cases}1 & \text{if } a > 0,\\ 0 & \text{if } a < 0.\end{cases}\end{aligned}$$

This concludes the proof of (2.4.44).

Second we prove:

$$\sqrt{n}\left(\hat{V}_n^{\Pi}(x) - V_n^{(\underline{x},\bar{x})}(x)\right) = o_p(1). \tag{2.4.45}$$

By the definitions of $\hat{V}_n^{\Pi}$ and $V_n^{(\underline{x},\bar{x})}$ in (2.3.12) and (2.3.14):

$$\sqrt{n}\left(\hat{V}_n^{\Pi}(x) - V_n^{(\underline{x},\bar{x})}(x)\right) = \frac{1}{\bar{x}-\underline{x}}\left\{\sqrt{n}\left(U_n^*(\bar{x}) - U_n(\bar{x})\right) + \sqrt{n}\left(U_n^*(\underline{x}) - U_n(\underline{x})\right)\right\}.$$

Both terms on the right tend to zero by Lemma 2.4.9.

We conclude by giving the asymptotic distributions. By the Central Limit Theorem:

$$\sqrt{n}\left(V_n^{(\underline{x},\bar{x})}(x) - V(x)\right) \rightsquigarrow N(0, \sigma^2_{\underline{x},\bar{x}}). \tag{2.4.46}$$

The estimators $V_n^{\Pi}(x)$ and $\hat{V}_n^{\Pi}(x)$ have the same limiting behaviour by Slutsky's lemma and (2.4.44)-(2.4.45). □

The proof Theorem 2.3.1.c. is deferred to the next section because it relies on Lemma 2.4.10, Lemma 2.4.11 and Proposition 2.3.2, which can be understood only in the context of Section 2.3.3.

Proof of Proposition 2.3.2. Define:

$$\begin{aligned}J_{n,i} &:= \left(1 - \frac{1}{c_{t_n}}\right) + \frac{1}{g_V(Z_i)}\int_{\{s\in[Z_i,M],\,(1+t_nh_0(s))\leq 0\}} c_{t_n}^{-1}\left(1 + t_nh_0(s)\right)\frac{dV(s)}{\pi\sqrt{s-Z_i}},\\ \Delta_{n,i} &:= \frac{1}{c_{t_n}\sqrt{n}}\left(k^{(\underline{x},\bar{x})}(Z_i) - \mathrm{E}k^{(\underline{x},\bar{x})}(Z)\right).\end{aligned} \tag{2.4.47}$$

Using the definition of h_0:

$$\frac{t_n}{c_{t_n}}\int_{Z_i}^{M} -\frac{h_0(s)v(s)}{\pi\sqrt{s-Z_i}}\,ds = t\Delta_{n,i}\,g_V(Z_i). \tag{2.4.48}$$

Now note that for any i, using $1 - \frac{(1+x)_+}{c} = -\frac{x}{c} + 1 - \frac{1}{c} + \frac{(1+x)_-}{c}$ and (2.4.47)-(2.4.48):

$$\frac{g_{V_{t_n}}(Z_i)}{g_V(Z_i)} - 1 = \frac{1}{g_V(Z_i)}\left(\int_{Z_i}^{M}\left(1 - \frac{(1+t_nh_0(s))_+}{c_{t_n}}\right)\frac{dV(s)}{\pi\sqrt{s-Z_i}}\right) \tag{2.4.49}$$

$$= -\frac{t_n}{c_{t_n} g_V(Z_i)} \int_{Z_i}^{M} \frac{h_0(s)v(s)}{\pi\sqrt{s - Z_i}}\, ds + J_{n,i} = t\Delta_{n,i} + J_{n,i}.$$

Through the Taylor expansion $\log\{1 + x\} = x - \frac{x^2}{2} + x^2 R(2x)$ where $R(x) \to 0$ as $x \to 0$, we can express the log-likelihood ratio:

$$\sum_{i=1}^{n} \log \frac{g_{V_{t_n}}(Z_i)}{g_V(Z_i)} \tag{2.4.50}$$
$$= \sum_{i=1}^{n} \left\{ t\Delta_{n,i} + J_{n,i} - \frac{1}{2}(t\Delta_{n,i} + J_{n,i})^2 + (t\Delta_{n,i} + J_{n,i})^2 R(2(t\Delta_{n,i} + J_{n,i})) \right\}.$$

We start by showing that:

$$c_{t_n} = -\frac{2}{\pi} \int_0^M \sqrt{s}\,(1 + t_n h_0(s))_+\, dV(s) = 1 + o\left(\frac{1}{n}\right). \tag{2.4.51}$$

Let $S \sim \mu_V$ for μ_V defined in (2.4.59). Since $\int h_0\, d\mu_V = 0$:

$$c_{t_n} \geq -\frac{2}{\pi} \int_0^M \sqrt{s}\,(1 + t_n h_0(s))\; dV(s) = 1.$$

Now note:

$$c_{t_n} = -\frac{2}{\pi} \int_{\{s\in[0,M],\,(1+t_n h_0(s))>0\}} \sqrt{s}\,(1 + t_n h_0(s))\, dV(s) = 1 + \frac{2}{\pi} \int_{\{s\in[0,M],\,(1+t_n h_0(s))\leq 0\}} \sqrt{s}\,(1 + t_n h_0(s))\, dV(s)$$
$$= 1 + \mathrm{E}_{\mu_V}\left[|1 + t_n h_0(S)|\, \mathbf{1}_{\{t_n h_0(S)\leq -1\}}\right].$$

If $t_n h_0(s) \leq -1$ then: $|1 + t_n h_0(s)| \leq |t_n h_0(s)| \leq t_n^2 (h_0(s))^2$, so that:

$$\frac{1}{t_n^2} \mathbf{1}_{\{t_n h_0(s)\leq -1\}} |1 + t_n h_0(s)| \leq (h_0(s))^2 \mathbf{1}_{\{t_n h_0(s)\leq -1\}}. \tag{2.4.52}$$

Now $\mathrm{E}_{\mu_V}\left[h_0^2(S)\right] \leq \mathrm{E}_{\mu_V}\left[h^2(S)\right]$, which by (2.4.67) is bounded by (2.3.24), which is finite by assumption. Therefore:

$$\limsup_{n\to\infty} \frac{1}{t_n^2} \mathrm{E}_{\mu_V}\left[|1 + t_n h_0(S)|\, \mathbf{1}_{\{t_n h_0(S)\leq -1\}}\right] = 0, \tag{2.4.53}$$

by the Dominated Convergence theorem. This concludes the proof of (2.4.51) as $t_n^2 = t^2/n$.

By (2.4.51) and the Central Limit theorem: $\sum_{i=1}^n \Delta_{n,i} \rightsquigarrow N(0, \sigma^2_{\underline{x},\bar{x}})$. Now note that because dV is a nonpositive measure (since V is decreasing), $\forall\, i$:

$$\frac{1}{g_V(Z_i)} \int_{\{s\in[Z_i,M],\,(1+t_n h_0(s))\leq 0\}} c_{t_n}^{-1}\,(1 + t_n h_0(s))\, \frac{dV(s)}{\pi\sqrt{s - Z_i}} \geq 0, \qquad \text{a.s.} \tag{2.4.54}$$

Again by (2.4.51), by Markov's inequality and Fubini's theorem, $\sum_{i=1}^n J_{n,i} = o_p(1)$ because $\forall\, \varepsilon > 0$:

$$\mathrm{P}\left(\sum_{i=1}^{n} \frac{1}{g_V(Z_i)} \int_{\{s\in[Z_i,M],\,(1+t_n h_0(s))\leq 0\}} c_{t_n}^{-1}\,(1 + t_n h_0(s))\, \frac{dV(s)}{\pi\sqrt{s - Z_i}} > \varepsilon \right)$$

$$
\begin{aligned}
&\leq \frac{n}{\varepsilon}\int_0^M\int_z^M \mathbf{1}_{\{(1+t_nh_0(s))\leq 0\}}\, c_{t_n}^{-1}\,(1+t_nh_0(s))\,\frac{v(s)}{\pi\sqrt{s-z}}\,ds\,dz\\
&= \frac{n}{\varepsilon\pi}\int \mathbf{1}_{\{(1+t_nh_0(s))\leq 0\}}\, c_{t_n}^{-1}\,(1+t_nh_0(s))\,v(s)\int_0^s \frac{1}{\sqrt{s-z}}\,dz\,ds\\
&= \frac{n}{\varepsilon c_{t_n}}\mathrm{E}_{\mu_v}\big[|1+t_nh_0(S)|\,\mathbf{1}_{\{t_nh_0(S)\leq -1\}}\big] \overset{(2.4.53)}{=} o(1). \qquad (2.4.55)
\end{aligned}
$$

All together this proves, for Δ_n as in (2.3.28): $\sum_{i=1}^n(t\Delta_{n,i}+J_{n,i}) = t\Delta_n + o_p(1)$, proving the behaviour of the linear term.

For the quadratic term, if $\max_{1\leq i\leq n}|t\Delta_{n,i}+J_{n,i}| \xrightarrow{\mathrm{P}} 0$, by property of the function R, the sequence $\max_{1\leq i\leq n}|R(t\Delta_{n,i}+J_{n,i})| \xrightarrow{\mathrm{P}} 0$ as well. By the triangle inequality:

$$
\max_{1\leq i\leq n}|t\Delta_{n,i}+J_{n,i}| \leq |t|\max_{1\leq i\leq n}|\Delta_{n,i}| + \sum_{i=1}^n |J_{n,i}|.
$$

The first term tends to zero in probability because for $\{X_i,\ i\geq 1\}$ i.i.d. random variables with finite variance: $\max_{1\leq i\leq n}\left\{\frac{|X_i|}{\sqrt{n}}\right\} \xrightarrow{\mathrm{P}} 0$. The second term tends to 0 by (2.4.51) and (2.4.55).

By the Law of Large numbers, $\sum_{i=1}^n \Delta_{n,i}^2 \to \sigma^2_{\underline{x},\bar{x}}$ in probability. Therefore we are left with showing:

$$
\sum_{i=1}^n \Delta_{n,i}J_{n,i} \xrightarrow{\mathrm{P}} 0 \qquad \text{and} \qquad \sum_{i=1}^n (J_{n,i})^2 \xrightarrow{\mathrm{P}} 0. \qquad (2.4.56)
$$

The second follows because $\sum_{i=1}^n (J_{n,i})^2 \leq \max_{1\leq i\leq n}|J_{n,i}|\sum_{i=1}^n J_{n,i}$ and the first next follows with the help of the Cauchy-Schwarz inequality. □

For the proof of the LAM (locally asymptotically minimax) theorem, the first statement of Theorem 2.3.2, and for the proof of Theorem 2.3.1.c., we need the derivative of the map we are interested in estimating, in our case $V(x)$ (cf. definition 1.10 in [7]). This is given in the following lemma.

Lemma 2.4.10. *Along the path given in* (2.3.27)*, as* $n\to\infty$*,*

$$
\sqrt{n}\,(V_{t_n}(x) - V(x)) \to t\,\sigma^2_{\underline{x},\bar{x}},
$$

where $\sigma^2_{\underline{x},\bar{x}}$ *is as in* (2.3.20).

Proof. Because $\int_x^s \frac{\sqrt{w-x}}{\sqrt{s-w}}\,dw = \frac{\pi}{2}(s-x)$ for every x, we have, $\forall\, x\in[\underline{x},\bar{x}]$ by Fubini's theorem:

$$
\begin{aligned}
&\int_0^M 2\frac{\sqrt{(w-\underline{x})_+} - \sqrt{(w-\bar{x})_+}}{\bar{x}-\underline{x}}\,g_v(w)\,dw\\
&= -\frac{2}{\pi}\frac{1}{\bar{x}-\underline{x}}\left(\int_{\underline{x}}^M\int_{\underline{x}}^s \frac{\sqrt{w-\underline{x}}}{\sqrt{s-w}}\,dw\,dV(s) - \int_{\underline{x}}^M\int_{\bar{x}}^s \frac{\sqrt{w-\bar{x}}}{\sqrt{s-w}}\,dw\,dV(s)\right)\\
&= -\frac{2}{\pi}\frac{1}{\bar{x}-\underline{x}}\int_{\underline{x}}^M \frac{\pi}{2}(\bar{x}-\underline{x})\,dV(s) = V(\underline{x}) = V(x). \qquad (2.4.57)
\end{aligned}
$$

Using the same computations as done for (2.4.49) we have:

$$(g_{V_t}(z) - g_V(z)) = -\frac{t_n}{c_{t_n}} \int_z^M \frac{h_0(s)v(s)}{\pi\sqrt{s-z}}\, ds + \left(1 - \frac{1}{c_{t_n}}\right) \int_z^M \frac{dV(s)}{\pi\sqrt{s-z}} + \int_{\{s\in[z,M],\,(1+t_n h_0(s))\le 0\}} c_{t_n}^{-1}\,(1 + t_n h_0(s))\, \frac{dV(s)}{\pi\sqrt{s-z}}.$$

Because $(\mathrm{E}k^{(\underline{x},\bar{x})} \int_0^M (g_{V_t}(w) - g_V(w))\, dw = 0$, using (2.4.57), we obtain:

$$\begin{aligned}
\sqrt{n}\,(V_{t_n}(\underline{x}) - V(\underline{x})) &= \sqrt{n} \int_0^M \left(k^{(\underline{x},\bar{x})}(z) - \mathrm{E}k^{(\underline{x},\bar{x})}\right)\left(g_{V_{t_n}}(z) - g_V(z)\right) dz \\
&= \frac{t}{c_{t_n}} \int_0^M \left(k^{(\underline{x},\bar{x})}(z) - \mathrm{E}k^{(\underline{x},\bar{x})}\right)\left(-\int_z^M \frac{h_0(s)v(s)}{\pi\sqrt{s-z}}\, ds\right) dz \\
&\quad + \sqrt{n}\left(\frac{1}{c_{t_n}} - 1\right) \underbrace{\int_0^M \left(k^{(\underline{x},\bar{x})}(z) - \mathrm{E}k^{(\underline{x},\bar{x})}\right) g_V(z)\, dz}_{=0} \\
&\quad + \sqrt{n} \int_0^M \left(k^{(\underline{x},\bar{x})}(z) - \mathrm{E}k^{(\underline{x},\bar{x})}\right) \int_{\{s\in[z,M],\,(1+t_n h_0(s))\le 0\}} c_{t_n}^{-1}\,(1 + t_n h_0(s))\, \frac{dV(s)}{\pi\sqrt{s-z}}\, dz \\
&\xrightarrow{n\to\infty} t \int_0^M \left(k^{(\underline{x},\bar{x})}(w) - \mathrm{E}k^{(\underline{x},\bar{x})}\right)^2 g_V(w)\, dw = t\,\sigma^2_{\underline{x},\bar{x}}.
\end{aligned}$$

by (2.4.48) and (2.4.53), where we use that $k^{(\underline{x},\bar{x})}(z) - \mathrm{E}k^{(\underline{x},\bar{x})}$ is bounded on $[0, M]$. □

The proof of Theorem 2.3.1.c., is based on the fact that the isotonic estimator is regular on the submodel $t \mapsto G_{V_t}$, i.e. the limit distribution of the sequence $\sqrt{n}(\hat{V}_n(x) - V_{G_{V_{t_n}}}(x))$ under $G_{V_{t_n}}$ is the same, for all $t \in \mathbb{R}$.

Lemma 2.4.11. *The IIE is regular in the submodel defined in* (2.3.27).

Proof. For clarity of notation, in this proof we denote $U(x)$ as: $U_V(x) = \int_0^x V(y)\, dy$ and thus $U_{V_{t_n}}(x) = \int_0^x V_{t_n}(y)\, dy$. We show, under $G_{V_{t_n}}$: $\sqrt{n}(\hat{V}_n(x) - V_{t_n}(x)) \rightsquigarrow L_x$, which is independent of t. We use the same steps as in the proof of Theorem 2.2.4. Using the switch relation, we study:

$$\begin{aligned}
\inf\{s \ge 0 : (U_n(s) - U_n(x) - U_{V_{t_n}}(s) + U_{V_{t_n}}(x)) + \\
+ (U_{V_{t_n}}(s) - U_{V_{t_n}}(x) - V_{t_n}(x)(s-x)) - (as)/\sqrt{n} \text{ is maximal}\} \le x.
\end{aligned}$$

Using the same proof of Lemma 2.2.1 it follows that, under $G_{V_{t_n}}$:

$$\sqrt{n}(U_n(s) - U_n(x) - U_{V_{t_n}}(s) + U_{V_{t_n}}(x) \;:\; s \ge 0) \rightsquigarrow \mathbb{Z}_x \quad \text{in } \ell^\infty[0,\infty).$$

Furthermore:

$$\begin{aligned}
U_{V_{t_n}}(s) - U_{V_{t_n}}(x) - V_{t_n}(x)(s-x) &= \int_x^s (V_{t_n}(y) - V_{t_n}(x))\, dy \\
&= \begin{cases} 0, & s, x \in [\underline{x}, \bar{x}], \\ < 0, & x \leqslant \bar{x} < s, \quad s < \underline{x} \leqslant x. \end{cases}
\end{aligned}$$

Now for $s = \bar{x} + \varepsilon_n$ and $\varepsilon_n \downarrow 0$ as in the proof of Theorem 2.2.4 and using the same reasoning as in the proof of (2.4.51), we obtain:

$$\begin{aligned}
&\sqrt{n}\int_{\bar{x}}^{\bar{x}+\varepsilon_n}(V_{t_n}(y) - V_{t_n}(\bar{x}))\,dy \\
&= -\sqrt{n}\int_{\bar{x}}^{\bar{x}+\varepsilon_n}\int_{\bar{x}}^{y} c_{t_n}^{-1}\left(1 + \frac{t}{\sqrt{n}}h_0(s)\right)_+ dV(s)dy \\
&= -\sqrt{n}\int_{\bar{x}}^{\bar{x}+\varepsilon_n}\int_{\bar{x}}^{y} c_{t_n}^{-1}\left(1 + \frac{t}{\sqrt{n}}h_0(s)\right)dV(s)dy + o\left(\frac{1}{\sqrt{n}}\right) \to -\infty,
\end{aligned}$$

since by the proof of Theorem 2.2.4 we have: $\sqrt{n}\int_{\bar{x}}^{\bar{x}+\varepsilon_n} V(y) - V(\bar{x})\,dy \to -\infty$. The same reasoning holds also for: $\sqrt{n}\int_{\underline{x}-\varepsilon_n}^{\underline{x}} (V_{t_n}(y) - V_{t_n}(\underline{x}))\,dy \to -\infty$. Using further the same computations as in the proof of Theorem 4 in [42] we conclude:

$$\mathrm{P}\Big(\sqrt{n}(\hat{V}_n(x) - V_{t_n}(x)) \le a\Big) \overset{n\to\infty}{\longrightarrow} \mathrm{P}\Big(\underset{s\in[\underline{x},\bar{x}]}{\arg\max}\{\mathbb{Z}_x(s) - as\} \le x\Big).$$

□

Proof of Theorem 2.3.1.c. (c.f. Theorem 1 in Section 2.3 of [2]). The efficient estimator satisfies:

$$\sqrt{n}(V_n^{(\underline{x},\bar{x})}(x) - V_0(x)) = \Delta_n + o_p(1)$$

where $V_0 \equiv V$ but this notation clarifies that it coincides with the perturbed V_t in (2.3.27) evaluated at $t = 0$. $\Delta_n \rightsquigarrow N(0, \sigma^2_{\underline{x},\bar{x}})$ is the linear term in (2.3.28) in the expansion of $\Lambda_n(t) := \sum_{i=1}^n \log\frac{g_{V_{t_n}}(Z_i)}{g_V(Z_i)}$. The joint sequence

$$(\sqrt{n}(\hat{V}_n(x) - V_n^{(x,x)}(x)),\ \sqrt{n}(V_n^{(x,x)}(x) - V_0(x)))$$

is uniformly tight (because marginally so). It suffices to show that every joint limit point under V_0 has independent coordinates. A joint limit point takes the form $(T - \Delta, \Delta)$ for (T, Δ) a joint limit of $\Big(\sqrt{n}\Big(\hat{V}_n(x) - V_0(x)\Big), \Delta_n\Big)$ under V_0. We have:

$$\Big(\sqrt{n}\Big(\hat{V}_n(x) - V_0(x)\Big), \Lambda_n(t)\Big) \overset{V_0}{\rightsquigarrow} \Big(T, t\Delta - \frac{1}{2}t^2\sigma^2_{\underline{x},\bar{x}},\Big).$$

Therefore by Le Cam's 3rd lemma (e.g. Theorem 6.6 in [102]):

$$\sqrt{n}\Big(\hat{V}_n(x) - V_0(x)\Big) \overset{V_{t/\sqrt{n}}}{\rightsquigarrow} L_t, \qquad L_t(B) = \mathrm{E}\mathbf{1}_B(T)e^{t\Delta - \frac{1}{2}t^2\sigma^2_{\underline{x},\bar{x}}}$$

because $\sqrt{n}\Big(V_{t/\sqrt{n}}(x) - V_0(x)\Big) \overset{n\to\infty}{\longrightarrow} t\sigma^2_{\underline{x},\bar{x}}$ (see Lemma 2.4.10) we conclude:

$$\sqrt{n}\Big(\hat{V}_n(x) - V_{t/\sqrt{n}}(x)\Big) \overset{V_{t/\sqrt{n}}}{\rightsquigarrow} \bar{L}_t, \qquad \bar{L}_t(B) = L_t\Big(B - t\sigma^2_{\underline{x},\bar{x}}\Big).$$

By regularity of the IIE (see Lemma 2.4.11): $\bar{L}_t = \bar{L}_0$, independent of t, so $\forall\, t, u$

$$\int e^{iuy}d\bar{L}_0(y) = \mathrm{E}e^{iu\left(T - t\sigma^2_{\underline{x},\bar{x}}\right)}e^{t\Delta - \frac{1}{2}t^2\sigma^2_{\underline{x},\bar{x}}}.$$

Choose $t = -iu$ to find: $\int e^{iuy} d\bar{L}_0(y) = \mathrm{E}e^{iu(T-\Delta)} e^{-\frac{1}{2}u^2\sigma^2_{\underline{x},\bar{x}}}$. Choose $t = -iu + iv$ to find: $\int e^{iuy} d\bar{L}_0(y) = \mathrm{E}e^{iu(T-\Delta)+iv\Delta} e^{\frac{1}{2}(v^2-u^2)\sigma^2_{\underline{x},\bar{x}}}$. Then:

$$\mathrm{E}e^{iu(T-\Delta)+iv\Delta} = \mathrm{E}e^{iu(T-\Delta)} e^{-\frac{1}{2}v^2\sigma^2_{\underline{x},\bar{x}}} = \mathrm{E}e^{iu(T-\Delta)} \mathrm{E}e^{iv\Delta}.$$

□

Proof of Theorem 2.3.2. The proof is a direct application of Theorems 25.20 and 25.21 from [102], using Proposition 2.3.2 and Lemma 2.4.10. The variance of the limiting random variable in (2.3.29) is given by $\dot{\nu}(\sigma^2_{\underline{x},\bar{x}})^{-1}\dot{\nu} = \sigma^2_{\underline{x},\bar{x}}$, where $\dot{\nu} = \sigma^2_{\underline{x},\bar{x}}$ is the derivative of the functional of interest and $(\sigma^2_{\underline{x},\bar{x}})^{-1}$ is the Fisher information for t, given in Proposition 2.3.2. □

2.4.4 Efficiency Theory Constrained Space

In this subsection we derive the least favourable submodel (2.3.27) from the semiparametric score calculus, as in [66] (or Chapter 25 of [102]).

The parameter V belongs to the class (c.f. [52]), for $\underline{x} < \bar{x}$:

$$\mathscr{V}_{\underline{x},\bar{x}} = \left\{ V \in \mathcal{V}_{\underline{x},\bar{x}}, : \int_0^\infty \sqrt{s}\, dV(s) = -\frac{\pi}{2}, \text{ zero on } [M, \infty) \right\}. \tag{2.4.58}$$

Because dV is not a probability measure, it is convenient to work with its associated probability measure μ_V defined for every measurable $A \subseteq [0, \infty)$ by:

$$\mu_V(A) := \int_A -\frac{2}{\pi} \sqrt{s}\, dV(s). \tag{2.4.59}$$

We consider the associated space of "underlying" probability measures $\mathscr{M}_{\underline{x},\bar{x}} := \{\mu_V : V \in \mathscr{V}_{\underline{x},\bar{x}}\}$ and the space of the "observed" probability measures $\mathscr{G}_{\underline{x},\bar{x}} := \{g_V : \mu_V \in \mathscr{M}_{\underline{x},\bar{x}}\}$. The tangent set $\dot{\mathscr{M}}_{\underline{x},\bar{x}}$ consists of all functions $h \in \mathbb{L}_2(\mu_V)$ such that $\int h\, d\mu_V = 0$. The derivative of the map $\mu_V \mapsto g_V$ (the score operator) maps score functions in the tangent set $\dot{\mathscr{M}}_{\underline{x},\bar{x}}$ into scores in the tangent set $\dot{\mathscr{G}}_{\underline{x},\bar{x}}$. For any $h \in \mathbb{L}_1(\mu_V)$ such that $-\int_z^M \frac{|h(s)|}{\sqrt{s-z}}\, dV(s) < \infty$, define:

$$(Th)(z) := \int_z^M \frac{h(s)}{\sqrt{s-z}}\, dV(s). \tag{2.4.60}$$

Then the score operator $A_V : \dot{\mathscr{M}}_{\underline{x},\bar{x}} \to \dot{\mathscr{G}}_{\underline{x},\bar{x}}$, for any $h \in \dot{\mathscr{M}}_{\underline{x},\bar{x}}$, is given by:

$$A_V h(z) = \frac{\int_z^M \frac{h(s)}{\sqrt{s-z}}\, dV(s)}{\int_z^M \frac{dV(s)}{\sqrt{s-z}}} = \frac{(Th)(z)}{(T1)(z)}. \tag{2.4.61}$$

For instance, for bounded $h \in \dot{\mathscr{M}}_{\underline{x},\bar{x}}$ define the following perturbation:

$$dV_t(s) = (1 + th(s))\, dV(s). \tag{2.4.62}$$

Then:

$$\frac{d}{dt} \log \left\{ -\int_z^M \frac{(1 + th(s))}{\pi\sqrt{s-z}} dV(s) \right\}\Big|_{t=0} = \frac{\int_z^M \frac{h(s)}{\sqrt{s-z}} dV(s)}{\int_z^M \frac{dV(s)}{\sqrt{s-z}}} =: A_V h(z).$$

We check that A_v is a linear operator between the $\mathbb{L}_2$ spaces claimed above. Linearity is clear. We can check by using the Cauchy-Schwarz inequality and that $h \in \mathbb{L}^2(\mu_v)$:

$$\int (A_v h)^2 dG_v = \int_0^\infty \left(\frac{\int_z^\infty \frac{h(s)dV(s)}{\sqrt{s-z}}}{\int_z^\infty \frac{dV(s)}{\sqrt{s-z}}} \right)^2 \left(-\frac{1}{\pi} \int_z^\infty \frac{dV(s)}{\sqrt{s-z}} \right) dz \tag{2.4.63}$$

$$\leqslant \int_0^\infty \left(\frac{\sqrt{\int_z^\infty -\frac{h^2(s)}{\sqrt{s-z}} dV(s)} \sqrt{\int_z^\infty -\frac{1}{\sqrt{s-z}} dV(s)}}{-\int_z^\infty \frac{dV(s)}{\sqrt{s-z}}} \right)^2 \left(-\frac{1}{\pi} \int_z^\infty \frac{dV(s)}{\sqrt{s-z}} \right) dz$$

$$= -\frac{1}{\pi} \int_0^\infty \int_z^\infty \frac{h^2(s)}{\sqrt{s-z}} dV(s) dz = -\frac{2}{\pi} \int_0^\infty h^2(s) \sqrt{s} dV(s) = \int h^2 \, d\mu_v < \infty.$$

The function $k^{(\underline{x},\bar{x})}$ is the influence function of the (regular) estimator (2.3.14). To show that this estimator is asymptotically efficient, it suffices to show that $k^{(\underline{x},\bar{x})}$ is contained in the closed linear span of the set of all score functions $A_v h$. We shall show that in fact $k^{(\underline{x},\bar{x})} - \mathrm{E}k^{(\underline{x},\bar{x})} = A_v h_0$ for h_0 given in (2.3.26). Equivalently, we show that h_0 solves

$$\begin{cases} (Th_0) = \left(k^{(\underline{x},\bar{x})} - \mathrm{E}k^{(\underline{x},\bar{x})} \right) \cdot (T1) \\ \int_0^M -|h_0(s)| \, dV(s) < \infty, \qquad \int_0^M \sqrt{s} h_0(s) \, dV(s) = 0. \end{cases} \tag{2.4.64}$$

Proposition 2.4.1. *Under Assumption 2.3.3, the solution to* (2.4.64) *is given by* (2.3.26)*.*

***Proof of Proposition** 2.4.1.* We show that the function h in (2.3.25) solves: $Th = q$, for $q = k^{(\underline{x},\bar{x})} \cdot T1$. Then the solution h_0 is found by centering h, i.e. $h_0(z) = h(z) - m$ (since $\int_0^M \sqrt{s} \, dV(s) = -\frac{\pi}{2}$) where m is such that:

$$\int_0^M \sqrt{s} \, (h(s) - m) \, dV(s) = 0 \quad \Longrightarrow \quad m = -\frac{2}{\pi} \int_0^M \sqrt{s} h(s) \, dV(s)$$

The equation $Th = q$, for $q = k^{(\underline{x},\bar{x})} \cdot T1$ is the following Abel's integral equation:

$$\int_z^M \frac{h(s)v(s)}{\sqrt{s-z}} \, ds = 2 \frac{\sqrt{(z-\underline{x})_+} - \sqrt{(z-\bar{x})_+}}{\bar{x} - \underline{x}} \int_z^M \frac{v(s)}{\sqrt{s-z}} \, ds$$

By Theorem 2.1 in [85] this equation is solvable for $hv \in \mathbb{L}_1[0, M]$ iff:

$$z \mapsto \int_z^M \frac{q(s)}{\sqrt{s-z}} \, ds$$

is absolutely continuous. Furthermore the unique in $\mathbb{L}_1$ solution is given by $h(z)v(z) = -\frac{d}{dz} \int_z^M \frac{q(s)}{\sqrt{s-z}} \, ds$, where the derivative is understood in the sense of absolute continuity (AC). Moreover, if $q \in \mathrm{AC}[0, M]$, then this condition is satisfied and for $z \notin [\underline{x}, \bar{x}]$:

$$h(z)v(z) = \frac{1}{\sqrt{\pi}} \left(\frac{q(M)}{\sqrt{M-z}} - \int_z^M \frac{q'(s)}{\sqrt{s-z}} \, ds \right). \tag{2.4.65}$$

Now we prove that under the given conditions $k^{(\underline{x},\bar{x})} \cdot T1 = q \in \mathrm{AC}[0, M]$. Because $\sqrt{(z-x)_+} = \int_0^z \frac{1}{2\sqrt{s-x}} \mathbf{1}_{s>x} \, ds$ we see that $k^{(\underline{x},\bar{x})} \in \mathrm{AC}[0, M]$. $T1$ is the $1/2$-fractional integral of v. Because

$v \in \mathrm{AC}[\underline{x}, \bar{x}]^c$ and $v = 0$ on $[\underline{x}, \bar{x}]$, we know that $T1 \in \mathrm{AC}[0, M]$ (c.f. property 3.2.(6).(b) pp. 209 in [43]). The product of absolutely continuous functions is absolutely continuous.

Finally we show:

$$\mathrm{E}k^{(\underline{x},\bar{x})} = -\frac{2}{\pi}\int_0^M \sqrt{s}h(s)v(s)\,ds.$$

For the given h in (2.3.25) we just proved:

$$\int_z^M -\frac{h(s)v(s)}{\pi\sqrt{s-z}}\,ds = 2\frac{\sqrt{(z-\underline{x})_+} - \sqrt{(z-\bar{x})_+}}{\bar{x}-\underline{x}}g_v(z). \tag{2.4.66}$$

By integrating both sides between 0 and M we get that the r.h.s. is $\mathrm{E}k^{(\underline{x},\bar{x})}$, whereas the l.h.s.

$$\int_0^M\int_z^M -\frac{h(s)v(s)}{\pi\sqrt{s-z}}\,ds\,dz = \int_0^M\int_0^s -\frac{h(s)v(s)}{\pi\sqrt{s-z}}\,dz\,ds = -\frac{2}{\pi}\int_0^M \sqrt{s}h(s)v(s)\,ds.$$

All together this proves that h_0 solves (2.4.64). □

Lemma 2.4.12. *For a fixed $\eta > 0$, there exist constants $c_1, c_2, c_3 > 0$ such that, for $z \in [\underline{x}, \bar{x}]^c$:*

$$|h(z)v(z)| \le c_1 + c_2\log\left(\frac{1}{\underline{x}-z}\right)\mathbf{1}_{\{0<\underline{x}-z\le\eta\}} + c_3\log\left(\frac{1}{z-\bar{x}}\right)\mathbf{1}_{\{0<z-\bar{x}\le\eta\}}. \tag{2.4.67}$$

Proof. Because $\lim_{z\to M}(T1)(z) = 0$, the function q in (2.4.65) satisfies $q(M) = 0$. Hence, for $z \in [\underline{x}, \bar{x}]^c$:

$$h(z) = -\frac{1}{\sqrt{\pi}v(z)}\Bigg(\underbrace{\int_z^M \frac{(k^{(\underline{x},\bar{x})})'(T1)(s)}{\sqrt{s-z}}\,ds}_{(1)} + \underbrace{\int_z^M \frac{(k^{(\underline{x},\bar{x})})(T1)'(s)}{\sqrt{s-z}}\,ds}_{(2)}\Bigg)$$

where:

$$(k^{(\underline{x},\bar{x})})'(s) = \frac{1}{(\bar{x}-\underline{x})}\left(\frac{1}{\sqrt{(s-\underline{x})_+}} - \frac{1}{\sqrt{(s-\bar{x})_+}}\right).$$

We prove (2) is bounded. By the same proof of Corollary 2.1 in [85] (pp. 32), since $(T1) \in \mathrm{AC}[0, M]$ (see proof of Proposition 2.4.1), $\lim_{z\to M} v(z) = 0$, and denoting by $\mathrm{I}^{\frac{1}{2}}(f)$ the half–integral of f (see [85] for an introduction to fractional calculus):

$$\frac{d}{dz}\int_z^M \frac{v(s)\,ds}{\sqrt{s-z}} = \int_z^M \frac{v'(s)\,ds}{\sqrt{s-z}} - \lim_{s\to M}\frac{v(s)}{\sqrt{s-z}} = \mathrm{I}^{\frac{1}{2}}(v'),$$

where the derivative is understood in the sense of absolute continuity (AC). Thus (2) coincides with $\mathrm{I}^{\frac{1}{2}}(k^{(\underline{x},\bar{x})}\,\mathrm{I}^{\frac{1}{2}}(v'))$. By Theorem 3.6 in [85], $\mathrm{I}^{\frac{1}{2}} : \mathbb{L}_p \mapsto H^{1/2-1/p}$, where H^α denotes the space of Hölder continuous functions of degree α. Thus if $k^{(\underline{x},\bar{x})}\,\mathrm{I}^{\frac{1}{2}}(v') \in \mathbb{L}_p$ for $p > 2$ we obtain the claim. Because $k^{(\underline{x},\bar{x})}$ is bounded we show $\mathrm{I}^{\frac{1}{2}}(v') \in \mathbb{L}_p$ for $p > 2$. By theorem 3.5 in [85], $\mathrm{I}^{\frac{1}{2}}(v') \in \mathbb{L}_{q/(1-q/2)}$ if $v' \in \mathbb{L}_q$. Because v is Lipschitz on $[\underline{x}, \bar{x}]^c$ and 0 on $[\underline{x}, \bar{x}]$, $\exists\ q > 1$ such that $v' \in \mathbb{L}_q$. As $q > 1 \iff q/(1-q/2) > 2$, $\mathrm{I}^{\frac{1}{2}}(v') \in \mathbb{L}_p$ for some $p > 2$ and we obtain the claim.

Regarding the absolute value of (1), using the fact that $(T1)(s) = -\frac{\pi}{2} g_v(s)$, which is bounded as well as v:

$$\Big| -\frac{1}{\sqrt{\pi}} \int_{z \vee \underline{x}}^{M} \frac{(T1)(s)}{\sqrt{(s-\underline{x})(s-z)}}\, ds \Big| \lesssim \Big| \int_{z \vee \underline{x}}^{M} \frac{ds}{\sqrt{(s-\underline{x})(s-z)}} \Big|.$$

For $z > \bar{x}$ the previous display is bounded. For $z < \underline{x}$ it is upper bounded (up to constants) by:

$$\left| \sin^{-1}\left(\sqrt{\frac{M-z}{\underline{x}-z}} \right) \right| = 2 \log \sqrt{\frac{M-z}{\underline{x}-z}} + O(1) + O\left(\frac{\underline{x}-z}{M-z} \right) \quad \text{as} \quad z \uparrow \underline{x}.$$

The integral $\int_{z \vee \bar{x}}^{M} \frac{(T1)(s)}{\sqrt{(s-\bar{x})(s-z)}}\, ds$ can be handled analogously.

□

2.5 Bibliographic remarks

There is a vast literature on Wicksell's problem, see for instance [19, 27, 48, 49, 52, 53, 56, 62, 92, 99, 114]. For a detailed introduction to Wicksell's problem we refer the reader to [99]. The setting of Wicksell's problem has several modern applications, as mentioned above. In these applications, it is common to see the use of the so-called Saltykov method or methods based on numerical discretization. These methods are far from being efficient and in some cases not even consistent, making our work of practical relevance. The estimation of F given data from g is of interest not only for stereological procedures, but also from the mathematical point of view. Consider the class of inverse problems that entails the estimation of a functional: $\Psi_\alpha(x; g) = \int_x^\infty (z-x)^{-\alpha} g(z)\, dz$, $0 < \alpha < 1$ based on samples from g. This type of functional estimation emerges in a vast variety of applications. For $\alpha < \frac{1}{2}$, the above gives a regular statistical functional (cf. Theorem 4 in [69]), whereas for $\alpha > \frac{1}{2}$ gives instances of irregular functionals. Wicksell's problem entails the boundary case for $\alpha = 1/2$, which makes the problem mathematically intriguing, both because of the unusual rate of convergence and because of the non-standard efficiency theory that in this section is developed.

The first time the naive plug-in estimator was studied is in [114]. Later in [52] the proof of the asymptotic normality of this estimator was simplified considerably. However, viewed as a function, the naive plug-in estimator is severely ill-behaved as we already pointed out, and this led to the definition of the IIE. Naturally, one may wonder why not use the Nonparametric Maximum Likelihood Estimator. For Wicksell's problem, the usual definition of the NPMLE is also ill-posed, as the log-likelihood can attain value infinity. However, work has been done, for instance in [62] and [19], to make the NPMLE an employable estimator. For results about confidence intervals in Wicksell's problem, we mention: [27, 92]. On the other hand for KDE-based estimators, we refer to [49, 56]. Finally, regarding efficient estimation, it is worth mentioning that one of the authors of [49] did some further work on Wicksell's problem in [48] where it was shown how to attain efficient smoothing for the plug-in estimator under square losses.

Chapter 3

Wicksell's problem: the Bayesian nonparametric approach

3.1 Semiparametric Bernstein-von Mises phenomenon via Isotonized Posterior

3.1.1 Introduction

Before diving into the results obtained using Bayesian nonparametric modelling approaches, we recall to the reader the main formulas that are useful to understand fully the following results. We start by formally recalling the main mathematical objects of interest in the material science setting and in the astronomical setting, respectively.

The setting of the opaque material having globular microstructure is formalized by assuming that the embedded particle squared radii are i.i.d. realizations from a random variable $X^* \sim F_0^*$ and our observations consist of cross sections of the medium (i.e. "planes cutting the medium"). Note that the spheres with bigger radii are more likely to be sampled. Thus the distribution of the squared radii of the spheres actually cut is not F_0^* but a biased version of it; we write for the squared radius of the sphere actually cut: $X \sim F_0$. Since we assume that the samples are gathered uniformly, by the Pythagorean Theorem, the observable squared circle radius Z is related to X via: $Z = (1 - U^2)X = YX$, where $U \sim U[0, 1]$ so $Y \sim \mathrm{Be}(1, 1/2)$, with density $y \mapsto \frac{1}{2}(1 - y)^{-1/2}\mathbf{1}_{(0,1)}(y)$. So, the observed squared radii Z are a random fraction of the true squared radii depending on where the plane cuts through the sphere. Consequently, since both X and Y are nonnegative, a version of the density g_0 of Z can, for $z \geq 0$, be written as:

$$g_0(z) = \frac{d}{dz}\int_{(0,\infty)} \mathrm{P}(Y \leq z/x)\, dF_0(x) = \int_z^\infty \frac{dF_0(x)}{2\sqrt{x^2 - xz}} \tag{3.1.1}$$

The problem in the astronomical setting can be modelled as follows: let $\mathbf{Z} = (Z_{p_1}, Z_{p_2}, Z_{p_3})$ be a spherically symmetric random vector (the three-dimensional position of a star in a galaxy) of which only (Z_{p_1}, Z_{p_2}) can be observed. Under the assumption that the joint of $\mathbf{Z}$ is a function of the distance to the center of the galaxy, astronomers are interested in estimating the distribution of the squared distance of a star to the center of the globular cluster or galaxy $X = Z_{p_1}^2 + Z_{p_2}^2 + Z_{p_3}^2 \sim F_0$ using $Z = Z_{p_1}^2 + Z_{p_2}^2$. From the assumed spherical symmetry, $\mathbf{Z} \sim O\mathbf{Z}$ for all orthogonal matrices $O \in \mathrm{O}(3)$. Therefore $\mathbf{U} := \mathbf{Z}/\|\mathbf{Z}\|$ (note $X = \|\mathbf{Z}\|^2$) is uniform on

the unit sphere $\mathbb{S}^2$. It follows that the random variable Z_{p_3} (the third coordinate) is distributed as: $\sqrt{X} \cdot U_3$ where $U_3 \sim U[-1, 1]$ independent of X. Note also that $Y \equiv 1 - U_3^2 \sim \text{Be}(1, 1/2)$ as above. Because:

$$\text{P}(Z \leq z) = \text{P}(Z_{p_1}^2 + Z_{p_2}^2 + Z_{p_3}^2 \leq z + Z_{p_3}^2) = \text{P}(X \leq z + X \cdot U_3^2) = \text{P}(YX \leq z),$$

it follows that both for the application in materials science and the one in astronomy, the density of the observations $Z \sim G_0$ is the one given on the right-hand side of (3.1.1).

By using Abel-integral equation's theory, we can invert (3.1.1) and obtain (c.f. [92]):

$$F_0(x) = 1 - \frac{2}{\pi} \sqrt{x}\, V_0(x) - \frac{2}{\pi} \int_x^\infty \frac{V_0(s)}{2\sqrt{s}}\, ds, \tag{3.1.2}$$

where:

$$V_0(x) := \int_x^\infty \frac{g_0(z)}{\sqrt{z - x}}\, dz. \tag{3.1.3}$$

(The proof of (3.1.2) is provided in Lemma 3.3.1 in the Appendix). In Corollary 3.1.1, we formally show that the functional $x \mapsto -\frac{2}{\pi} \int_x^\infty V_0(s)/(2\sqrt{s})\, ds$ can be estimated at the rate $\sqrt{n}$, whereas the functional $V_0(x)$ can be estimated (nonparametrically) only at the slower rate $\sqrt{n}/\sqrt{\log n}$. This can be intuitively seen because, by plugging (3.1.3) into the second term of (3.1.2) and using Fubini's theorem to exchange order of integration:

$$\int_x^\infty \frac{V_0(s)}{2\sqrt{s}}\, ds = \frac{\pi}{2} - \int_0^\infty \sin^{-1} \sqrt{1 \wedge \frac{x}{z}}\, g_0(z)\, dz \tag{3.1.4}$$

and $\text{Var}[\sin^{-1} \sqrt{1 \wedge (x/Z)}] < \infty$, for $Z \sim G_0$, in contrast with $\text{Var}[(Z - x)^{-1/2}] = \infty$. It thus follows that the estimation of V_0 and F_0 are essentially equivalent. Furthermore, as shown in [52], the relation $F_0^*(x) = 1 - \frac{V_0(x)}{V_0(0)}$ holds, which implies that also in the materials science setting, the primary target of estimation is V_0. Accordingly, our focus in this chapter will first be on estimating $V_0(x)$, and we will subsequently show that these results naturally extend to the estimation of $F_0(x)$. The latter will be the main object of interest in the astronomical application discussed in the simulation study.

We aim to develop an easy-to-compute and statistically efficient nonparametric Bayesian method for estimating $V_0(x)$ at a given point $x \geq 0$. We begin by placing a Dirichlet Process prior directly on the distribution function of the observables: $G \sim \text{DP}(\alpha)$. Given an observed sample $Z_1, \ldots, Z_n$ from cdf G_0 (associated with density g_0), the posterior distribution is (c.f. [34], Theorem 4.6 in [42]):

$$G \mid Z_1, \ldots, Z_n \sim \text{DP}(\alpha + n\mathbb{G}_n), \qquad \mathbb{G}_n := \frac{1}{n} \sum_{i=1}^n \delta_{Z_i}. \tag{3.1.5}$$

This choice offers computational simplicity due to the conjugacy of the posterior. Leveraging the inverse relations inherent to the problem, we can estimate the functional of interest using (3.1.2) by

$$V_G(x) := \int_x^\infty \frac{dG(z)}{\sqrt{z - x}}. \tag{3.1.6}$$

This initial step yields the *Naive Bayes Posterior* (NBP). We show below that this satisfies a BvM phenomenon with asymptotic variance $g_0(x)$. However, the realisations V_G of this plug-in posterior are not necessarily monotone functions. To remedy this, we project them into the $\mathbb{L}_2$ subspace of decreasing and right-continuous functions. It turns out that this not only corrects the non-monotonicity, but also improves the estimator. We show that the resulting *Isotonized Inverse Posterior* (IIP) $\hat{V}_G(x)$ satisfies a BvM theorem with improved variance that depends on the local smoothness of V_0. The smoothness at x is measured through a parameter $\gamma > \frac{1}{2}$ that can be loosely understood as the degree of Hölder continuity at x. Using (3.1.2), we can define the IIP to target the estimation of $F_0(x)$, which we will denote by $\hat{F}_G(x)$. The formal definitions of $\hat{V}_G(x)$ and $\hat{F}_G(x)$, being quite technical, are given in (3.1.16) and (3.1.18) below, respectively.

Thus the approach in this chapter has three advantages:

(i) **Computational speed**: The Dirichlet posterior (3.1.5) is explicit and its projection can be effectively computed by the Pool-Adjacent-Violators Algorithm (PAVA, c.f. [4]).

(ii) **Efficiency and adaptation**: The IIP achieves the asymptotic variance of the efficient estimators introduced in [43] and [49] and it is adaptive to the level of local smoothness of V_0.

(iii) **Uncertainty quantification**: The Bayesian framework inherently provides uncertainty quantification without the need to estimate γ or other parameters. We show below that this is asymptotically correct in the frequentist sense.

The Naive Bayes Posterior (NBP) is analogous to the naive estimator introduced in [52] and [114] which is constructed as follows: given a sample $Z_1, \dots, Z_n$ from the density g_0 and for $\mathbb{G}_n$ the empirical distribution function based on this sample,

$$V_n(x) := \int_x^\infty \frac{d\mathbb{G}_n(z)}{\sqrt{z-x}}. \tag{3.1.7}$$

In [43], the authors showed that the projection of the naive estimator onto the set of monotone functions, the *Isotonized Inverse Estimator* (IIE) $\hat{V}_n$, is asymptotically efficient and the attained asymptotic variance depends on the local smoothness of V_0 in the same way as for the IIP.

Projecting the naive posterior distribution departs from a classical Bayesian framework which might assign a Dirichlet Process (DP) prior to F and employ a hierarchical model to sample from the density of the observations. A Bernstein-von Mises (BvM) result for this approach is currently unavailable. Moreover, the available algorithms for posterior sampling rely on Gibbs sampling procedures, which are much slower than the method we propose here.

From a practical perspective, the Bernstein-von Mises (BvM) result established in this chapter assures practitioners that the Bayesian methodology based on DP prior provides all the efficiency benefits of the Isotonic Inverse Estimator (IIE) of [43], but without requiring the estimation of difficult-to-access nuisance quantities such as the density of the observations $g_0(x)$ or the local Hölder smoothness $\gamma > 1/2$. In applications, where one aims to recover the distribution of squared distances of stars from their projected two-dimensional positions or the distribution of the embedded particles' squared radii, this is especially valuable: the underlying smoothness of these distributions (and thus of V_0) may vary across radii, and the BvM result holds for all local smoothness levels $\gamma > 1/2$. This significantly improves over existing bootstrap-based uncertainty guarantees of [92], which only hold under the stronger assumption that V_0 is differentiable at the estimation point x (which is even stronger than $\gamma = 1$).

3.1.2 Main results

We start by proving a BvM result for the *Naive Bayes Posterior* (NBP), analogous to the result obtained for the naive estimator (c.f. Theorem 2 in [52]). For our first result, we need our version of g_0 to be well-defined and bounded, i.e.:

$$\|g_0\|_\infty := \sup_{z\geq 0}\left|\int_z^\infty \frac{dV_0(s)}{\pi\sqrt{s-z}}\right| < \infty. \tag{3.1.8}$$

Assumption (3.1.8) asserts that projected 2D star positions do not become arbitrarily concentrated. From a mathematical perspective, without this assumption the inverse problem becomes highly unstable: small errors in the data can cause arbitrarily large errors in estimating V_0. For instance, if g_0 is unbounded at x, this leads to instability in estimating $V_0(x)$, reflected in an infinite asymptotic variance. Thus, boundedness of g_0 ensures stability of the inverse mapping. Mathematically, if F_0 is Hölder continuous with degree $\gamma > 1/2$, then g_0 is bounded (cf. Lemma A.3 in [43]).

In equation (3.1.9) and (3.1.22) below we use the notion of conditional weak convergence in distribution (in probability). For its precise definition we refer the reader to (3.3.1)-(3.3.2) in Appendix A.

Theorem 3.1.1 (BvM for NBP)**.** *Let $x \geq 0$ and suppose that* (3.1.8) *holds true. Let $G \sim \mathrm{DP}(\alpha)$, with $\int_x^\infty (s-x)^{-1/2}\, d\alpha(s) < \infty$. Then, in probability under G_0:*

$$\frac{\sqrt{n}}{\sqrt{\log n}}(V_G(x) - V_n(x)) \mid Z_1,\ldots,Z_n \rightsquigarrow N(0, g_0(x)). \tag{3.1.9}$$

Proof. For a measure P and a function f, we denote by Pf the integral $\int f\, dP$, this holding also for empirical measures like $\mathbb{G}_n = \frac{1}{n}\sum_{i=1}^n \delta_{Z_i}$.

Since $G \mid Z_1,\ldots,Z_n \sim \mathrm{DP}(\alpha + n\mathbb{G}_n)$ by proposition G.10 in [42], for $Q \sim \mathrm{DP}(\alpha)$, $\mathbb{B}_n \sim \mathrm{DP}(n\mathbb{G}_n)$, $V_n \sim \mathrm{Be}(|\alpha|, n)$ and $f(z) = 1/\sqrt{(z-x)_+}$, $x \geq 0$, conditionally on $Z_1,\ldots,Z_n$, we have the following representation in distribution:

$$Gf = V_n Qf + (1 - V_n)\mathbb{B}_n f, \tag{3.1.10}$$

where Q, $\mathbb{B}_n$, V_n are independent. First note that by assumption Qf is a.s. well-defined (c.f. Remark 4.4 in [42]). By Proposition G.2 in [42] for $\varepsilon_i \overset{\text{i.i.d.}}{\sim} \mathrm{Exp}(1)$ and $W_{ni} := \varepsilon_i/\bar{\varepsilon}_n$, we have $(W_{n1},\ldots,W_{nn}) \sim \mathrm{Dir}(n; 1/n,\ldots,1/n)$. Since $\mathbb{B}_n \sim \mathrm{DP}(n\mathbb{G}_n)$, for δ_{Z_i} Dirac measure at Z_i, we have the following representation in distribution of $\mathbb{B}_n$ (c.f. Example 3.7.9 in [103]):

$$\mathbb{B}_n = \frac{1}{n}\sum_{i=1}^n W_{ni}\delta_{Z_i}.$$

Because of this representation (c.f. section 3.7.2. in [103]), $\mathbb{B}_n$ is also known as *Bayesian bootstrap*. Conclude that:

$$\frac{\sqrt{n}}{\sqrt{\log n}}(\mathbb{B}_n - \mathbb{G}_n)f = \frac{1}{\sqrt{n\log n}}\sum_{i=1}^n (W_{ni} - 1)\, f(Z_i).$$

Because $|\bar{\varepsilon}_n - 1| = O_p(1/\sqrt{n})$, we study: $(n\log n)^{-1/2}\sum_{i=1}^n (\varepsilon_i - 1)\, f(Z_i)$. Now note that by Lemma 3.3.6 in the Appendix:

$$\frac{1}{\sqrt{n\log n}}\sum_{i=1}^n \frac{(\varepsilon_i - 1)}{\sqrt{Z_i - x}}\mathbf{1}_{\{Z_i > x\}} \mid Z_1,\ldots,Z_n \rightsquigarrow N(0, g_0(x)) \quad \text{under } G_0. \tag{3.1.11}$$

Because Qf is a.s. well-defined, we obtain the claim by Slutsky's lemma upon noticing $\frac{\sqrt{n}}{\sqrt{\log n}} V_n Qf \overset{G_0}{\to} 0$. □

We consider a projection-type posterior by isotonizing draws from the obtained naive posterior (3.1.6) as follows. First define, for $G \sim \mathrm{DP}(\alpha + n\mathbb{G}_n)$ and $\mathbb{G}_n$, G_0 as above:

$$U_G(x) := \int_0^x V_G(y)\,dy = 2\int_0^\infty \sqrt{z}\,dG(z) - 2\int_x^\infty \sqrt{z-x}\,dG(z), \tag{3.1.12}$$

$$U_n(x) := \int_0^x V_n(y)\,dy = 2\int_0^\infty \sqrt{z}\,d\mathbb{G}_n(z) - 2\int_x^\infty \sqrt{z-x}\,d\mathbb{G}_n(z), \tag{3.1.13}$$

$$U_0(x) := \int_0^x V_0(y)\,dy = 2\int_0^\infty \sqrt{z}\,dG_0(z) - 2\int_x^\infty \sqrt{z-x}\,dG_0(z). \tag{3.1.14}$$

Furthermore, for a continuous function $k : [0,\infty) \to [0,\infty)$, the Least Concave Majorant (LCM) and its right-hand side derivative, are given for $x \in [0,\infty)$ by:

$$k^*(x) := \inf\{f(x) \,:\, f \geq k \text{ on } [0,\infty),\ f \text{ concave on } [0,\infty)\},$$
$$(k^*)'_+(x) := \lim_{h\to 0^+} \frac{k^*(x+h) - k^*(x)}{h}.$$

We denote by $\hat{V}_n$ (IIE) the right-hand side derivative of U_n^*, the LCM of U_n. Similarly, $\hat{V}_G$ is the right-hand side derivative of U_G^*, the LCM of U_G, i.e., the isotonization of a draw from the naive Bayes posterior. Thus, for G as in (3.1.5):

$$\hat{V}_n(x) := (U_n^*)'_+(x), \tag{3.1.15}$$

$$\hat{V}_G(x) := (U_G^*)'_+(x). \tag{3.1.16}$$

By Lemma 2 in [52], $\hat{V}_n$ and $\hat{V}_G$ can be viewed as (equivalent to) $\mathbb{L}_2$-projections of the naive estimators V_n and V_G into the space of nonnegative, decreasing and right-continuous functions on $[0,\infty)$. We can finally define the Naive posterior and the IIP to target the estimation of F_0 as follows:

$$F_G(x) := 1 - \frac{2}{\pi}\sqrt{x}\,V_G(x) - \frac{2}{\pi}\int_x^\infty \frac{V_G(s)}{2\sqrt{s}}\,ds, \tag{3.1.17}$$

$$\hat{F}_G(x) := 1 - \frac{2}{\pi}\sqrt{x}\,\hat{V}_G(x) - \frac{2}{\pi}\int_x^\infty \frac{\hat{V}_G(s)}{2\sqrt{s}}\,ds, \tag{3.1.18}$$

and analogously we define $F_n(x)$, with V_n instead of V_G in (3.1.17) and $\hat{F}_n(x)$, with $\hat{V}_n$ instead of $\hat{V}_G$ in (3.1.18).

We now state two additional conditions needed for Theorem 3.1.2. First, a natural assumption is a moment constraint for Z, (which can be translated into a moment condition on X c.f. [43]):

$$\int_0^\infty z\,g_0(z)\,dz < \infty. \tag{3.1.19}$$

This is a mild assumption, which is practically always satisfied, given that materials studied in materials science are of bounded size and globular clusters typically do not extend indefinitely. Additionally, assume $\exists\, K > 0$ and $\gamma > \frac{1}{2}$ such that

$$\int_0^1 (V_0(x) - V_0(x+u\delta))\,du \sim \mathrm{sgn}(\delta)|\delta|^\gamma K, \qquad \text{as } \delta \to 0, \tag{3.1.20}$$

where $\sim$ means that the ratio of the left-hand side and of the right-hand side goes to 1 as $\delta \to 0$. The constant γ indicates the level of smoothness at x of V_0. This is analogous to equation (5) from [43]. Note that it is possible to see what this condition means in terms of the underlying F_0, using (3.1.2), we obtain that $\exists\, K > 0$ and $\gamma > \frac{1}{2}$ such that as $\delta \to 0$:

$$\int_0^1 (\sqrt{x}\, V_0(x) - \sqrt{x + u\delta}\, V_0(x + u\delta))\, du \sim \operatorname{sgn}(\delta)|\delta|^\gamma K - \frac{V_0(x)}{4\sqrt{x}} \operatorname{sgn}(\delta)|\delta|, \tag{3.1.21}$$

(upon using a Taylor expansion for $\int_0^1 (\sqrt{x} - \sqrt{x + u\delta})\, du$). Because $\frac{2}{\pi} \int_x^\infty \frac{V_0(s)}{2\sqrt{s}}\, ds$ is Lipschitz continuous, the local smoothness of F_0 at x is $\min(\gamma, 1)$. Therefore for the most relevant levels of smoothness, i.e. $\gamma \in (\frac{1}{2}, 1]$, the smoothness of V_0 carries over to F_0. Because $F_0^*(x) = 1 - \frac{V_0(x)}{V_0(0)}$, equation (5) from [43] is equivalent to (3.1.20) (note that in that paper F denotes the current F_0^*). Thus, for a better mathematical understanding of the role of γ, we refer the interested reader to the discussion given in [43], in particular Example 2.1, 2.2, 2.3 and Remark 2.1.

From a physical point of view, the role of γ is also interpretable. In a globular cluster, different regions (e.g., dense core vs. sparse halo) may correspond to different levels of smoothness of F_0, and the semiparametric theory reflects this. The local adaptivity of the Bayesian estimator is encoded in the limiting variance of the Bernstein-von Mises mirrors this phenomenon. If there is a sharp increase in the concentration of stars with size close to x (i.e., γ close to $1/2$), then the 2D observations are much noisier. This makes the estimation more difficult and results in a higher asymptotic variance.

Theorem 3.1.2 (BvM for IIP)**.** *Let $x \geq 0$, suppose that* (3.1.8)*,* (3.1.19)*, and* (3.1.20) *hold true. Moreover let $G \sim \mathrm{DP}(\alpha)$, where the base measure α has bounded density. Then, in probability under G_0:*

$$\frac{\sqrt{n}}{\sqrt{\log n}} \left(\hat{V}_G(x) - \hat{V}_n(x) \right) \mid Z_1, \ldots, Z_n \rightsquigarrow N\left(0, \frac{g_0(x)}{2\gamma}\right). \tag{3.1.22}$$

Proof. The proof essentially consists of three parts. The first part is to rewrite the event

$$\left\{ \sqrt{\frac{n}{\log n}} \left(\hat{V}_G(x) - \hat{V}_n(x) \right) \leq a \right\}$$

given $Z_1, \ldots, Z_n$ in terms of the location of the maximum (arg max) of a stochastic process $t \mapsto \tilde{Z}_n^G(t) - at$. It turns out that the process $\tilde{Z}_n^G(t)$ can be decomposed into three essential parts. The second step consists in studying these processes. One process, which can be interpreted as the Bayesian bootstrap counterpart of the process studied in the proof of Theorem 2.1 in [43] (c.f. eq. (14) in [43]). This process converges conditionally in distribution to a Gaussian process $\mathbb{Z}$, which we show can be written as $\mathbb{Z}(t) = tX$, where $X \sim N(0, g_0(x)/(2\gamma))$. A second process (see (3.1.29)), which is a deterministic process, whose behavior is controlled by the local smoothness of V_0 at x and which converges to $-|t|^{\gamma+1}K$ (with γ and K as in (3.1.20)). Lastly, a third remainder process which converges uniformly to zero with respect to the topology of uniform convergence on compacta. The final step is to prove that from the conditional convergence in distribution of $t \mapsto \tilde{Z}_n^G(t) - at$ to $\mathbb{Z}(t) - at$ it follows that the arg max of $(\tilde{Z}_n^G(t) - at)$ converges in distribution to the arg max of $\mathbb{Z}(t) - at$. Since the arg max functional is certainly not continuous on the space of continuous functions equipped with the uniform norm, the simplest form of the continuous mapping theorem cannot be applied. For

this reason, we prove in Lemma 3.3.9 specifically for this application: a conditional argmax continuous mapping theorem (extension of Theorem 3.2.2 in [103]). From (3.1.36) below, it follows that the sequence of locations of the maximum of $(\tilde{Z}_n(t) - at)$ is conditionally uniformly tight. This enables us to use Lemma 3.3.9. Moreover, since the limiting process turns out almost surely to have a unique maximum, the asserted conditional convergence in distribution in (3.1.22) follows.

For sequences $\delta_n = \sqrt{\log n}/\sqrt{n}$ and $\delta_n^* = \delta_n^{1/\gamma}$ where $\gamma > 1/2$ satisfies condition (3.1.20), define

$$Z_n^{G_0}(t) := \delta_n^{-1}(\delta_n^*)^{-1}\{U_n(x+\delta_n^* t) - U_n(x) - U_0(x+\delta_n^* t) + U_0(x)\} \tag{3.1.23}$$

$$= 2\delta_n^{-1}(\delta_n^*)^{-1}\int \left\{\sqrt{(z-x)_+} - \sqrt{(z-x-\delta_n^* t)_+}\right\} d(\mathbb{G}_n - G_0)(z). \tag{3.1.24}$$

By Lemma 3.3.2 in Appendix A it suffices to show, for $X \sim N\left(0, \frac{g_0(x)}{2\gamma}\right)$ and every $a \in \mathbb{R}$:

$$\Pi_n\left(\sqrt{\frac{n}{\log n}}\left(\hat{V}_G(x) - V_0(x)\right) - Z_n^{G_0}(1) \le a \mid Z_1, \ldots, Z_n\right) \xrightarrow{G_0} \mathrm{P}(X \le a). \tag{3.1.25}$$

Define the process T_G for $a > 0$:

$$T_G(a) := \inf\{t \ge 0 : U_G(t) - at \text{ is maximal}\}.$$

By the switch relation (see Lemma 3.2 in [53]):

$$T_G(a) \le x \iff \hat{V}_G(x) \le a.$$

Therefore, for sufficiently large n, conditionally on $Z_1, \ldots, Z_n$:

$$\delta_n^{-1}(\hat{V}_G(x) - V_0(x)) - Z_n^{G_0}(1) \le a \iff (\delta_n^*)^{-1}\left(T_G\left(V_0(x) + \delta_n(a + Z_n^{G_0}(1))\right) - x\right) \le 0.$$

Because the location of a maximum of a function is equivariant under translations and invariant under multiplication by a positive number and addition of a constant:

$$\begin{aligned}
&(\delta_n^*)^{-1}\left(T_G\left(V_0(x) + \delta_n(a + Z_n^{G_0}(1))\right) - x\right) \\
&= (\delta_n^*)^{-1}\big(\inf\{x + t \ge 0 : U_G(x+t) - V_0(x)(x+t) \\
&\qquad - \delta_n(a + Z_n^{G_0}(1))(x+t) \text{ is max}\} - x\big) \\
&= \inf\{t \ge -(\delta_n^*)^{-1}x : U_G(x+\delta_n^* t) - U_G(x) - U_n(x+\delta_n^* t) + U_n(x) \\
&\quad + U_0(x+\delta_n^* t) - U_0(x) - V_0(x)\delta_n^* t - \delta_n\delta_n^*(at + Z_n^{G_0}(1)t - Z_n^{G_0}(t)) \text{ is max}\} \qquad (3.1.26) \\
&= \inf\{t \ge -(\delta_n^*)^{-1}x : \tilde{Z}_n^G(t) - at \text{ is max}\}.
\end{aligned}$$

where we define the localized process $\tilde{Z}_n^G$ by,

$$\tilde{Z}_n^G(t) := \delta_n^{-1}(\delta_n^*)^{-1}\{U_G(x+\delta_n^* t) - U_G(x) - U_n(x+\delta_n^* t) + U_n(x)\} \tag{3.1.27}$$

$$+ \delta_n^{-1}(\delta_n^*)^{-1}\{U_0(x+\delta_n^* t) - U_0(x) - V_0(x)\delta_n^* t\} + Z_n^{G_0}(t) - tZ_n^{G_0}(1) \tag{3.1.28}$$

for $t \in I_x$, where $I_x := [0, \infty)$ if $x = 0$, and $I_x := \mathbb{R}$ if $x > 0$.

We determine the asymptotic behavior of the process $\tilde{Z}_n^G$ by studying separately the components in (3.1.27) and (3.1.28), and next we conclude by applying Slutsky's lemma. First

note that $|Z_n^{G_0}(1)t - Z_n^{G_0}(t)| = o_{G_0}(1)$ uniformly in t in compacta by the proof of Theorem 2.1 in [43] (c.f. display after equation (16)). Furthermore, using assumption (3.1.20) (as in equation (13) in [43]), uniformly in t in compacta:

$$\delta_n^{-1}\left(\delta_n^*\right)^{-1}\left\{U_0\left(x+\delta_n^* t\right) - U_0(x) - V_0(x)\delta_n^* t\right\} = -|t|^{\gamma+1}(K+o(1)). \tag{3.1.29}$$

This determines the behavior of (3.1.28). Now we study the posterior behaviour of the process in (3.1.27), which can be interpreted as a *Bayesian bootstrap* counterpart of the process (3.1.23) studied in the proof of Theorem 2.1 in [43] (c.f. eq. (14) in [43]):

$$\begin{aligned} Z_n^G(t) &:= \delta_n^{-1}\left(\delta_n^*\right)^{-1}\left\{U_G\left(x+\delta_n^* t\right) - U_G(x) - U_n\left(x+\delta_n^* t\right) + U_n(x)\right\} \\ &= 2\delta_n^{-1}\left(\delta_n^*\right)^{-1}\int\left\{\sqrt{(z-x)_+} - \sqrt{(z-x-\delta_n^* t)_+}\right\} d\left(G-\mathbb{G}_n\right)(z) \end{aligned}$$

As in proof of Theorem 3.1.1 in equation (3.1.10), we have by Proposition G.10 in [42], for independent $Q \sim \mathrm{DP}(\alpha)$, $\mathbb{B}_n \sim \mathrm{DP}(n\mathbb{G}_n)$, and $V_n \sim \mathrm{Be}(|\alpha|, n)$, the following representation in distribution:

$$Gf_t^n = V_n Q f_t^n + (1-V_n)\mathbb{B}_n f_t^n, \tag{3.1.30}$$

where $f_t^n(z) := 2(\delta_n^*)^{-1}\{\sqrt{(z-x)_+} - \sqrt{(z-x-\delta_n^* t)_+}\}$ and thus the following representation in distribution of the process $Z_n^G(t)$:

$$Z_n^G(t) = \delta_n^{-1} V_n(Q-\mathbb{G}_n)f_t^n + \delta_n^{-1}(1-V_n)(\mathbb{B}_n - \mathbb{G}_n)f_t^n.$$

This representation is crucial for the asymptotic analysis of the process $Z_n^G(t)$. The first thing to do is to identify which parts of the process $Z_n^G(t)$ do not give any contribution to limiting dstribution of the process, as they converge conditionally to zero in probability. The second part is to identify the limiting distribution of the remaining parts of the process, which will be a Gaussian process. In Appendix A, we give the proof that uniformly in t in compacta:

$$Z_n^G(t) = \mathbb{Z}_n^*(t) + o_p(1), \tag{3.1.31}$$

where for $\varepsilon_i \overset{\text{i.i.d.}}{\sim} \mathrm{Exp}(1)$, the process $\mathbb{Z}_n^*$ is defined as:

$$\mathbb{Z}_n^*(t) := \frac{1}{\sqrt{n\log n}}\sum_{i=1}^n (\varepsilon_i - 1)\, f_t^n(Z_i). \tag{3.1.32}$$

Then by Slutsky's lemma, if we determine the conditional weak limit of $(\mathbb{Z}_n^*(t) : t \in I_x)$, then we determine the one of $(Z_n^G(t) : t \in I_x)$. We now study the marginals of $\mathbb{Z}_n^*$. Note that by Lemma 3.3.7 for fixed $t \in I_x$, in probability under G_0:

$$\mathbb{Z}_n^*(t) \mid Z_1, \ldots, Z_n \rightsquigarrow N\left(0, t^2 g_0(x)/(2\gamma)\right), \tag{3.1.33}$$

and again by Lemma 3.3.7 that:

$$\mathrm{Cov}\left(\mathbb{Z}_n^*(t), \mathbb{Z}_n^*(s) \,\middle|\, Z_1, \ldots, Z_n\right) = \frac{1}{2\gamma} g_0(x) st\left(1 - O_{G_0}\left(\frac{\log\log n}{\log n}\right)\right) + O_{G_0}\left(\frac{1}{\log n}\right), \tag{3.1.34}$$

By the conditional Chebyshev inequality and (3.1.34) we conclude $\forall\, \varepsilon > 0$:

$$\mathrm{P}\left(|\mathbb{Z}_n^*(t) - t\mathbb{Z}_n^*(1)| > \varepsilon \mid Z_1, \ldots, Z_n\right) \le \frac{1}{\varepsilon^2}\mathrm{Var}(\mathbb{Z}_n^*(t) - t\mathbb{Z}_n^*(1) \mid Z_1, \ldots, Z_n) \overset{G_0}{\to} 0.$$

By (3.1.31) conclude that (3.1.33) holds true with $\mathbb{Z}_n^*$ replaced by Z_n^G. Therefore, the finite-dimensional distributions of Z_n^G converge conditionally in distribution to the finite-dimensional distributions of the process: $\mathbb{Z}(t) = tX$, where $X \sim N(0, g_0(x)/(2\gamma))$.

Lemma 3.3.8 gives the conditional asymptotic equicontinuity of the process $\mathbb{Z}_n^*$. The conditional asymptotic equicontinuity together with the conditional convergence in distribution of the finite dimensional distributions, implies in probability under G_0 as $n \to \infty$:

$$\mathbb{Z}_n^* \mid Z_1, \ldots, Z_n \rightsquigarrow \mathbb{Z} \quad \text{in} \quad \ell^\infty(K). \tag{3.1.35}$$

for every compact $K \subset I_x$ and ℓ^∞ being the space of bounded functions on K equipped with the supremum norm. For a review of the conditional weak convergence in ℓ^∞, see [103] (c.f. Example 1.5.1 and Theorem 1.5.4). By (3.1.31) and (3.1.33) conclude that (3.1.35) holds true with $\mathbb{Z}_n^*$ replaced by Z_n^G, i.e. the conditional weak limit of Z_n^G relative to the topology of uniform convergence on compacta is the same Gaussian process. Altogether, recalling also (3.1.29), we can conclude that conditionally on $Z_1, \ldots, Z_n$ the weak limit of $(\tilde{Z}_n^G(t) - at \,:\, t \in K)$ in ℓ^∞ is given by $(tX - |t|^{\gamma+1}K - at \,:\, t \in K)$ for every compact $K \subset I_x$.

In order to apply Lemma 3.3.9, notice that $\mathbb{Z}$ has a.s. continuous sample paths and a.s. unique maxima (see Lemma 2.6 in [65]) and that $\hat{t}_n^G := \inf\{t \geq -(\delta_n^*)^{-1}x : \tilde{Z}_n^G(t) - at \text{ is max}\}$ is an argmax process conditionally on $Z_1, \ldots, Z_n$, satisfying the second condition on Lemma 3.3.9. It is thus enough to show that there exists a conditionally stochastically bounded sequence $(\tilde{M}_n)$ that satisfies:

$$\Pi_n\left((\delta_n^*)^{-1}\left|T_G\left(V_0(x) + \delta_n(a + Z_n^{G_0}(1))\right) - x\right| \leq \tilde{M}_n \mid Z_1, \ldots, Z_n\right) \xrightarrow{G_0} 1, \tag{3.1.36}$$

The proof of (3.1.36) is given in Appendix A. Thus by Lemma 3.3.9 we can conclude:

$$\begin{aligned}&\Pi_n\left(\sqrt{\frac{n}{\log n}}\left(\hat{V}_G(x) - V_0(x)\right) - Z_n^{G_0}(1) \leq a \mid Z_1, \ldots, Z_n\right)\\ &\xrightarrow{G_0} \mathrm{P}\left\{\arg\max_{t \in I_x}\left\{tX - |t|^{\gamma+1}K - at\right\} \leq 0\right\} = \mathrm{P}\{X \leq a\},\end{aligned}$$

because $t \mapsto t(X - a) - |t|^{\gamma+1}K$ attains its maximum on $(-\infty, 0]$ iff $X - a \leq 0$. □

We conclude this section by showing the implications of the above results for the uncertainty quantification of the posterior. Under the conditions of Theorem 3.1.2 (c.f. [43])

$$\frac{\sqrt{n}}{\sqrt{\log n}}\left(\hat{V}_n(x) - V_0(x)\right) \rightsquigarrow N\left(0, \frac{g_0(x)}{2\gamma}\right).$$

Thus as consequence of (3.1.2) and the analogous definition of $\hat{F}_n(x)$, we have that (see proof of Corollary 3.1.1 in the Appendix):

$$\frac{\sqrt{n}}{\sqrt{\log n}}\left(\hat{F}_n(x) - F_0(x)\right) \rightsquigarrow N\left(0, \frac{2x}{\pi^2\gamma}g_0(x)\right). \tag{3.1.37}$$

Together with (3.1.39) below, this will imply that the credible intervals constructed from the Isotonized Inverse Posterior (IIP) of $\hat{F}_G(x)$ are (asymptotic) frequentist confidence sets. The following Corollary 3.1.1, whose proof is given in the Appendix, summarizes formally the above discussion. Note that because the uncertainty quantification based on the IIP is always preferable to the one based on the Naive Bayes Posterior, we only propose uncertainty quantification using the latter.

Corollary 3.1.1 (BvM for $\hat{F}_G$ and F_G and UQ for F_0)**.** *Let F_G as in* (3.1.17) *and $\hat{F}_G$ as in* (3.1.18) *and analogously let $F_n(x)$, with V_n instead of V_G in* (3.1.17) *and $\hat{F}_n(x)$, with $\hat{V}_n$ instead of $\hat{V}_G$ in* (3.1.18)*. Under the same conditions of Theorem 2.1 in [47], in probability under G_0 as $n \to \infty$:*

$$\frac{\sqrt{n}}{\sqrt{\log n}} \left(F_G(x) - F_n(x)\right) \mid Z_1, \ldots, Z_n \rightsquigarrow N\left(0, \frac{4x}{\pi^2} g_0(x)\right). \tag{3.1.38}$$

Under the same conditions of Theorem 2.2 in [47], in probability under G_0 as $n \to \infty$:

$$\frac{\sqrt{n}}{\sqrt{\log n}} \left(\hat{F}_G(x) - \hat{F}_n(x)\right) \mid Z_1, \ldots, Z_n \rightsquigarrow N\left(0, \frac{2x}{\pi^2 \gamma} g_0(x)\right). \tag{3.1.39}$$

Further, if we define the posterior quantiles for isotonized posterior $\hat{F}_G$, for $\alpha \in [0,1]$, as:

$$q_\alpha := \inf\{t \in \mathbb{R} \,:\, \Pi_n(\hat{F}_G(x) \leq t \mid Z_1, \ldots, Z_n) \geq \alpha\}$$

then under G_0 as $n \to \infty$:

$$\mathrm{P}\left(F_0(x) \in [q_{\alpha/2}, q_{1-\alpha/2}]\right) \to 1 - \alpha. \tag{3.1.40}$$

Proof. We start by proving (3.1.39), as the proof of (3.1.38) is analogous but simpler. First note that by the inverse relation (3.1.2) inherent to Wicksell's problem:

$$\hat{F}_G(x) - \hat{F}_n(x) = -\frac{2}{\pi}\sqrt{x}\left\{\hat{V}_G(x) - \hat{V}_n(x)\right\} - \frac{2}{\pi}\int_x^\infty \frac{\hat{V}_G(u) - \hat{V}_n(u)}{2\sqrt{u}}\, du.$$

From Theorem 2.2 in [47], we obtain the weak conditional limit of $\frac{\sqrt{n}}{\sqrt{\log n}}\left\{\hat{V}_G(x) - \hat{V}_n(x)\right\} \mid Z_1, \ldots, Z_n$. Integrating by parts, the integral on the last display is at most

$$\begin{aligned}\frac{\left|U_G^*(x) - U_n^*(x)\right|}{2\sqrt{x}} + \frac{1}{4}\left|\int_x^\infty \frac{U_G^*(u) - U_n^*(u)}{u^{3/2}}\, du\right| &\leq \frac{\left\|U_G^* - U_n^*\right\|_\infty}{2\sqrt{x}} + \left\|U_G^* - U_n^*\right\|_\infty \frac{1}{2\sqrt{x}} \\ &= \frac{\left\|U_G^* - U_n^*\right\|_\infty}{\sqrt{x}} \leq \frac{\|U_G - U_n\|_\infty}{\sqrt{x}}.\end{aligned}$$

where in the last inequality we applied a generalized version of Marshall's inequality (c.f. Lemma 2.2 in [29]). Now note that $\sqrt{n}\,\|U_G - U_n\|_\infty \mid Z_1, \ldots, Z_n$ has a conditional weak limit by Theorem 12.2 in [42] combined with the Continuous Mapping Theorem. We can apply Theorem 12.2 because the class of functions $\mathcal{F} := \{z \mapsto \sqrt{(z)_+} - \sqrt{(z-s)_+} \,:\, s \geq 0\}$ is Donsker by Lemma 2.1 in [43]. Conclude thus by Slutsky's lemma upon noticing $\frac{\sqrt{n}}{\sqrt{\log n}}\,\|U_G - U_n\|_\infty \mid Z_1, \ldots, Z_n \to 0$ under G_0.

(3.1.38) follows using the fact that $\sqrt{n}\,\|U_G - U_n\|_\infty \mid Z_1, \ldots, Z_n$ has a conditional weak limit and the rewriting:

$$F_G(x) - F_n(x) = -\frac{2}{\pi}\sqrt{x}\left\{V_G(x) - V_n(x)\right\} - \frac{2}{\pi}\int_x^\infty \frac{V_G(u) - V_n(u)}{2\sqrt{u}}\, du.$$

We now show the third claim (3.1.40). By (3.1.39) and Lemma 21.2 in [102], in probability under G_0 as $n \to \infty$:

$$\widetilde{q}_\alpha := \inf\left\{t \in \mathbb{R} \,:\, \Pi_n\left(\frac{\sqrt{n}}{\sqrt{\log n}}\left(\hat{F}_G(x) - \hat{F}_n(x)\right) \leq t \mid Z_1, \ldots, Z_n\right) \geq \alpha\right\}$$

$$\xrightarrow{G_0} \sqrt{\frac{2xg_0(x)}{\pi^2\gamma}}\Phi^{-1}(\alpha),$$

where Φ^{-1} is the quantile function of the standard normal distribution. Then note:

$$\begin{aligned} q_\alpha &= \inf_{t\in\mathbb{R}}\left\{\Pi_n\left(\frac{\sqrt{n}}{\sqrt{\log n}}\left(\hat{F}_G(x)-\hat{F}_n(x)\right)\le\frac{\sqrt{n}}{\sqrt{\log n}}\left(t-\hat{F}_n(x)\right)\mid Z_1,\dots,Z_n\right)\ge\alpha\right\} \\ &= \hat{F}_n(x)+\frac{\sqrt{\log n}}{\sqrt{n}}\widetilde{q}_\alpha = \hat{F}_n(x)+\sqrt{\frac{\log n}{n}\cdot\frac{2xg_0(x)}{\pi^2\gamma}}\,\Phi^{-1}(\alpha)+o_{G_0}\left(\sqrt{\frac{\log n}{n}}\right). \end{aligned}$$

Note that

$$\frac{\sqrt{n}}{\sqrt{\log n}}\left(\hat{F}_n(x)-F_0(x)\right)\rightsquigarrow N\left(0,\frac{2x}{\pi^2\gamma}g_0(x)\right) \tag{3.1.41}$$

follows by the same rewriting in (3.1.2) once we notice, by Marshall's inequality (c.f. Exercise 3.1 in [53]) and Lemma 2.1 in [43], that as $n\to\infty$:

$$\left|\frac{\sqrt{n}}{\sqrt{\log n}}\int_x^\infty\frac{\hat{V}_n(u)-V_0(u)}{\pi\sqrt{u}}\,du\right|\le\frac{\sqrt{n}}{\sqrt{\log n}}\frac{\|U_n-U_0\|_\infty}{\sqrt{x}}\xrightarrow{G_0}0.$$

Using (3.1.41) and the symmetry of the normal distribution, for $X\sim N(0,(2xg_0(x))/(\pi^2\gamma))$

$$\begin{aligned} \mathrm{P}\left(F_0(x)\in[q_{\alpha/2},q_{1-\alpha/2}]\right) &= \mathrm{P}\left(\sqrt{\frac{n}{\log n}}\left(\hat{F}_n(x)-F_0(x)\right)\in[-\widetilde{q}_{1-\alpha/2},-\widetilde{q}_{\alpha/2}]\right) \\ &\to \mathrm{P}\left(X\in\sqrt{\frac{2xg_0(x)}{\pi^2\gamma}}[\Phi^{-1}(\alpha/2),\Phi^{-1}(1-\alpha/2)]\right)=1-\alpha, \end{aligned}$$

where we assumed that $\alpha\in[0,1]$. □

3.1.3 Simulation Study and Discussion about Uncertainty Quantification

In this section, we present simulations to illustrate the practical behavior of the IIP in comparison to the IIE studied in [43]. Furthermore we present a simulation study that compares the IIP-uncertainty quantification to the one proposed in [92].



$n = 200$



$n = 2000$



Zoom for $n = 2000$



Histograms



QQ-plot

Figure 3.1: *Synthetic* setting 1: uncertainty for different sample sizes and analysis of normality at $x = 1.5$, based on 300 draws from the posterior.

We consider three distinct settings. The first two are *synthetic* settings where the true underlying distribution function F_0 is known. The first two simulation settings serve as valuable benchmarks for assessing the quality of our methodology. In particular, the first one verifies that the variance attained by the IIP coincides with the theoretical one that we derived

in Theorem 3.1.1. The second *synthetic* setting compares the uncertainty quantification of our methodology to the bootstrap-based one derived in [92]. The last simulation setting is a *real-data* application involving the positions of stars in the Globular Cluster Messier 62. In the last case, the ground truth distribution of the stars is unknown. Also in this setting, we compare our methodology with the bootstrap of [92].

In Figure 3.1, we analyze the setting where the underlying cdf F_0 is exponential with rate 1.2. Samples $Z_1, \ldots, Z_n$ used to compute the projection posterior of the IIP are generated hierarchically: $X_1, \ldots, X_n \sim F_0$, $Y_1, \ldots, Y_n \sim \text{Be}(1, 1/2)$, with $Z_i = X_i \cdot Y_i$ for all i. The plots use the following color scheme: green for estimates of F based on prior draws from $G \sim \text{DP}(M\alpha)$ (parameters M and α specified in the plots); purple for draws from the isotonized posterior; red for the average of these draws (approximating the posterior mean); yellow for the Isotonic Inverse Estimator (IIE) of [43]; and orange for F_0.

The first row of Figure 3.1 compares the behavior of the IIP and the IIE as the sample size goes from 200 to 2000. With $n = 200$, the IIE and the average of the IIP draws already behave very similarly and with $n = 2000$, the uncertainty significantly reduces and the two estimation procedures essentially coincide. The average IIP appears to be somewhat smoother, which will typically be seen as an advantage. The middle row focuses on $n = 2000$, providing a zoomed-in view and highlighting $x = 1.5$, the point used for subsequent analysis of the normality. The final row examines normality ($n = 2000$): histograms, QQ-plots, and statistical tests strongly confirm normality of the gathered samples. Furthermore, the Bayesian-estimated standard deviation aligns closely with the theoretical value attained by the frequentist procedure (c.f. [43], note that $\gamma = 1$ because F_0 is Lipschitz). In all the plots we employ 300 posterior draws.

In Figure 3.2, we analyze six different settings where at $x = 5$ the underlying cdfs F_0 have varying degrees of smoothness, making them locally Hölder smooth of degree $\gamma \in \{0.55, 0.6, 0.65, 0.7, 0.8, 1.5\}$. In particular, for an appropriate constant $K > 0$, $F_0(y) = \frac{1}{2} - K(5 - y)^\gamma$, for $y \in [0, 5]$ and $F_0(y) = \frac{1}{2} + K(y - 5)^\gamma$, for $y \in [5, 10]$. The plots use the following color scheme: pink for the Bayesian-based credible sets; red for the approximated posterior mean; light blue for the bootstrap-based confidence sets of [92]; yellow for the Isotonic Inverse Estimator (IIE) of [43]; and lime green for F_0. The prior base measure used for these simulation is the same used in the previous simulation study. In all cases $n = 2000$ and we employed 300 posterior-based and bootstrap-based draws. This comparison confirms that the IIP delivers valid uncertainty quantification for varying degrees of smoothness and that the uncertainty quantification delivered by [92] behaves closely, even though the authors formally showed it works only for a differentiable underlying truth F_0.

In Figure 3.3, we apply the methodology to star position data from globular cluster Messier 62. Let $\mathbf{Z} = (Z_{p_1}, Z_{p_2}, Z_{p_3})$ represent a star's three-dimensional position, where only (Z_{p_1}, Z_{p_2}) are observed. Here, $X = Z_{p_1}^2 + Z_{p_2}^2 + Z_{p_3}^2 \sim F_0$, and $Z = Z_{p_1}^2 + Z_{p_2}^2$. Beyond serving as a compelling application of our methodology, this example highlights a few key conclusions. First of all, also for real data the IIE closely resembles the IIP. However, because in this case the true γ is unknown, it is not immediately clear how the frequentist uncertainty quantification compares with the Bayesian one. From the work of [92], we know that bootstrap procedures work for Wicksell's problem, at least when the underlying truth is differentiable at x. We thus compared our methodology with the one proposed in [92] and we observed that they behave very similarly with 1000 or more samples. In Figure 3.3 the reader can observe an instance with $n = 1400$.



Figure 3.2: *Synthetic* setting 2: comparison uncertainty quantification IIP and bootstrap method from [92].

	$\gamma = 0.55$	$\gamma = 0.6$	$\gamma = 0.65$	$\gamma = 0.7$	$\gamma = 0.8$	$\gamma = 1.5$
Bootstrap CI width at $x = 5$	0.116	0.110	0.104	0.108	0.100	0.073
Bayesian CI width at $x = 5$	0.114	0.121	0.103	0.101	0.099	0.077



Figure 3.3: *Real-data* application: $n = 1400$ projected star positions. On the left-hand side, the IIE from [43] and the IIP with 300 posterior samples; on the right-hand side, a comparison between the 95% bootstrap confidence set for the IIE from [92] and the 95% credible sets for the IIP.

Discussion about Uncertainty Quantification

The main limitation of the approach in [43] is the lack of uncertainty quantification, which is essential for practical applications. The asymptotic variance of the limiting distribution achieved by the IIE involves two nuisance parameters: the local smoothness γ and the density $g_0(x)$ of the observed data at the evaluation point x. While $g_0(x)$ may be estimable (for most practical applications the rate of convergence would be slower than $\sqrt{n}/\sqrt{\log n}$), estimating γ is particularly challenging, as it requires recovering the local smoothness of the target distribution function F_0 at a specific point x using samples from g_0, effectively a separate inverse problem. To our knowledge, no practical methods currently exist for this task. Moreover, because of the method proposed in this chapter, estimating γ offers little practical benefit and thus makes this research direction of limited value. This highlights the key advantage of the Bayesian approach proposed in this chapter: it bypasses the need to estimate either γ or $g_0(x)$, as the posterior distribution inherently provides valid uncertainty quantification over a wide range of local smoothness levels, i.e. all $\gamma > 1/2$. For $\gamma \leq 1/2$, the inverse problem becomes severely ill-behaved due to the unboundedness of the density of the observations g_0. This might even lead to a slower rate of convergence both for the naive

plug-in estimator and for the IIE. Therefore, for all practically relevant cases with $\gamma > 1/2$, practitioners can apply our method without needing to estimate nuisance parameters or make strong assumptions about the form of F_0, thanks to its adaptive nature and the automatic uncertainty quantification.

The results displayed in Figure 3.2 suggest that the bootstrap method of [92] behaves very closely to our method. We can thus make the empirically grounded hypothesis that the bootstrap method proposed in [92] might work under the less stringent assumptions of the current chapter. The value of this simulation study has to be taken carefully, as a formal mathematical proof is still needed. However, proving this hypothesis is nontrivial and we therefore leave this as potential furure work.

From a practitioner's perspective, thus, the key advantage of our Bayesian methodology lies in its provable robustness across a broad class of smoothness levels: it provides valid uncertainty quantification for all $\gamma > 1/2$. In contrast, the bootstrap approach in [92] is provably consistent only when F_0 is differentiable at x, a substantially stronger condition (stricter than $\gamma = 1$). This distinction is crucial, since determining whether F_0 is differentiable at a given point based solely on samples from g_0 constitutes a challenging inverse problem both theoretically and in practice. By avoiding this requirement, our method offers a more broadly applicable and practitioner-friendly solution.

3.1.4 Conclusion

We have introduced a novel Bayesian approach for nonparametric estimation in Wicksell's problem, leveraging a Dirichlet Process prior on the distribution of observables rather than unobservables. This deviation from the classical Bayesian framework in inverse problems allows for computational efficiency through posterior conjugacy. By projecting the posterior draws onto the $\mathbb{L}_2$ subspace of increasing, right-continuous functions, we obtained the Isotonized Inverse Bayes Posterior (IIP) and we verified that it allows for a Bernstein-von Mises theorem with minimax asymptotic variance, reflecting the Hölder continuity of the true distribution. This proves that our methodology gives asymptotically correct uncertainty quantification under the frequentist point of view and it highlights its potential for Bayesian nonparametrics in inverse problems.

Regarding potential future work, several interesting directions emerge. First, it would be important to verify whether the bootstrap procedure of [92] remains valid for all degrees of smoothness of the underlying truth, not just for F_0 differentiable at x. Second, an open question is whether a Bernstein-von Mises (BvM) result can be obtained for Wicksell's problem within the classical Bayesian framework that is, by placing a Dirichlet Process (DP) prior F on F_0. Formally obtaining a nonparametric hierarchical Bayesian model for the observations $Z_1, \ldots, Z_n$. In the following section we will illustrate the obtained results with this hierarchical Bayesian approach.

We conclude by remarking that the work of the present chapter introduces a fresh perspective on Bayesian inverse problems by employing the concept of the *projection posterior*, first proposed in [17]. The methods proposed here are also significantly computationally faster if compared to the standard hierarchical Bayes model illustrated above, because they do not rely on computationally expensive Gibbs samplers. Finally, a valuable direction for future research is to identify other problems where this approach could be effectively utilized.

3.2 Posterior Contraction via Dirichlet-Beta Hierarchical Modelling

3.2.1 Introduction

We closed the previous section by mentioning that one of the most interesting directions for future research is the study of Wicksell's problem within the classical Bayesian framework that is, by placing a Dirichlet Process (DP) prior F on F_0. In particular, obtaining a Bernstein-von Mises result in this setting, remains an open question. However, in this section we show consistency and posterior contraction results obtained for this modelling approach.

We perform statistical inference in the setting of Wicksell's problem using the following nonparametric hierarchical Bayesian model for the observations $Z_1, \ldots, Z_n$:

(i) A probability distribution F on $\mathbb{R}_+$ is generated from the Dirichlet Process prior $\mathrm{DP}(\alpha)$.

(ii) An i.i.d. sample $X_1, \ldots, X_n$ is generated from F.

(iii) An i.i.d. sample $Y_1, \ldots, Y_n$ is generated from a density $\psi(y) = \frac{1}{2\sqrt{1-y}}$, $y \in (0,1)$. of a beta distribution $\mathrm{Be}(1, 1/2)$, independent of the other samples $X_1, \ldots, X_n$.

(iv) The observations are $Z_i = Y_i \cdot X_i$, for $i = 1, \ldots, n$, with density as in (3.1.1).

In the above model, the posterior is a Mixture of Dirichlet Processes (MDP), with distribution and mean, for any measurable A:

$$F \mid Z_1, \ldots, Z_n \sim \int \mathrm{DP}\Big(\alpha + \sum_{i=1}^{n} \delta_{X_i}\Big)\, d\mathrm{P}(X_1, \ldots, X_n \mid Z_1, \ldots, Z_n), \tag{3.2.1}$$

$$\mathrm{E}[F(A) \mid Z_1, \ldots, Z_n] = \tag{3.2.2}$$

$$= \int \Big(\frac{|\alpha|}{|\alpha| + n}\overline{\alpha}(A) + \frac{1}{|\alpha| + n}\sum_{i=1}^{n} \delta_{X_i}(A)\Big)\, d\mathrm{P}(X_1, \ldots, X_n \mid Z_1, \ldots, Z_n),$$

where (3.2.1) follows directly from Corollary 3.1 in [3] and (3.2.2) follows from (3.2.1), Fubini's theorem and Proposition 4.2 in [42].

3.2.2 Main results

Given two probability distributions G and G_0 with densities g and g_0, the Kullback-Leiber discrepancy and corresponding variance are denoted by

$$K(g_0, g) := \int \log(g_0/g)\, dG_0, \quad K_2(g_0, g) := \int (\log(g_0/g))^2\, dG_0.$$

We write $g_0 \in \mathrm{KL}(\Pi)$ if, with positive prior probability, $\forall\, \varepsilon > 0$:

$$K(g_0; g) < \varepsilon.$$

We will consider three notions of recovering the truth: the bounded Lipschitz distance d_{BL} on probability distributions on $\mathbb{R}$, the Hellinger metric and the Wasserstein metric. The d_{BL}

distance metrizes the topology of weak convergence on the space of probability measures over a separable metric space and it is formally defined as:

$$d_{BL}(G, G_0) := \sup_{f \in BL_1} \left| \int f dG - \int f dG_0 \right|, \tag{3.2.3}$$

The Hellinger distance on the space of densities is given by

$$h(g_0, g) := \left(\int \left(g_0^{1/2}(z) - g^{1/2}(z) \right)^2 dz \right)^{1/2}$$

We are going to use several times that:

$$h^2(g_0, g) \leq \|g_0 - g\|_1 \leq 2h(g_0, g), \tag{3.2.4}$$

The third notion of recovery consists in measuring recovery of the underlying truth via the Wasserstein metric

$$W_k(G, G_0) := \inf_{\pi \in \Pi^*(G, G_0)} \left(\int |x - y|^k d\pi(x, y) \right)^{1/k},$$

where $\Pi^*(G, G_0)$ is the collection of all couplings π of G and G_0 into a bivariate measure with marginals G and G_0 (i.e. if $(x, y) \sim \pi$, then $x \sim G$ and $y \sim G_0$), and $k \geq 1$.

Consistency

We assume that the distribution F is supported inside a finite interval $[0, L]$. In practice, the endpoint will be known (e.g. the maximal length of the studied surface) and hence we can choose a base measure of the $\mathrm{DP}(\alpha)$ prior to have support at least $[0, L]$. In short:

$$\mathrm{supp}(F_0) \subseteq [0, L] \subset \mathrm{supp}(\alpha) \quad \text{where:} \quad F \sim \mathrm{DP}(\alpha), \tag{3.2.5}$$

where L is not an atom of F_0.

Theorem 3.2.1 (Consistency)**.** *Let $F \sim \Pi = \mathrm{DP}(\alpha)$ be the Dirichlet process prior and g_F as in* (3.1.1)*. Suppose that $\mathrm{E}_0 \left| \log g_0(Z) \right| \vee \mathrm{E}_0 \left| \log F_0(\max(Z, 1), L) \right| < \infty$ and that* (3.2.5) *holds true, then the following assertions are true:*

a. $g_0 \in \mathrm{KL}(\Pi)$ and the posterior distribution is consistent relative to the weak topology at g_0, i.e. $\forall\, \varepsilon > 0$ as $n \to \infty$:

$$\Pi_n(F : d_{BL}(G_F, G_0) > \varepsilon \mid Z_1, \ldots, Z_n) \to 0, \qquad G_0^n \text{ a.s.}$$

b. If $\overline{\alpha}([0, n^{-\beta}]) \leq e^{-Cn}$ for some $C > 0$ and $\beta > 0$, then the posterior is consistent relative to the $\mathbb{L}_1$ norm at g_0, i.e. $\forall\, \varepsilon > 0$ as $n \to \infty$:

$$\Pi_n(F : \|g_F - g_0\|_1 > \varepsilon \mid Z_1, \ldots, Z_n) \to 0, \qquad G_0^n \text{ a.s.}$$

c. If the condition of b. holds, then the posterior is consistent relative to the weak topology at F_0, i.e. $\forall\, \varepsilon > 0$ as $n \to \infty$:

$$\Pi_n(F : d_{BL}(F, F_0) > \varepsilon \mid Z_1, \ldots, Z_n) \to 0, \qquad G_0^n \text{ a.s.}$$

d. If the condition of b. holds, then the posterior is consistent relative to the weak topology at F_0^*, i.e. $\forall\, \varepsilon > 0$ as $n \to \infty$:

$$\Pi_n\left(F : d_{BL}\left(F^*, F_0^*\right) > \varepsilon \mid Z_1, \ldots, Z_n\right) \to 0, \qquad G_0^n \text{ a.s.}$$

The proof of Theorem 3.2.1 is deferred to Section 3.3.2.

Contraction rates

In this section we will focus on the contraction rate of the three fundamental functionals for the application in material sciences: g_0, V_0 and F_0^*. However we will use three different notions of recovery for the three different functionals: for the first one in Hellinger distance, for the second one pointwise convergence at a fixed point $x \geq 0$ and for the third one the Wasserstein metric. The reason for this choice is that the first one will help to prove the second one, and the first and the second will help to prove the third one. Again for mathematical convenience, we are going to make use of a slightly more stringent assumption than (3.2.5), i.e. for $0 < \delta < L$:

$$\operatorname{supp}(F_0) \subseteq [\delta, L] \subset \operatorname{supp}(\alpha) \quad \text{where:} \quad F \sim \mathrm{DP}(\alpha). \tag{3.2.6}$$

Theorem 3.2.2 (Contraction rates)**.** *Let g_F as in* (3.1.1) *and $F \sim \Pi = \mathrm{DP}(\alpha)$ be the Dirichlet process prior with center measure α that has a Lebesgue density bounded away from 0 and ∞ on its bounded support that satisfies* (3.2.6)*. Further, let F_0 have density f_0 bounded away from zero and infinity, $\|1/g_0\|_1 < \infty$ and $Z_1, \ldots, Z_n \overset{i.i.d.}{\sim} g_0$. Then:*

a. there exists a constant M such that, in G_0-probability:

$$\Pi_n\Big(F : h(g_F, g_0) \geq M n^{-1/4} (\log n)^{1/4} \mid Z_1, \ldots, Z_n\Big) \to 0,$$

b. for every $x \geq 0$, there exists a constant M such that, in G_0-probability:

$$\Pi_n\left(F : |V_F(x) - V_0(x)| \geq M n^{-1/4} (\log n)^{3/4} \mid Z_1, \ldots, Z_n\right) \to 0.$$

c. for every $k \geq 1$, there exists a constant M such that, in G_0-probability:

$$\Pi_n\left(F : W_k\left(F^*, F_0^*\right) \geq M n^{-1/(2k+4)} (\log n)^{\frac{4k+3}{2k+4}} \mid Z_1, \ldots, Z_n\right) \to 0.$$

The proof of Theorem 3.2.2 is deferred to Section 3.3.2. The contraction rate $n^{-1/4}$ for the Hellinger distance is equal to the $\mathbb{L}_2$-minimax rate for estimating a density in a Sobolev space of order $1/2$ (see [32]). Because the densities g_0 are in $H^{1/2}[0, L]$ if F_0 possesses a bounded density, this suggests that the rate $n^{-1/4}$ is close to optimal. The rate with respect to the Wasserstein distance remains unknown. The pointwise contraction rate for the functional V_0 follows from the contraction rate in Hellinger distance, combined with a proof technique based on a local perturbation. This perturbation, which is believed to correspond to the efficient influence function for the functional of interest, is essential for handling arguments involving $1/g_F(z)$, which is unbounded as $z \to L$. However, this localization comes at the cost of contracting at the Hellinger distance rate. Based on the findings of [45] and the results of our simulation study, we believe that this rate is suboptimal and that a Bernstein-von Mises phenomenon may hold.

3.2.3 Simulation study

In this section, we present simulations to illustrate the practical behavior of the presented methodology. We compare it also with the Inverse Isotonic Estimator (IIE) studied in [43]. We consider two distinct settings. The first is a *synthetic* setting where the true underlying distribution function F_0 is known. The first simulation setting serves as a valuable benchmark for assessing the quality of our methodology. The second is a *real-data* application involving

the positions of stars in the Globular Cluster Messier 62. In the second case, the ground truth distribution of the stars is unknown.



Figure 3.4: Comparison for $n = 200$ isotonic estimator and bayesian Dirichlet-based estimator.



Figure 3.5: Comparison for $n = 112$ and $n = 450$ projected star positions between isotonic estimator and bayesian Dirichlet-based estimator.



$n = 200$ $n = 1000$

Figure 3.6: Analysis normality.

In Figure 3.4 and 3.6, we analyze the setting where the underlying cdf F_0 is exponential with rate 1.2. Samples $Z_1, \ldots, Z_n$ used to compute the posterior are generated hierarchically: $X_1, \ldots, X_n \sim F_0$, $Y_1, \ldots, Y_n \sim \mathrm{Be}(1, 1/2)$, with $Z_i = X_i \cdot Y_i$ for all i. The plots use the following color scheme: green for prior draws $F \sim \mathrm{DP}(M\alpha)$ (parameters M and α specified in the plots); purple for draws from the posterior; red for the average of these draws (approximating the posterior mean); yellow for the Isotonic Inverse Estimator (IIE) of [43]; and orange for F_0.

In Figure 3.5, we apply the methodology to star position data from globular cluster Messier 4. Let $\mathbf{Z} = (Z_{p_1}, Z_{p_2}, Z_{p_3})$ represent a star's three-dimensional position, where only (Z_{p_1}, Z_{p_2}) are observed. Here, $X = Z_{p_1}^2 + Z_{p_2}^2 + Z_{p_3}^2 \sim F_0$, and $Z = Z_{p_1}^2 + Z_{p_2}^2$. Beyond serving as a

compelling application of our methodology, this example highlights that also in real data applications the IIE of [43] behaves very closely to the present methodology.

The first column of Figure 3.6 shows that with $n = 200$. The average of the posterior appears to be somewhat smoother, which will typically be seen as an advantage. By increasing the sample size to $n = 1000$ the uncertainty significantly reduces. The two bottom rows examine normality: histograms, QQ-plots, and statistical tests strongly confirm normality of the gathered samples. Furthermore, the Bayesian-estimated standard deviation aligns closely with the theoretical value attained by the frequentist procedure, suggesting that a Bernstein-von Mises phenomenon is satisfied. In all the plots we employ 300 posterior draws.

3.2.4 Conclusion

In this section we studied Wicksell's problem within the classical nonparametric Bayesian framework, placing a Dirichlet Process prior directly on F_0. We established consistency of the posterior in several metrics (Theorem 3.2.1) and derived contraction rates for the key functionals g_0, V_0 and F_0^* (Theorem 3.2.2).

The contraction rate $n^{-1/4}$ for the Hellinger distance equals the $\mathbb{L}_2$-minimax rate for estimating a density in a Sobolev space of order $1/2$ (see [32]). Since $g_0 \in H^{1/2}[0,\infty)$ whenever F_0 possesses a bounded density, this suggests that $n^{-1/4}$ is close to optimal. The pointwise contraction rate for V_0 follows from the Hellinger contraction rate, combined with a Laplace transform rewriting of the posterior as in equation (2.13) of [16], a local perturbation and a LAN expansion as in [43]. This perturbation, believed to correspond to the efficient influence function for the functional of interest, is essential for controlling terms involving $1/g_F(z)$, which is unbounded as $z \to L$. However, the local nature of the perturbation in Wicksell's problem comes at the cost of contracting at the Hellinger rate h rather than at the faster rate $h^2 = O_p(\sqrt{\log n}/\sqrt{n})$ that arises in the proof style of [16]. Despite significant efforts, overcoming this limitation remains an open problem.

The question of whether a Bernstein-von Mises theorem holds in this setting is also open. The simulation study presents encouraging evidence: the posterior concentrates around the truth at the expected frequentist rate, and normality of the centered and rescaled posterior draws is strongly supported by histograms, QQ-plots and formal tests. Moreover, the Bayesian standard deviation closely matches the theoretical frequentist value, suggesting that efficiency is attained. Establishing this rigorously, however, remains a challenging open problem.

From a practical standpoint, the hierarchical Bayesian approach presented here is considerably more demanding to implement than the projection-based method of the previous section. The latter requires no Markov Chain Monte Carlo: isotonized posterior draws are obtained analytically via the Greatest Convex Majorant, making computation fast and straightforward. The present approach, by contrast, requires MCMC sampling from the Mixture of Dirichlet Processes posterior (3.2.1), which is computationally slow and harder to tune. Unless the full posterior distribution over F is of interest, the projection approach is therefore preferable in practice.

3.3 Appendix

Measure-theoretical framework: in this chapter, the sample space is $(\mathbb{R}^+, \mathscr{B}(\mathbb{R}^+))$, and $\mathfrak{M}$ (with Borel σ-field $\mathscr{M}$ for the weak topology) is the collection of all Borel probability measures on $(\mathbb{R}^+, \mathscr{B}(\mathbb{R}^+))$. The Dirichlet process prior is a probability measure on $\mathfrak{M}$. A prior on the set of probability measures $\mathfrak{M}$ is a probability measure on a sigma-field $\mathscr{M}$ of subsets of $\mathfrak{M}$. Alternatively, it can be viewed as map from some probability space $(\Omega, \mathscr{U}, \Pr)$ into $(\mathfrak{M}, \mathscr{M})$. We can think of the hierarchy of the spaces as follows: $(\Omega, \mathscr{U}, \Pr) \to (\mathfrak{M}, \mathscr{M}) \to (\mathbb{R}^+, \mathscr{B}(\mathbb{R}^+))$. We put a Dirichlet process prior over G and write $G \sim \Pi = \mathrm{DP}(\alpha)$ for some base measure α and prior precision $|\alpha|$. This means that for any measurable set $B \in \mathscr{M}$ we have $\Pi(B) = \Pr(G \in B)$. Throughout the chapter, we use the notations P and E, which should be understood as the outer probability measure and outer expectation, respectively, whenever issues of measurability in the considered processes arise (see [103] for a rigorous treatment of these topics).

Furthermore, we use the notion of conditional convergence in distribution. This happens if the bounded Lipschitz distance between the conditional law of Borel measurable X_n given $\mathcal{B}_n$ and a Borel probability measure L tends to zero, where the convergence can be in outer probability. The sequence X_n tends to L conditionally given $\mathcal{B}_n$ in outer probability if

$$d_{BL}\left(\mathcal{L}\left(X_n \mid \mathcal{B}_n\right), L\right) := \sup_{f \in BL_1} \left| \mathrm{E}\left(f\left(X_n\right) \mid \mathcal{B}_n\right) - \int f dL \right| \xrightarrow{\mathrm{P}} 0, \tag{3.3.1}$$

where BL_1 is the space of bounded Lipschitz functions with Lipschitz constant 1.

In particular if $Z_1, \ldots, Z_n \overset{\text{i.i.d.}}{\sim} \mathrm{P}_Z$, then $X_n \mid Z_1, \ldots, Z_n \rightsquigarrow N(0, \sigma^2)$ means:

$$d_{BL}\left(\mathrm{P}(X_n \in \cdot \mid Z_1, \ldots, Z_n), N(0, \sigma^2)\right) \xrightarrow{\mathrm{P}_Z} 0. \tag{3.3.2}$$

3.3.1 Proofs of complemental statements for Semiparametric BvM

Lemma 3.3.1.

$$F_0(x) = 1 - \frac{2}{\pi}\sqrt{x}\, V_0(x) - \frac{2}{\pi}\int_x^\infty \frac{V_0(s)}{2\sqrt{s}}\, ds, \tag{3.3.3}$$

where:

$$V_0(x) := \int_x^\infty \frac{g_0(z)}{\sqrt{z - x}}\, dz. \tag{3.3.4}$$

Proof. We will prove (3.3.3) using the materials science application setting, which is the same that Wicksell [115] considered. In this way we will use his result. However the same formula can be obtained with similar reasonings in the astronomical application setting (c.f. eq. (1.3) and proof of Proposition 2.2 in [92]).

Recall that the embedded particle squared radii are i.i.d. realizations from a random variable $X^* \sim F_0^*$. However, when sampling, two opposing effects shape the model. First, a circular cross-section cannot exceed the radius of its parent sphere. Second, larger spheres are more likely to be intersected—sampling is proportional to radius. As a result, the observed distribution is not F_0^* but its size-biased version:

$$F_0(x) = \frac{1}{m_F}\int_0^x \sqrt{v}\, dF_0^*(v), \tag{3.3.5}$$

where $0 < m_F = \int_0^\infty \sqrt{y}\, dF_0^*(y) < \infty$ is the expected radius under F^*.

By plugging this formula into (3.1.1) we obtain the equivalent formula for g_0 as in [43]:

$$g_0(z) = \frac{1}{2m_F} \int_z^\infty \frac{dF_0^*(x)}{\sqrt{x-z}}.$$

Wicksell [115] inverted this equation, by recognising an Abel-type integral, and found an expression for F_0^* in terms of g_0:

$$F_0^*(x) = 1 - \frac{\int_x^\infty (z-x)^{-1/2} g_0(z)\, dz}{\int_0^\infty z^{-1/2} g_0(z)\, dz}, \quad x > 0 \tag{3.3.6}$$

and $F_0^*(x) = 0$ for all $x \le 0$. Using (3.3.5), by defining $c_F = \int_0^\infty z^{-1/2} g_0(z)\, dz$, and integrating by parts we obtain:

$$\begin{aligned} F_0(x) \cdot m_F &= \sqrt{v} F_0^*(v)\Big|_0^x - \int_0^x F_0^*(v) \frac{1}{2\sqrt{v}}\, dv = \sqrt{x} F_0^*(x) - \int_0^x F_0^*(v) \frac{1}{2\sqrt{v}}\, dv \\ &= \sqrt{x}\left(1 - \frac{\int_x^\infty (z-x)^{-1/2} g_0(z) dz}{\int_0^\infty z^{-1/2} g_0(z) dz}\right) - \int_0^x \left(1 - \frac{\int_v^\infty (z-v)^{-1/2} g_0(z) dz}{\int_0^\infty z^{-1/2} g_0(z) dz}\right) \frac{1}{2\sqrt{v}}\, dv \\ &= -\sqrt{x} \frac{1}{c_F} \int_x^\infty (z-x)^{-1/2} g_0(z)\, dz + \frac{1}{c_F} \int_0^x \int_v^\infty (z-v)^{-1/2} g_0(z)\, dz\, \frac{1}{2\sqrt{v}}\, dv, \end{aligned}$$

since: $\frac{2m_F}{\pi} = (\int_0^\infty \frac{g_0(z)}{\sqrt{z}} dz)^{-1} \Rightarrow \frac{2m_F}{\pi} = \frac{1}{c_F}$, therefore: $m_F c_F = \frac{\pi}{2}$, and $V_0(x) = \int_x^\infty (z-x)^{-1/2} g_0(z)\, dz$ we obtain:

$$F_0(x) = -\frac{2}{\pi} \sqrt{x} V_0(x) + \frac{2}{\pi} \int_0^x \frac{V_0(v)}{2\sqrt{v}}\, dv$$

because the density g_0 integrates to 1 (c.f. [52] p. 1519), it follows by integration by parts:

$$1 = -\frac{2}{\pi} \int_0^\infty \sqrt{u}\, dV_0(u) = \frac{2}{\pi} \int_0^\infty \frac{V_0(u)}{2\sqrt{u}}\, du = \frac{2}{\pi} \int_0^x \frac{V_0(v)}{2\sqrt{v}}\, dv + \frac{2}{\pi} \int_x^\infty \frac{V_0(v)}{2\sqrt{v}}\, dv,$$

thus the claim in (3.3.3) follows. □

Lemma 3.3.2. (3.1.25) *implies* (3.1.22).

Proof. By Theorem 1.13.1 (i) and (ii) in [103] the claim (3.1.22) is obtained as soon as we show (c.f. (3.3.1)), for $X \sim N\left(0, \frac{g_0(x)}{2\gamma}\right)$:

$$\sup_{f \in BL_1} \left| \mathrm{E}\left(f\Big(\frac{\sqrt{n}}{\sqrt{\log n}} (\hat{V}_G(x) - \hat{V}_n(x)) \Big) \mid Z_1, \ldots, Z_n \right) - \mathrm{E}\left(f(X)\right) \right| \xrightarrow{G_0} 0.$$

By Lemma 2.1 given in the Online Supplement [46], for any bounded 1-Lipschitz function $f : \mathbb{R} \mapsto \mathbb{R}$ and $Z_n^{G_0}$ as in (3.1.23):

$$\left| \mathrm{E}\Big[f\Big(\frac{\sqrt{n}}{\sqrt{\log n}} (\hat{V}_G(x) - \hat{V}_n(x)) \Big) - f\Big(\frac{\sqrt{n}}{\sqrt{\log n}} (\hat{V}_G(x) - V_0(x)) - Z_n^{G_0}(1) \Big) \mid Z_1, \ldots, Z_n \Big] \right|$$

$$\leq \left| \frac{\sqrt{n}}{\sqrt{\log n}} \left(\hat{V}_n(x) - V_0(x) \right) - Z_n^{G_0}(1) \right| \xrightarrow{G_0} 0.$$

This means that the claim is obtained as soon as we show:

$$\sup_{f \in BL_1} \left| \mathrm{E} \left(f\Big(\frac{\sqrt{n}}{\sqrt{\log n}} (\hat{V}_G(x) - V_0(x)) - Z_n^{G_0}(1) \Big) \mid Z_1, \ldots, Z_n \right) - \mathrm{E}\left(f(X)\right) \right| \xrightarrow{G_0} 0.$$

By Theorem 1.13.1 (iii) in [103] the last display is implied by (3.1.25). □

Proof of (3.1.31). First note that by assumption Qf is a.s. well-defined (c.f. Remark 4.4 in [42]). Moreover note that $\mathrm{E}V_n = |\alpha|/(|\alpha| + n)$ and $|f_n^t(Z)| \leq 2\sqrt{|t|/\delta_n^*}$ and $\forall\, \varepsilon > 0$, by independence, that:

$$\begin{aligned} \mathrm{P}\left(\delta_n^{-1} \left| V_n(Q - \mathbb{G}_n) f_t^n \right| > \varepsilon \mid Z_1, \ldots, Z_n \right) &\leq \delta_n^{-1} \varepsilon^{-1} \mathrm{E}V_n \, \mathrm{E}\left[(Q + \mathbb{G}_n)|f_t^n| \mid Z_1, \ldots, Z_n \right] \\ &\leq 4\delta_n^{-1}\varepsilon^{-1} \frac{|\alpha|}{|\alpha| + n} \frac{\sqrt{|t|}}{\sqrt{\delta_n^*}} = O\left(\delta_n^{-1} n^{-1} (\delta_n^*)^{-1/2} \right). \end{aligned}$$

As $\gamma > \frac{1}{2}$, note that $\delta_n^{-1} n^{-1} (\delta_n^*)^{-1/2} = n^{-1}(n/\log n)^{\frac{1}{2} + \frac{1}{4\gamma}} \ll n^{-1}(\sqrt{n}/\sqrt{\log n})^2 \to 0$. Similarly: $\mathrm{P}\left(\delta_n^{-1} \left| V_n(\mathbb{B}_n - \mathbb{G}_n) f_t^n \right| > \varepsilon \mid Z_1, \ldots, Z_n \right) \overset{G_0}{\to} 0$.

By Proposition G.2 in [42] for $\varepsilon_i \overset{\text{i.i.d.}}{\sim} \mathrm{Exp}(1)$ and $W_{ni} := \varepsilon_i / \bar{\varepsilon}_n$, we have $(W_{n1}, \ldots, W_{nn}) \sim \mathrm{Dir}(n; 1/n, \ldots, 1/n)$. Since $\mathbb{B}_n \sim \mathrm{DP}(n\mathbb{G}_n)$, for δ_{Z_i} Dirac measure at Z_i, we have the following representation in distribution of $\mathbb{B}_n$ (c.f. Example 3.7.9 in [103]):

$$\mathbb{B}_n = \frac{1}{n} \sum_{i=1}^{n} W_{ni} \delta_{Z_i}.$$

Because of this representation (c.f. section 3.7.2. in [103]), $\mathbb{B}_n$ is also known as *Bayesian bootstrap*. Conclude that:

$$\frac{\sqrt{n}}{\sqrt{\log n}} (\mathbb{B}_n - \mathbb{G}_n) f_t^n = \frac{1}{\sqrt{n \log n}} \sum_{i=1}^{n} (W_{ni} - 1)\, f_t^n(Z_i).$$

Because $|\bar{\varepsilon}_n - 1| = O_p(1/\sqrt{n})$, for almost every sequence $Z_1, \ldots, Z_n$ conclude that, uniformly in t in compacta:

$$\frac{1}{\sqrt{n \log n}} \sum_{i=1}^{n} (W_{ni} - 1)\, f_t^n(Z_i) = \frac{1}{\sqrt{n \log n}} \sum_{i=1}^{n} (\varepsilon_i - 1)\, f_t^n(Z_i) + o_p(1), \tag{3.3.7}$$

conclude by recalling the definition of $\mathbb{Z}_n^*$ in (3.1.32). □

Lemma 3.3.3. *For fixed $x \geq 0$, $\gamma > \frac{1}{2}$ which satisfies* (3.1.20)*, let $\hat{V}_n$ be the Inverse Isotonic Estimator (IIE) (see also Theorem 2 in [43]). Then for $n \to \infty$:*

$$\frac{\sqrt{n}}{\sqrt{\log n}} \left(\hat{V}_n(x) - V_0(x) \right) = Z_n^{G_0}(1) + o_{G_0}(1).$$

where $Z_n^{G_0}(1)$ *is given, for sequences* $\delta_n = \sqrt{\log n}/\sqrt{n}$ *and* $\delta_n^* = \delta_n^{1/\gamma}$ *where* $\gamma > 1/2$ *satisfies condition* (3.1.20)*,*

$$Z_n^{G_0}(t) := \delta_n^{-1}(\delta_n^*)^{-1}\left\{U_n(x+\delta_n^* t) - U_n(x) - U_0(x+\delta_n^* t) + U_0(x)\right\} \tag{3.3.8}$$

$$= 2\delta_n^{-1}(\delta_n^*)^{-1}\int\left\{\sqrt{(z-x)_+} - \sqrt{(z-x-\delta_n^* t)_+}\right\} d\left(\mathbb{G}_n - G_0\right)(z). \tag{3.3.9}$$

Proof. Fix $x \geq 0$ such that the conditions stated above are satisfied. Define, for U_n as above and $a > 0$, the process T_n as follows:

$$T_n(a) := \inf\left\{t \geq 0 : U_n(t) - at \text{ is maximal }\right\}.$$

The relation between T_n and $\hat{V}_n$ is given by the switch relation (Lemma 3.2, [53]):

$$T_n(a) \leq x \quad \Leftrightarrow \quad \hat{V}_n(x) \leq a. \tag{3.3.10}$$

From this relation, it follows that for all n sufficiently large:

$$\delta_n^{-1}\left(\hat{V}_n(x) - V_0(x)\right) \leq a + Z_n^{G_0}(1) \;\Leftrightarrow\; (\delta_n^*)^{-1}\left(T_n\left(a_0 + \delta_n\left(a + Z_n^{G_0}(1)\right)\right) - x\right) \leq 0.$$

We aim to study the probability of the event on the left-hand side. We write a_0 for $V_0(x)$, which is fixed. Using the same steps as in the proof of Theorem 2 in [43] we obtain:

$$\begin{aligned}&(\delta_n^*)^{-1}\left(T_n\left(a_0 + \delta_n\left(a + Z_n^{G_0}(1)\right)\right) - x\right)\\ &\quad = \inf\left\{t \geq -(\delta_n^*)^{-1}x : \tilde{Z}_n(t) - \left(a + Z_n^{G_0}(1)\right)t \text{ is max}\right\},\end{aligned}$$

where:

$$\tilde{Z}_n(t) - at := \delta_n^{-1}(\delta_n^*)^{-1}\left\{U_n(x+\delta_n^* t) - U_n(x) - U_0(x+\delta_n^* t) + U_0(x)\right\} + \tag{3.3.11}$$

$$+ \delta_n^{-1}(\delta_n^*)^{-1}\left\{U_0(x+\delta_n^* t) - U_0(x) - V_0(x)\delta_n^* t\right\} - at. \tag{3.3.12}$$

Again by the proof of Theorem 2 in [43] we have:

$$\tilde{Z}_n(t) = Z_n^{G_0}(t) - |t|^{\gamma+1}(K + o(1)), \tag{3.3.13}$$

moreover: $Z_n^{G_0}(t) - tZ_n^{G_0}(1) \xrightarrow{G_0} 0$. From these facts, we conclude by applying Theorem 2.7 from [65] to the process $\tilde{Z}_n$:

$$(\delta_n^*)^{-1}\left(T_n\left(a_0 + \delta_n\left(a + Z_n^{G_0}(1)\right)\right) - x\right) \rightsquigarrow \underset{t\in I_x}{\arg\max}\left\{-|t|^{\gamma+1}K - at\right\}. \tag{3.3.14}$$

Since:

$$\begin{cases} \underset{t\in I_x}{\arg\max}\left\{-|t|^{\gamma+1}K - at\right\} < 0 & \text{if } a > 0,\\ \underset{t\in I_x}{\arg\max}\left\{-|t|^{\gamma+1}K - at\right\} > 0 & \text{if } a < 0.\end{cases}$$

We conclude then:

$$\begin{aligned}&\mathrm{P}\left\{\delta_n^{-1}\left(\hat{V}_n(x) - V_0(x)\right) - Z_n^{G_0}(1) \leq a\right\}\\ &\qquad \overset{(3.3.14)}{\longrightarrow} \mathrm{P}\left\{\underset{t\in I_x}{\arg\max}\left\{-|t|^{\gamma+1}K - at\right\} \leq 0\right\} = \begin{cases} 1 & \text{if } a > 0,\\ 0 & \text{if } a < 0,\end{cases}\end{aligned}$$

which gives the claim. □

Lemma 3.3.4. *If* $g_0(x) = \int_x^\infty (4s(s-x))^{-1/2}\, dF_0(s) < \infty$, *then as* $t \downarrow 0$:

$$\mathrm{E}\left\{\frac{1}{Z-x}\mathbf{1}_{\{Z\geq x+t\}}\right\} \sim \log\left(\frac{1}{t}\right) g_0(x), \tag{3.3.15}$$

where $\sim$ *means that the ratio of the left-hand side and of the right-hand side goes to 1 as* $t \downarrow 0$. *Moreover let* $Z \perp\!\!\!\perp \Lambda \sim \mathrm{Exp}(1)$, *then it is also true:*

$$\mathrm{E}\left\{\frac{(\Lambda-1)^2}{Z-x}\mathbf{1}_{\{Z\geq x+t(\Lambda-1)^2\}}\right\} \sim \log\left(\frac{1}{t}\right) g_0(x). \tag{3.3.16}$$

Proof. We start with (3.3.15). Consider independent $X \sim F_0$ and $V \sim \mathrm{Be}(1, 1/2)$, so that $Z \sim VX$. Consider $t \downarrow 0$, we study:

$$\begin{aligned}
&\mathrm{E}\left\{\frac{1}{Z-x}\mathbf{1}_{\{Z>x+t\}}\right\} = \mathrm{E}\left\{\frac{1}{VX-x}\mathbf{1}_{\{VX>x+t\}}\right\} = \iint \frac{1_{\{vs>x+t\}}}{vs-x}\frac{1_{v\in[0,1]}}{2\sqrt{1-v}}\, dv\, dF_0(s) \\
&= \mathrm{E}\mathbf{1}_{\{X>x+t\}} \int_{\frac{x+t}{X}}^1 \frac{1}{vX-x}\frac{1}{2\sqrt{1-v}} dv \overset{(\star)}{=} \mathrm{E}\left\{\int_{\frac{t}{X-x}}^1 \frac{1}{z}\frac{1}{2\sqrt{1-z}} dz \frac{\mathbf{1}_{\{X-x>t\}}}{\sqrt{X}\sqrt{X-x}}\right\}
\end{aligned}$$

where in $(\star)$ we changed variable by using $vX - x = z(X-x)$, which means $1 - v = \frac{X-x}{X}(1-z)$. Define now $X_x := X - x$ and $\mathcal{J}(t) := \int_t^1 \frac{1}{z}\frac{1}{\sqrt{1-z}} dz$. We will therefore show:

$$\mathrm{E}\left\{\mathcal{J}\left(\frac{t}{X_x}\right)\frac{1}{2\sqrt{X_x}\sqrt{X_x+x}}\mathbf{1}_{\{X_x>t\}}\right\} \sim \log\left(\frac{1}{t}\right)\mathrm{E}\left\{\frac{1}{2\sqrt{X_x}\sqrt{X_x+x}}\mathbf{1}_{\{X_x>0\}}\right\}$$

In order to show the above, we first prove as $t \downarrow 0$:

$$\mathcal{J}(t) = \int_t^1 \frac{1}{z}\frac{1}{\sqrt{1-z}} dz \sim \log\frac{1}{t}$$

and $\forall\, 0 < t < 1$, it holds $\mathcal{J}(t) \lesssim \log\left(\frac{1}{t}\right) + 1$.

For any $b \in [t, 1]$, write: $\mathcal{J}(t) = \int_t^b \frac{1}{z}\frac{1}{\sqrt{1-z}} dz + \int_b^1 \frac{1}{z}\frac{1}{\sqrt{1-z}} dz$. For $r > 1$:

$$\begin{aligned}
\int_b^1 \frac{1}{z}\frac{1}{\sqrt{1-z}} dz &\leq \left(\log\frac{1}{b}\right)^r \int_b^1 \frac{1}{z(\log\frac{1}{z})^r}\frac{1}{\sqrt{1-z}} dz \\
&\leq \left(\log\frac{1}{b}\right)^r \int_0^1 \frac{1}{z(\log\frac{1}{z})^r}\frac{1}{\sqrt{1-z}} dz \lesssim \left(\log\frac{1}{b}\right)^r.
\end{aligned}$$

Furthermore:

$$\left|\int_t^b \frac{1}{z}\frac{1}{\sqrt{1-z}} dz - \int_t^b \frac{1}{z} dz\right| \leq \int_t^b \frac{1}{z} dz \left(\frac{1}{\sqrt{1-b}} - 1\right) \lesssim \left(\log\frac{b}{t}\right)\frac{b}{(1-b)^{3/2}}.$$

Together the above gives:

$$\left|\frac{\int_t^1 \frac{1}{z}\frac{1}{\sqrt{1-z}} dz}{\log\frac{b}{t}} - 1\right| = \left|\frac{\int_t^b \frac{1}{z}\frac{1}{\sqrt{1-z}} dz + \int_b^1 \frac{1}{z}\frac{1}{\sqrt{1-z}} dz - \log\frac{b}{t}}{\log\frac{b}{t}}\right|$$

$$= \left| \frac{\int_t^b \frac{1}{z\sqrt{1-z}}dz - \int_t^b \frac{1}{z}dz + \int_b^1 \frac{1}{z\sqrt{1-z}}dz}{\log \frac{b}{t}} \right| \lesssim \frac{b}{(1-b)^{3/2}} + \left| \frac{1}{\log \frac{b}{t}} \right| \left| \log \frac{b}{t} \right|^r .$$

Take $b = 1/\sqrt{\log \frac{1}{t}}$ to see that

$$\left| \frac{\mathcal{J}(t)}{\log \frac{1}{t} - \frac{1}{2}\log\log \frac{1}{t}} - 1 \right| \lesssim \frac{1}{\sqrt{\log \frac{1}{t}}}(1 + o(1)) + \frac{\left(\log\log \frac{1}{t}\right)^r}{\log \frac{1}{t} - \frac{1}{2}\log\log \frac{1}{t}} \longrightarrow 0$$

This gives $\mathcal{J}(t) \sim \log \frac{1}{t}$. Also $\mathcal{J}(t) \lesssim \mathcal{J}(\varepsilon) \lesssim 1$, for $t \geq \varepsilon$, $\forall\, \varepsilon > 0$ and hence $\mathcal{J}(t) \lesssim \log \frac{1}{t} + 1$.

Now in order to prove the claim we need to be able to apply the Dominated Convergence Theorem to the following:

$$\frac{1}{\log \frac{1}{t} + 1} \mathrm{E}\left\{ \mathcal{J}\left(\frac{t}{X_x}\right) \frac{1}{2\sqrt{X_x}\sqrt{X_x + x}} \mathbf{1}_{\{X_x > t\}} \right\}$$

To this end note that since $\log \frac{X_x}{t} \leq \log X_x \mathbf{1}_{X_x > 1} + \log \frac{1}{t}$:

$$\frac{\mathcal{J}\left(\frac{t}{X_x}\right)\mathbf{1}_{X_x > t}}{\log \frac{1}{t} + 1} = \frac{\mathcal{J}\left(\frac{t}{X_x}\right)\mathbf{1}_{X_x > t}}{\log \frac{X_x}{t} + 1} \frac{\log \frac{X_x}{t} + 1}{\log \frac{1}{t} + 1} \leq \frac{\log \frac{X_x}{t} + 1}{\log \frac{1}{t} + 1} \mathbf{1}_{X_x > t} \leq \log X_x \mathbf{1}_{X_x > 1} + \mathbf{1}_{X_x > 0}$$

note that $\mathrm{E}\{\mathbf{1}_{X_x > 0} + (\log X_x \mathbf{1}_{X_x > 1})\} < \infty$, which means we can apply the Dominated Convergence Theorem upon noticing:

$$\frac{\mathcal{J}\left(\frac{t}{X_x}\right)}{\log \frac{1}{t} + 1} \mathbf{1}_{X_x > t} \xrightarrow{t \to 0} \mathbf{1}_{X_x > 0} \text{ a.s.}$$

We proceed now with (3.3.16). Recall $Z \perp\!\!\!\perp \Lambda \sim \mathrm{Exp}(1)$, $Z = XV$ and $X \perp\!\!\!\perp V \sim \mathrm{Be}(1, 1/2)$. Then for $t \downarrow 0$ we study:

$$\mathrm{E}\left[\mathbf{1}_{\{Z - x > (\Lambda-1)^2 t\}} \frac{(\Lambda-1)^2}{Z - x} \right] = \mathrm{E}\left[\mathbf{1}_{\{X > x + (\Lambda-1)^2 t\}} \int_{\frac{x + (\Lambda-1)^2 t}{X}}^1 \frac{(\Lambda-1)^2}{vX - x} \frac{1}{2\sqrt{1-v}} dv \right]$$

using $vX - x = z(X - x)$ and $1 - v = \frac{X-x}{X}(1 - z)$ the above display equals:

$$\mathrm{E}\left[\mathbf{1}_{\{X > x + (\Lambda-1)^2 t\}} \int_{\frac{(\Lambda-1)^2 t}{X-x}}^1 \frac{(\Lambda-1)^2}{z(X-x)} \frac{1}{2\sqrt{1-z}} \sqrt{\frac{X-x}{X}} \frac{X-x}{X} dz \right]$$
$$= \mathrm{E}\left[\mathbf{1}_{\{X > x + (\Lambda-1)^2 t\}} \mathcal{J}\left(\frac{(\Lambda-1)^2 t}{X - x} \right) \frac{(\Lambda-1)^2}{2\sqrt{X}\sqrt{X-x}} \right]$$

where again $\mathcal{J}(t) = \int_t^1 \frac{1}{z} \frac{1}{\sqrt{1-z}} dz$, which satisfies $\mathcal{J}(t) \sim \log \frac{1}{t}$ and $\mathcal{J}(t) \lesssim \log \frac{1}{t} + 1$ for $t \in (0, 1)$. Thus:

$$\frac{\mathcal{J}(\frac{(\Lambda-1)^2 t}{X-x})}{\log \frac{1}{t}} \frac{(\Lambda-1)^2}{2\sqrt{X}\sqrt{X-x}} \mathbf{1}_{\{X > x + (\Lambda-1)^2 t\}}$$

$$\sim \frac{\log\left(\frac{X-x}{(\Lambda-1)^2 t}\right)}{\log(\frac{1}{t})} \frac{(\Lambda-1)^2}{2\sqrt{X}\sqrt{X-x}} \mathbf{1}_{\{X>x+(\Lambda-1)^2 t\}} \xrightarrow{t\downarrow 0} \frac{(\Lambda-1)^2}{2\sqrt{X}\sqrt{(X-x)_+}} \text{ a.s.}$$

Moreover:

$$\begin{aligned}
&\left| \frac{\mathcal{J}(\frac{(\Lambda-1)^2 t}{X-x})}{\log\frac{1}{t}} \frac{(\Lambda-1)^2}{2\sqrt{X}\sqrt{X-x}} \mathbf{1}_{\{X>x+(\Lambda-1)^2 t\}} \right| \\
&\leq \frac{\left(\mathcal{J}(t) + \left(1 + \log\left(\frac{X-x}{(\Lambda-1)^2 t}\right) \mathbf{1}_{\{\frac{(\Lambda-1)^2}{X-x}<1\}}\right)\right)}{\log(\frac{1}{t})} \frac{(\Lambda-1)^2}{2\sqrt{X}\sqrt{X-x}} \mathbf{1}_{\{X>x+(\Lambda-1)^2 t\}} \\
&\lesssim \frac{(\Lambda-1)^2}{\sqrt{X}\sqrt{X-x}} \mathbf{1}_{\{X>x\}} + \sup_{u>1} \frac{\log u}{\sqrt{u}} \cdot \frac{|\Lambda-1|}{\sqrt{X}} \mathbf{1}_{\{X>x\}}
\end{aligned}$$

Conclude again by applying the Dominated Convergence Theorem. □

Lemma 3.3.5. *Let $\xi_{n1}, \ldots, \xi_{nn}$ be a triangular array of independent random variables that satisfy, $\sum_{i=1}^n \mathrm{E}|\xi_{ni}| = O(1)$, and $\forall\, \varepsilon > 0$, $\sum_{i=1}^n \mathrm{E}|\xi_{ni}| \mathbf{1}_{|\xi_{ni}|>\varepsilon} \to 0$. as $n \to \infty$. Then: $\sum_{i=1}^n (\xi_{ni} - \mathrm{E}\xi_{ni}) \xrightarrow{\mathrm{P}} 0$.*

Proof. Fix $\varepsilon > 0$

$$\mathrm{Var}\left(\sum_{i=1}^n \xi_{ni} \mathbf{1}_{|\xi_{ni}|\leq\varepsilon}\right) = \sum_{i=1}^n \mathrm{Var}\left(\xi_{ni} \mathbf{1}_{|\xi_{ni}|\leq\varepsilon}\right) \lesssim \sum_{i=1}^n \varepsilon\, \mathrm{E}|\xi_{ni}| \lesssim \varepsilon$$

So, $\forall\, \varepsilon_n \to 0$:

$$\sum_{i=1}^n (\xi_{ni} \mathbf{1}_{|\xi_{ni}|\leq\varepsilon_n} - \mathrm{E}\xi_{ni} \mathbf{1}_{|\xi_{ni}|\leq\varepsilon_n}) \xrightarrow{\mathrm{P}} 0.$$

Now note that $\forall\, \varepsilon > 0$,

$$\mathrm{E}\left(\left|\sum_{i=1}^n (\xi_{ni} - \mathrm{E}\xi_{ni}) - \sum_{i=1}^n (\xi_{ni} \mathbf{1}_{|\xi_{ni}|\leq\varepsilon} - \mathrm{E}\xi_{ni} \mathbf{1}_{|\xi_{ni}|\leq\varepsilon})\right|\right) \leq 2 \sum_{i=1}^n \mathrm{E}|\xi_{ni}| \mathbf{1}_{|\xi_{ni}|>\varepsilon} \to 0,$$

hence also for some $\varepsilon_n \to 0$. □

Lemma 3.3.6. *For $\varepsilon_i \overset{i.i.d.}{\sim} Exp(1)$ and $Z_i \overset{i.i.d.}{\sim} G_0$:*

$$\frac{1}{\sqrt{n \log n}} \sum_{i=1}^n \frac{(\varepsilon_i - 1)}{\sqrt{Z_i - x}} \mathbf{1}_{\{Z_i>x\}} \mid Z_1, \ldots, Z_n \rightsquigarrow N(0, g_0(x)) \quad \textit{under } G_0. \tag{3.3.17}$$

Proof. We apply Theorem 6.16 in [63] (with $\xi_{ni} := \frac{1}{\sqrt{n \log n}} \frac{\varepsilon_i - 1}{\sqrt{Z_i - x}} \mathbf{1}_{\{Z_i>x\}}$) conditionally on $Z_1, \ldots, Z_n$. Therefore we need to show:

$$\sum_{i=1}^n \mathrm{P}\left((\varepsilon_i - 1)^2 \geq \delta\, n \log n\, |Z_i - x|,\, Z_i > x \;\middle|\; Z_1, \ldots, Z_n\right) \xrightarrow{G_0} 0, \quad \forall\, \delta > 0, \tag{3.3.18}$$

$$\frac{1}{\sqrt{n \log n}} \sum_{i=1}^n \mathrm{E}\left(\frac{(\varepsilon_i - 1)}{\sqrt{Z_i - x}} \mathbf{1}_{\{(\varepsilon_i-1)^2 < n \log n\, |Z_i - x|,\, Z_i > x\}} \;\middle|\; Z_1, \ldots, Z_n\right) \xrightarrow{G_0} 0, \tag{3.3.19}$$

and:

$$\frac{1}{n\log n}\sum_{i=1}^{n}\mathrm{Var}\left(\frac{(\varepsilon_i-1)}{\sqrt{Z_i-x}}\mathbf{1}_{\{(\varepsilon_i-1)^2<n\log n\,|Z_i-x|,\,Z_i>x\}}\,\middle|\,Z_1,\ldots,Z_n\right)\overset{G_0}{\to} g_0(x). \tag{3.3.20}$$

First note that (3.3.18) and (3.3.19) are implied by convergence in mean (which means unconditional probability and expectation). So (3.3.18) is implied by:

$$\begin{aligned} n\,\mathrm{P}\left((\varepsilon-1)^2\geq\delta\,n\log n\,|Z-x|,\,Z>x\right) &= n\int_0^\infty\int_x^{x+\frac{(\lambda-1)^2}{\delta n\log n}} e^{-\lambda}g_0(z)\,dz\,d\lambda \\ &\leq n\,\|g_0\|_\infty\frac{1}{\delta n\log n}\int_0^\infty(\lambda-1)^2\,e^{-\lambda}\,d\lambda\to 0. \end{aligned}$$

Because $\mathrm{E}(\varepsilon_i-1)=0$, $\forall\, i$, the expected value of (3.3.19) goes to zero because:

$$\begin{aligned} &\mathrm{E}\left|\frac{1}{\sqrt{n\log n}}\sum_{i=1}^{n}\mathrm{E}\left(\frac{(\varepsilon_i-1)}{\sqrt{Z_i-x}}\mathbf{1}_{\{(\varepsilon_i-1)^2\geq n\log n\,|Z_i-x|,\,Z_i>x\}}\,\middle|\,Z_1,\ldots,Z_n\right)\right| \\ &\leq\frac{\sqrt{n}}{\sqrt{\log n}}\int_0^\infty\int_x^{x+\frac{(\lambda-1)^2}{n\log n}}\frac{|\lambda-1|}{\sqrt{z-x}}e^{-\lambda}g_0(z)\,dz\,d\lambda \\ &\leq\frac{\sqrt{n}}{\sqrt{\log n}}\|g_0\|_\infty\int_0^\infty\sqrt{\frac{(\lambda-1)^2}{n\log n}}|\lambda-1|e^{-\lambda}\,d\lambda\to 0. \end{aligned}$$

In order to prove (3.3.20) we proceed in steps. First note that (3.3.20) equals:

$$\frac{1}{n\log n}\sum_{i=1}^{n}\Bigg\{\mathrm{E}\left(\frac{(\varepsilon_i-1)^2}{Z_i-x}\mathbf{1}_{\{(\varepsilon_i-1)^2<n\log n\,|Z_i-x|,\,Z_i>x\}}\,\middle|\,Z_1,\ldots,Z_n\right) \tag{3.3.21}$$

$$-\left(\mathrm{E}\left(\frac{(\varepsilon_i-1)}{\sqrt{Z_i-x}}\mathbf{1}_{\{(\varepsilon_i-1)^2<n\log n\,|Z_i-x|,\,Z_i>x\}}\,\middle|\,Z_1,\ldots,Z_n\right)\right)^2\Bigg\}. \tag{3.3.22}$$

First we show the term in (3.3.22) goes to zero. Define:

$$\begin{aligned} &\mathcal{J}(u):=\mathrm{E}\,(\varepsilon-1)\,\mathbf{1}_{\{|\varepsilon-1|<u\}} \\ &=\begin{cases}\mathrm{E}\,(\varepsilon-1)\,\mathbf{1}_{\{|\varepsilon-1|>u\}}=\int_{1+u}^\infty(\lambda-1)e^{-\lambda}\,d\lambda\lesssim e^{-u/2},\ u>1\\ \int_{1-u}^{1+u}(\lambda-1)e^{-\lambda}\,d\lambda\leq\int_{1-u}^{1+u}(\lambda-1)\,d\lambda=u^2,\qquad 0\leq u\leq 1\end{cases}\lesssim u^2\wedge e^{-u/2},\quad\text{all } u. \end{aligned}$$

Therefore for (3.3.22) we have:

$$\begin{aligned} &\mathrm{E}\frac{1}{n\log n}\sum_{i=1}^{n}\left(\mathrm{E}\left(\frac{\varepsilon_i-1}{\sqrt{Z_i-x}}\mathbf{1}_{\{|\varepsilon_i-1|^2<n\log n|Z_i-x|,\,Z_i\geq x\}}\,\middle|\,Z_1,\ldots,Z_n\right)\right)^2 \\ &\leq\frac{1}{\log n}\mathrm{E}\left(\frac{\mathbf{1}_{Z>x}}{Z-x}\mathcal{J}^2\left(\sqrt{n\log n\,(Z-x)_+}\right)\right) \\ &\lesssim\frac{1}{\log n}\mathrm{E}\left(\frac{\mathbf{1}_{Z>x}}{Z-x}\left((Z-x)\,n\log n\wedge e^{-\sqrt{(Z-x)\,n\log n}/2}\right)^2\right) \end{aligned}$$

$$\leq \frac{1}{\log n}\mathrm{E}\left(\mathbf{1}_{Z>x}\left((Z-x)(n\log n)^2\mathbf{1}_{\{Z-x<\frac{\log\log n}{n\log n}\}}\right)+\frac{1}{Z-x}e^{-\sqrt{\log\log n}}\mathbf{1}_{\{Z-x>\frac{\log\log n}{n\log n}\}}\right)$$
$$\leq \frac{(n\log n)^2}{\log n}\int_x^{x+\frac{\log\log n}{n\log n}}(z-x)g_0(z)\,dz+\frac{e^{-\sqrt{\log\log n}}}{\log n}\mathrm{E}\frac{1}{Z-x}\mathbf{1}_{\{Z-x>\frac{\log\log n}{n\log n}\}}$$
$$\leq \|g_0\|_\infty\frac{1}{2}\left(\frac{\log\log n}{\log n}\right)^2+\frac{e^{-\sqrt{\log\log n}}}{\log n}\log\left(\frac{n\log n}{\log\log n}\right)(g_0(x)+O(1)).$$

where in the last line we used Lemma 3.3.4. Conclude that the last display goes to 0. Now we proceed with (3.3.21).

$$(3.3.21)=\frac{1}{n\log n}\sum_{i=1}^n\mathrm{E}\left(\frac{(\varepsilon_i-1)^2}{Z_i-x}\mathbf{1}_{\{|\varepsilon_i-1|^2<n\log n|Z_i-x|,\,Z_i>x\}}\,\middle|\,Z_1,\ldots,Z_n\right)$$
$$=\frac{1}{n\log n}\sum_{i=1}^n\frac{\mathbf{1}_{\{Z_i>x\}}}{Z_i-x}T(\sqrt{n\log n}\sqrt{Z_i-x})$$

where $T(u):=\mathrm{E}(\varepsilon-1)^2\mathbf{1}_{|\varepsilon-1|<u}\asymp u^3\wedge 1$. Define

$$\xi_{n,i}:=\frac{\mathbf{1}_{\{Z_i>x\}}}{Z_i-x}\cdot\frac{T(\sqrt{n\log n}\sqrt{Z_i-x})}{n\log n}$$

note $0\leq\xi_{n,i}\leq 1$, because $\frac{T(u)}{u^2}\leq 1$. Now we verify the conditions of Lemma 3.3.5 and:

$$\sum_{i=1}^n\mathrm{E}\xi_{ni}\to g_0(x). \tag{3.3.23}$$

First note that (3.3.23) and the fact that $\xi_{ni}\geq 0$, $\forall\, i,n$ imply the fact that $\sum_{i=1}^n\mathrm{E}|\xi_{ni}|=O(1)$. Now:

$$\sum_{i=1}^n\mathrm{E}|\xi_{ni}|\mathbf{1}_{|\xi_{ni}|>\varepsilon}\leq\sum_{i=1}^n\mathrm{P}(|\xi_{ni}|>\varepsilon)\leq n\,\mathrm{P}\left(\frac{1}{n\log n}\frac{\mathbf{1}_{Z>x}}{Z-x}>\varepsilon\right)$$
$$\leq n\,\mathrm{P}\left(0<Z-x<\frac{1}{\varepsilon n\log n}\right)\leq n\,\|g_0\|_\infty\frac{1}{\varepsilon n\log n}\to 0.$$

Finally note that the left hand side of (3.3.23) is equal to:

$$\frac{1}{\log n}\mathrm{E}\frac{(\varepsilon-1)^2}{Z-x}\mathbf{1}_{\{|\varepsilon-1|^2<n\log n|Z-x|,\,Z>x\}}\sim\frac{\log(n\log n)}{\log n}g_0(x).$$

where in the last line we used Lemma 3.3.4. □

Lemma 3.3.7. *For $\varepsilon_i\overset{i.i.d.}{\sim}Exp(1)$, $Z_i\overset{i.i.d.}{\sim}G_0$, the process $\mathbb{Z}_n^*$ is defined as:*

$$\mathbb{Z}_n^*(t):=\frac{1}{\sqrt{n\log n}}\sum_{i=1}^n(\varepsilon_i-1)\,f_t^n(Z_i). \tag{3.3.24}$$

where $f_t^n(z):=2(\delta_n^*)^{-1}\{\sqrt{(z-x)_+}-\sqrt{(z-x-\delta_n^*t)_+}\}$, $\delta_n=\sqrt{\log n}/\sqrt{n}$ and $\delta_n^*=\delta_n^{1/\gamma}$ where $\gamma>1/2$ satisfies condition (3.1.20).

Fix $t \in I_x$, where $I_x := [0,\infty)$ if $x = 0$, and $I_x := \mathbb{R}$ if $x > 0$. Then, in probability under G_0:

$$\frac{1}{\sqrt{n \log n}} \sum_{i=1}^{n} (\varepsilon_i - 1)\, f_t^n(Z_i) \mid Z_1, \ldots, Z_n \rightsquigarrow N\left(0, t^2 g_0(x)/(2\gamma)\right), \tag{3.3.25}$$

Furthermore:

$$Cov\left(\mathbb{Z}_n^*(t), \mathbb{Z}_n^*(s) \,\middle|\, Z_1, \ldots, Z_n\right) = \frac{1}{2\gamma} g_0(x) st\left(1 - O_{G_0}\left(\frac{\log\log n}{\log n}\right)\right) + O_{G_0}\left(\frac{1}{\log n}\right). \tag{3.3.26}$$

Proof. Without loss of generality set $t = 1$. We apply Theorem 6.16 in [63] (with $\xi_{ni} = ((\varepsilon_i - 1) f_1^n(Z_i))/\sqrt{n \log n}$) conditionally on $Z_1, \ldots, Z_n$. Therefore we need to show:

$$\sum_{i=1}^{n} \mathrm{P}\left(|\varepsilon_i - 1|\, f_1^n(Z_i) \geq \sqrt{\delta\, n \log n} \;\middle|\; Z_1, \ldots, Z_n\right) \overset{G_0}{\to} 0, \quad \forall\, \delta > 0, \tag{3.3.27}$$

$$\frac{1}{\sqrt{n \log n}} \sum_{i=1}^{n} \mathrm{E}\left((\varepsilon_i - 1) f_1^n(Z_i) \mathbf{1}_{\{|\varepsilon_i - 1| f_1^n(Z_i) \geq \sqrt{\delta\, n \log n}\}} \;\middle|\; Z_1, \ldots, Z_n\right) \overset{G_0}{\to} 0, \tag{3.3.28}$$

and:

$$\frac{1}{n \log n} \sum_{i=1}^{n} \mathrm{Var}\left((\varepsilon_i - 1) f_1^n(Z_i) \mathbf{1}_{\{|\varepsilon_i - 1| f_1^n(Z_i) \geq \sqrt{\delta\, n \log n}\}} \;\middle|\; Z_1, \ldots, Z_n\right) \overset{G_0}{\to} \frac{g_0(x)}{2\gamma}. \tag{3.3.29}$$

First note that (3.3.27) and (3.3.28) are implied by convergence in mean (which means unconditional probability and expectation). In what follows, we will often use that $|f_n^1(Z)| \leq 2/\sqrt{\delta_n^*}$. So (3.3.27) is implied by:

$$\begin{aligned}
&n\,\mathrm{P}\left((f_n^1(Z))^2 > \frac{\delta\, n \log n}{(\varepsilon - 1)^2}\right) \lesssim n\,\mathrm{P}\left(\frac{1}{\delta_n^*} > \frac{\delta\, n \log n}{(\varepsilon - 1)^2}\right) = n\,\mathrm{P}\left(|\varepsilon - 1| > \sqrt{\delta\, \delta_n^*\, n \log n}\right) \\
&= n \int_{|\lambda - 1| > \sqrt{\delta_n^*\, n \log n\, \delta}} e^{-\lambda} d\lambda = n \int_{\lambda > 1 + \sqrt{\delta_n^*\, n \log n\, \delta}} e^{-\lambda} d\lambda = n \int_{1 + \sqrt{\delta_n^*\, n \log n\, \delta}}^{\infty} e^{-\lambda} d\lambda \\
&= n\, e^{-1 - \sqrt{\delta_n^*\, n \log n\, \delta}} \to 0,
\end{aligned}$$

because $\left(\frac{\log n}{n}\right)^{1/2\gamma} \cdot n \log n \gg n^{1 - 1/2\gamma} \log^2 n$, since $\delta_n^* = \left(\frac{\log n}{n}\right)^{1/2\gamma}$, $\quad \gamma > 1/2$.

Because $\mathrm{E}(\varepsilon_i - 1) = 0$ and $\mathrm{E} f_n^1(Z) = O(1)$ (see (A9) in [43]), the expected value of (3.3.28) goes to zero because:

$$\begin{aligned}
&\mathrm{E}\left|\frac{1}{\sqrt{n \log n}} \sum_{i=1}^{n} \mathrm{E}\Big((\varepsilon_i - 1) f_n^1(Z_i) \mathbf{1}_{\{|\varepsilon_i - 1| f_1^n(Z_i) > \sqrt{\delta\, n \log n}\}} \Big| Z_1, \ldots, Z_n\Big)\right| \\
&\lesssim \sqrt{\frac{n}{\log n}}\, \mathrm{E} f_n^1(Z) \mathrm{E}(\varepsilon - 1) \mathbf{1}_{\{|\varepsilon - 1| > \sqrt{\delta\, \delta_n^*\, n \log n}\}} \lesssim \sqrt{\frac{n}{\log n}} \frac{(1 + \sqrt{\delta_n^* n \log n})}{e^{\sqrt{\delta\, \delta_n^*\, n \log n} + 1}},
\end{aligned}$$

which goes to zero as $\gamma > 1/2$ and thus $\left(\frac{\log n}{n}\right)^{1/2\gamma} \cdot n \log n \gg n^{1 - 1/2\gamma} \log^2 n \gtrsim n^{\eta}$ some $\eta > 0$.

In order to prove (3.3.29) we proceed in steps. First note that (3.3.29) equals:

$$\frac{1}{n \log n} \sum_{i=1}^{n} \left\{ \mathrm{E}\left((f_1^n(Z_i))^2 (\varepsilon_i - 1)^2 \mathbf{1}_{\{|\varepsilon_i - 1| f_1^n(Z_i) < \sqrt{\delta\, n \log n}\}} \;\middle|\; Z_1, \ldots, Z_n\right) \right. \tag{3.3.30}$$

$$- \left(\mathrm{E}\left(f_1^n(Z_i)(\varepsilon_i - 1)\mathbf{1}_{\{|\varepsilon_i - 1| f_1^n(Z_i) < \sqrt{\delta\, n \log n}\}} \,\middle|\, Z_1, \ldots, Z_n \right) \right)^2 \Big\}. \tag{3.3.31}$$

First we show the term in (3.3.31) goes to zero. Because $\mathrm{E}(\varepsilon - 1) = 0$ and $\varepsilon \perp\!\!\!\perp Z$:

$$\begin{aligned}
&\mathrm{E}\left(\frac{1}{n \log n} \sum_{i=1}^n \mathrm{E}\left((\varepsilon_i - 1) f_n^1(Z_i) \mathbf{1}_{\{|\varepsilon_i - 1| f_n^1(Z_i) > \sqrt{\delta\, n \log n}\}} \,\middle|\, Z_1, \ldots, Z_n \right) \right)^2 \\
&\leq \frac{1}{\log n} \mathrm{E}(\varepsilon - 1)^2 (f_n^1(Z))^2 \mathbf{1}_{\{|\varepsilon - 1| f_n^1(Z) > \sqrt{\delta\, n \log n}\}} \\
&\leq \frac{1}{\log n} \mathrm{E}(\varepsilon - 1)^2 (f_n^1(Z))^2 \mathbf{1}_{\{|\varepsilon - 1| > \sqrt{\delta\, \delta_n^*\, n \log n}\}} \\
&= \frac{1}{\log n} \mathrm{E}(f_n^1(Z))^2 \mathrm{E}(\varepsilon - 1)^2 \mathbf{1}_{\{|\varepsilon - 1| > \sqrt{\delta\, \delta_n^*\, n \log n}\}} \to 0,
\end{aligned}$$

because by (A7) in [43] with $s = t = 1$, $\frac{1}{\log n} \mathrm{E}(f_n^1(Z))^2 = O(1)$ and for all n big,

$$\begin{aligned}
\mathrm{E}(\varepsilon - 1)^2 \mathbf{1}_{\{|\varepsilon - 1| > \sqrt{\delta\, \delta_n^*\, n \log n}\}} &\lesssim \int_{1 + \sqrt{\delta\, \delta_n^*\, n \log n}}^{\infty} (\lambda - 1)^2 e^{-\lambda} d\lambda \\
&= \left(\sqrt{\delta\, \delta_n^*\, n \log n}\, (\sqrt{\delta\, \delta_n^*\, n \log n} + 2) + 2 \right) e^{-\sqrt{\delta\, \delta_n^*\, n \log n} - 1} \to 0
\end{aligned} \tag{3.3.32}$$

again because $\delta_n^*\, n \log n \sim n^{1 - 1/2\gamma} \log^{1 + \frac{1}{2\gamma}}(n)$, $\gamma > 1/2$.

Now we show the behavior of (3.3.30). Denote in what follows by E_ε the expectation only under ε, thus conditionally on $Z_1, \ldots, Z_n$. Using that $\mathrm{E}(\varepsilon - 1)^2 = 1$ and $Z \perp\!\!\!\perp \varepsilon$:

$$\begin{aligned}
(3.3.30) &= \frac{1}{n \log n} \sum_{i=1}^n \mathrm{E}_\varepsilon (\varepsilon_i - 1)^2 (f_n^1(Z_i))^2 \mathbf{1}_{\{|\varepsilon_i - 1| f_n^1(Z_i) < \sqrt{\delta\, n \log n}\}} \\
&= \frac{1}{n \log n} \sum_{i=1}^n \mathrm{E}_\varepsilon (\varepsilon_i - 1)^2 (f_n^1(Z_i))^2 \{1 - \mathbf{1}_{\{|\varepsilon_i - 1| f_n^1(Z_i) \geq \sqrt{\delta\, n \log n}\}} \} \\
&= \frac{1}{n \log n} \sum_{i=1}^n (f_n^1(Z_i))^2 - \frac{1}{n \log n} \sum_{i=1}^n \mathrm{E}_\varepsilon (\varepsilon_i - 1)^2 (f_n^1(Z_i))^2 \mathbf{1}_{\{|\varepsilon_i - 1| f_n^1(Z_i) > \sqrt{\delta\, n \log n}\}}.
\end{aligned}$$

First we show $\frac{1}{n \log n} \sum_{i=1}^n (f_n^1(Z_i))^2 \overset{G_0}{\to} g_0(x)/2\gamma$. We use again Lemma 3.3.5 with $\xi_{ni} = (f_n^1(Z_i))^2 / n \log n$. Now $\forall\, \varepsilon > 0$ and all n big enough:

$$\begin{aligned}
\sum_{i=1}^n \mathrm{E}|\xi_{ni}| \mathbf{1}_{\{|\xi_{ni}| > \varepsilon\}} &= \frac{1}{n \log n} \sum_{i=1}^n \mathrm{E}(f_n^1(Z_i))^2 \mathbf{1}_{\{(f_n^1(Z_i))^2 > \varepsilon n \log n\}} \\
&\leq \frac{1}{n \log n} \sum_{i=1}^n \mathrm{E}(f_n^1(Z_i))^2 \mathbf{1}_{\{\frac{1}{\delta_n^*} > \varepsilon n \log n\}} = 0,
\end{aligned}$$

because $\frac{1}{\delta_n^*} = (\frac{n}{\log n})^{1/2\gamma} \ll n \log n$ as $\gamma > 1/2$. Finally:

$$\sum_{i=1}^n \mathrm{E}\xi_{ni} = \frac{1}{\log n} \int (f_n^1(z))^2 g_0(z) dz \to \frac{1}{2\gamma} g_0(x)$$

where we use (A7) of [43] with $s = t = 1$. The fact that ξ_{ni} are nonnegative and $\sum_{i=1}^n \mathrm{E}\xi_{ni} \to g_0(x)/2\gamma$ imply $\sum_{i=1}^n \mathrm{E}|\xi_{ni}| = O(1)$. Using (3.3.32) we conclude:

$$\begin{aligned}
&\frac{1}{n\log n}\sum_{i=1}^n \mathrm{E}_\varepsilon(\varepsilon_i - 1)^2 (f_n^1(Z_i))^2 \mathbf{1}_{\{|\varepsilon_i - 1| f_n^1(Z_i) > \sqrt{\delta\, n\log n}\}} \\
&\leq \frac{1}{n\log n}\sum_{i=1}^n (f_n^1(Z_i))^2 \mathrm{E}_\varepsilon(\varepsilon - 1)^2 \mathbf{1}_{\{|\varepsilon - 1| > \sqrt{\delta\, \delta_n^* n\log n}\}} \to 0
\end{aligned}$$

since $\frac{1}{n\log n}\sum_{i=1}^n (f_n^1(Z_i))^2 = O_{G_0}(1)$.

We now prove (3.3.26). Fix $s, t \in I_x$. Since:

$$\mathrm{E}\left(\mathbb{Z}_n^*(t) \mid Z_1, \ldots, Z_n\right) = \frac{1}{\sqrt{n\log n}}\sum_{i=1}^n \mathrm{E}\left((\varepsilon_i - 1)\, f_t^n(Z_i) \mid Z_1, \ldots, Z_n\right) = 0. \tag{3.3.33}$$

Thus using (3.3.33), the above relations, that $Z \perp\!\!\!\perp \varepsilon$ and that $\varepsilon \overset{\text{i.i.d.}}{\sim} \mathrm{Exp}(1)$:

$$\mathrm{Cov}\left(\mathbb{Z}_n^*(t), \mathbb{Z}_n^*(s) \,\middle|\, Z_1, \ldots, Z_n\right) = \frac{1}{n\log n}\sum_{i=1}^n f_t^n(Z_i) f_s^n(Z_i) \underbrace{\mathrm{E}\,(\varepsilon_i - 1)^2}_{=1}. \tag{3.3.34}$$

Now we apply Lemma 3.3.5 with $\xi_{ni} := \frac{f_t^n(Z_i) f_s^n(Z_i)}{n\log n}$ to show:

$$\frac{1}{n\log n}\left\{\sum_{i=1}^n f_t^n(Z_i) f_s^n(Z_i) - n\,\mathrm{E}\, f_t^n(Z) f_s^n(Z)\right\} \overset{G_0}{\to} 0.$$

First, by using (A7) from [43] and that $\delta_n^* = (\log n/n)^{1/2\gamma}$:

$$\begin{aligned}
\frac{1}{\log n}\mathrm{E}\, f_t^n(Z_i) f_s^n(Z_i) &= -st g_0(x)\frac{\log \delta_n^*}{\log n} + O\left(\frac{1}{\log n}\right) \\
&= \frac{1}{2\gamma} g_0(x) st\left(1 - O\left(\frac{\log\log n}{\log n}\right)\right) + O\left(\frac{1}{\log n}\right).
\end{aligned} \tag{3.3.35}$$

This implies together with the nonnegativity of the functions f_s^n that $\sum_{i=1}^n \mathrm{E}|\xi_{ni}| = O(1)$. Moreover:

$$\sum_{i=1}^n \mathrm{E}\left|\frac{f_t^n(Z_i) f_s^n(Z_i)}{n\log n}\right| \mathbf{1}_{\left|\frac{f_t^n(Z_i) f_s^n(Z_i)}{n\log n}\right| > \varepsilon} \leq \sum_{i=1}^n \mathrm{E}\left|\frac{f_t^n(Z_i) f_s^n(Z_i)}{n\log n}\right| \mathbf{1}_{\left|\frac{\sqrt{|st|}}{\delta_n^*}\right| > \varepsilon n\log n} = 0,$$

for all n big enough, because $(n/\log n)^{1/2\gamma} \ll n\log n$ as $\gamma > 1/2$. Conclude that the behavior of (3.3.34) is given by (3.3.35). □

Lemma 3.3.8. *The process $\mathbb{Z}_n^*$ is conditionally asymptotically equicontinuous, i.e. $\forall\, \eta > 0$ as $n \to \infty$ followed by $\delta > 0$:*

$$\mathrm{P}\left\{\sup_{|s-t|<\delta} \left|\mathbb{Z}_n^*(s) - \mathbb{Z}_n^*(t)\right| > \eta \mid Z_1, \ldots, Z_n\right\} \xrightarrow{G_0} 0.$$

Proof. Take the class of functions for $\delta > 0,\ M > 0$:

$$\mathscr{F}_{n,\delta}^{M} := \{(\log n)^{-1/2}(\varepsilon - 1)(f_s^n - f_t^n)(z) :\ s,t \in I_x, |s-t| < \delta, \max(|s|,|t|) < M\}. \tag{3.3.36}$$

and the class of functions $\mathscr{H}_{n,\delta}^{M}$, whose functions for $f \in \mathscr{F}_{n,\delta}^{M}$ are given by $f/(\varepsilon - 1)$, thus without the $\varepsilon - 1$. These classes have square integrable natural envelopes $F_{n,\delta}$ and $H_{n,\delta}$ that satisfy for all $\delta > 0$ small (cf. Lemma 5 in [43]):

$$\|F_{n,\delta}\|_2^2 = \mathrm{E}(\varepsilon - 1)^2 \|H_{n,\delta}\|_2^2 = \|H_{n,\delta}\|_{2,G_0}^2 \leq g_0(x + \delta_n^* s \wedge t)\, \delta^2 (\log n)^{-1} (\log(\delta \delta_n^*)^{-2}).$$

For fixed $s,t \in I_x$, we prove conditional asymptotic equicontinuity of $\mathbb{Z}_n^*$, i.e. $\forall\, \eta > 0$ as $n \to \infty$ followed by $\delta > 0$, the right-hand side of the display below converges to 0 under G_0:

$$\mathrm{P}\Big\{ \sup_{|s-t|<\delta} \big|\mathbb{Z}_n^*(s) - \mathbb{Z}_n^*(t)\big| > \eta \mid Z_1, \ldots, Z_n \Big\} \leq \frac{1}{\eta} \mathrm{E}\Big[\sup_{|s-t|<\delta} \big|\mathbb{Z}_n^*(s) - \mathbb{Z}_n^*(t)\big| \mid Z_1, \ldots, Z_n \Big].$$

By the maximal inequality in Theorem 2.14.1 in [103]:

$$\mathrm{E}\Big[\sup_{|s-t|<\delta} \big|\mathbb{Z}_n^*(s) - \mathbb{Z}_n^*(t)\big| \Big] \lesssim J(1) \|F_{n,\delta}\|_2 \lesssim \sqrt{\|g_0\|_\infty\, \delta^2 (\log n)^{-1} (\log(\delta \delta_n^*)^{-2})} \to 0,$$

where $J(1) := J(1, \mathscr{F}_{n,\delta}^{M}, \mathbb{L}_2) = \sup_Q \int_0^1 \sqrt{\log N(\eta \|F_{n,\delta}\|_2, \mathscr{F}_{n,\delta}^{M}, \mathbb{L}_2(Q))}\, d\eta$ and where the square $\mathbb{L}_2(Q)$-distance is given for s,t and $s',t' \in I_x$ by: $\frac{1}{\log n} \int ((f_s^n - f_t^n)(z) - (f_{s'}^n - f_{t'}^n)(z))^2\, dQ(z)$. (Note that we did not take $\varepsilon - 1$ into account as $\mathrm{E}(\varepsilon - 1)^2 = 1$). Furthermore, we used that $J(1, \mathscr{F}_{n,\delta}^{M}, \mathbb{L}_2) < \infty$ as a consequence of Lemma 2.1 in [43], where it is shown that the function classes under consideration are VC of uniform bounded index. □

In the following Lemma, we make use of the concept of weak convergence of a process in $\ell^\infty(K)$ for every compact $K \subset \mathbb{R}$ and ℓ^∞ being the space of bounded functions on K equipped with the supremum norm. For a review of the conditional weak convergence in ℓ^∞, see [103] (c.f. Example 1.5.1 and Theorem 1.5.4).

Lemma 3.3.9 (Conditional argmax continuous mapping)**.** *Let $\mathbb{Z}_n, \mathbb{Z}$ be stochastic processes indexed by $H \subseteq \mathbb{R}$ such that in probability given a sequence of random variables $Z_1, \ldots, Z_n$:*

$$\mathbb{Z}_n \mid Z_1, \ldots, Z_n \rightsquigarrow \mathbb{Z} \quad \textit{in} \quad \ell^\infty(K)$$

for every compact $K \subset H$. Suppose that almost all sample paths $t \mapsto \mathbb{Z}(t)$ are upper semicontinuous and possess a unique maximum at a (random) point $\hat{t}$, which as a random map in H is tight. If $\forall\, n \in \mathbb{N}$, $\mathbb{Z}_n(\hat{t}_n) = \sup_t \mathbb{Z}_n(t)$ a.s. and there exist a compact set $\tilde{K} \subset H$ such that, under P*:*

$$\mathrm{P}\left(\hat{t}_n \notin \tilde{K} \mid Z_1, \ldots, Z_n\right) \xrightarrow{\mathrm{P}} 0,$$

then $\hat{t}_n \mid Z_1, \ldots, Z_n \rightsquigarrow \hat{t}$ in probability in H.

Proof. By the continuous mapping theorem, for every closed set F and A, B arbitrary sets of H: $\sup_{t \in F \cap A} \mathbb{Z}_n(t) - \sup_{t \in B} \mathbb{Z}_n(t)$ converges conditionally on $Z_1, \ldots, Z_n$ in distribution to $\sup_{t \in F \cap A} \mathbb{Z}(t) - \sup_{t \in B} \mathbb{Z}(t)$, where $\mathbb{Z}(t)$ has a.s. continuous sample paths and a.s. unique maxima. Thus using the assumption on $\hat{t}_n$:

$$\limsup_{n\to\infty} \mathrm{P}(\hat{t}_n \in F \cap A \mid Z_1, \ldots, Z_n) \leq \limsup_{n\to\infty} \mathrm{P}\left(\sup_{t \in F \cap A} \mathbb{Z}_n(t) > \sup_{t \in B} \mathbb{Z}_n(t) \mid Z_1, \ldots, Z_n \right)$$

$$= \mathrm{P}\left(\sup_{t\in F\cap A} \mathbb{Z}(t) > \sup_{t\in B} \mathbb{Z}(t) \right) \leq \mathrm{P}(\hat{t} \in F \cup B^c),$$

where in the last inequality we use that $\mathbb{Z}(\hat{t}) > \sup_{t\notin F^c, t\in A} \mathbb{Z}(t)$ and thus the event

$$\left\{ \sup_{t\in F\cap A} \mathbb{Z}(t) > \sup_{t\in B} \mathbb{Z}(t) \right\} \subseteq \{\hat{t} \in F \cup B^c\}.$$

Because Z_n converges in distribution conditionally on $Z_1, \ldots, Z_n$ to Z in $\ell^\infty(K)$ for every compact K, take now $A = B = K$. Then for every open set G containing $\hat{t}$, it holds

$$\mathbb{Z}(\hat{t}) > \sup_{t\notin G, t\in K} \mathbb{Z}(t),$$

because of the continuity of $\mathbb{Z}$ and the uniqueness of $\hat{t}$. Thus, the above yields:

$$\limsup_{n\to\infty} \mathrm{P}(\hat{t}_n \in F \mid Z_1, \ldots, Z_n) \leq \limsup_{n\to\infty} \mathrm{P}(\hat{t}_n \in F \cap K \mid Z_1, \ldots, Z_n)$$
$$+ \limsup_{n\to\infty} \mathrm{P}(\hat{t}_n \in K^c \mid Z_1, \ldots, Z_n) = \mathrm{P}(\hat{t} \in F \cup K^c).$$

Note: $\mathrm{P}(\hat{t} \in F \cup K^c) \leq \mathrm{P}(\hat{t} \in F) + \mathrm{P}(\hat{t} \in K^c)$, where by the stochastic boundedness of $\hat{t}_n \mid Z_1, \ldots, Z_n$ and the choice of K^c, the terms $\mathrm{P}(\hat{t} \in K^c)$ can be made arbitrarily small and $\limsup_{n\to\infty} \mathrm{P}(\hat{t}_n \in K^c \mid Z_1, \ldots, Z_n)$ converges to 0. □

Lemma 3.3.10 (Consistency)**.** *Assume* (3.1.20) *holds true. The IIP is consistent, i.e.* $\forall\, \varepsilon > 0$ *as* $n \to \infty$*:*

$$\Pi_n(|\hat{V}_G(x) - V_0(x)| > \varepsilon \mid Z_1, \ldots, Z_n) \xrightarrow{G_0} 0. \tag{3.3.37}$$

Moreover:

$$\Pi_n\left(\left| T_G\left(V_0(x) + \delta_n(a + Z_n^{G_0}(1)) \right) - x \right| > \varepsilon \mid Z_1, \ldots, Z_n \right) \xrightarrow{G_0} 0. \tag{3.3.38}$$

Proof. We begin by showing (3.3.37). Let $\psi_x(z) := 2\sqrt{z} - 2\sqrt{(z-x)_+}$, using Proposition 4.3 in [42] with ψ_x we obtain:

$$\mathrm{E}\,(U_G(x) \mid Z_1, \ldots, Z_n) = \frac{1}{|\alpha| + n} \int \psi_x \, d\left(\alpha + \sum_{i=1}^n \delta_{Z_i} \right),$$
$$\mathrm{Var}\,(U_G(x) \mid Z_1, \ldots, Z_n) \lesssim \frac{1}{1 + |\alpha| + n} \frac{1}{|\alpha| + n} \int \psi_x^2 \, d\left(\alpha + \sum_{i=1}^n \delta_{Z_i} \right).$$

Thus, using the conditional Chebyshev inequality:

$$\Pi_n\left(|U_G(x) - U_0(x)| > \varepsilon \mid Z_1, \ldots, Z_n \right) \to 0, \quad \text{a.s.}$$

Note now that the functions U_G are non-decreasing, U_0 is bounded continuous non-decreasing, thus $\forall\, \varepsilon > 0$ by Exercise 3.2 in [53] we conclude:

$$\Pi_n\left(\|U_G - U_0\|_\infty > \varepsilon \mid Z_1, \ldots, Z_n \right) \to 0, \quad \text{a.s.}$$

Thus, by Marshall's inequality (c.f. Exercise 3.1 in [53]), with U_G^* least concave majorant of U_G:

$$\Pi_n\left(\|U_G^* - U_0\|_\infty > \varepsilon \mid Z_1, \ldots, Z_n\right) \overset{G_0}{\to} 0.$$

Now fix $\varepsilon > 0$ and $h > 0$ such that $|(U_0(x) - U_0(x-h))/h - V_0(x)| < \varepsilon$. By concavity of U_G^* we have:

$$V_G(x) \leq \frac{U_G^*(x) - U_G^*(x-h)}{h}.$$

Thus by Lemma 3.1 in [53]:

$$\begin{aligned}
&\Pi_n\left(V_G(x) \geq V_0(x) + 2\varepsilon \mid Z_1, \ldots, Z_n\right) \\
&\leq \Pi_n\left(\frac{U_G^*(x) - U_G^*(x-h)}{h} \geq \frac{U_0(x) - U_0(x-h)}{h} + \varepsilon \mid Z_1, \ldots, Z_n\right) \to 0, \quad \text{a.s.}
\end{aligned}$$

Similarly $V_G(x) \geq (U_G^*(x+h) - U_G^*(x))/h$ gives $\Pi_n\left(V_G(x) \leq V_0(x) - 2\varepsilon \mid Z_1, \ldots, Z_n\right) \to 0$, a.s. This being true for every $\varepsilon > 0$ gives (3.3.37).

From (3.3.37) follows consistency of T_G; by the switch relation, $\forall\, \varepsilon > 0$:

$$\begin{aligned}
&\Pi_n\left(\left|T_G\left(V_0(x) + \delta_n(a + Z_n^{G_0}(1))\right) - x\right| > \varepsilon \mid Z_1, \ldots, Z_n\right) \\
&\leq \Pi_n\left(T_G\left(V_0(x) + \delta_n(a + Z_n^{G_0}(1))\right) > x + \varepsilon \mid Z_1, \ldots, Z_n\right) \\
&\quad + \Pi_n\left(T_G\left(V_0(x) + \delta_n(a + Z_n^{G_0}(1))\right) < x - \varepsilon \mid Z_1, \ldots, Z_n\right) \\
&\leq \Pi_n\left(\hat{V}_G(x+\varepsilon) > V_0(x) + \delta_n(a + Z_n^{G_0}(1)) \mid Z_1, \ldots, Z_n\right) \\
&\quad + \Pi_n\left(\hat{V}_G(x-\varepsilon) < V_0(x) + \delta_n(a + Z_n^{G_0}(1)) \mid Z_1, \ldots, Z_n\right) \overset{G_0}{\to} 0, \qquad (3.3.39)
\end{aligned}$$

where by (3.3.37) above $\hat{V}_G(x \pm \varepsilon) \mid Z_1, \ldots, Z_n \to V_0(x \pm \varepsilon)$, $Z_n^{G_0}(1) \rightsquigarrow \mathbb{Z}(1)$ and $\delta_n \downarrow 0$, thus $\delta_n(a + Z_n^{G_0}(1)) \overset{G_0}{\to} 0$ and V_0 is strictly decreasing at x thus we have the strict inequalities $V_0(x+\varepsilon) < V_0(x) < V_0(x-\varepsilon)$. □

Proof of (3.1.36). The claim is implied by the unconditional version of the convergence given in (3.1.36) and this entails proving that the rate for $\left|T_G\left(V_0(x) + \delta_n(a + Z_n^{G_0}(1))\right) - x\right|$ is $(\delta_n^*)^{-1}$. Because of Lemma 3.3.10, we can proceed by verifying the rest of the conditions of Theorem 3.2.5 in [103] (or see Theorem 3.4.1 for a more general version).

The argmax process we are interested in is $\tilde{T}_n := \arg\max_t \{\tilde{Z}_n^G(t) - at\}$ where $\tilde{Z}_n^G$ is given in (3.1.27)-(3.1.28). Thus note that we can rewrite it as: $\tilde{T}_n = (\delta_n^*)^{-1} \arg\max_t \{\mathbb{M}_n(t)\}$, where:

$$\mathbb{M}_n(t) := 2\int \left(\sqrt{(z-x)_+} - \sqrt{(z-x-t)_+} - \frac{1}{2}V_0(x)t\right) d(G - \mathbb{G}_n + G_0)(z) \qquad (3.3.40)$$

$$- 2\frac{t}{\delta_n^*}\int \left(\sqrt{(z-x)_+} - \sqrt{(z-x-\delta_n^*)_+}\right) d(\mathbb{G}_n - G_0)(z) \qquad (3.3.41)$$

$$+ 2\int \left(\sqrt{(z-x)_+} - \sqrt{(z-x-t)_+}\right) d(\mathbb{G}_n - G_0)(z) - at\delta_n. \qquad (3.3.42)$$

Moreover define the deterministic function:

$$\mathbb{M}(t) := 2\int \left(\sqrt{(z-x)_+} - \sqrt{(z-x-t)_+} - \frac{1}{2}V_0(x)t\right) dG_0(z).$$

Note that $\mathbb{M}_n(0) = \mathbb{M}(0) = 0$. In [43] (c.f. proof of (18)) it is shown that $\exists\, \delta_0 > 0$ such that $\forall\, |t| < \delta_0$:

$$\mathbb{M}(t) - \mathbb{M}(0) \leq -K\,|t|^{\gamma+1} \tag{3.3.43}$$

Following Theorem 3.2.5 in [103], define thus the metric $d(s,t) := |s-t|^{\frac{\gamma+1}{2}}$.

Consider the class of functions $h_t : [0,\infty) \to \mathbb{R}$:

$$\mathscr{H} := \{h_t \,:\, h_t(z) = 2\sqrt{(z-x)_+} - 2\sqrt{(z-x-t)_+} - V_0(x)t,\ t \geq 0\}$$

and, for $\eta > 0$, its subclass $\mathscr{H}_\eta := \{h_t \in \mathscr{H} : |t| \leq \eta\}$. In [52] it is shown that the envelope of $\mathscr{H}_\eta$, $H_\eta(z) := 2\sqrt{(z-x)_+} - 2\sqrt{(z-x-\eta)_+} - V_0(x)\eta$, satisfies for all η small enough (c.f. (11) in [52]):

$$\int H_\eta^2(z)\, g_0(z)\, dz \leq -2g_0(x)\eta^2 \log(\eta). \tag{3.3.44}$$

Consider now the class of functions $\widetilde{\mathscr{H}_n} := \{\tilde{h}_t^n \,:\, t \geq 0\}$ where, for any fixed $t \geq 0$:

$$\tilde{h}_t^n(z) := 2\Big(\frac{t}{\delta_n^*} - 1\Big)\sqrt{(z-x)_+} - 2\frac{t}{\delta_n^*}\sqrt{(z-x-\delta_n^*)_+} + 2\sqrt{(z-x-t)_+},$$

and, for $\eta > 0$, its subclass $\widetilde{\mathscr{H}_{n,\eta}} := \{\tilde{h}_t^n \in \widetilde{\mathscr{H}_n} : |t| \leq \eta\}$. Using the same computations as in [52], it follows that the envelope of $\widetilde{\mathscr{H}_{n,\eta}}$, $\tilde{H}_{n,\eta}(z) := \sup_{|t|<\eta} |\tilde{h}_t^n(z)|$, satisfies for all η small enough:

$$\int \tilde{H}_{n,\eta}^2(z)\, g_0(z)\, dz \leq -2g_0(x)\eta^2(\log(\eta) + \log(\delta_n^*)). \tag{3.3.45}$$

Using the representation (3.1.30) we have:

$$\begin{aligned}\mathbb{M}_n(t) - \mathbb{M}(t) = (1-V_n)\int h_t(z)\, d(\mathbb{B}_n - \mathbb{G}_n)(z) + V_n \int h_t(z)\, d(Q - \mathbb{G}_n)(z) - at\delta_n \\ - \int \tilde{h}_t^n(z)\, d(\mathbb{G}_n - G_0)(z),\end{aligned}$$

and, by applying the maximal inequality in Theorem 2.14.1 in [103] for sufficiently small $\delta > 0$, we conclude ($d(t,0) < \delta \iff |t| < \delta^{\frac{2}{1+\gamma}}$):

$$\begin{aligned}&\mathrm{E}\Big[\sup_{|t|<\delta^{\frac{2}{1+\gamma}}} \sqrt{n}|(\mathbb{M}_n - \mathbb{M})(t)|\Big] \lesssim J(1, \mathscr{H}_{\delta^{\frac{2}{1+\gamma}}}, \mathbb{L}_2)\sqrt{\int H^2_{\delta^{\frac{2}{1+\gamma}}}\, dG_0} + \sqrt{n}\frac{1}{n}\delta^{\frac{2}{1+\gamma}} \\ &+ |a|\delta^{\frac{2}{1+\gamma}}\delta_n\sqrt{n} + J(1, \widetilde{\mathscr{H}}_{n,\delta^{\frac{2}{1+\gamma}}}, \mathbb{L}_2)\sqrt{\int \tilde{H}^2_{n,\delta^{\frac{2}{1+\gamma}}}\, dG_0} \lesssim \delta^{\frac{2}{1+\gamma}}(\sqrt{\log\delta} + \sqrt{\log n}).\end{aligned}$$

The first term after applying the maximal inequality comes from the $(1-V_n)(\mathbb{B}_n - \mathbb{G}_n)$-term in the representation above of $\mathbb{M}_n - \mathbb{M}$; the second one comes from the $V_n(Q - \mathbb{G}_n)$-term, the third one from the $at\delta_n$-term and the last one from the $\mathbb{G}_n - G_0$-term (having used also for this one the maximal inequality). The last inequality is just a consequence of (3.3.44), (3.3.45)

and the facts that $J(1, \mathscr{H}_{\delta^{\frac{2}{1+\gamma}}}, \mathbb{L}_2) < \infty$ by the proof of Lemma 2.1 in [43] and by the same reasonings $J(1, \widetilde{\mathscr{H}}_{n,\delta^{\frac{2}{1+\gamma}}}, \mathbb{L}_2)$ is uniformly bounded.

The above computations and Theorem 3.2.5 in [103] imply that the rate for d is determined as follows: $\delta^{\frac{2}{1+\gamma}}(\sqrt{\log \delta} + \sqrt{\log n}) \leq \sqrt{n}\delta^2$, hence $\delta \gtrsim \left(\frac{\log n}{n}\right)^{\frac{1+\gamma}{4\gamma}}$. Because $d(s,t) = |s-t|^{\frac{\gamma+1}{2}}$, the rate for the distance $|\cdot|$ is given by:

$$\left(\frac{\log n}{n}\right)^{\frac{1+\gamma}{4\gamma}\cdot\frac{2}{1+\gamma}} = \left(\frac{\log n}{n}\right)^{\frac{1}{2\gamma}} \equiv \delta_n^*.$$

Altogether the above proves that the rate for $\arg\max_t \{\mathbb{M}_n(t)\}$ is $(\delta_n^*)^{-1}$. □

3.3.2 Proofs of complemental statements for Hierarchical Modelling

Proofs of the consistency results

***Proof of Theorem** 3.2.1.a.* Fix $\varepsilon > 0$, let F_ε be F_0 restricted and renormalized to $[\varepsilon, L]$. Then $g_{F_\varepsilon} > 0$ whenever $g_0 > 0$ and hence:

$$K(g_0, g_F) = K(g_0, g_{F_\varepsilon}) + \int \left(\log \frac{g_{F_\varepsilon}}{g_F}(z)\right) g_0(z)\, dz. \tag{3.3.46}$$

We shall show: (i) that the first term on the right tends to 0 as $\varepsilon \downarrow 0$ and (ii) that the lim sup of the second term is non-positive if $F \rightsquigarrow F_\varepsilon$ weakly, for any fixed $\varepsilon > 0$. Because the DP(α) prior puts positive mass on every weak neighborhood of F_ε (see Theorem 4.15 in [42]), together (i) and (ii) show $g_0 \in \mathrm{KL}(\Pi)$. Next the consistency is immediate from Example 6.20 in [42].

(i) Because $F_0[\varepsilon, L] \to 1$ as $\varepsilon \downarrow 0$, we have $g_{F_\varepsilon} \to g_0$ pointwise on $(0, \infty)$ and hence $\|g_{F_\varepsilon} - g_0\|_1 \to 0$ by Sheffe's Theorem. We strengthen this convergence in KL-convergence by providing bounds on $\log\left(\frac{g_0}{g_{F_\varepsilon}}\right)$.

First $\log \frac{g_0}{g_{F_\varepsilon}} \geq 1 - \frac{g_{F_\varepsilon}}{g_0}$ since $-\log x \geq 1 - x$ for $x > 0$ where $\frac{g_{F_\varepsilon}}{g_0}$ is integrable relative to g_0. Second: note we can bound g_{F_ε} from below as follows:

$$g_{F_\varepsilon}(z) = \int_{z\vee\varepsilon}^{L} \frac{1}{2\sqrt{x(x-z)}} \cdot \frac{dF_0(x)}{F_0[\varepsilon, L]} \geq \frac{F_0(\max(z,1), L)}{\sqrt{L^2 - Lz}} \geq \frac{F_0(\max(z,1), L)}{L}.$$

Now we check integrability:

$$\mathrm{E}_0 \left|\log L \frac{g_0(z)}{F_0(\max(Z,1), L)}\right| \lesssim \mathrm{E}_0 \log g_0(z) + \int_0^L g_0(z) \left|\log F_0(\max(z,1), L)\right| dz$$

which is integrable. This concludes the proof of (i).

(ii) For fixed $x, x_1 \in \mathbb{R}^+$, $\psi(z; x) := \frac{\mathbf{1}_{x>z}}{2\sqrt{x^2 - xz}}$:

$$\|\psi_x(\cdot) - \psi_{x_1}(\cdot)\|_1 = 2\frac{\sqrt{x_1 - x}}{\sqrt{x_1}} \mathbb{1}_{\{x \leq x_1\}} + 2\frac{\sqrt{x - x_1}}{\sqrt{x}} \mathbb{1}_{\{x_1 \leq x\}}. \tag{3.3.47}$$

Now for any partition $[0, L] = \bigcup_{i=0}^N I_i$ and points $x_i \in I_i$, using (3.3.47)

$$\|g_F - g_{F_\varepsilon}\|_1 = \left\| \sum_{i=1}^N \int_{I_i} (\psi_x - \psi_{x_i}) d(F - F_\varepsilon)(x) + \sum_{i=1}^N \psi_i (F - F_\varepsilon)(I_i) \right\|$$

$$\leq \sum_{i=1}^{N} \sup_{x\in I_i} \|\psi_x - \psi_{x_i}\|_1 (F + F_\varepsilon)(I_i) + \sum_{i=1}^{N} |(F - F_\varepsilon)(I_i)|$$

$$\leq 2(F + F_\varepsilon)(I_0) + \max_{i\geq 1} \sup_{x\in I_i} 2\sqrt{\frac{|x - x_i|}{x \vee x_i}} + \sum_{i=1}^{N} |(F - F_\varepsilon)(I_i)|.$$

We can choose a partition in intervals with boundary points continuity points of F_ε and $I_0 \subset [0, \varepsilon]$ so that $F_\varepsilon(I_0) = 0$. Then the first and last terms on the right tend to zero as $F \rightsquigarrow F_\varepsilon$ weakly. We can further choose the partition fine enough so that the second term is arbitrarily small. Thus $\|g_F - g_{F_\varepsilon}\|_1 \to 0$ as $F \rightsquigarrow F_\varepsilon$ weakly. It follows that $\log \frac{g_{F_\varepsilon}}{g_F} \to 0$ in measure.

If $F \rightsquigarrow F_\varepsilon$ weakly, by Portmanteau lemma $\exists\, \delta > 0$ and $0 < c \leq 1$ such that $\forall\, z \in [L - \delta, L)$:

$$\liminf F(z, L) \geq c F_\varepsilon(z, L).$$

Therefore we can choose neighborhoods of Fs with positive prior probability such that $\forall\, z \in [L - \delta, L)$:

$$g_F(z) = \int \frac{dF(x)}{2\sqrt{x^2 - xz}} \geq \frac{F(z, L)}{2L} \geq \frac{cF_\varepsilon(z, L)}{2L}.$$

Conclude:

$$\int_0^L g_0(z) \log\left(\frac{g_{F_\varepsilon}}{g_F}(z)\right) dz = \int_0^{L-\delta} g_0(z) \log \frac{g_{F_\varepsilon}}{g_F}(z) dz + \int_{L-\delta}^{L} g_0(z) \log \frac{g_{F_\varepsilon}}{g_F}(z) dz$$

$$\leq \int_0^{L-\delta} g_0(z) \log \frac{2g_0(z)L}{F_0(\varepsilon, L)F(z, L)} dz + \int_{L-\delta}^{L} g_0(z) \log \frac{g_0(z)2L}{cF_0(z, L)} dz$$

$$\lesssim \int_0^{L-\delta} g_0(z) \log g_0(z) dz + \int_{L-\delta}^{L} g_0(z) \log \frac{1}{F_0(z, L)} dz < \infty,$$

where we used that $F \rightsquigarrow F_\varepsilon$ and $F(L - \delta, L) > 0$ a.s. □

***Proof of Theorem** 3.2.1.b.* We verify the conditions of Theorem 6.23 from [42]. We also use the result derived above for the Kullback-Leibler property. Now consider the following partitions of the space $\mathfrak{X}$ where the X's take values: set $\mathfrak{X}_n = [\frac{1}{a_n}, L]$, for $a_n = n^\beta$. Then $\operatorname{diam}(\mathfrak{X}_n) = L - \frac{1}{a_n}$. For given $\varepsilon > 0$, for $N_n \sim \eta n / \log n$ and small $\eta > 0$ to be determined at the end of the proof, define $\mathcal{P}_n$ by:

$$\mathcal{P}_n = \left\{ \sum_{j=1}^{\infty} w_j \psi\left(\cdot; x_j\right) : \left(w_j\right) \in \mathbb{S}_\infty, \sum_{j>N_n} w_j < \frac{\varepsilon}{8}, x_1, \ldots, x_{N_n} \in \mathfrak{X}_n \right\}.$$

By the series representation (Sethuraman) $F = \sum_j W_j \delta_{X_j}$ of the Dirichlet process, the prior density g_F is contained in $\mathcal{P}_n$, unless the weights of the representation satisfy $\sum_{j>N_n} W_j \geq \varepsilon/8$ or one of the points $X_1, \ldots, X_{N_n}$ is not in $\mathfrak{X}_n$. It follows that

$$\Pi\left(g_F \notin \mathcal{P}_n\right) \leq \mathrm{P}\left(\sum_{j>N_n} W_j \geq \frac{\varepsilon}{8} \right) + N_n \overline{\alpha}(\mathfrak{X}_n^c).$$

The last term is exponentially small by assumption. The stick-breaking weights are $W_j = V_j \prod_{l=1}^{j-1} (1 - V_l)$, for $V_l \overset{\text{iid}}{\sim} \mathrm{Be}(1, |\alpha|)$ and $\sum_{j>N} W_j = \prod_{j=1}^{N} \left(1 - V_j\right)$. It can be checked that

$-\log\left(1-V_j\right)$ is exponentially distributed with parameter $|\alpha|$, so that $R_n := -\log\sum_{j>N_n} W_j$ possesses a gamma distribution with parameters N_n and $|\alpha|$. Therefore the first term is equal to:

$$\mathrm{P}\left(R_n \le \log\frac{8}{\varepsilon}\right) = \int_0^{\log(8/\varepsilon)} \frac{1}{(N_n-1)!} r^{N_n-1}|\alpha|^{N_n} e^{-|\alpha|r} dr \le$$
$$\le \frac{(\log(8/\varepsilon)|\alpha|)^{N_n}}{N_n!} \le \left(\frac{e\log(8/\varepsilon)|\alpha|}{N_n}\right)^{N_n}$$

The last inequality follows, since $x^N/N! \le e^x$, for every $x > 0$, including $x = N_n$. For large n and fixed $\varepsilon > 0$ the term within brackets is (easily, because $\log n \ll \sqrt{n}$) bounded above by $1/\sqrt{n}$ and hence the right side is bounded by $e^{-(N_n/2)\log N_n} \le e^{n\eta/2}$.

Given any infinite probability vector $w = (w_1, w_2, \ldots)$, the probability vector $\tilde{w}$ obtained by setting the coordinates with $j > N_n$ to zero and renormalizing the remaining coordinates to sum to 1 (hence $\tilde{w}_j = 0$ for $j > N_n$ and $\tilde{w}_j = w_j/\sum_{j\le N_n} w_j$, otherwise) satisfies $\|w-\tilde{w}\|_1 = 2\sum_{j>N_n} w_j$. By the triangle inequality and the fact that $\|\psi(\cdot;x)\|_1 = 1$, for $\sum_{j=1}^n w_j\psi\left(\cdot;x_j\right) \in \mathcal{P}_n$ it follows:

$$\left\|\sum_{j=1}^{\infty} w_j\psi\left(\cdot;x_j\right) - \sum_{j\le N_n}\tilde{w}_j\psi\left(\cdot;x_j\right)\right\|_1 \le \|w-\tilde{w}\|_1 = 2\sum_{j>N_n} w_j \le \frac{\varepsilon}{4}.$$

Therefore, the finite mixtures $\sum_{j=1}^{N_n} w_j\psi\left(\cdot;x_j\right)$, with $(w_1,\ldots,w_{N_n})$ an arbitrary probability vector, form an $\varepsilon/4$-net over $\mathcal{P}_n$ with respect to the $\mathbb{L}_1$-norm. Consequently, by the triangular inequality, a $3\varepsilon/4$-net over these finite mixtures is an ε-net over $\mathcal{P}_n$. To construct such a net we restrict $(w_1,\ldots,w_{N_n})$ to an $\varepsilon/4$-net over all probability vectors of length N_n, restrict $x_1,\ldots,x_{N_n}$ to an $\frac{\varepsilon^2}{16a_n}$-net over $\mathfrak{X}_n$. By the triangle inequality:

$$\left\|\sum_{j\le N_n} w_j\psi\left(\cdot;x_j\right) - \sum_{j\le N_n}\tilde{w}'_j\psi\left(\cdot;x'_j\right)\right\|_1 \le \left\|w-w'\right\|_1 + \max_{j\le N_n}\left\|\psi\left(\cdot;x_j\right)-\psi\left(\cdot;x'_j\right)\right\|_1$$
$$\le \frac{\varepsilon}{4} + 2\sqrt{a_n}\max_{j\le N_n}\sqrt{|x'_j-x_j|} \le \frac{\varepsilon}{4}+\frac{\varepsilon}{2} = \frac{3\varepsilon}{4}.$$

The second upper bound follows from (3.3.47). It follows that for every w, x there exist w', x' in the restricted set, so that the right side is smaller than $3\varepsilon/4$. By Proposition C.2 in [42] the cardinality of this net is bounded above as follows (denote by $\mathcal{P}_n^F$ the set $\mathcal{P}_n$ but composed by finite mixtures, which constitute as shown above a $\frac{3}{4}\varepsilon$-net over $\mathcal{P}_n$):

$$N\left(\frac{3\varepsilon}{4},\mathcal{P}_n^F,\|\cdot\|_1\right) \le N\left(\frac{\varepsilon}{4},\{w=(w_1,\ldots,w_{N_n})\in\mathbb{R}^{N_n} \;:\; \|w\|_1=1\},\|\cdot\|_1\right)\cdot$$
$$\cdot N\left(\frac{\varepsilon^2}{16a_n^2},\{x=(x_1,\ldots,x_{N_n})\in\mathbb{R}^{N_n} \;:\; \|x\|_\infty \le \mathrm{diam}(\mathfrak{X}_n)\},\|\cdot\|_\infty\right)$$
$$\le \left(\frac{12}{\varepsilon}\right)^{N_n} \times \left(\frac{3\cdot 16a_n^2\cdot\mathrm{diam}(\mathfrak{X}_n)}{\varepsilon^2}\right)^{N_n}$$

Together with the definition $N_n \sim n\eta/\log n$ and the fact that $\log a_n \le \beta\log n$ for $\beta > 0$, this shows that the log-cardinality of $\mathcal{P}_n$ for sufficiently large n: $\log N(\varepsilon,\mathcal{P}_n,\|\cdot\|_1) \le \frac{n\varepsilon^2}{2}$, provided η is chosen small enough. □

***Proof of Theorem** 3.2.1.c.* First note that by substituting $y = z/x$:

$$\int_0^L z^k \int_z^L \frac{1}{\sqrt{x^2 - xz}} dF(x)dz = \int_0^1 y^k \frac{1}{2} \frac{1}{\sqrt{1-y}} dy \int_0^L x^k dF(x)$$

And using the above derivation we conclude for $c_k = \int_0^1 y^k \frac{1}{2} \frac{1}{\sqrt{1-y}} dy$:

$$\begin{aligned}
&\lim_{n\to\infty} \Pi_n\left(\left|c_k \int_0^L x^k d(F - F_0)(x)\right| > \varepsilon \mid Z_1, \ldots, Z_n\right) \\
&\leq \lim_{n\to\infty} \Pi_n\left(\left| \int_0^L z^k g_F(z)dz - \int_0^L z^k g_0(z)dz\right| > \varepsilon \mid Z_1, \ldots, Z_n\right) \\
&\leq \lim_{n\to\infty} \Pi_n\left(\int_0^L |z|^k \left|g_F(z) - g_0(z)\right| dz > \varepsilon \mid Z_1, \ldots, Z_n\right) \\
&\leq \lim_{n\to\infty} \Pi_n(L^k \|g_F - g_0\|_1 > \varepsilon \mid Z_1, \ldots, Z_n) = 0, \qquad G_0^n \text{ a.s.}
\end{aligned}$$

By the above we conclude that for any polynomial p_k:

$$\lim_{n\to\infty} \Pi_n\left(\left| \int_0^L p_k(x)d(F - F_0)(x)\right| > \varepsilon \mid Z_1, \ldots, Z_n\right) = 0, \qquad G_0^n \text{ a.s.} \tag{3.3.48}$$

In order to prove the claim, by the Portmanteau Lemma, it is enough to show that $\forall\ f \in C_b([0, L])$:

$$\lim_{n\to\infty} \Pi_n\left(\left| \int_0^L f(x)d(F - F_0)(x)\right| > \varepsilon \mid Z_1, \ldots, Z_n\right) = 0, \qquad G_0^n \text{ a.s.}$$

Recall now that every continuous function on a compact interval can be approximated uniformly by polynomials using the Stone-Weierstrass theorem. This means that $\forall\ \varepsilon > 0$ there exists k_ε such that $\forall\ x \in [0, L]$ we have $|p_{k_\varepsilon}(x) - f(x)| < \varepsilon/4$. Since:

$$\begin{aligned}
\varepsilon < \left| \int_0^L f(x)d(F - F_0)(x)\right| \leq \left| \int_0^L f(x) - p_{k_\varepsilon}(x)d(F - F_0)(x)\right| + \\
+ \left| \int_0^L p_{k_\varepsilon}(x)d(F - F_0)(x)\right| \leq \frac{\varepsilon}{2} + \left| \int_0^L p_{k_\varepsilon}(x)d(F - F_0)(x)\right|
\end{aligned}$$

which means:

$$\left\{F : \left| \int_0^L f(x)d(F - F_0)(x)\right| > \varepsilon\right\} \subseteq \left\{F : \left| \int_0^L p_{k_\varepsilon}(x)d(F - F_0)(x)\right| > \frac{\varepsilon}{2}\right\}$$

Therefore we conclude by (3.3.48):

$$\begin{aligned}
&\lim_{n\to\infty} \Pi_n\left(\left| \int_0^L f(x)d(F - F_0)(x)\right| > \varepsilon \mid Z_1, \ldots, Z_n\right) \leq \\
&\leq \lim_{n\to\infty} \Pi_n\left(\left| \int_0^L p_{k_\varepsilon}(x)d(F - F_0)(x)\right| > \frac{\varepsilon}{2} \mid Z_1, \ldots, Z_n\right) = 0 \qquad G_0^n \text{ a.s.}
\end{aligned}$$

By Portmanteau Lemma (Theorem A.2 in [42]) and the fact that $\mathbb{R}$ is separable, it follows that the above is equivalent to $\forall\, \varepsilon > 0$:

$$\lim_{n\to\infty} \Pi_n \left(d_{BL}\left(F, F_0\right) > \varepsilon \mid Z_1, \ldots, Z_n\right) = 0 \qquad G_0^n \text{ a.s.}$$

where d_{BL} is the bounded Lipschitz metric on the space of probability measures. Again by Portmanteau Lemma (see proof of Proposition 6.2 from [42]) the above is equivalent to: $d_{BL}\left(\Pi_n\left(\cdot \mid Z_1, \ldots, Z_n\right), \delta_{F_0}(\cdot)\right) \to 0,\ G_0^n$ a.s.. □

***Proof of Theorem** 3.2.1.d.* Because $g_F \overset{\mathbb{L}_1}{\to} g_0$, then $g_F \rightsquigarrow g_0$, which implies pointwise convergence of the Fourier transforms: $\hat{g}_F(\lambda) \to \hat{g}_0(\lambda)$. But $\hat{g}_F(\lambda) = \hat{k}(\lambda) \cdot \hat{V}_F(\lambda)$, where $\hat{k}$ is the Fourier transform of $k(x) = -\mathbf{1}_{\{x<0\}}/\pi\sqrt{-x}$, since $g_F = k * V_F$. Similarly $\hat{g}_0 = \hat{k} \cdot \hat{V}_0$. Note that $\hat{k}$ is nonzero. Thus also: $\hat{V}_F(\lambda) \to \hat{V}_0(\lambda)$, meaning $V_F \rightsquigarrow V_0$.

Because $V_0(0) = \int \delta_0\, dV_0$ and it is possible to find a sequence of mollifiers $\{f_m\}_m$ (which are bounded continuous functions) such that $\int f_m\, dV_0 \to \int \delta_0\, dV_0$ as $m \to \infty$, we conclude that also pointwise convergence $V_F(0) \to V_0(0)$ holds true. Because $F_0^*(x) = 1 - V_0(x)/V_0(0)$ conclude $\forall\, f \in C_b([0, L])$

$$\left| \int_0^L f(x) d(F^* - F_0^*)(x) \right| \lesssim |V_F(0) - V_0(0)| + V_0^{-1}(0) \int_0^L f(x) d(V_F - V_0)(x),$$

which converges to zero as $V_F \rightsquigarrow V_0$. □

Proofs of the contraction rates results

***Proof of the contraction rate in Hellinger distance, Theorem** 3.2.2.a.* We verify the conditions of Theorem 8.11 from [42] with d the Hellinger distance. Let $\bar{\varepsilon}_n = \varepsilon_n = (1/n)^{1/4} \log^{1/4} n$. Correspondingly, we obtain the set $\mathcal{P}_n$ which is $\mathcal{P}_\varepsilon$ in (3.3.49) with $\varepsilon = \varepsilon_n$. Note that: $(1/\varepsilon_n)^2 \log(1/\varepsilon_n) \asymp n\varepsilon_n^2$ and that the conditions required by Proposition 3.3.2 are the most restrictive and imply the ones required by Lemma 3.3.11 and Proposition 3.3.1. The chosen ε_n, combined with Proposition 3.3.2 below implies condition (i) of Theorem 8.11 and Proposition 3.3.1 implies condition (ii). Using Lemma 3.3.11 we obtain condition (iii):

$$\frac{\Pi(g_F \notin \mathcal{P}_n)}{\Pi(F : K\left(g_F, g_0\right) \le \varepsilon_n^2, K_2\left(g_F, g_0\right) \le \varepsilon_n^2)} = o(e^{-2n\varepsilon_n^2}).$$

□

For $\varepsilon \downarrow 0$ and $\mathfrak{X}_\varepsilon$ to be specified later, define:

$$\mathcal{P}_\varepsilon = \left\{ \sum_{j=1}^{\infty} w_j \psi\left(\cdot; x_j\right) : \left(w_j\right) \in \mathbb{S}_\infty, \sum_{j>N_\varepsilon} w_j < \frac{\varepsilon^2}{32}, x_1, \ldots, x_{N_\varepsilon} \in \mathfrak{X}_\varepsilon \right\}. \tag{3.3.49}$$

Lemma 3.3.11 (Sieves)**.** *Assume $\exists \beta > 0$ such that $\overline{\alpha}[0, \varepsilon^\beta] \le e^{-C\frac{1}{\varepsilon^2}\log\frac{1}{\varepsilon}}$ for some constant $C > 0$. Then for $\mathcal{P}_\varepsilon$ as in* (3.3.49) *with $N_\varepsilon \asymp \varepsilon^{-2}$ and all ε sufficiently small,*

$$\log \Pi(g_F \notin \mathcal{P}_\varepsilon) \lesssim -\frac{1}{\varepsilon^2} \log\left(\frac{1}{\varepsilon}\right).$$

Proof. Let $\mathfrak{X}_\varepsilon = \operatorname{supp}(\alpha) \setminus [0, \varepsilon^\beta]$. By the series representation $F = \sum_j W_j \delta_{X_j}$ of the Dirichlet process, the prior density g_F is contained in $\mathcal{P}_\varepsilon$, unless the weights of the representation satisfy $\sum_{j>N_\varepsilon} W_j \geq \varepsilon^2/32$ or $X_1, \ldots, X_{N_\varepsilon} \notin \mathfrak{X}_\varepsilon$. It follows that

$$\Pi\left(g_F \notin \mathcal{P}_\varepsilon\right) \lesssim \mathrm{P}\left(\sum_{j>N_\varepsilon} W_j \geq \frac{\varepsilon^2}{32}\right) + N_\varepsilon \overline{\alpha}(\mathfrak{X}_\varepsilon^c).$$

The last term is exponentially small with exponent $-C\frac{1}{\varepsilon^2}$ by assumption and the choice of N_ε. The stick-breaking weights in the first term satisfy $W_j = V_j \prod_{l=1}^{j-1}(1 - V_l)$, for $V_l \overset{\text{iid}}{\sim} \mathrm{Be}(1, |\alpha|)$ and $\sum_{j>N} W_j = \prod_{j=1}^{N}\left(1 - V_j\right)$. It can be checked that $-\log\left(1 - V_j\right)$ is exponentially distributed with parameter $|\alpha|$, so that $R_\varepsilon := -\log \sum_{j>N_\varepsilon} W_j$ possesses a gamma distribution with parameters N_ε and $|\alpha|$. Therefore this term is bounded above by:

$$\begin{aligned}\mathrm{P}\left(R_\varepsilon < \log\frac{32}{\varepsilon^2}\right) &= \int_0^{\log\frac{32}{\varepsilon^2}} \frac{1}{(N_\varepsilon - 1)!} r^{N_\varepsilon - 1} |\alpha|^{N_\varepsilon} e^{-|\alpha| r} dr \\ &\leq \frac{(\log(32/\varepsilon^2)|\alpha|)^{N_\varepsilon}}{N_\varepsilon!} \leq \left(\frac{e\log(32/\varepsilon^2)|\alpha|}{N_\varepsilon}\right)^{N_\varepsilon} \asymp \exp\left\{-\frac{1}{\varepsilon^2}\log\left(\frac{1}{\varepsilon}\right)\right\}.\end{aligned}$$

The second inequality follows, since $x^N/N! \leq e^x$, for every $x > 0$, including $x = N_\varepsilon$. □

Proposition 3.3.1 (Entropy). *Let $\mathcal{P}_\varepsilon$ as in* (3.3.49) *with $\mathfrak{X}_\varepsilon = [\varepsilon^\beta, L]$ for some $\beta > 0, L > 0$ as in Lemma 3.3.11 and $N_\varepsilon \asymp \varepsilon^{-2}$. Then:*

$$\log N(\varepsilon, \mathcal{P}_\varepsilon, h) \lesssim \left(\frac{1}{\varepsilon}\right)^2.$$

Proof. Given any infinite probability vector $w = (w_1, w_2, \ldots)$, the probability vector $\tilde{w}$ obtained by setting the coordinates with $j > N_\varepsilon$ to zero and renormalizing the remaining coordinates to sum to 1 (hence $\tilde{w}_j = 0$ for $j > N_\varepsilon$ and $\tilde{w}_j = w_j / \sum_{j\leq N_\varepsilon} w_j$, otherwise) satisfies $\|w - \tilde{w}\|_1 = 2\sum_{j>N_\varepsilon} w_j$. By the triangle inequality and the fact that $\|\psi_x\|_1 = 1$, it follows that for $F = \sum_{j=1}^{\infty} w_j \delta_{x_j}$ and $F_1 = \sum_{j\leq N_\varepsilon} \tilde{w}_j \delta_{x_j}$:

$$h^2(g_F, g_{F_1}) \leq \left\|\sum_{j=1}^{\infty} w_j \psi_{x_j} - \sum_{j\leq N_\varepsilon} \tilde{w}_j \psi_{x_j}\right\|_1 \leq \|w - \tilde{w}\|_1 = 2\sum_{j>N_\varepsilon} w_j \leq \frac{\varepsilon^2}{16}.$$

Therefore, the finite mixtures $\sum_{j\leq N_\varepsilon} w_j \psi_{x_j}$, with $(w_1, \ldots, w_{N_\varepsilon})$ an arbitrary probability vector, form an $\varepsilon/4$-net over $\mathcal{P}_\varepsilon$ with respect to the Hellinger distance. Consequently, by the triangle inequality, a $3\varepsilon/4$-net over these finite mixtures is an ε-net over $\mathcal{P}_\varepsilon$. To construct such a net we restrict $(w_1, \ldots, w_{N_\varepsilon})$ to an $\varepsilon^2/16$-net over all probability vectors of length N_ε, and restrict $x_1, \ldots, x_{N_\varepsilon}$ to an $\frac{\varepsilon^{4+\beta}}{16}$-net over $\mathfrak{X}_\varepsilon$. If $\bar{F} = \sum_{j\leq N_\varepsilon} w_j \delta_{x_j}$ and $F_2 = \sum_{j\leq N_\varepsilon} w'_j \delta_{x'_j}$ then by the triangle inequality:

$$\begin{aligned}h^2(g_{\bar{F}}, g_{F_2}) &\leq \left\|\sum_{j\leq N_\varepsilon} w_j \psi_{x_j} - \sum_{j\leq N_\varepsilon} w'_j \psi_{x'_j}\right\|_1 \leq \left\|w - w'\right\|_1 + \max_{j\leq N_\varepsilon}\left\|\psi_{x_j} - \psi_{x'_j}\right\|_1 \\ &\leq \frac{\varepsilon^2}{16} + 2\max_{j\leq N_\varepsilon}\sqrt{\frac{|x'_j - x_j|}{x'_j \vee x_j}} \leq \frac{\varepsilon^2}{16} + \frac{\varepsilon^2}{2}.\end{aligned}$$

The second upper bound follows from (3.3.47). It follows that for every w, x there exist w', x' in the restricted set, so that $h(g_F, g_{F_2}) \le 3\varepsilon/4$. By Proposition C.2 in [42] the cardinality of this net is bounded above by

$$N\left(\frac{\varepsilon^2}{16}, \mathbb{S}_{N_\varepsilon}, \|\cdot\|_1\right) \cdot N\left(\frac{\varepsilon^{4+\beta}}{16}, \mathfrak{X}_\varepsilon^{N_\varepsilon}, \|\cdot\|_\infty\right) \le \left(\frac{48}{\varepsilon^2}\right)^{N_\varepsilon} \cdot \left(\frac{3 \cdot 16 \cdot \operatorname{diam}(\mathfrak{X}_\varepsilon)}{\varepsilon^{4+\beta}}\right)^{N_\varepsilon}.$$

Since $N_\varepsilon \asymp \varepsilon^{-2}$, the logarithm of this is bounded by a multiple of $\left(\frac{1}{\varepsilon}\right)^2$. □

Lemma 3.3.12 (Finite approximation)**.** *Let the probability measure F_0 be supported on $[\delta, L]$ for $0 < \delta < L$ with density f_0 bounded away from zero and infinity and $\|1/g_0\|_1 < \infty$. Then there exists a discrete measure F_1 with at most $N \asymp \varepsilon^{-2}$ support points such that for g_{F_1} as in* (3.1.1)*:*

$$h(g_0, g_{F_1}) \le \varepsilon$$

Proof. Using Lemma A.1 in [40] and Lemma 3.3.13, let F_1 be a discrete measure on $[\delta, L]$ such that for $m \ge 1$:

$$\int x^{-j} d\left(F_1 - F_0\right)(x) = 0, \quad \forall\ 0 \le j \le m,$$
$$\int x^{j+1/4} d\left(F_1 - F_0\right)(x) = 0, \quad \forall\ 0 \le j \le 2m-1.$$

then F_1 can be chosen to have at most $3m$ support points. Using the power series:

$$\frac{\mathbf{1}_{\{x>z\}}}{2\sqrt{x^2 - xz}} = \frac{1}{2x} \frac{\mathbf{1}_{\{x>z\}}}{\sqrt{1 - \frac{z}{x}}} = \frac{1}{2x} \sum_{k=0}^{\infty} \binom{-1/2}{k} \left(-\frac{z}{x}\right)^k \mathbf{1}_{x>z},$$

and using the way F_1 is constructed, we obtain:

$$\begin{aligned}
h^2(g_0, g_{F_1}) &= \frac{1}{2} \int_0^L (g_0^{1/2}(z) - g_{F_1}^{1/2}(z))^2 \, dz \le \frac{1}{4} \int_{[0,\infty)} \frac{\left(g_{F_1}(z) - g_0(z)\right)^2}{g_0(z)} \, dz \\
&= \int_0^L \frac{1}{g_0(z)} \left(\int_z^\infty \frac{1}{2x} \sum_{k=m+1}^{\infty} \binom{-1/2}{k} \left(-\frac{z}{x}\right)^k d\left(F_0 - F_1\right)(x) \right)^2 dz \\
&\lesssim \int_0^L \frac{1}{g_0(z)} \left(\int_z^\infty \frac{1}{2x} \sum_{k=m+1}^{\infty} \binom{-1/2}{k} \left(-\frac{z}{x}\right)^k dF_0(x) \right)^2 dz \\
&\quad + \int_0^L \frac{1}{g_0(z)} \left(\int_z^\infty \frac{1}{2x} \sum_{k=m+1}^{\infty} \binom{-1/2}{k} \left(-\frac{z}{x}\right)^k dF_1(x) \right)^2 dz \\
&\lesssim \max\{1, \|f_0\|_\infty\} \int_0^L \frac{1}{g_0(z)} \left(\sum_{k=m+1}^{\infty} \binom{-1/2}{k} (-z)^k \int_z^\infty \frac{1}{x^{k+1}} \, dx \right)^2 dz \qquad (3.3.50) \\
&\lesssim \left(\sum_{k=m+1}^{\infty} \frac{1}{\sqrt{k}} \frac{L}{k+1} \right)^2 \lesssim \left(\sum_{k=m+1}^{\infty} \frac{1}{k^{3/2}} \right)^2 \le \left(\int_m^\infty \frac{1}{x^{3/2}} \, dx \right)^2 \asymp \frac{1}{m}.
\end{aligned}$$

In (3.3.50) we used the fact that for all k big enough:

$$\left|\binom{-1/2}{k}\right| = \frac{\sqrt{\pi}}{|\Gamma(\frac{1}{2}-k)\Gamma(k+1)|} \asymp \frac{1}{\sqrt{k}},$$

Moreover, it is possible to see that for $k > 0$, $\binom{-1/2}{k}(-1)^k > 0$, so for fixed z the functions $\binom{-1/2}{k}(-1)^k \frac{z^k}{x^{k+1}}$ are decreasing in x and nonnegative. Therefore using the integral test for series and the fact that $F_1\{x_i\} \in [0,1]$:

$$\sum_{x_i>z} \frac{1}{2x_i} \sum_{k=m+1}^{\infty} \binom{-1/2}{k}\left(-\frac{z}{x_i}\right)^k F_1\{x_i\} \leq \int_z^{\infty} \frac{1}{2x} \sum_{k=m+1}^{\infty} \binom{-1/2}{k}\left(-\frac{z}{x}\right)^k dx.$$

Similarly, we can upper bound the other integral by using $\|f_0\|_\infty < \infty$.

By choosing $m \asymp \varepsilon^{-2}$ we obtain the claim. □

Lemma 3.3.13. *Let F_0 as in Lemma 3.3.12. Then there exists a discrete probability measure F on $[0, L]$ with at most $N+1$ support points such that*

$$\int \frac{1}{x^j}\, dF(x) = \int \frac{1}{x^j}\, dF_0(x) \quad \text{for } j = 1, \ldots, N.$$

Proof. This lemma is an adaptation of Lemma A.1 in [40].

Consider the set $C = \left\{\left(1/x, \ldots, 1/x^N\right) : x \in (0, L]\right\}$ and its convex hull $\mathrm{conv}(C)$. Then for every probability measure F_0 on $[0, L]$ with finite inverse moments $\int_0^L \frac{1}{x^k}\, dF_0(x) < \infty$ for $k = 0, 1, 2, \ldots$ as in Lemma 3.3.12,

$$v_0 = \left(\int \frac{1}{x}\, dF_0(x), \ldots, \int \frac{1}{x^N}\, dF_0(x)\right)^T \in \mathrm{conv}(C)$$

(see, e.g., Lemma 3 p. 74 of [35]). Since $\mathrm{conv}(C) \subset \mathbb{R}^N$, any element of $\mathrm{conv}(C)$ may be written as a convex combination of at most $N+1$ elements of C (see, e.g., [90]). Thus there exist $\lambda_1, \ldots, \lambda_{N+1} \geq 0, \sum_{j=1}^{N+1} \lambda_j = 1, x_1 \ldots, x_{N+1} \in K$ such that

$$v_0 = \sum_{j=1}^{N+1} \lambda_j \left(\frac{1}{x_j}, \ldots, \frac{1}{(x_j)^N}\right)^T = \left(\int \frac{1}{x}\, dF(x), \ldots, \int \frac{1}{x^N}\, dF(x)\right)^T,$$

where $F = \sum_{j=1}^{N+1} \lambda_j \delta_{x_j}$. □

Proposition 3.3.2 (Prior mass)**.** *Let the density g_0 be bounded and let the conditions of Lemma 3.3.12 hold true. Let Π be the Dirichlet process $\mathrm{DP}(\alpha)$ with precision $|\alpha|$ and center measure α that has a Lebesgue density bounded away from 0 and ∞ and bounded support, that satisfies $\mathrm{supp}(F_0) \subseteq [\delta, L] \subset \mathrm{supp}(\alpha)$. For all sufficiently small $\varepsilon > 0$,*

$$\log \Pi\Big(F : K(g_F, g_0) \leq \varepsilon^2, K_2(g_F, g_0) \leq \varepsilon^2\Big) \gtrsim -\left(\frac{1}{\varepsilon}\right)^2 \log\left(\frac{1}{\varepsilon}\right). \tag{3.3.51}$$

Proof. By Lemma 3.3.12 there exists a discrete measure F_1 with $N \lesssim \varepsilon^{-2}$ support points such that $h(g_{F_1}, g_0) \leq \varepsilon$. We can take the support points x_j of F_1 such that $\forall\, j$ there exists an interval I_j with $\sup_{x \in I_j} \sqrt{|x_j - x|/(x_j \vee x)} \lesssim \varepsilon^2$ and $\bigcup_{j=1}^N I_j \subset [0, L]$, $F_1(x : x \notin I_j, \forall\, j) \leq \varepsilon^2$.

If not, we take a maximal set of points in the support points of F_1 which respect this property, and replace F_1 by the discrete measure obtained by relocating the masses of F_1 to the nearest points in the net which fulfill the above requirement. Call the obtained measure F_2. Then $h(g_{F_2}, g_{F_1}) \lesssim \varepsilon$, as seen in the proof of Proposition 3.3.1.

By Lemma B.2 in [42] we have:

$$\max\left(K\left(g_0, g_F\right), K_2\left(g_0, g_F\right)\right) \lesssim h^2\left(g_0, g_F\right) \log_-^2 h\left(g_0, g_F\right) \tag{3.3.52}$$

where $\log_- x = \max(-\log x, 0)$. In order to conclude we need a further step. We need to upper bound the $\mathbb{L}_1$ distance between g_0 and g_F. Let $F_1 = \sum_{i=1}^N w_j \delta_{x_j}$, with the support points $x_j \in I_j$ for every j as set in the beginning of the proof. Next note:

$$\begin{aligned}
&|g_F(z) - g_{F_1}(z)| = \left| \int \psi_x^z dF(x) - \int \psi_x^z dF_1(x) \right| \\
&\leq \int_{x \notin I_j,\, \forall j} \psi_x^z d(F + F_1)(x) + \sum_{j=1}^N \int_{I_j} |\psi_x^z - \psi_{x_j}^z| dF(x) + \sum_{j=1}^N \psi_{x_j}^z |F(I_j) - w_j|
\end{aligned}$$

Now note that: $\left\|\psi_x - \psi_{x_j}\right\|_1 = 2\sqrt{\frac{|x - x_j|}{x_j \vee x}}$, that $\left\|\psi_x^z\right\|_1 = 1$, that by construction:

$$\sup_{x \in I_j} \sqrt{|x_j - x|/(x_j \vee x)} \lesssim \varepsilon^2$$

and that:

$$F(x \,:\, x \notin I_j,\ \forall j) = 1 - \sum_{j=1}^N F(I_j) \leq \sum_{j=1}^N |F(I_j) - w_j|$$

Altogether, this gives:

$$\begin{aligned}
&\left\|g_F - g_{F_1}\right\|_1 \\
&\leq F(x \,:\, x \notin I_j,\ \forall\ j) + 2\sum_{j=1}^N \int_{I_j} \sqrt{\frac{|x_j - x|}{x_j \vee x}} dF(x) + \sum_{j=1}^N |F(I_j) - w_j| + \varepsilon^2 \\
&\leq 2\sum_{j=1}^N |F(I_j) - w_j| + 2\sum_{j=1}^N \sup_{x \in I_j} \sqrt{\frac{|x_j - x|}{x_j \vee x}} F(I_j) \lesssim 2\sum_{j=1}^N |F(I_j) - w_j| + 3\varepsilon^2.
\end{aligned} \tag{3.3.53}$$

Thus there exist constants d_1, d_2 such that for $F_1 = \sum_{i=1}^N w_j \delta_{x_j}$, with the support points $x_j \in I_j$, $\forall j$,

$$\begin{aligned}
\left\{F : \max\left(K, K_2\right)\left(g_0, g_F\right) < d_1\varepsilon^2\right\} &\overset{(3.3.52)}{\supset} \left\{F : h\left(g_0, g_F\right) \leq 2\varepsilon/\log_- \varepsilon\right\} \\
&\overset{\Delta \text{ in.}}{\supset} \left\{F : h\left(g_{F_1}, g_F\right) \leq \varepsilon/\log_- \varepsilon\right\} \\
&\overset{(3.2.4)}{\supset} \left\{F : \left\|g_F - g_{F_1}\right\|_1 \leq d_2\varepsilon^2/\log_-^2 \varepsilon\right\} \\
&\overset{(3.3.53)}{\supset} \left\{F : \sum_{j=1}^N \left|F(I_j) - w_j\right| \leq \varepsilon^2/\log_-^2 \varepsilon\right\}.
\end{aligned}$$

By Lemma A.2 in [40], since the base measure α has density bounded away from zero and infinity on its support by assumption, we have

$$\log \Pi\Big(F : \sum_{j=1}^{N} \big|F(I_j) - w_j\big| \leq \varepsilon^2 / \log_-^2 \varepsilon\Big) \gtrsim -N \log\left(\frac{\log_- \varepsilon}{\varepsilon}\right).$$

The proposition follows upon combining the preceding. □

***Proof of pointwise contraction rate for V, Theorem** 3.2.2.b.* Let $\mathcal{F}_n$ be such that $\Pi_n(F \in \mathcal{F}_n \mid Z_1, \ldots, Z_n) \xrightarrow{G_0} 1$. Without loss of generality, let the $F \in \mathcal{F}_n$ satisfy Theorem 3.2.2.a.. Denote by $\|\cdot\|_{\mathcal{F}_n} = \sup_{F \in \mathcal{F}_n} |\cdot|$. Now note for $\eta_n = \frac{1}{\sqrt{\log n}}$:

$$\begin{aligned}
|V_F(x) - V_0(x)| &= \left| \int_x^\infty \frac{1}{\sqrt{z-x}} (g_F - g_0)(z) dz \right| \\
&= \left| \left(\int_x^{x+\eta_n} + \int_{x+\eta_n}^\infty \right) \frac{1}{\sqrt{z-x}} (g_F - g_0)(z) dz \right| \\
&\leq \left| \int_x^{x+\eta_n} \frac{1}{\sqrt{z-x}} (g_F - g_0)(z) dz \right| + \frac{2}{\sqrt{\eta_n}} h(g_F, g_0)
\end{aligned}$$

Then define $V_F^{\eta_n}(x) = \int_x^{x+\eta_n} \frac{1}{\sqrt{z-x}} g_F(z) dz$ and note:

$$\begin{aligned}
&\Pi_n\left(\frac{\sqrt[4]{n}}{(\log n)^{3/4}} |V_F(x) - V_0(x)| > M \mid Z_1, \ldots, Z_n \right) \\
&\leq \Pi_n\left(\frac{\sqrt[4]{n}}{(\log n)^{3/4}} |V_F^{\eta_n}(x) - V_0^{\eta_n}(x)| > M \mid Z_1, \ldots, Z_n \right) \\
&\quad + \Pi_n\left(\frac{\sqrt[4]{n}}{(\log n)^{3/4}} \cdot \sqrt{\log n} \cdot h(g_F, g_0) > M \mid Z_1, \ldots, Z_n \right)
\end{aligned}$$

where the second term goes to 0 under G_0. By using Lemma 14 in [85] the claim is proven if we show that $\forall\, t$ in a neighborhood of 0, $r_n = \frac{\sqrt[4]{n}}{(\log n)^{3/4}}$:

$$\mathrm{E}\left[e^{t r_n (V_F^{\eta_n}(x) - V_0^{\eta_n}(x))} \mid Z_1, \ldots, Z_n, F \in \mathcal{F}_n \right] \xrightarrow{G_0} 1 = \int e^{ts} d\delta_0(s). \tag{3.3.54}$$

The first step to prove the claim consists of the definition of the appropriate perturbations of F. Let $\eta_n^* = \frac{1}{\sqrt{n}}$ and define the function

$$h_n^x(u) := \frac{2\sqrt{u}}{\pi f_0(u)} \frac{1}{u-x} \mathbf{1}_{\{\eta_n^* \leq u - x \leq \eta_n\}}$$

then $\|h_n^x\|_\infty = O(\frac{1}{\eta_n^*})$. Define for $t_n = t r_n / n$.

$$dF_{t_n}(u) := D_{t_n}^{-1} \left(1 + t_n h_n^x(u)\right) dF(u), \quad D_{t_n} := \int \left(1 + t_n h_n^x(u)\right) dF(u) \tag{3.3.55}$$

It follows that:

$$D_{t_n} g_{F_{t_n}}(z) = \int_z^\infty \frac{D_{t_n} dF_{t_n}(v)}{2\sqrt{v^2 - vz}} = g_F(z) + t_n \int_z^L \frac{1}{\pi \sqrt{u-z}} \frac{\mathbf{1}_{\{\eta_n^* \leq u - x \leq \eta_n\}}}{u-x} \frac{dF(u)}{f_0(u)}$$

The proof's second step involves rewriting the left-hand side of (3.3.54) as:

$$\frac{\int_{\mathcal{F}_n} e^{tr_n(V_F(x)-V_0(x))+\sum_{i=1}^n\left(\log g_F(Z_i)-\log g_{F_{-t_n}}(Z_i)\right)}\prod_{i=1}^n g_{F_{-t_n}}(Z_i)\,d\Pi(F)}{\int_{\mathcal{F}_n}\prod_{i=1}^n g_F(Z_i)\,d\Pi(F)} \tag{3.3.56}$$

where F_{-t_n} is as in (3.3.55) (note that the inserted perturbation has negative sign). The proof's third step consists of studying the sum of the log-likelihood ratios:

$$\sum_{i=1}^n \log\frac{g_{F_{-t_n}}}{g_F}(Z_i) = \sum_{i=1}^n \log\left\{1-\frac{t_n}{D_{t_n}}Y_{ni}^F\right\} \tag{3.3.57}$$

where $Y_{ni}^F = V_{ni}^F - \mathrm{E}V_{ni}^F$, where E denotes the expectation under g_F and:

$$V_{ni}^F = \frac{1}{g_F(Z_i)}\int_{Z_i}^L \frac{1}{\sqrt{u-Z_i}}\cdot\frac{\mathbf{1}_{\eta_n^*\le u-x\le\eta_n}}{\pi(u-x)}\frac{dF(u)}{f_0(u)}.$$

Now, using the Taylor expansion of the logarithm $\log\{1-\frac{tY}{1-tK}\}$ at $t=0$ in (3.3.57) at $t=0$, we obtain for some $\tilde{t}$ between 0 and t:

$$(3.3.57) = \sum_{i=1}^n\left\{\frac{-t_nY_{ni}^F}{D_{\tilde{t}_n}(D_{\tilde{t}_n}-\tilde{t}_nY_{ni}^F)}+t_nY_{ni}^F-t_nY_{ni}^F\right\}.$$

The supremum over $F\in\mathcal{F}_1$ of the above expression can be upper bounded as follows:

$$n\max_{i\le i\le n}\left\|\frac{t_nY_{ni}^F}{D_{\tilde{t}_n}(D_{\tilde{t}_n}-\tilde{t}_nY_{ni}^F)}\right\|_{\mathcal{F}_1}\left\|D_{\tilde{t}_n}^2-\tilde{D}_{\tilde{t}_n}\tilde{t}_nY_{ni}^F-1\right\|_{\mathcal{F}_1}\le$$
$$n\max_{i\le i\le n}\underbrace{\|t_nY_{ni}^F\|_{\mathcal{F}_1}}_{(I)}\underbrace{\max_{i\le i\le n}\left\|\frac{1}{D_{\tilde{t}_n}(D_{\tilde{t}_n}-\tilde{t}_nY_{ni}^F)}\right\|_{\mathcal{F}_1}}_{(II)}\underbrace{\max_{i\le i\le n}\|D_{\tilde{t}_n}^2-D_{\tilde{t}_n}\tilde{t}_nY_{ni}^F-1\|_{\mathcal{F}_1}}_{(III)}$$

which we prove goes to zero under G_0 in probability.

(I) Since $t_n = tr_n/n$ we show: $\max_{i\le i\le n}\|t_nY_{ni}^F\|_{\mathcal{F}_1} = O_{G_0}\left(\frac{r_n}{n}\frac{1}{\sqrt{\eta_n^*}}\right)$. First note:

$$\begin{aligned}
|\mathrm{E}V_{ni}^F| &= \left|\int_0^L\int_0^\infty \frac{1}{\sqrt{u-z}}\cdot\frac{\mathbf{1}_{\eta_n^*\le u-x\le\eta_n}}{\pi(u-x)}\frac{dF(u)}{f_0(u)}dz\right| \\
&\le \left\|\frac{1}{f_0}\right\|_\infty\left|\int_0^L\int_0^\infty\frac{1}{\sqrt{u-z}}\cdot\frac{\mathbf{1}_{\eta_n^*\le u-x\le\eta_n}}{\pi(u-x)}\,dF(u)\,dz\right| \\
&\le \left\|\frac{1}{f_0}\right\|_\infty\int_0^L\int_0^\infty\frac{1}{\sqrt{u-z}}\cdot\frac{1}{\pi(u-x)}\,du\,dz \lesssim \left\|\frac{1}{f_0}\right\|_\infty\int_x^L\frac{1}{\sqrt{z-x}}dz = O(1).
\end{aligned}$$

Now, using Lemma 8 from [43] and $\|\sup_{z\in[x,x+\eta_n]}\frac{1}{g_F(z)}\|_{\mathcal{F}_1}<\infty$, for all $1\le i\le n$ uniformly over $\mathcal{F}_1$:

$$|V_{ni}^F| \le \left\|\frac{1}{f_0}\right\|_\infty\frac{1}{g_F(Z_i)}\int_{Z_i}^L\frac{1}{\sqrt{u-Z_i}}\cdot\frac{\mathbf{1}_{\eta_n^*\le u-x\le\eta_n}}{\pi(u-x)}\,dF(u)$$

$$\lesssim \frac{1}{g_F(Z_i)} \int_{Z_i}^{L} \frac{1}{\sqrt{u - Z_i}} \cdot \frac{\mathbf{1}_{\eta_n^* \le u - x \le \eta_n}}{\pi(u - x)}\, du = O_{G_0}\left(\frac{1}{\sqrt{\eta_n^*}}\right).$$

(II) Uniformly for $F \in \mathcal{F}_1$, we have:

$$|D_{\tilde{t}_n}| \le 1 + |\tilde{t}| \frac{r_n}{n} \sup_{F \in \mathcal{F}_1} \int h_n^x(u) dF(u) \le 1 + O\left(\frac{r_n}{n} \log\left(\frac{1}{\eta_n^*}\right)\right),$$

because we can use the integral test for series as $(u - x)^{-1}$ is decreasing and nonnegative on $[x + \eta_n^*, x + \eta_n]$. We have:

$$\begin{aligned}\int h_n^x(u) dF(u) &= \int \frac{2\sqrt{u}}{\pi f_0(u)} \cdot \frac{\mathbf{1}_{\eta_n^* \le u - x \le \eta_n}}{u - x}\, dF(u) \\ &\le \left\| \frac{1}{f_0} \right\|_\infty \frac{2\sqrt{x + \eta_n}}{\pi} \int_{x + \eta_n^*}^{x + \eta_n} \frac{du}{u - x} \sim \log\left(\frac{1}{\eta_n^*}\right).\end{aligned}$$

Using also the results derived for (I) we conclude $(II) = 1 + O_{G_0}(1)$ because:

$$\begin{aligned}\|D_{\tilde{t}_n}(D_{\tilde{t}_n} - \tilde{t}_n Y_{ni}^F)\|_{\mathcal{F}_1} &\le \|D_{\tilde{t}_n}\|_{\mathcal{F}_1}^2 + \|D_{\tilde{t}_n}\|_{\mathcal{F}_1} \|\tilde{t}_n Y_{ni}^F\|_{\mathcal{F}_1} \\ &= 1 + O\left(\frac{r_n}{n} \log \frac{1}{\eta_n^*}\right) + O\left(\frac{r_n^2}{n^2} \log^2 \frac{1}{\eta_n^*}\right) + O_{G_0}\left(\frac{r_n}{n} \frac{1}{\sqrt{\eta_n^*}}\right).\end{aligned}$$

(III) First note that $D_{\tilde{t}_n}^2 - 1 = 2 \int \tilde{t}_n h_n^x(u) dF(u) + \left(\int \tilde{t}_n h_n^x(u) dF(u)\right)^2$. Thus:

$$\begin{aligned}\max_{i \le i \le n} \|D_{\tilde{t}_n}^2 - 1 - D_{\tilde{t}_n} \tilde{t}_n Y_{ni}^F\|_{\mathcal{F}_1} &\le \|D_{\tilde{t}_n}^2 - 1\|_{\mathcal{F}_1} + \|D_{\tilde{t}_n}\|_{\mathcal{F}_1} \|\tilde{t}_n Y_{ni}^F\|_{\mathcal{F}_1} \\ &= O_{G_0}\left(\frac{r_n}{n}\left(\frac{1}{\sqrt{\eta_n^*}} + \log\left(\frac{1}{\eta_n^*}\right)\right)\right).\end{aligned}$$

We conclude that we can reduce the study of the numerator of (3.3.56) to:

$$\int_{\mathcal{F}_n} e^{t r_n (V_F^{\eta_n}(x) - V_0^{\eta_n}(x)) + \sum_{i=1}^n \left\{ \frac{t r_n}{n} Y_{ni}^F \right\} \pm t r_n \mathrm{E}_0 Y_{ni}^F} \prod_{i=1}^n g_{F_{-t_n}}(Z_i)\, d\Pi(F)$$

where E_0 denotes the expectation under g_0. The proof's fourth step consists in studying the exponent given in the integral in the above display. We now prove:

$$r_n \left\| V_F^{\eta_n}(x) - V_0^{\eta_n}(x) + \mathrm{E}_0 Y_{ni}^F \right\|_{\mathcal{F}_n} = o(1), \tag{3.3.58}$$

$$r_n \left\| \frac{1}{n} \sum_{i=1}^n Y_{ni}^F - \mathrm{E}_0 Y_{ni}^F \right\|_{\mathcal{F}_n} = o_{G_0}(1). \tag{3.3.59}$$

Denote for $k_n^x(u) = \frac{\mathbf{1}_{\eta_n^* \le u - x \le \eta_n}}{\pi(u - x)}$, its fractional $1/2$ integral by $(I^{1/2} k_n^x)(z) = \int_z^\infty \frac{k_n^x(u)}{\sqrt{u - z}}\, du$ and analogously $(I_{F/f_0}^{1/2} k_n^x)(z) = \int_z^\infty \frac{k_n^x(u)}{\sqrt{u - z}} \frac{dF(u)}{f_0(u)}$. Then:

$$(3.3.58) = r_n \left\| \int_0^L (g_F - g_0)(z) \left(\frac{\mathbf{1}_{x \le z \le x + \eta_n}}{\sqrt{z - x}} - \frac{1}{g_F(z)} (I_{F/f_0}^{1/2} k_n^x)(z) \right) \right\|_{\mathcal{F}_n}$$

$$\leq r_n \left\| \int_0^L (g_F - g_0)(z) \frac{1}{\sqrt{z-x}} \mathbf{1}_{x \leq z \leq x+\eta_n^*} dz \right\|_{\mathcal{F}_n} \tag{3.3.60}$$

$$+ r_n \left\| \int_0^L \frac{\mathbf{1}_{\eta_n^* \leq z-x \leq \eta_n}}{\sqrt{z-x}} (g_0 - g_F)(z) \left(1 - \frac{g_0}{g_F}(z)\right) dz \right\|_{\mathcal{F}_n} \tag{3.3.61}$$

$$+ r_n \left\| \int_0^L \left(1 - \frac{g_0}{g_F}(z)\right) \left(\frac{g_0(z)}{\sqrt{z-x}} \mathbf{1}_{\eta_n^* \leq z-x \leq \eta_n} - (I^{1/2}_{F/f_0} k_n^x)(z) \right) dz \right\|_{\mathcal{F}_n} \tag{3.3.62}$$

We consider first (3.3.60). Using the substitution $vs - x = y(s-x)$ thus $(1-v) = (1-y)\frac{s-x}{s}$:

$$\begin{aligned}
\int_x^{x+\eta_n^*} \frac{1}{\sqrt{z-x}} g_0(z) dz &= \int_0^L \int_{0<vs-x<\eta_n^*} \frac{\mathbf{1}_{0<v<1}}{\sqrt{vs-x}\sqrt{1-v}} \, dv \, dF_0(s) \\
&= \int_0^{x+\eta_n^*} \int_0^{\frac{\eta_n^*}{s-x} \wedge 1} \frac{s-x}{s\sqrt{(y(s-x))}} \frac{\sqrt{s}}{2\sqrt{(1-y)(s-x)}} \, dy \, dF_0(s) \\
&= \int_x^{x+\eta_n^*} \frac{1}{\sqrt{s}} \int_0^{\frac{\eta_n^*}{s-x} \wedge 1} \frac{1}{2\sqrt{y(1-y)}} \, dy \, dF_0(s) \leq \frac{\pi}{2} \int_x^{x+\eta_n^*} \frac{f_0(s)}{\sqrt{s}} \, ds \\
&\leq \pi \, \|f_0\|_\infty \, (\sqrt{x+\eta_n^*} - \sqrt{x}) \leq \pi \sqrt{\eta_n^*} \, \|f_0\|_\infty
\end{aligned}$$

where we used that: $\mathrm{Be}(\frac{\eta_n^*}{s-x} \wedge 1; \frac{1}{2}, \frac{1}{2}) = -\arcsin\sqrt{1 - \frac{\eta_n^*}{s-x} \wedge 1} + \frac{\pi}{2} \leq \frac{\pi}{2}$ and that since $x \mapsto \frac{1}{\sqrt{x}}$ is decreasing, by the integral test for series:

$$\int_x^{x+\eta_n^*} \frac{dF(s)}{2\sqrt{s}} = \sum_{i \,:\, 0<x_i-x<\eta_n^*} \frac{F\{x_i\}}{2\sqrt{x_i}} \leq \sum_{i \,:\, 0<x_i-x<\eta_n^*} \frac{1}{2\sqrt{x_i}} \leq \int_x^{x+\eta_n^*} \frac{ds}{2\sqrt{s}} \leq \sqrt{\eta_n^*}$$

thus (3.3.60) $= O(r_n \sqrt{\eta_n^*}) = o(1)$.

Now note that because $\sup_{z \in [x, x+\eta_n^*]} \left|(1 + \frac{g_0}{g_F}(z))^2\right| < \infty$ uniformly for $F \in \mathcal{F}_1$ and using the contraction rate given in Theorem 3.2.2.a. for the Hellinger distance:

$$(3.3.61) \leq \frac{r_n}{\eta_n^*} \sup_{F \in \mathcal{F}_1} \sup_{z \in [x, x+\eta_n^*]} \left| \left(1 + \frac{g_0}{g_F}(z)\right)^2 \right| \sup_{F \in \mathcal{F}_n} h^2(g_F, g_0) = o(1).$$

The quantity in (3.3.62) can be upper bounded by the following quantity:

$$r_n \left(\int_0^{x+\eta_n} \left| \frac{g_0}{g_F}(z) - 1 \right|^2 dz \right)^{\frac{1}{2}} \left(\int_0^L \left(\frac{\mathbf{1}_{\eta_n^* \leq z-x \leq \eta_n}}{\sqrt{z-x}} g_0(z) - I^{1/2}_{F/f_0} k_n^x(z) \right)^2 dz \right)^{\frac{1}{2}}$$

and by using Lemma 8 from [43]:

$$\begin{aligned}
\left(\int \left(I^{1/2} k_n^x(z) \right)^2 dz \right)^{1/2} &\leq \left(\int \left(\frac{1}{\sqrt{z-x}} \mathbf{1}_{\eta_n^* \leq z-x \leq \eta_n} \right)^2 dz \right)^{1/2} + O(\eta_n^{-1}) \\
&= O\left(\sqrt{\log \frac{\eta_n}{\eta_n^*}} + \eta_n^{-1} \right).
\end{aligned}$$

Moreover since the function $u \to \frac{1}{\sqrt{u-z}} \frac{1}{u-x} \mathbf{1}_{\eta_n^* \leq u-x \leq \eta_n}$ is decreasing and nonnegative in u, we can thus use the integral test for series thus $\forall\, z \in [0, L]$: $\left| I^{1/2}_{F/f_0} k_n^x(z) \right| \leq \left| I^{1/2} k_n^x(z) \right|$ and conclude:

$$\left(\int \left(I^{1/2}_{F/f_0} k_n^x(z) \right)^2 dz \right)^{1/2} = O\left(\sqrt{\log \frac{\eta_n}{\eta_n^*}} + \eta_n^{-1} \right).$$

Finally since uniformly for $F \in \mathcal{F}_1$, $\sup_{z\in[0,x+\eta_n]} g_F^{-1/2}(z) < \infty$ and similarly $\sup_{z\in[0,x+\eta_n]} ((1 + \frac{g_0}{g_F}(z))^2)^{1/2} < \infty$, we conclude:

$$(3.3.62) \lesssim r_n h(g_F, g_0)\, O\left(\sqrt{\log \frac{\eta_n}{\eta_n^*}} + \eta_n^{-1}\right).$$

We now prove (3.3.59). We consider the class of functions:

$$\overline{\mathcal{V}}_n = \left\{ z \mapsto \frac{\mathbf{1}_{0\le z\le x+\eta_n}}{g_F(z)} \int_z^L \frac{1}{\sqrt{u-z}} \frac{\mathbf{1}_{\eta_n^*\le u-x\le\eta_n}}{\pi(u-x)} \frac{dF(u)}{f_0(u)} \ : \ F \in \mathcal{F}_n \right\}$$

where $D^* = \sup_{F\in\mathcal{F}_1} \int \frac{\mathbf{1}_{\eta_n^*\le u-x\le\eta_n}}{(u-x)\pi} \frac{dF(u)}{f_0(u)} = O(\log \frac{\eta_n}{\eta_n^*})$, then $\mathcal{J}(1, \overline{\mathcal{V}}_n, V_n^*) < \infty$, and V_n^* is the maximizer over support of the DP of the functions $V_n^F \in \overline{\mathcal{V}}_n$.

Here comes in the importance of the localization argument because now the function: $z \mapsto \int_z^L \frac{\mathbf{1}_{\eta_n^*\le u-x\le\eta_n}}{\pi(u-x)\sqrt{u-z}} \frac{dF(u)}{f_0(u)}$ is 0 for all $z \ge x + \eta_n$ by definition; this means that we can use: $\sup_{F\in\mathcal{F}_1} \sup_{z\in[0,x+\eta_n]} g_F^{-1}(z) < \infty$ if $x < L$.

The class $\overline{\mathcal{V}}_n$ is the product of two classes:

$$\mathcal{U}_n = \left\{ z \mapsto \frac{1}{D^*} \int_z^L \frac{1}{\sqrt{u-z}} \frac{\mathbf{1}_{\eta_n^*\le u-x\le\eta_n}}{\pi(u-x)} \frac{dF(u)}{f_0(u)} \ : \ F \in \mathcal{F}_n \right\}$$

and $\phi(\mathcal{G}_n)$ where $g \mapsto \phi(g)$ is the mapping $g(z) \mapsto g^{-1}(z)\mathbf{1}_{[0,x+\eta_n]}(z)$ and

$$\mathcal{G}_n = \left\{ z \mapsto g_F(z) = \sum_{i=1}^{\infty} \frac{F\{x_i\}}{\sqrt{x_i^2 - x_i z}} \ : \ F \in \mathcal{F}_n \right\}.$$

We study for $C_n = [0, x + \eta_n]$ and U_n^* the maximizer over the support of the DP of the functions $U_n^F \in \mathcal{U}_n$, analogously g_* means the maximizer over the support of the DP of the functions g_F:

$$\begin{aligned}
&\int_0^1 \sup_Q \sqrt{\log N(\varepsilon \|\overline{V}_n^*\|_{Q,2}, \overline{\mathcal{V}}_n, \mathbb{L}_2(Q))}\, d\varepsilon \\
&\quad \le \int_0^1 \sup_Q \sqrt{\log N\left(\frac{\varepsilon}{2} \left\| \frac{\mathbf{1}_{C_n}}{g_*} \right\|_{Q,2}, \phi(\mathcal{G}_n), \mathbb{L}_2(Q)\right)} d\varepsilon \\
&\qquad\qquad + \int_0^1 \sup_Q \sqrt{\log N\left(\frac{\varepsilon}{2} \|U_n^{F\star}\|_{Q,2}, \mathcal{U}_n, \mathbb{L}_2(Q)\right)} d\varepsilon.
\end{aligned}$$

Take now the class of functions: $\mathcal{S} := \left\{ z \mapsto \frac{\mathbf{1}_{x>z}}{2\sqrt{x^2-xz}} \ : \ x \in \mathbb{R}^+ \right\}$. We prove that this class of functions is VC. It is immediate using Lemma 2.6.16 and 2.6.18 from [103] that the class of functions $\{z \mapsto x(x-z) \ : \ x \in \mathbb{R}^+\}$ is VC. Next by Lemma 2.6.18 (ii) in [103] we see that the class $\{z \mapsto x(x-z)_+ \ : \ x \in \mathbb{R}^+\}$ is VC. Again by Lemma 2.6.18 (viii) in [103] we see that the class of functions $\left\{ z \mapsto (x(x-z)_+)^{-1/2} \ : \ x \in \mathbb{R}^+ \right\}$ is VC, as the square root of a function is a monotone function. Note that taking the reciprocal of a function is monotone except for one point, which is the point where the original function assumes value 0, and in that case we

assign to the reciprocal also value 0. This makes the reciprocal function monotone except at one point. We can view the resulting class as the union of two VC classes (one obtained by taking the inverse except at zero and the other assigning value zero when the original function gets value zero), thus still VC. This proves $\mathcal{S}$ is VC.

Since in $\mathcal{F}_n$ are contained realizations of the Dirichlet process with probability one, and these are almost surely discrete, we can look at the infinite mixtures of functions belonging to the set $\mathcal{S}$. We can therefore define $\mathcal{G}_n := \{z \mapsto g_F(z) = \sum_{i=1}^{\infty} \frac{\mathbf{1}_{x_i>z}}{2\sqrt{x_i^2 - x_i z}} F(x_i) \ : \ F \in \mathcal{F}_n\}$. Then $\mathcal{G}_n$ is a VC-hull of the class $\mathcal{S}$, as we showed previously that $\mathcal{S}$ is VC. Using Corollary 2.6.12 from [103] we have that for a constant K that depends only on the VC-index $V_m(\mathcal{G}_n)$:

$$\log N\left(\varepsilon \|g_*\|_{Q,2}, \mathcal{G}_n, \mathbb{L}_2(Q)\right) \leq K\left(\frac{1}{\varepsilon}\right)^{2-2V_m^{-1}(\mathcal{G}_n)} \tag{3.3.63}$$

Now consider the functions $f \mapsto \phi(f) := \frac{1}{f}$, then using (3.3.63) we have:

$$\begin{aligned} &\int_0^1 \sup_Q \sqrt{\log N\left(\varepsilon \|\mathbf{1}_{C_n} g_*^{-1}\|_{Q,2}, \phi(\mathcal{G}_n), \mathbb{L}_2(Q)\right)}\, d\varepsilon \\ &\leq \int_0^1 \sup_Q \sqrt{\log N\left(\varepsilon \|g_*\|_{Q,2}, \mathcal{G}_n, \mathbb{L}_2(Q)\right)}\, d\varepsilon < \infty \end{aligned} \tag{3.3.64}$$

where in the last inequality we used the fact that we can use Theorem 2.10.20 from [103] as $\sup_{z\in[0,x+\eta_n]} |g_F^{-4}(z)| < \infty$ uniformly over $\mathcal{F}_1$, and thus:

$$|\phi \cdot g_{F_1}(z) - \phi \cdot g_{F_2}(z)|^2 \leq \left\| \sup_{z\in[0,x+\eta_n]} |g_F^{-4}(z)| \right\|_{\mathcal{F}_1} |g_{F_1}(z) - g_{F_2}(z)|$$

We show:

$$J(1, \mathcal{U}_n, U_n^*) = \int_0^1 \sup_Q \sqrt{\log N\left(\varepsilon \|U_n^*\|_{Q,2}, \mathcal{U}_n, \mathbb{L}_2(Q)\right)}\, d\varepsilon < \infty. \tag{3.3.65}$$

Note that we can write any function $U_n^F \in \mathcal{U}_n$ as: $z \mapsto \frac{1}{D_*} \sum_{u_i>z} \frac{w(u_i)}{\sqrt{u_i - z}} F\{u_i\}$ for weights $w(u_i)$:

$$w(u_i) := \frac{1}{f_0(u_i)} \frac{\mathbf{1}_{\eta_n^* \leq u_i - x \leq \eta_n}}{\pi(u_i - x)}, \quad \text{and} \quad D_* = \sup_{F\in\mathcal{F}_1} \sum_i |w(u_i)| F(u_i)$$

where $\mathcal{F}_1$ denotes the weak support of the Dirichlet process prior. Note D_* is an appropriate renormalizing constant and that $D_* < \infty$. Because the $\mathbb{L}_1$ norm of the weights is bounded for every n, we conclude that the weights are absolutely summable in the sense: $(D^*)^{-1} \sum |w(u_i)| F\{u_i\} \leq 1$ for every n. (3.3.65) now follows from the same reasoning used for $\mathcal{G}_n$. Indeed the class of functions $\left\{z \mapsto ((u-z)_+)^{-1/2} \ : \ u \in \mathbb{R}^+\right\}$ is VC, thus taking its convex hull, using Corollary 2.6.12 from [103] and the fact that the weights are absolutely summable, gives (3.3.65).

We can now conclude by using the maximal inequality:

$$\mathrm{P}_0\left(r_n \left\| \frac{1}{n} \sum_{i=1}^n Y_{n,i}^F - E_0 Y_{n,i}^F \right\|_{\mathcal{F}_n} > \varepsilon \right)$$

$$= \mathrm{P}_0\left(\sqrt{n}\left\| \frac{1}{n}\sum_{i=1}^{n} \overline{V}^F_{n,i} - \mathrm{E}_0 \overline{V}^F_{n,i} \right\|_{\mathcal{F}_n} > \frac{\varepsilon\sqrt{n}}{r_n D^*} \right) \leq \frac{r_n}{\varepsilon\sqrt{n}} O\left(\sqrt{\log\frac{1}{\eta_n^*}} + \frac{1}{\eta_n} \right),$$

because

$$\begin{aligned} \|V_n^*\|^2_{G_0,2} &= \frac{1}{(D^*)^2}\int_0^L \frac{g_0(z)}{g_F^2(z)}\left(\int_z^L \frac{1}{\sqrt{u-z}} \frac{\mathbf{1}_{\eta_n^* \leq u-x \leq \eta_n}}{(u-x)} \frac{dF(u)}{f_0(u)} \right)^2 dz \\ &= \frac{1}{(D^*)^2}\left(O\left(\log\frac{1}{\eta_n^*} \right) + O(\eta_n^{-2}) \right). \end{aligned}$$

The last display converges to zero under G_0 proving (3.3.59).

The last step of the proof consists in showing $\forall\, t$ in a neighborhood of 0:

$$\frac{\displaystyle\int_{\mathcal{F}_n} \prod_{i=1}^{n} g_{F_{t\eta}}(Z_i)\, d\Pi(F)}{\displaystyle\int_{\mathcal{F}_n} \prod_{i=1}^{n} g_F(Z_i)\, d\Pi(F)} \xrightarrow{G_0} 1. \tag{3.3.66}$$

This condition entails the so-called "invariance" under changing the prior from $F \sim \Pi$ (in the denominator) to $F \circ \phi^{-1}$ (in the numerator) for $\phi : g_F \mapsto g_{F_{t\eta}}$.

If $F \sim \mathrm{DP}(\alpha)$, F is the normalized Gamma process, a *completely random measure*. So if $F = \frac{\Gamma}{|\Gamma|}$ the:

$$dF_t = \frac{(1+th^x)\, dF}{\int (1+th^x)\, dF} = \frac{(1+th^x)\, d\Gamma}{\int (1+th^x)\, d\Gamma} \tag{3.3.67}$$

Hence F_t is the normalized process $\overline{\Gamma}$ defined, for any measurable A, by:

$$\overline{\Gamma}(A) = \int_A (1+th^x)\, d\Gamma$$

Since F has base measure α, we prove that the process $\overline{\Gamma}$ has intensity measure:

$$\nu(ds, dv) = s^{-1} \exp\left\{ -\frac{s}{(1+th^x(v))} \right\} ds\, d\alpha(v)$$

A CRM is characterized by its intensity measure. For a CRM $\overline{\Gamma}$ without fixed atoms, this is the measure on $(0,\infty)\times\mathfrak{X}$ so that:

$$-\log \mathrm{E}\, e^{-\int f d\overline{\Gamma}} = \int_{\mathfrak{X}}\int_0^\infty \left(1 - e^{-sf(v)}\right) \nu(ds, dv)$$

for any measurable f which is integrable with respect to $\overline{\Gamma}$ almost surely. For the Gamma Process $\nu(ds, dv) = s^{-1}e^{-s}\, ds\, d\alpha(v)$. Because $\int f d\overline{\Gamma} = \int f(1+sh) d\Gamma$ it follows, by substituting $s(1+th) = u$:

$$-\log \mathrm{E}\, e^{-\int f d\overline{\Gamma}} = \iint \left(1 - e^{-sf(v)(1+th^x(v))}\right) s^{-1}e^{-s}\, ds\, d\alpha(v)$$

$$= \iint \left(1 - e^{-uf(v)}\right) \frac{(1 + th^x(v))}{u} \exp\left\{-\frac{u}{(1 + th^x(v))}\right\} \frac{du}{1 + th^x(v)}\, d\alpha(v)$$
$$= \iint \left(1 - e^{-sf(v)}\right) s^{-1} \exp\left\{-\frac{s}{(1 + th^x(v))}\right\} ds\, d\alpha(v)$$

which proves the claim.

Now for $X_1, \ldots, X_n \mid F \sim F_t$, for any measurable functions $\varphi_1, \ldots, \varphi_n$ (taken to be $\varphi(Z_i, \cdot)$ in our case):

$$\mathrm{E}\Big(\prod_{i=1}^{n} \varphi_i(X_i)\Big) = \mathrm{E}\,\mathrm{E}\Big(\prod_{i=1}^{n} \varphi_i(X_i) \mid F\Big) = \mathrm{E}\prod_{i=1}^{n} \int \varphi_i\, dF_t$$

therefore in our case, the functions $\varphi(Z_i, \cdot)$ are given as follows:

$$\frac{\int_{\mathcal{F}_n} \prod_{i=1}^{n} g_{F_{t_n}}(Z_i)\, d\Pi(F)}{\int_{\mathcal{F}_n} \prod_{i=1}^{n} g_F(Z_i)\, d\Pi(F)} = \frac{\int_{\mathcal{F}_n} \prod_{i=1}^{n} \int_{Z_i}^{\infty} \frac{dF_{t_n}(u)}{2\sqrt{(u)^2 - uZ_i}}\, d\Pi(F)}{\int_{\mathcal{F}_n} \prod_{i=1}^{n} \int_{Z_i}^{\infty} \frac{dF(u)}{2\sqrt{(u)^2 - uZ_i}}\, d\Pi(F)}$$
$$= \frac{\mathrm{E}\left[\prod_{i=1}^{n} \frac{\mathbf{1}_{X_{i,n} > Z_i}}{2\sqrt{(X_{i,n})^2 - X_{i,n}Z_i}}\right]}{\mathrm{E}\left[\prod_{i=1}^{n} \frac{\mathbf{1}_{X_i > Z_i}}{2\sqrt{(X_i)^2 - X_i Z_i}}\right]} \tag{3.3.68}$$

Let $\mathscr{S}_n$ stand for the collection of all partitions of $\{1, \ldots, n\}$ that emerge from the ties obtained by sampling from a Dirichlet process, and write a partition as $\mathcal{S} = \{S_1, \ldots, S_N\}$, for $N = \#\mathcal{S}$ the number of partitioning sets. Then using Proposition 4.7 from [42] the denominator of (3.3.68) can be written as:

$$\frac{1}{|\alpha|^{[n]}} \sum_{\mathcal{S} \in \mathscr{S}_n} \prod_{j=1}^{\#\mathcal{S}} \left[\Big(\#S_j - 1\Big)! \int \prod_{l \in S_j} \varphi_l(v) d\alpha(v)\right]. \tag{3.3.69}$$

Now we are going to give both an upper bound and lower bound for the numerator in (3.3.68) and we are going to show that both the upper bound and the lower bound, divided by (3.3.69), go to 1.

For $X_1, \ldots, X_n \mid F \sim F_t$, for any measurable functions $\varphi_1, \ldots, \varphi_n$:

$$\mathrm{E}\prod_{i=1}^{n} \varphi_i(x_i) = \tag{3.3.70}$$
$$\sum_{\mathcal{S} \in \mathscr{S}_n} \int \cdots \int \prod_{j=1}^{k} \prod_{l \in S_j} \varphi_l\left(v_j\right) \int_0^{\infty} \frac{\lambda^{n-1}}{\Gamma(n)} e^{-\psi(\lambda)} \prod_{j=1}^{k} \tau_{|s_j|}\left(\lambda \mid v_j\right) d\lambda d\alpha(v_1) \cdots d\alpha(v_k)$$

where:

$$\psi(\lambda) = \int \log\left(1 + \lambda(1 + th^x(v))\right) d\alpha(v)$$

$$\tau_{|N|}(\lambda \mid v) = \Gamma(N)\left(\frac{1+th^x(v)}{1+\lambda(1+th^x(v))}\right)^N$$

The idea is that the realizations of F_t are discrete, whence $X_1, \ldots, X_n$ can be grouped in sets of equal values. The joint distribution of the random partitions and values is given by Lemma 14.62 in [42]. In order to prove (3.3.70) by using Lemma 14.62 in [42] we need to compute:

$$\begin{aligned}
\psi(\lambda) &= \iint \left(1-e^{-\lambda s}\right)\nu(dv,ds) = \\
&= \iint \left(1-e^{-\lambda s}\right)s^{-1}e^{-\frac{s}{1+th^x(v)}}\,ds\,d\alpha(v) \\
&= \iint \left(1-e^{-\lambda(1+th^x(v))s}\right)s^{-1}e^{-s}\,ds\,d\alpha(v) = \\
&= \int \log(1+\lambda(1+th^x(v)))\,d\alpha(v)
\end{aligned}$$

where by example J.7 in [42], for $\nu^c(dv,ds) = s^{-1}e^{-s}\,ds\,d\alpha(v)$ we have for $\theta > 0$:

$$\iint (1-e^{-\theta s f(v)})\,\nu^c(dv,ds) = \int \log(1+\theta f(v))\,d\alpha(v).$$

Moreover using $\int_0^\infty s^{n-1}e^{-(c+k)s}\,ds = \Gamma(n)(c+k)^{-n}$:

$$\begin{aligned}
\tau_N(\lambda \mid v) &= \int s^N e^{-\lambda s}\,\rho(ds \mid v) = \int s^N e^{-\lambda s}s^{-1}\exp\left\{-\frac{s}{(1+th^x(v))}\right\}ds \\
&= \Gamma(N)\frac{1}{(\lambda+\frac{1}{(1+th^x(v))})^N}.
\end{aligned}$$

We conclude that (3.3.70) holds true and as a special case for $t = 0$ we retrieve $F_t = F \sim \mathrm{DP}(\alpha)$, which gives $\psi(\lambda) = (\log(1+\lambda))|\alpha|$ and $\tau_N(\lambda \mid v) = \Gamma(N)/(1+\lambda)^N$ and it is possible to prove that (3.3.70) reduces to (3.3.69).

Now for $t > 0$ small, using that $\frac{1}{1+\lambda(1-r)} \le \frac{1+r}{(1-r)(1+\lambda(1+r))}$ we have:

$$\begin{aligned}
&\int_0^\infty \lambda^{n-1}e^{-\psi(\lambda)}\prod_{j=1}^k \frac{\tau_{|S_j|}\left(\lambda \mid v_j\right)}{\Gamma\left(\left|S_j\right|\right)}d\lambda = \\
&= \int_0^\infty \lambda^{n-1}e^{-\int \log(1+\lambda(1+th^x(v)))\,d\alpha(v)}\prod_{j=1}^k\left(\frac{1+th^x\left(v_j\right)}{1+\lambda\left(1+th^x\left(v_j\right)\right)}\right)^{|S_j|}d\lambda \\
&\le \int_0^\infty \lambda^{n-1}\frac{1}{(1+\lambda\,(1-t\|h\|_\infty))^{|\alpha|}}\left(\frac{1+t\|h\|_\infty}{1+\lambda\,(1+t\|h\|_\infty)}\right)^n d\lambda \\
&\le \left(\frac{1+t\|h\|_\infty}{1-t\|h\|_\infty}\right)^{|\alpha|}\int_0^\infty \frac{\lambda^{n-1}\,(1+t\|h\|_\infty)^n}{(1+\lambda\,(1+t\|h\|_\infty))^{|\alpha|+n}}d\lambda \\
&= \left(\frac{1+t\|h\|_\infty}{1-t\|h\|_\infty}\right)^{|\alpha|}\frac{\Gamma(|\alpha|)\Gamma(n)}{\Gamma(|\alpha|+n)}.
\end{aligned}$$

The lower bound is given similarly by:

$$\int_0^\infty \lambda^{n-1}e^{-\psi(\lambda)}\prod_{j=1}^k \frac{\tau_{|S_j|}\left(\lambda \mid v_j\right)}{\Gamma\left(\left|S_j\right|\right)}d\lambda \ge \left(\frac{1-t\|h\|_\infty}{1+t\|h\|_\infty}\right)^{|\alpha|}\frac{\Gamma(|\alpha|)\Gamma(n)}{\Gamma(|\alpha|+n)}.$$

Therefore we can upper-bound (and similarly lower-bound) (3.3.70) by:

$$\left(\frac{1+t\|h\|_\infty}{1-t\|h\|_\infty}\right)^{|\alpha|}\frac{1}{|\alpha|^{[n]}}\sum_{\mathcal{S}\in\mathscr{S}_n}\prod_{j=1}^{\#\mathcal{S}}\Big[\int\prod_{l\in S_j}\varphi_l(v)\underbrace{\prod_{i}^{N}\Gamma(|S_j|)}_{=(\#S_j-1)!}d\alpha(v)\Big].$$

because t was chosen to depend on n in the way: $t_n = tr_n/n$ and $\|h_n^x\|_\infty = O(\sqrt{n})$, we conclude that: $(1-O(1/\sqrt[4]{n}))^{|\alpha|} \le (3.3.68) \le (1+O(1/\sqrt[4]{n}))^{|\alpha|}$ which proves (3.3.66).

□

□

***Proof of contraction rate in Wasserstein metric, Theorem** 3.2.2.c.* We follow the structure of an analogous result given in [39]. The first thing is to upper bound the Wasserstein distance in terms of the Hellinger distance and this is done in Lemma 3.3.14 below.

We establish the contraction rate for the "discrepancy" $d(g_F, g_0) = \gamma(h(g_F, g_0))$ for $\varepsilon \mapsto \gamma(\varepsilon) = D_k\varepsilon^{1/(k+2)}[\log(C_k/\varepsilon)]^{\frac{k+1/2}{k+2}}$. Note that $\gamma^{-1}(d) \le h(g_F, g_0)$ and it possesses the properties (a)-(d) in Section 9 of [39] and the balls of d are convex. We can therefore apply Theorem 3 from [39] to obtain a rate of posterior contraction relative to d and thus relative to $h^{1/(k+2)}(\log(1/h))^{\frac{k+1/2}{k+2}}$. We apply the theorem with the same set $\mathcal{P}_n$ as in the proof of Theorem 3.2.2.a.. Thus by Proposition 3.3.1:

$$\log N(\varepsilon,\mathcal{P}_n,d) = \log N\Big(\varepsilon^{\frac{1}{k+2}}\Big(\log\frac{1}{\varepsilon}\Big)^{\frac{k+1/2}{k+2}},\mathcal{P}_n,h\Big) \lesssim \Big(\frac{1}{\varepsilon}\Big)^{\frac{2}{k+2}}\Big(\log\frac{1}{\varepsilon}\Big)^{\frac{2k+1}{k+2}}.$$

Therefore condition (9.1) from Theorem 3 in [39] holds for the rate $\varepsilon_n \gtrsim n^{-\gamma}$ and any $\gamma < (k+2)/(2k+4)$. The prior mass condition proved in Proposition 3.3.2 is satisfied with the rate $\varepsilon_n \asymp (\log n/n)^{1/4}$. The slowest of the two rates is $\varepsilon_n \asymp (\log n/n)^{1/4}$ which, by substitution, yields a rate:

$$\varepsilon_n^{2/(k+2)}(\log(1/\varepsilon_n))^{\frac{2k+1}{k+2}} \asymp n^{-1/(2k+4)}(\log n)^{(4k+3)/(2k+4)}.$$

Theorem 3.2.2.b. gives the contraction rate of $|V_F(0)-V_0(0)|$ and it is faster than the contraction rate for d. The latter is thus the dominating contraction rate. □

Lemma 3.3.14. *For probability measures F and F_0 supported on $[0,L]$, and $g_F = \int\psi\,dF$ for ψ density as in* (3.1.1) *and for $x \ge 0$, $F^*(x) = 1 - V_F(x)/V_F(0)$ and similarly $F_0^*(x)$ as in* (3.1.2) *with $V_0(0) > 0$, then for any $k \ge 1$,*

$$W_k\left(F^*,F_0^*\right) \lesssim h\left(g_F,g_0\right)^{1/(k+2)}\left(\log\frac{C_k}{h\left(g_F,g_0\right)}\right)^{\frac{(k+1/2)}{(k+2)}} + |V_F(0)-V_0(0)|.$$

where C_k is a constant such that the map $\varepsilon \mapsto \varepsilon\left[\log(C_k/\varepsilon)\right]^{k+1/2}$ is monotone on $(0,2]$.

Proof. The proof of this lemma follows very closely the proof of Lemma 7 in [39], with some adaptations for our case. The strategy consists of approximating F^* and F_0^* by convolving them with a mollifier. Using Theorem 6.15 from [110], $W_k\left(F^*,F_0^*\right) \lesssim \left(\int|x|^k\,d|F^*-F_0^*|(x)\right)^{1/k}$, where $|F^*-F_0^*|$ is the total variation measure of $F^*-F_0^*$. For a measure $\mu = \mu^+-\mu^-$, $|\mu| = \mu^++\mu^-$,

where respectively μ^+ and μ^- are the positive and negative variation of μ. We apply the above upper bound to the convolutions of F^* and F_0^* with the Gaussian mollifier $\phi_\delta(x) = \frac{1}{\delta\sqrt{2\pi}} e^{-\frac{x^2}{2\delta^2}}$ (denote by Φ_δ the normal distribution). For any $M > 0$ and Z standard normal variable:

$$\begin{aligned} W_k(F * \Phi_\delta,\, F_0 * \Phi_\delta)^k &\lesssim \int |x|^k \, d\left|\left(F^* - F_0^*\right) * \Phi_\delta\right|(x) \\ &\leq \frac{|V_F(0) - V_0(0)|}{|V_F(0)V_0(0)|} \int |x|^k \, d|V_F * \Phi_\delta|(x) + \frac{1}{V_0(0)} \int |x|^k \, d\,|(V_F - V_0) * \Phi_\delta|\,(x) \\ &\lesssim |V_F(0) - V_0(0)| + \int |x|^k \, d\,|(V_F - V_0) * \Phi_\delta|\,(x) \end{aligned}$$

where we used that $V_F(0), V_0(0) > 0$ and that $\int |x|^k \, d|V_F * \Phi_\delta|(x) < \infty$, $\forall\, \delta > 0$. Now we analyze separately the second expression in the inequality above:

$$\begin{aligned} \int |x|^k \, d\,|(V_F - V_0) * \Phi_\delta|\,(x) &= \int |x|^k \;|(V_0 - V_F) * \phi_\delta(x)| \; dx \\ &\leq \left(\int_{-M}^{M} x^{2k} dx \int_{-M}^{M} |(V_0 - V_F) * \phi_\delta(x)|^2 \, dx \right)^{1/2} \\ &\quad + e^{-M} \int_{|x|>M} |x|^k e^{|x|} \, |(V_0 - V_F) * \phi_\delta(x)| \, dx \\ &\lesssim M^{k+\frac{1}{2}} \, \|(V_0 - V_F) * \phi_\delta\|_2 + e^{-M} \max\{1, e^{2|L|}\} \mathrm{E} e^{2|\delta Z|}, \end{aligned}$$

where we used that the measures V_F and V_0 are supported on $[0, L]$, the fact for differentiable functions f and g in $\mathbb{L}_1$, $\frac{d}{dx}(f * g)(x) = (\frac{d}{dx} f(x)) * g(x)$ and the fact for a measure that is absolutely continuous with respect to the Lebesgue measure we have $\frac{d}{d\lambda}|\mu| = |\frac{d\mu}{d\lambda}|$, where λ is the Lebesgue measure and $|\mu|$ the total variation measure. Note also that the number $\max\{1, e^{2|L|}\}\mathrm{E} e^{2|\delta Z|}$ is uniformly bounded. The reason why we upper bounded the distance $W_k(F^* * \Phi_\delta, F_0^* * \Phi_\delta)$ with an expression in terms of V_0 and V_F is that we can express the density of the observations g_F as a convolution with respect to the measure V_F: $g_F = k * V_F$ for $k(x) = -\mathbf{1}_{\{x<0\}}/\pi\sqrt{-x}$. This will allow us to use tools like the convolution theorem for Fourier transforms and Plancherel's theorem. We denote the Fourier transform $\mathcal{F}$ as follows $f \mapsto \mathcal{F}(f) = \hat{f}$. Then:

$$\begin{aligned} \|(V_0 - V_F) * \phi_\delta\|_2^2 &= \int \left|\hat{V}_0 - \hat{V}_F\right|^2 (\lambda) \hat{\phi}_\delta^2(\lambda) d\lambda \\ &= \int \left|\hat{f}(\hat{V}_0 - \hat{V}_F)\right|^2 (\lambda) \frac{\hat{\phi}_\delta^2}{|\hat{f}|^2}(\lambda) \, d\lambda = \left\|(g_F - g_0) * \phi_\delta * \mathcal{F}(\hat{f}^{-1})\right\|_2^2 \\ &\leq 4 d_{TV}^2(g_F, g_0) \, \|\phi_\delta * \mathcal{F}(\hat{f}^{-1})\|_2^2 \lesssim h^2\,(g_F, g_0)\, \delta^{-2}, \end{aligned}$$

where d_{TV} is the total variation distance. Above we used a known fact (cf. [77] pp. 391) that for an arbitrary (signed) measure μ on $\mathbb{R}$ and a function $g \in \mathbb{L}_2(\mathbb{R})$, there holds $\|\mu * g\|_2 \leq |\mu| \|g\|_2$ ($|\mu|$ total variation measure). By Plancherel's theorem and that $|\hat{f}(\lambda)|$ is proportional to $1/\sqrt{|\lambda|}$,

$$\int \frac{\hat{\phi}_\delta^2(\lambda)}{|\hat{f}(\lambda)|^2} d\lambda \lesssim \int \lambda e^{-\delta^2\lambda^2} \, d\lambda \lesssim \delta^{-2}.$$

If $U \sim F^*$ is independent of $Z \sim N(0,1)$, then $(U, U + \delta Z)$ gives a coupling of F^* and $F^* * \phi_\delta$. Under this coupling $\mathrm{E}|U + \delta Z - U|^k = O(\delta^k)$ which implies $W_k\,(F^*, F^* * \phi_\delta)^k \leq \mathrm{E}|\delta Z|^k \lesssim \delta^k$.

By combining the preceding inequalities we see that, for any $M > 0$:

$$\begin{aligned} W_k\left(F^*, F_0^*\right)^k &\leq W_k(F^*, F_0^* * \phi_\delta)^k + W_k(F_0^* * \phi_\delta, F^* * \phi_\delta)^k + W_k(F^* * \phi_\delta, F_0^*)^k \\ &\lesssim M^{k+1/2} h\left(g_F, g_0\right) \delta^{-2} + e^{-M} + \delta^k \end{aligned}$$

As done in [39], the lemma follows by choosing for $\varepsilon = h\left(g_F, g_0\right)$ the optima $M = k/(k + 1/2) \log\left(C_k/\varepsilon\right)$ and $\delta = \left(M^{k+2}\varepsilon\right)^{1/(k+1/2)}$, which are eligible by assumption. □

3.4 Bibliographic remarks

There is a vast literature on Wicksell's problem, see for instance [19, 27, 43, 48, 49, 52, 53, 56, 62, 92, 99, 114]. Wicksell's problem has present-day applications, where it is common to see the use of the so-called Saltykov method [22], and numerical discretization methods [72], often lacking theoretical guarantees. These methods are far from being efficient and in some cases not even consistent. The estimation of V_0 given data from g_0 is of interest not only for stereological procedures, but also from the mathematical point of view, both because of the unusual rates of convergence and because of the non-standard efficiency theory.

In the nonparametric frequentist approach to Wicksell's problem, the standard rate of convergence for pointwise asymptotic normality in estimating V_0 is $(\log n)^{-1/2} \sqrt{n}$ (cf. [43, 52, 114]). The first time the naive plug-in estimator was studied is in [114]. Later in [52] the proof of the asymptotic normality of this estimator was simplified considerably. However, viewed as a function, the naive plug-in estimator is severely ill-behaved due to infinite discontinuities and non-monotonicity, and this led to the definition of the IIE in [43, 52]. The Nonparametric Maximum Likelihood Estimator is studied in [19, 62]. For Wicksell's problem, the usual definition of the NPMLE is also ill-posed, as the log-likelihood can attain value infinity. However, work has been done, in [19] and [62], to obtain likelihood based estimators. Confidence interval results can be found in [27, 92]. The results of [27] provide confidence bands by developing a methodology for isonization procedures, which often lead to nuisance parameters in the limiting distributions, that tackles this issue. However, they do not cover the setting considered in the present chapter. Extending their results to our case appears non-trivial, as their proof techniques rely on at least the differentiability of the estimated object, an assumption we do not make. The bootstrap results in [92] are more directly relevant and will be discussed in detail in the simulation section, including a comprehensive numerical comparison. The main advantage of the methodology proposed in this chapter, if compared to the bootstrap methods of [92], is that it relies on significantly less stringent assumptions. In particular, our results do not depend on the assumption that the underlying distribution function F_0 is differentiable at x. Instead, our methodology is provably consistent for all local smoothness levels $\gamma > 1/2$ of V_0, where γ is defined in condition (3.1.20) below.

In [49], the authors investigate efficient estimation in the classical Wicksell problem, focusing on the scenario where the distribution function F_0^* of interest belongs to a Hölder class of smoothness with degree $\gamma > \frac{1}{2}$. They propose a kernel density-based estimator that, under appropriate bandwidth selection, achieves asymptotic efficiency. However, a notable limitation is that the estimator requires prior knowledge or guesstimation of the smoothness parameter γ, making it non-adaptive. Moreover, efficiency is only guaranteed under global smoothness assumptions, i.e., the entire distribution function must belong to a global Hölder class. Despite these limitations, their work remains the first foundational contribution to semiparametric efficiency in Wicksell's problem. [43] builds upon earlier work of [49] and

[52] by addressing both the adaptivity and locality of efficient estimation in Wicksell's problem. The key innovation consists in showing that the Isotonized Inverse Estimator (IIE) achieves asymptotic efficiency under the same local smoothness assumptions on the target function F_0^* of the present chapter. Specifically, instead of assuming that $F_0^* \in \Lambda_\gamma$ globally as in [49], the analysis in [43] only requires that F_0^* satisfies local Hölder-type smoothness at the evaluation point x and at the origin. The estimator adapts automatically to the local regularity of F_0^*, offering a more flexible framework for practitioners. Additionally, [43] provides a local asymptotic minimax lower bound, showing that the IIE attains the asymptotically optimal risk in this refined setting. The estimator thus achieves the same efficiency as the KDE-based approaches, but without requiring tuning of smoothness parameters, and without imposing strong global assumptions. Finally, other results concerning efficient estimation in Wicksell's problem are contained in [44] and [48]. [44] shows that in the case the underlying cdf is constant on a *known* interval of strictly positive length the IIE evaluated at x is not efficient, for x in the interval of constancy. However, it behaves in practice very closely to informed estimators that make use of the fact that the cdf is constant on that interval. Finally, one of the authors of [49] did some further work in [48] where it was shown how to attain efficient smoothing for the plug-in estimator under squared losses.

While Bayesian methods for inverse problems are widely adopted, their theoretical guarantees remain scarce, particularly in infinite-dimensional settings. Although recent years have seen significant advances in this area [14–16, 57, 66, 67, 76, 78, 79, 84, 85, 88], the technical frameworks developed there do not readily apply to the specific inverse problem and Dirichlet process prior setting of this chapter. We provide a detailed discussion of this related work in Supplement [46]. A similar methodology to the one proposed in this chapter — using an unconstrained conjugate prior for the distribution function of the observables and projecting the posterior samples into the space of monotone functions — has been very recently adopted in isotonic regression settings (c.f. [17, 112, 113]), isotonic density estimation (c.f. [18]) and ODEs settings (c.f. [5, 6]). [5, 6] develop Bayesian two-step methods for parameter inference in ODE models by embedding them in a nonparametric regression framework with B-spline priors. The parameter posterior is obtained by minimizing a discrepancy between the estimated regression function and the ODE solution, either via derivative matching or Runge-Kutta approximations. [6] further establishes a Bernstein-von Mises theorem for the Runge-Kutta-based method, showing that the resulting Bayes estimator is asymptotically efficient. However, their framework relies on a very different type of prior modelling and the projection method does not entail isotonization. In [17], the authors study a nonparametric Bayesian regression model for a response variable Y with respect to a predictor variable $X \in [0, 1]$, given by $Y = f(X) + \varepsilon$, where f is a monotone increasing function and ε is a mean-zero random error. They propose a "projection-posterior" estimator f^*, constructed using a finite random series of step functions with normal basis coefficients as a prior for f. Their main finding is that when f^* is centered at the maximum likelihood estimator (MLE) and rescaled appropriately, the Bernstein-von Mises (BvM) theorem does not hold. However, when f^* is instead centered at the true function f_0, a BvM result is obtained. This phenomenon resonates with the known inconsistency of the bootstrap for the Grenander estimator in isotonic regression (c.f. [93]) and represents a key difference with the present work. In our case, the "projection-posterior" IIP $\hat{V}_G$, when centered at the IIE $\hat{V}_n$, is shown to converge in distribution. This may also not be surprising given the existing results in [92], where bootstrap methods are shown to be consistent for Wicksell's problem.

Recent research contributions [14–16, 57, 66, 67, 76, 78, 79, 84, 85, 88] have introduced proof techniques aimed at establishing theoretical guarantees for inference based on posterior

distributions in a variety of models. These techniques rely on appropriately chosen priors and are applicable across some contexts. [14] and [15] made significant advances in understanding the Bernstein–von Mises (BvM) phenomenon in nonparametric Bayesian analysis. Specifically, [14] provided foundational results in the Gaussian white noise model, while [15] extended these findings to more general settings, introducing innovative tools such as multiscale spaces, which naturally accommodate certain nonparametric priors and posteriors. However, their results do not cover the setting of the current chapter, as the theoretical tools there developed do not naturally accommodate DP priors in inverse settings. [16] extended the BvM theorem to semiparametric models, focusing on the asymptotic behavior of posterior distributions for smooth functionals of the parameter of interest. However, their methodology is not applicable to our setting for two key reasons. First, the primary functional of interest in our work, V_0, which is essentially the $1/2$-integral of g_0, is not a smooth functional of g_0. This is reflected in the fact that, since their framework applies to smooth functionals, they obtain $\sqrt{n}$ convergence rates, whereas we obtain the slower $\sqrt{n}/\sqrt{\log n}$ rate. Second, the study of the so-called *least favorable direction* in our context, as conducted in [43], does not lead to the classical Laplace expansion of the posterior (cf. Eq. (2.13) in [16]). This discrepancy arises because the LAN expansion for Wicksell's model involves a local functional, which does not yield the global $\mathbb{L}_2$ contractions needed. [57] provides a comprehensive treatment of quantile estimation under the DP prior; however, their results do not involve any inverse setting. [66] and [67] establish semiparametric and nonparametric BvM results in the proportional hazards and right-censoring models, respectively. While they develop highly useful techniques for handling DP process priors in survival analysis, these methods are difficult to generalize beyond that framework. In addition, their approaches do not employ projection-based posterior methodologies, as used in the present chapter. [76] investigates the statistical inverse problem of recovering a function f from general X-ray transform measurements corrupted by Gaussian noise. The proof techniques of this chapter rely on the fact that the noise is Gaussian and that the used prior is a Gaussian process, making it difficult to extend them to our setting (with beta noise and DP process prior). [78] considers the inverse problem of estimating the potential in a Schrödinger equation while the solution of the equation is corrupted by Gaussian noise, while [79] considers the inverse problem of recovering the parameters governing compound Poisson processes. In both settings, the BvM results are based on ad-hoc priors constructed with wavelet basis functions, which is far from the DP modelling we use in the current chapter. [84] establishes a BvM result in the Gaussian white model with priors constructed with wavelet basis functions and [85] for the Average Causal Effect in a causal setting, using Gaussian process priors. The latter uses techniques closely related to [16], which cannot be applied to our setting. Finally [88] also uses wavelet-based priors to derive semiparametric BvM results for linear functionals of densities. While the approach of this chapter constitutes a relevant advancement in the semiparametric BvM theory, the inverse problem treated in the current work cannot be framed in the context of [88].

One of the key advantages our approach shares with the works in [5, 6, 17, 112, 113] is the computational efficiency, driven by posterior conjugacy, as well as the relative simplicity of the asymptotic analysis. However, unlike in [17], our method ensures that the limiting distribution in the Bernstein-von Mises (BvM) theorem coincides with that of the asymptotically efficient estimator — the IIE of [43].

Chapter 4

Semiparametric Uncertainty Quantification via Isotonized Posterior for Deconvolutions

4.1 Introduction

In many statistical applications, one encounters situations where the observed data Z_i arises from the sum of two independent components: a variable of interest X_i, which is usually the quantity we are interested in, and another variable Y_i whose distribution is assumed to be known or to be inferred using expert knowledge.

This setup applies to many different areas of research. For example, X_i might represent the true underlying quantity we aim to measure in an experiment or measurement, while Y_i captures the random measurement error introduced during the process — and which is assumed to be additive in nature. In epidemiology, X_i could correspond to the actual time of infection, and Y_i to the incubation period. This framework plays a key role in approaches like back-calculation, which has been used in HIV/AIDS research to estimate historical infection rates based on observed diagnoses [4]. The statistical properties of Y_i vary significantly depending on the context: measurement errors are often modeled with symmetric distributions over the entire real line, such as the normal distribution, while durations — like incubation periods or the time until a component fails — are typically modeled with skewed, non-negative distributions, such as the Gamma or Weibull distributions.

This method of separating the distribution of X_i from the one of Y_i, which is called deconvolution, is also key in other fields. For example, in image processing, a blurry image Z can be thought of as the true, sharp image X distorted by a blur effect Y caused by the camera or optics. Methods like the Richardson-Lucy algorithm are used to try and recover the original sharp image X [73, 87]. Similarly, in astronomical spectroscopy, the spectrum of light observed from a star or galaxy Z is the object's actual spectrum X that has been spread out or blurred by the telescope's instruments represented by additive noise Y. Deconvolution helps scientists recover the true details about the celestial object [8]. Another application is in seismology, where the seismic waves recorded from underground Z are a combination of the Earth's rock layer structure X and the seismic signal Y that was sent out or generated. Deconvolution is thus a fundamental step to get clearer images of the Earth's subsurface [89].

More formally, throughout this chapter we confine attention to the case where both X and

Y are nonnegative random variables, i.e. we work on the half-line $[0, \infty)$. Let $X_1, X_2, \ldots, X_n$ denote a sample from an unknown distribution with distribution function F_0 supported on $[0, \infty)$ and, independent of that sample, $Y_1, Y_2, \ldots, Y_n$ a sample from a known distribution with density k on $[0, \infty)$. Consider the problem of estimating F_0 based on the sample $Z_1, Z_2, \ldots, Z_n$, where $Z_i = X_i + Y_i$. The density g_0 of Z_i is the convolution of k and F_0 in the following sense:

$$g_0(z) = \int_0^\infty k(z - x)\, dF_0(x) =: k * dF_0(z).$$

For this reason, this estimation problem is known as a deconvolution problem.

Suppose that, given a kernel k, we have a function p living on $[0, \infty)$, solving the integral equation

$$(p * k)(x) := \int_0^x p(x - y)k(y)\, dy = (\mathbf{1} * \mathbf{1})(x) = x\mathbf{1}(x), \tag{4.1.1}$$

where the function $\mathbf{1}$ is defined by

$$\mathbf{1}(x) = \mathbf{1}_{[0,\infty)}(x).$$

Then we can write, for each $x > 0$ and a Z having density function $g_0 = k * dF_0$,

$$\mathrm{E}[p(x-Z)] = (p*g_0)(x) = (p*k*dF_0)(x) = (\mathbf{1}*\mathbf{1}*dF_0)(x) = \int_0^x F_0(s)\, ds =: H_0(x). \tag{4.1.2}$$

A crucial question concerning integral equation (4.1.1) is whether a function p exists that satisfies it. in the Appendix (Section 4.5) 4.5.2, we give a review of the literature on this topic.

We start by placing a Dirichlet Process prior directly on the distribution function of the observables: $G \sim \mathrm{DP}(\alpha)$. Given an observed sample $Z_1, \ldots, Z_n$ from cdf G_0 (associated with density g_0), the posterior distribution is (c.f. [34], Theorem 4.6 in [42]):

$$G \mid Z_1, \ldots, Z_n \sim \mathrm{DP}(\alpha + n\mathbb{G}_n), \qquad \mathbb{G}_n := \frac{1}{n}\sum_{i=1}^n \delta_{Z_i}. \tag{4.1.3}$$

This choice offers computational simplicity due to the conjugacy of the posterior. Leveraging the inverse relations inherent to the problem, we define, by analogy with (4.1.2), the functional H_G (with G in place of G_0) as

$$H_G(x) = \int_{[0,x)} p(x - z)\, dG(z). \tag{4.1.4}$$

Similarly, the empirical counterpart of the left-hand side of (4.1.2) is given by a sample mean:

$$H_n(x) = \int_{[0,x)} p(x - z)\, d\mathbb{G}_n(z) = \frac{1}{n}\sum_{i=1}^n p\,(x - Z_i)$$

This function H_n is an estimator for H_0.

Taking the derivative of some smoothed version of H_n or H_G (H_n and H_G will in general not be differentiable) would therefore yield methodologies for the estimation of F_0. We call such a methodology an inverse methodology, since it is based on the inverse relation

$$F_0(x) = \frac{d}{dx} p * g_0(x) \quad \text{a.e.}$$

which follows from (4.1.2). However, by using general smoothing techniques, for example kernel estimation, we would not use the information that H_0 is convex (which follows from (4.1.2) and the monotonicity of F_0). Consequently, inverse estimators based on standard smoothing techniques will in general yield non-monotone estimators.

To remedy this, we isotonize the posterior draws using the following procedure. For $T \in (0, \infty]$ and a continuous function $h : [0, \infty) \to [0, \infty)$, the Greatest Convex Minorant (GCM) of h on $[0, T]$ and its right-hand side derivative, are given for $x \in [0, T]$ by:

$$h^\star(x) := \sup\{f(x) \,:\, f \leq h \text{ on } [0,T],\ f \text{ convex on } [0,T]\},$$
$$(h^\star)'_+(x) := \lim_{t \to 0^+} \frac{h^\star(x+t) - h^\star(x)}{t}.$$

For any $x \in [0, T]$, we define the isotonizations of the draws of the posterior as the right derivative of $H_G^\star$ evaluated at x,

$$\hat{F}_G(x) = (H_G^\star)'_+(x). \tag{4.1.5}$$

The posterior draws $\hat{F}_G$ are now by construction monotone (isotonic with respect to natural ordering on $\mathbb{R}$), and therefore we call the posterior *Isotonic Inverse Posterior*. Using the same procedure we define the *Isotonic Inverse Estimator* of [109], for any $x \in [0, T]$, as:

$$\hat{F}_n(x) = (H_n^\star)'_+(x). \tag{4.1.6}$$

One possible choice for T in the definition of $\hat{F}_G$ or $\hat{F}_n$ is $T = \infty$. As we will see in Section 4.2, we need finiteness of T in order to prove our asymptotic distribution result for a large class of densities k. If k is assumed monotone on $[0, \infty)$, our result does allow the choice $T = \infty$. For practical purposes, there is no difference between finite (but large) and infinite T.

In the domain of nonparametric inference, frequentist methods often yield estimators with well-characterized asymptotic distributions (e.g., the Chernoff distribution for isotonic regression, c.f. [53]). These distributions are crucial for constructing asymptotic confidence intervals and understanding the estimator's behavior. However, a significant practical limitation arises because these limiting distributions generally depend on unknown nuisance parameters. For instance, in our deconvolution problem of estimating $F_0(x)$ from samples $Z_1, \ldots, Z_n \sim g_0 = k * dF_0$, the limiting distribution depends on quantities such as $g_0(x)$ and, more critically, $f_0(x)$ (in a similar way as in the isotonic regression setting where the limit distribution often involves the derivative of the true underlying function). The necessity of estimating these nuisance parameters complicates the direct application of these asymptotic results for uncertainty quantification. This practical difficulty highlights the desirability of methods that can provide uncertainty quantification without relying on the estimation of such nuisance parameters. Bayesian nonparametric procedures offer a promising alternative in this regard. As explored in approaches in the "projection-posterior" literature (c.f. [17, 112, 113]), a key advantage is that the resulting credible intervals can achieve asymptotic frequentist coverage that is free of these nuisance parameters. The limiting coverage in such Bayesian frameworks can depend only on the chosen credibility level, not on characteristics of the true unknown function like its derivative $f_0(x)$. This circumvents the harder problem of nuisance parameter estimation, leading to more direct and more robust inferential procedures for quantifying uncertainty in nonparametric models. Furthermore, another possible Bayesian approach would consist in placing a DP process prior directly on F. However this procedure has some clear computational disadvantages due to the needed Gibbs sampling procedures to sample from the posterior.

Thus the approach in this chapter has two advantages:

(i) **Computational speed**: The Dirichlet posterior (4.1.3) is explicit and its projection can be effectively computed by the Pool-Adjacent-Violators Algorithm (PAVA, c.f. [4]).

(ii) **Uncertainty quantification**: The Bayesian framework inherently provides uncertainty quantification without the need to estimate nuisance parameters. We show below that this is asymptotically correct in the frequentist sense, after an appropriate calibration.

Advantage (ii) is the core contribution of this chapter, as currently there are no nuisance-free methods in the literature for semiparametric uncertainty quantification of $F_0(x)$ for deconvolution problems as treated in this chapter. Projecting the unconstrained posterior distribution departs from a classical Bayesian framework which might assign a Dirichlet Process (DP) prior to F and employ a hierarchical model to sample from the density of the observations. Uncertainty quantification results for this approach are currently unavailable and, at the same time, this approach is not advantageous from a computational prespective, being slower as described above.

4.2 Main results

In this section, we present the main result of this chapter, which establishes the asymptotic distribution of the isotonized posterior of $\hat{F}_G$, and we compare it with the asymptotic distribution attained by the isotonized inverse estimator (IIE) $\hat{F}_n$ (c.f. [109]) for a fixed $x \in [0, T)$, where T is a positive constant. We show in particular an interesting analogy. The results of [109] show that the IIE $\hat{F}_n$ attains at cube n root rate an asymptotic distribution, which, up to constants, is the Chernoff distribution. Symmetrically, we show that under the same conditions, the isotonized posterior of $\hat{F}_G$ attains at cube root rate an asymptotic distribution which, up to constants, is the Bayesian Chernoff distribution (4.2.17), as introduced in [17]. This curious phenomenon both confirms and extends the vision of [17], who introduced the Bayesian Chernoff distribution (4.2.17), proved its key properties (symmetry, monotone CDF, and the resulting recalibration procedure), and predicted that this distribution would play a central role in uncertainty quantification in nonparametric Bayesian inference far beyond the regression setting they originally considered. Indeed, also in the current setting, with a very different inverse problem and different prior, the Bayesian Chernoff distribution arises as the limiting distribution of the isotonized posterior of $\hat{F}_G$, confirming that the concept and the framework of [17] have a broad reach.

Integral equation (4.1.1) is a Volterra equation of the first kind, of convolution type. The function p is sometimes called the *resolvent of the first kind* of k (see [51], page 158). In all what follows, we restrict our attention to the case where F_0 has support contained in $[0, \infty)$. Furthermore, to prove consistency of the posterior $\hat{F}_G$, we have to impose a condition on p. The asymptotic distribution is derived under more stringent conditions (the same as in [109]) that the kernel k satisfies certain regularity conditions, which we will specify below.

Condition 1. On bounded intervals, the function p has only finitely many discontinuities. All these discontinuities are finite in size.

For the cube root asymptotics of $\hat{F}_G(x)$ for $x < T$, we need a slightly stronger condition on p

Condition 2. The function p is Hölder continuous of order $\gamma > 1/2$ on $(0, \infty)$ and $0 < p(0) < \infty$ and right continuous at zero.

A crucial question concerning integral equation (4.1.1) is whether a function p exists that satisfies it. in the Appendix (Section 4.5) 4.5.2, we give a review of the literature on this topic, showing that such a function p exists for many kernels k of interest. We also report a lemma, Lemma 4.5.8, that provides a sufficient condition for the existence of such a function p that satisfies the main condition required by the results in [109] and thus by our results, Condition 2.

For convenience of the reader, we report here without proof the main result of [109] for the IIE $\hat{F}_n$ (c.f. Theorem 2 in [109]).

Theorem 4.2.1 (Theorem 2 in [109])**.** *Let p satisfy Condition 2, $0 < T < \infty$ and let $x \in [0, T)$ be fixed and F_0 be such that F_0 has a continuous strictly positive derivative f_0 in a neighborhood of x. Then $n^{1/3}(\hat{F}_n(x) - F_0(x))$ converges in distribution as $n \to \infty$; specifically, $\forall\, z \in \mathbb{R}$:*

$$\mathrm{P}\left(n^{1/3}(\hat{F}_n(x) - F_0(x)) \le z\right) \xrightarrow{d} \mathrm{P}\left(\frac{2^{2/3} f_0(x)^{1/3} g_0(x)^{1/3}}{k(0)^{2/3}} \operatorname*{arg\,min}_{t\in\mathbb{R}}\{\mathbb{W}_1(t) + t^2\} \le z\right) \tag{4.2.1}$$

where $\mathbb{W}_1$ is two-sided Brownian motion originating from 0.

Finally, we proceed to the main result of this chapter, which establishes the asymptotic distribution of the isotonized posterior of $\hat{F}_G$. By stating it next to Theorem 4.2.1, we highlight the analogy between the asymptotic distribution of the IIP and that of the IIE.

Theorem 4.2.2. *Suppose $G \sim \mathrm{DP}(\alpha)$, where the base measure α has bounded density, and $\hat{F}_G$ as in* (4.1.5)*. Let p satisfy Condition 2, $0 < T < \infty$ and let $x \in [0, T)$ be fixed and F_0 be such that F_0 has a continuous strictly positive derivative f_0 in a neighborhood of x. Then $n^{1/3}(\hat{F}_G(x) - F_0(x))$ converges in distribution conditionally on $Z_1, \ldots, Z_n$ as $n \to \infty$; specifically, $\forall\, z \in \mathbb{R}$:*

$$\Pi\left(n^{1/3}(\hat{F}_G(x) - F_0(x)) \le z \mid Z_1, \ldots, Z_n\right) \tag{4.2.2}$$

$$\xrightarrow[n\to\infty]{d} \mathrm{P}\left(\frac{2^{2/3} f_0(x)^{1/3} g_0(x)^{1/3}}{k(0)^{2/3}} \operatorname*{arg\,min}_{t\in\mathbb{R}}\{\mathbb{W}_1(t) + \mathbb{W}_2(t) + t^2\} \le z \mid \mathbb{W}_1\right), \tag{4.2.3}$$

where $\mathbb{W}_1$ and $\mathbb{W}_2$ are two independent Brownian motions on $\mathbb{R}$ originating at 0.

Proof. Consider for $a \in (0, 1)$ and $\tau \in [0, T)$ the event $T_G(a) > \tau$ where:

$$T_G(a) := \inf\{t \in [0, T] : H_G(t) - at \text{ is minimal}\}. \tag{4.2.4}$$

This event takes place if and only if the maximal affine function with slope a that is bounded above by H_G coincides with H_G at a point $t_0 > \tau$, while for all $t \le \tau$, it stays strictly beneath H_G. This is equivalent to the inequality $\hat{F}_G(\tau) < a$. So $\forall\, \tau \in [0, T)$ and $a \in (0, 1)$:

$$T_G(a) \le \tau \iff \hat{F}_G(\tau) \ge a.$$

Fix $x \in (0, T)$. Then, by Lemma 4.5.1, for $a \in \mathbb{R}$ and n sufficiently large

$$n^{1/3}(\hat{F}_G(x) - F_0(x)) < a \iff \inf\{t \in [-xn^{1/3}, (T - x)n^{1/3}] : Z_G(t) - at \text{ is minimal}\} > 0,$$

where

$$Z_G(t) := n^{2/3}\left(H_G(x + tn^{-1/3}) - H_G(x) - F_0(x)tn^{-1/3}\right). \tag{4.2.5}$$

This process can be decomposed as

$$Z_G(t) = W_G(t) + W_n(t) + n^{2/3}(H_0(x + tn^{-1/3}) - H_0(x) - F_0(x)tn^{-1/3})$$

where:

$$W_G(t) := n^{2/3}(H_G(x + n^{-1/3}t) - H_G(x) - H_n(x + n^{-1/3}t) + H_n(x)),$$
$$W_n(t) := n^{2/3}(H_n(x + n^{-1/3}t) - H_n(x) - H_0(x + n^{-1/3}t) + H_0(x)).$$

We can rewrite $W_n(t)$ by identifying two processes: one of them converging weakly and a remainder term that converges to zero in probability. By defining the γ-Hölder continuous, $\gamma > 1/2$, function $\tilde{p} := p - p(0)\mathbf{1}_{[0,\infty)}$, and

$$\mathbb{W}_n(t) := n^{2/3}p(0)\int_0^\infty (\mathbf{1}_{[0,x+n^{-1/3}t)}(z) - \mathbf{1}_{[0,x)}(z))\, d(\mathbb{G}_n - G_0)(z),$$
$$R_n(t) := n^{2/3}\int_0^\infty (\tilde{p}(x + n^{-1/3}t - z) - \tilde{p}(x - z))\, d(\mathbb{G}_n - G_0)(z),$$

we can write:

$$W_n(t) = \mathbb{W}_n(t) + R_n(t). \tag{4.2.6}$$

Using the fact $\tilde{p} = p - p(0)\mathbf{1}_{[0,\infty)}$ is Hölder continuous of degree $\gamma > 1/2$ we have that as $n \to \infty$:

$$\sup_{|t|\in K} |R_n(t)| \xrightarrow{G_0} 0 \tag{4.2.7}$$

for all compact sets $K \subset \mathbb{R}$ (see the Appendix (Section 4.5) in [109] for a proof). By example 3.2.14. in [103], for every compact $K \subset \mathbb{R}$:

$$\mathbb{W}_n \rightsquigarrow \frac{\sqrt{g_0(x)}}{k(0)}\mathbb{W}_1 \quad \text{in} \quad \ell^\infty(K) \tag{4.2.8}$$

where $\mathbb{W}_1$ is a two sided standard Brownian Motion originating at 0. Now, using Taylor's theorem, for $n \to \infty$

$$n^{2/3}(H_0(x + n^{-1/3}t) - H_0(x) - F_0(x)n^{-1/3}t) = \frac{1}{2}f_0(x)t^2 + o(1), \tag{4.2.9}$$

uniformly for t in compacta.

We now turn to studying W_G. Define $f_t^n(z) = n^{1/3}(p(x + n^{-1/3}t - z)\mathbf{1}_{[0,x+n^{-1/3}t)}(z) - p(x - z)\mathbf{1}_{[0,x)}(z))$ then, for independent $Q \sim \mathrm{DP}(\alpha)$, $\mathbb{B}_n \sim \mathrm{DP}(n\mathbb{G}_n)$ and $V_n \sim \mathrm{Be}(|\alpha|, n)$:

$$Gf_t^n = V_nQf_t^n + (1 - V_n)\mathbb{B}_nf_t^n$$

and thus we study the process:

$$W_G(t) = n^{1/3}(G - \mathbb{G}_n)f_t^n = n^{1/3}V_n(Q - \mathbb{G}_n)f_t^n + n^{1/3}(1 - V_n)(\mathbb{B}_n - \mathbb{G}_n)f_t^n.$$

By Lemma 4.5.2 (eq. (4.5.3)-(4.5.4)), the terms premultiplied by V_n converge to zero in probability under G_0, so we can focus on the study of:

$$W_n^*(t) := n^{1/3}(\mathbb{B}_n - \mathbb{G}_n)f_t^n.$$

By Proposition G.2 in [42] for $\varepsilon_i \overset{\text{i.i.d.}}{\sim} \text{Exp}(1)$ and $W_{ni} := \varepsilon_i/\bar{\varepsilon}_n$, we have $(W_{n1}, \ldots, W_{nn}) \sim \text{Dir}(n; 1/n, \ldots, 1/n)$. Since $\mathbb{B}_n \sim \text{DP}(n\mathbb{G}_n)$, denoting by δ_{Z_i} Dirac measure at Z_i, we have the following representation in distribution of $\mathbb{B}_n$ (c.f. Example 3.7.9 in [103]):

$$\mathbb{B}_n = \frac{1}{n}\sum_{i=1}^{n} W_{ni}\delta_{Z_i}.$$

Because of this representation (c.f. section 3.7.2. in [103]), $\mathbb{B}_n$ is also known as *Bayesian bootstrap*. Conclude, by Lemma 4.5.3, that uniformly for t in compacta:

$$W_n^*(t) = n^{1/3}(\mathbb{B}_n - \mathbb{G}_n)f_t^n = \frac{1}{n^{2/3}}\sum_{i=1}^{n}(W_{ni} - 1)f_t^n(Z_i) = \mathbb{W}_n^*(t) + o_p(1). \tag{4.2.10}$$

where for $\varepsilon_i \overset{\text{i.i.d.}}{\sim} \text{Exp}(1)$, the process $\mathbb{W}_n^*$, for $\tilde{f}_t^n(z) := n^{1/3}p(0)\mathbf{1}_{[x,x+n^{-1/3}t)}(z)$, is defined as:

$$\mathbb{W}_n^*(t) = \frac{1}{\sqrt{n}}\sum_{i=1}^{n}\frac{(\varepsilon_i - 1)}{\sqrt[6]{n}}\tilde{f}_t^n(Z_i). \tag{4.2.11}$$

Using the results in the Appendix (Section 4.5) A, Lemma 4.5.4, which shows that $\mathbb{W}_n^*$ converges conditionally in $\ell^\infty(K)$ to $\frac{\sqrt{g_0(x)}}{k(0)}\mathbb{W}_2$, together with Lemma 4.5.2 and (4.2.10), we have in probability under G_0, as $n \to \infty$,

$$W_G \mid Z_1, \ldots, Z_n \rightsquigarrow \frac{\sqrt{g_0(x)}}{k(0)}\mathbb{W}_2 \quad \text{in} \quad \ell^\infty(K), \tag{4.2.12}$$

for every compact $K \subset \mathbb{R}$. Hence, by (4.2.6), (4.2.7), (4.2.8), (4.2.12) and Slutsky's lemma, for every compact $K \subset \mathbb{R}$:

$$\begin{aligned}(W_G(t) + W_n(t) + n^{2/3}(H_0(x + tn^{-1/3}) - H_0(x) - F_0(x)tn^{-1/3}) : t \in K) \mid Z_1, \ldots, Z_n \\ \rightsquigarrow \left(\frac{\sqrt{g_0(x)}}{k(0)}\mathbb{W}_1(t) + \frac{\sqrt{g_0(x)}}{k(0)}\mathbb{W}_2(t) + \frac{1}{2}f_0(x)t^2 : t \in K\right) \mid \mathbb{W}_1.\end{aligned} \tag{4.2.13}$$

Let us elaborate on why the unconditional convergence $W_n \rightsquigarrow \frac{\sqrt{g_0(x)}}{k(0)}\mathbb{W}_1$ in $\ell^\infty(K)$ (equation (4.2.8)) gives rise to a *conditional* limit involving $\mathbb{W}_1$ in (4.2.13). The key observation is that W_G is driven exclusively by the Bayesian bootstrap weights $(\varepsilon_i)_{i=1}^n$, which are independent of the data $(Z_1, \ldots, Z_n)$. Hence W_G and W_n are conditionally independent given the data. As $n \to \infty$: the orbit of W_n (a measurable function of the data alone) converges to $\frac{\sqrt{g_0(x)}}{k(0)}\mathbb{W}_1$, and $W_G|Z_1, \ldots, Z_n$ converges independently to $\frac{\sqrt{g_0(x)}}{k(0)}\mathbb{W}_2$. In the joint limit, the realization of $\mathbb{W}_1$ (coming from W_n) plays the role of a fixed, conditioning object, while $\mathbb{W}_2$ remains a fresh independent Brownian motion. Conditional on the data, an application of Slutsky's lemma in the conditional setting then yields (4.2.13) with the right-hand side conditioned on $\mathbb{W}_1$.

Now, in order to apply Lemma A.3 in [47], we need to prove that there exists a stochastically bounded sequence $\tilde{M}_n$ such that:

$$\Pi_n\left(\inf\{t \in [-xn^{1/3}, (T-x)n^{1/3}] : Z_G(t) - at \text{ is minimal}\} \leq \tilde{M}_n \mid Z_1, \ldots, Z_n\right) \xrightarrow{G_0} 1 \tag{4.2.14}$$

The proof of (4.2.14) is given in the Appendix (Section 4.5) A, Lemma 4.5.6. An application of Lemma A.3 in [47], together with (4.2.13) and (4.2.14) gives:

$$\begin{aligned}&\inf\{t \in \mathbb{R} : Z_G(t) - at \text{ is minimal}\} \mid Z_1, \ldots, Z_n \\ &\rightsquigarrow \arg\min_{t\in\mathbb{R}} \left(\frac{\sqrt{g_0(x)}}{k(0)} \mathbb{W}_1(t) + \frac{\sqrt{g_0(x)}}{k(0)} \mathbb{W}_2(t) + \frac{1}{2} f_0(x) t^2 - at \right) \mid \mathbb{W}_1.\end{aligned}$$

Using that $\forall\, c > 0$, $b \in \mathbb{R}$ that as process the standard Brownian Motion satisfies $W(\cdot/a^2) \overset{d}{=} W(\cdot)/a$:

$$\operatorname{argmin}_{t\in\mathbb{R}}\{c(\mathbb{W}_1(t) + \mathbb{W}_2(t)) + (t-b)^2\} \overset{d}{=} c^{2/3} \operatorname{argmin}_{t\in\mathbb{R}}\{\mathbb{W}_1(t) + \mathbb{W}_2(t) + t^2\} + b.$$

Note that this distributional equality holds *jointly in the pair* (argmin, $\mathbb{W}_1$): the time-rescaling $t \mapsto c^{1/3}t$ is applied simultaneously to both $\mathbb{W}_1$ and $\mathbb{W}_2$, so the rescaled $\mathbb{W}_1(c^{1/3}\,\cdot)/c^{1/6}$ has the same law as $\mathbb{W}_1$, and a joint rescaling argument shows that the equality in distribution is in the space of (argmin, $\mathbb{W}_1$). Thus conditioning on $\mathbb{W}_1$ on the right-hand side corresponds to conditioning on the same Brownian trajectory (up to the distributional scaling identity), and the conditional distribution of the argmin given $\mathbb{W}_1$ is the same on both sides of the $\overset{d}{=}$. Using the change of variable $t \to \sqrt{\frac{2}{f_0(x)}}t$ we get

$$\begin{aligned}&\arg\min_{t\in\mathbb{R}} \left(\frac{\sqrt{g_0(x)}}{k(0)} \mathbb{W}_1(t) + \frac{\sqrt{g_0(x)}}{k(0)} \mathbb{W}_2(t) + \frac{1}{2} f_0(x) t^2 - at \right) \\ &\overset{d}{=} \frac{g_0(x)^{1/3}}{k(0)^{2/3}} \left(\frac{2}{f_0(x)} \right)^{1/6+1/2} \arg\min_{t\in\mathbb{R}} \{\mathbb{W}_1(t) + \mathbb{W}_2(t) + t^2\} + \frac{a}{f_0(x)},\end{aligned}$$

which proves the claim, since:

$$\begin{aligned}&\mathrm{P}\left(\arg\min_{t\in\mathbb{R}} \left(\frac{\sqrt{g_0(x)}}{k(0)} \mathbb{W}_1(t) + \frac{\sqrt{g_0(x)}}{k(0)} \mathbb{W}_2(t) + \frac{1}{2} f_0(x) t^2 - at \right) > 0 \mid \mathbb{W}_1 \right) \\ &= \mathrm{P}\left(\frac{g_0(x)^{1/3}}{k(0)^{2/3}} \left(\frac{2}{f_0(x)} \right)^{1/6+1/2} \arg\min_{t\in\mathbb{R}} \{\mathbb{W}_1(t) + \mathbb{W}_2(t) + t^2\} > -\frac{a}{f_0(x)} \mid \mathbb{W}_1 \right) \\ &= \mathrm{P}\left(\frac{2^{2/3} f_0(x)^{1/3} g_0(x)^{1/3}}{k(0)^{2/3}} \arg\min_{t\in\mathbb{R}} \{\mathbb{W}_1(t) + \mathbb{W}_2(t) + t^2\} \le a \mid \mathbb{W}_1 \right).\end{aligned}$$

□

The limiting distribution established in Theorem 4.2.2 is fundamental for establishing the asymptotic coverage of credible intervals. For any given $n \ge 1$ and $\tau \in [0, 1]$, the credible interval is represented by $I_{n,\tau} := [Q_{n,1-\tau/2}, Q_{n,\tau/2}]$. Here, $Q_{n,\tau}$ signifies the $(1-\tau)$-quantile of the projection-posterior distribution associated with $\hat{F}_G(x)$:

$$Q_{n,\tau} := \inf\{z \in \mathbb{R} \,:\, \Pi(\hat{F}_G(x) \le z \mid Z_1, \ldots, Z_n) \ge 1 - \tau\}.$$

Consequently, $F_0(x) \le Q_{n,\tau}$ is true if and only if $\Pi(\hat{F}_G(x) \le F_0(x) \mid Z_1, \ldots, Z_n) \le 1 - \tau$. The limiting coverage of the interval $I_{n,\tau}$ as $n \to \infty$ can be determined. To begin, observe that

$$\mathrm{P}(F_0(x) \le Q_{n,\tau}) = \mathrm{P}(\Pi(n^{1/3}(\hat{F}_G(x) - F_0(x)) \le 0 \mid Z_1, \ldots, Z_n) \le 1 - \tau) \tag{4.2.15}$$

$$\to \mathrm{P}\Big(\mathrm{P}(\operatorname{argmin}_{t\in\mathbb{R}}\{\mathbb{W}_1(t)+\mathbb{W}_2(t)+t^2\}\le 0 \mid \mathbb{W}_1)\le 1-\tau\Big). \tag{4.2.16}$$

This can be written as $\mathrm{P}(Z_B \le 1-\tau)$, where Z_B is specified as

$$Z_B := \mathrm{P}(\operatorname{argmin}_{t\in\mathbb{R}}\{\mathbb{W}_1(t)+\mathbb{W}_2(t)+t^2\}\le 0 \mid \mathbb{W}_1). \tag{4.2.17}$$

The random variable Z_B, having distribution known as the Bayes–Chernoff distribution, plays a pivotal role in understanding the limiting coverage. While the distribution of Z_B is analogous to the Chernoff distribution for the MLE, Z_B takes values on the unit interval and exhibits symmetry around $1/2$ ($Z_B = 1 - Z_B$ in distribution). Employing (4.2.15), we arrive at the primary conclusion regarding the limiting coverage of $I_{n,\tau}$ as $n \to \infty$:

$$\mathrm{P}(F_0(x) \in I_{n,\tau}) \to \mathrm{P}(\tau/2 \le Z_B \le 1-\tau/2).$$

Therefore, the limiting coverage of a $(1-\tau)$-credible interval does not necessarily equal $(1-\tau)$. It can, however, be computed using the distribution of Z_B. Let us define $A(u) = \mathrm{P}(Z_B \le u)$ for $u \in [0,1]$. The function A is continuous, strictly increasing, and maps the interval $[0,1]$ onto itself. Consequently, the inverse function A^{-1} is well-defined and also strictly increasing. Additionally, the function A features a symmetry property, which is formalized in Lemma 3.5 in [17] and reported here for completeness.

Lemma 4.2.1 (Lemma 3.5 in [17])**.** *The random variable $Z_B \in [0,1]$ is symmetrically distributed about $1/2$, and hence $A(1-u) = 1-A(u)$, for all $u \in [0,1]$, $A^{-1}(1-v) = 1-A^{-1}(v)$ for all $v \in [0,1]$, and $A(1/2) = 1/2 = A^{-1}(1/2)$.*

Using these properties, the limiting coverage $\mathrm{P}(\tau/2 \le Z_B \le 1-\tau/2)$ can be rewritten as $A(1-\tau/2) - A(\tau/2) = A(1-\tau/2) - (1-A(1-\tau/2)) = 2A(1-\tau/2) - 1$. A recalibration technique allows us to achieve any desired asymptotic coverage, say $1-\beta$. For a two-sided credible interval $I_{n,\tau}$, Theorem 4.2.2 and Lemma 4.2.1 imply that its asymptotic coverage is $2A(1-\tau/2)-1$. To obtain a desired coverage of $1-\beta$, we set $2A(1-\tau/2)-1 = 1-\beta$. This implies $1-\tau/2 = A^{-1}(1-\beta/2)$, or $\tau/2 = 1 - A^{-1}(1-\beta/2) = A^{-1}(\beta/2)$. Thus, we choose $\tau = 2A^{-1}(\beta/2)$ (c.f. Corollary 3.6 in [17]).

Corollary 4.2.1. *For any $0 < \beta < 1$, $\mathrm{P}(F_0(x) \in I_{n,2A^{-1}(\beta/2)}) \to 1-\beta$.*



Figure 4.1: The Bayes–Chernoff distribution Z_B and its inverse function A^{-1}, with the identity line for reference (the computations are based on 20000 gathered samples of Z_B).

The back-calculation of the required credibility level $1 - \tau = 1 - 2A^{-1}(\beta/2)$ to achieve a coverage level $1 - \beta$ can be obtained from a Table 2 in [17], which presents values of the function A^{-1}. For convenience of the reader and for completeness, we report Table 2 from [17] here below, in Table 4.1 (we verified via our simulations its correctness). Numerical calculations suggest that $A(u) \geq u$ for all values of $u \geq 1/2$ where the computation has been numerically performed and the results are shown in Fig. 4.1. This implies that a credible interval typically has asymptotic coverage greater than its nominal coverage, which may be called a reverse Cox phenomenon.

Table 4.1: Table of the values of the function A^{-1}

v	0.700	0.750	0.800	0.850	0.900	0.910	0.920	0.930	0.940	0.950	0.960
$A^{-1}(v)$	0.677	0.723	0.771	0.820	0.874	0.885	0.897	0.909	0.922	0.934	0.946
v	0.965	0.970	0.975	0.980	0.985	0.990	0.995	0.996	0.997	0.998	0.999
$A^{-1}(v)$	0.952	0.960	0.966	0.973	0.980	0.986	0.993	0.995	0.996	0.997	0.999

4.3 Simulation study

In this section, we present simulations to illustrate the practical behavior of the IIP in comparison to the IIE studied in [109]. In this simulation study, we analyze the setting where the underlying cdf F_0 is exponential with rate 1.2. Samples $Z_1, \ldots, Z_n$ used to compute the projection posterior of the IIP are generated hierarchically: $X_1, \ldots, X_n \sim F_0$, $Y_1, \ldots, Y_n \sim k$, with $Z_i = X_i + Y_i$ for all i. The noise density k which crucially determines the solvability of the Volterra integral equation in (4.1.1), is chosen from the following set: $\{\text{Exp}(1), \text{Mix} : \frac{1}{4}\text{Erlang}(2) - \frac{3}{4}\text{Exp}(2), \text{HalfNorm}(0, 1), \text{HalfCauchy}(0, 2), \text{Lomax}(10, 1), \text{HalfLogistic}\}$. We recall the associated kernels k in Fig. 4.2 and we plot the numerical solutions p (and the analytical solutions when available) of the Volterra integral equation in (4.1.1) for each of these noise distributions.

The reasons why we chose this set of distributions are two: on the one hand, for our methodology to work we need $k(0) \neq 0$, with k living on the positive real line. On the other hand, we want to illustrate the performance of the IIP in a variety of settings, including both light-tailed and heavy-tailed noise distributions. Thus we included a good variety of distributions, that go from light-tailed (e.g. exponential and half-normal) to heavy-tailed (e.g. Lomax with $c = 10$, with tails proportional to y^{-11} and which is a special case of Pareto) to very heavy-tailed (e.g. half-Cauchy with tails proportional to y^{-2}). Because the underlying cdf F_0 is always the same, our simulations show that the IIP is able to recover the underlying cdf F_0 in all these settings, even when the noise distribution is very heavy-tailed. In particular, in Figure 4.3 we show the results of the IIP deconvolution methodology compared to the IIE, visualizing the isonotized posterior draws, for the same kernels of Figure 4.2. We observe that the Bayesian methodology agrees with the frequentist one. In Figure 4.4, we provide in the same simulation setting the recalibrated credible sets according to the recalibration $2A^{-1}(0.975) - 1 = 0.932$. The numerical solutions of the Volterra integral equation are computed using the trapezoidal rule, as described in detail here below. The plots use the following color scheme: green for draws of $\hat{F}_G$ based on prior draws from $G \sim \text{DP}(M\alpha)$ (parameters M and α a pdf specified in the plots not necessarily in the above mentioned set); purple for draws from the isotonized posterior; red for the average of these draws (approximating the posterior mean); yellow for the Isotonic Inverse Estimator (IIE); and orange for F_0.



Figure 4.2: Resolvent of proposed kernels.



Figure 4.3: Combined deconvolution plots for $n = 200$. In green, we plot draws of $\hat{F}_G$ based on prior draws from $G \sim \mathrm{DP}(M\alpha)$ with $\alpha \sim \mathrm{Ga}(2, 2)$ and prior precision $M = 10$; in purple, draws from the isotonized posterior; in red, the average of these draws (approximating the posterior mean); in yellow, the (IIE); and in orange F_0.



Figure 4.4: Combined deconvolution plots with calibrated uncertainty quantification for $n = 200$. In green, we plot draws of $\hat{F}_G$ based on prior draws from $G \sim \mathrm{DP}(M\alpha)$ with $\alpha \sim \mathrm{Ga}(2, 2)$ and prior precision $M = 10$; in pink, the calibrated credible bands from the isotonized posterior; in red, the average of these draws (approximating the posterior mean); in yellow, the (IIE); and in orange F_0.

Overall, these simulations demonstrate that the IIP reliably recovers the underlying distribution F_0 across a wide spectrum of noise distributions, ranging from light-tailed to extremely heavy-tailed. Despite substantial variation in the shape and tail behavior of the kernels k, the numerical solutions of the associated Volterra equations remain stable, and the resulting posterior draws closely track the frequentist IIE benchmark. The agreement between methods, together with the well-calibrated credible sets, highlights the robustness of the IIP and its

practical effectiveness for deconvolution in diverse settings.

Numerical Solver of the Volterra Equation

In order to compute the estimators in case p is not explicitly known, we use a numerical approximation. The presented simulations need the unknown function $p(x)$ solving the specific Volterra integral equation of the first kind:

$$x\mathbf{1}(x) = \int_0^x k(x-t)p(t)\,dt, \quad x \geq 0. \tag{4.3.1}$$

The kernel k is a known function. The solver uses a numerical method based on the trapezoidal rule. A key requirement for this method is $k(0) \neq 0$.

We begin by discretizing the domain $[0, T]$ into N equally spaced points, $x_n = nh$, $p_n := p(x_n)$ and $k_n := k(x_n)$, for $n = 0, 1, \ldots, N-1$, with step size $h := T/(N-1)$. At each grid point x_n, Equation (4.3.1) becomes:

$$x_n = nh = \int_0^{x_n} k(x_n - t)p(t)\,dt. \tag{4.3.2}$$

We approximate the integral using the composite trapezoidal rule over the points $t_j = jh$ for $j = 0, \ldots, n$:

$$nh \approx h\left[\frac{1}{2}k(x_n - t_0)p(t_0) + \sum_{j=1}^{n-1} k(x_n - t_j)p(t_j) + \frac{1}{2}k(x_n - t_n)p(t_n)\right]. \tag{4.3.3}$$

Let $k(x_n - t_j) := k((n-j)h) = k_{n-j}$ and $k(x_n - t_n) := k(0) = k_0$, then dividing both sides by the step size h:

$$n = \frac{1}{2}k_n p_0 + \sum_{j=1}^{n-1} p_j k_{n-j} + \frac{1}{2}k_0 p_n. \tag{4.3.4}$$

We can now rearrange this equation to solve for the unknown term p_n, assuming all previous values $(p_0, \ldots, p_{n-1})$ are known. This gives the recurrence relation for p_n:

$$p_n = \frac{2}{k_0}\left[n - \frac{1}{2}k_n p_0 - \sum_{j=1}^{n-1} p_j k_{n-j}\right]. \tag{4.3.5}$$

To start the recurrence, we need the value of $p_0 = p(0)$. This can be found by differentiating the original integral equation using the Leibniz integral rule, which yields the relation $1 = k(0)p(0)$.

The algorithm implemented in the simulation proceeds as follows:

1. Discretize the domain $[0, T]$ into N points with step size h. Pre-compute the kernel values k_n for $n = 0, \ldots, N-1$.
2. Calculate the exact initial value p_0: $p_0 = 1/k_0$.
3. For each step $n = 1, 2, \ldots, N-1$, sequentially compute p_n using the recurrence relation from Equation (4.3.5).

This procedure yields a stable solution for all examples in Figure 4.2.

4.4 Conclusion

This work introduces and evaluates a nonparametric Bayesian method for uncertainty quantification in a specific class of statistical deconvolution problems. To address the challenge of nuisance-parameter-dependent confidence intervals in frequentist methods, we propose the "Isotonic Inverse Posterior." This approach combines a Dirichlet Process prior on the distribution of the observed data with an inverse operator and an isotonic regression step to produce valid estimates for the signal's cumulative distribution function, F_0. The procedure is computationally fast and, following a straightforward recalibration, its credible sets achieve asymptotically correct frequentist coverage without relying on nuisance parameter estimation. As demonstrated through simulations, the method performs effectively across a variety of noise distributions.

4.5 Appendix

4.5.1 Complementary results

Measure-theoretical framework: in this chapter, the sample space is $(\mathbb{R}^+, \mathscr{B}(\mathbb{R}^+))$, and $\mathfrak{M}$ (with Borel σ-field $\mathscr{M}$ for the weak topology) is the collection of all Borel probability measures on $(\mathbb{R}^+, \mathscr{B}(\mathbb{R}^+))$. The Dirichlet process prior is a probability measure on $\mathfrak{M}$. A prior on the set of probability measures $\mathfrak{M}$ is a probability measure on a sigma-field $\mathscr{M}$ of subsets of $\mathfrak{M}$. Alternatively, it can be viewed as map from some probability space $(\Omega, \mathscr{U}, \Pr)$ into $(\mathfrak{M}, \mathscr{M})$. We can think of the hierarchy of the spaces as follows: $(\Omega, \mathscr{U}, \Pr) \to (\mathfrak{M}, \mathscr{M}) \to (\mathbb{R}^+, \mathscr{B}(\mathbb{R}^+))$. We put a Dirichlet process prior over G and write $G \sim \Pi = \mathrm{DP}(\alpha)$ for some base measure α and prior precision $|\alpha|$. This means that for any measurable set $B \in \mathscr{M}$ we have $\Pi(B) = \Pr(G \in B)$. Throughout the chapter, we use the notations P and E, which should be understood as the outer probability measure and outer expectation, respectively, whenever issues of measurability in the considered processes arise (see [103] for a rigorous treatment of these topics).

Furthermore, we use the notion of conditional convergence in distribution. This happens if the bounded Lipschitz distance between the conditional law of Borel measurable X_n given $\mathcal{B}_n$ and a Borel probability measure L tends to zero, where the convergence can be in outer probability. The sequence X_n tends to L conditionally given $\mathcal{B}_n$ in outer probability if

$$d_{BL}\left(\mathcal{L}\left(X_n \mid \mathcal{B}_n\right), L\right) := \sup_{f \in BL_1} \left| \mathrm{E}\left(f\left(X_n\right) \mid \mathcal{B}_n\right) - \int f dL \right| \xrightarrow{\mathrm{P}} 0, \tag{4.5.1}$$

where BL_1 is the space of bounded Lipschitz functions with Lipschitz constant 1.

In particular if $Z_1, \ldots, Z_n \overset{\text{i.i.d.}}{\sim} \mathrm{P}_Z$, then $X_n \mid Z_1, \ldots, Z_n \rightsquigarrow N(0, \sigma^2)$ means:

$$d_{BL}\left(\mathrm{P}(X_n \in \cdot \mid Z_1, \ldots, Z_n), N(0, \sigma^2)\right) \xrightarrow{\mathrm{P}_Z} 0. \tag{4.5.2}$$

Lemma 4.5.1. *Let $x \in (0, T)$, $\hat{F}_G$ as in* (4.1.5) *and F_0 true underlying cdf. Then for $a \in \mathbb{R}$ and n sufficiently large*

$$n^{1/3}(\hat{F}_G(x) - F_0(x)) < a \iff \inf\{t \in [-xn^{1/3}, (T-x)n^{1/3}] : Z_G(t) - at \text{ is minimal}\} > 0,$$

where, for H_G as in (4.1.4)*,*

$$Z_G(t) := n^{2/3}\left(H_G(x + tn^{-1/3}) - H_G(x) - F_0(x)tn^{-1/3}\right).$$

Proof. Fix $x \in (0, T)$. Then for $a \in \mathbb{R}$ and n sufficiently large

$$\begin{aligned}
n^{1/3}(\hat{F}_G(x) - F_0(x)) < a &\iff \hat{F}_G(x) < F_0(x) + an^{-1/3} \\
&\overset{(4.2.4)}{\iff} T_G(F_0(x) + an^{-1/3}) > x \\
&\iff \inf\{x + tn^{-1/3} \in [0, T] : H_G(x + tn^{-1/3}) - H_G(x) \\
&\qquad - F_0(x)tn^{-1/3} - atn^{-2/3} \text{ is minimal}\} > x \\
&\iff \inf\{t \in [-xn^{1/3}, (T-x)n^{1/3}] : n^{2/3}(H_G(x + tn^{-1/3}) \\
&\qquad - H_G(x) - F_0(x)tn^{-1/3}) - at \text{ is minimal}\} > 0.
\end{aligned}$$

□

Lemma 4.5.2. *Let* $f_t^n(z) := n^{1/3}(p(x+n^{-1/3}t-z)\mathbf{1}_{[0,x+n^{-1/3}t)}(z) - p(x-z)\mathbf{1}_{[0,x)}(z))$ *then, for independent* $Q \sim \mathrm{DP}(\alpha)$, $\mathbb{B}_n \sim \mathrm{DP}(n\mathbb{G}_n)$ *and* $V_n \sim \mathrm{Be}(|\alpha|, n)$. *As* $n \to \infty$, *under* G_0 *in probability:*

$$\mathrm{P}(n^{1/3}|V_n(Q-\mathbb{G}_n)f_t^n| > \varepsilon \mid Z_1,\dots,Z_n) \xrightarrow{G_0} 0. \tag{4.5.3}$$

and

$$\mathrm{P}(n^{1/3}|V_n(\mathbb{B}_n-\mathbb{G}_n)f_t^n| > \varepsilon \mid Z_1,\dots,Z_n) \xrightarrow{G_0} 0. \tag{4.5.4}$$

Proof. Using the fact that p is Hölder continuous on $(0,\infty)$ of degree $\gamma > 1/2$:

$$\begin{aligned}
|f_t^n(z)| &= n^{1/3}|p(x+n^{-1/3}t-z)\mathbf{1}_{[0,x)}(z) - (x-z)\mathbf{1}_{[0,x)}(z)| \\
&\quad + |p(x+n^{-1/3}t-z)\mathbf{1}_{[x,x+n^{-1/3}t)}(z)| \\
&\lesssim n^{1/3}|(x+n^{-1/3}t-z)-(x-z)|^\gamma \mathbf{1}_{[0,x)}(z) \\
&\quad + n^{1/3}\sup_{z\in[x,x+n^{-1/3}t]}|p(x+n^{-1/3}t-z)| \\
&\lesssim n^{1/3}|n^{-1/3}t\mathbf{1}_{[0,x)}(z)|^\gamma + n^{1/3}\sup_{z\in[x,x+n^{-1/3}t]}|x+n^{-1/3}t-z|^\gamma \\
&\lesssim n^{1/3}p(0) + n^{(1-\gamma)/3}|t|^\gamma.
\end{aligned}$$

Note that $\mathrm{E}V_n = |\alpha|/(|\alpha|+n)$, thus $\forall\,\varepsilon > 0$ by independence, $\gamma > 1/2$ as $n \to \infty$

$$\begin{aligned}
\mathrm{P}(n^{1/3}|V_n(Q-\mathbb{G}_n)f_t^n| > \varepsilon \mid Z_1,\dots,Z_n) &\lesssim n^{1/3}\varepsilon^{-1}\mathrm{E}V_n\,\mathrm{E}[(Q+\mathbb{G}_n)|f_t^n| \mid Z_1,\dots,Z_n] \\
&\lesssim n^{(2-\gamma)/3}\varepsilon^{-1}\frac{|\alpha|}{|\alpha|+n}|t|^\gamma \to 0.
\end{aligned}$$

Similarly $\mathrm{P}(n^{1/3}|V_n(\mathbb{B}_n-\mathbb{G}_n)f_t^n| > \varepsilon \mid Z_1,\dots,Z_n) \xrightarrow{G_0} 0$. □

Lemma 4.5.3. *For* W_n^* *as in* (4.2.10), *in probability under* G_0, *as* $n \to \infty$:

$$W_n^*(t) = \mathbb{W}_n^*(t) + o_p(1). \tag{4.5.5}$$

where for $\varepsilon_i \overset{i.i.d.}{\sim} \mathrm{Exp}(1)$, *the process* $\mathbb{W}_n^*$ *is defined as:*

$$\mathbb{W}_n^*(t) = \frac{1}{\sqrt{n}}\sum_{i=1}^n \frac{(\varepsilon_i - 1)}{\sqrt[6]{n}}\tilde{f}_t^n(Z_i). \tag{4.5.6}$$

and $\tilde{f}_t^n(z) := n^{1/3}p(0)\mathbf{1}_{[x,x+n^{-1/3}t)}(z)$.

Proof. First recall the decomposition:

$$W_n^*(t) = n^{1/3}(\mathbb{B}_n-\mathbb{G}_n)f_t^n = \frac{1}{n^{2/3}}\sum_{i=1}^n (W_{ni}-1)f_t^n(Z_i),$$

where $W_{ni} = \frac{\varepsilon_i}{\bar{\varepsilon}_n}$. Because $|\bar{\varepsilon}_n - 1| = O_p(1/\sqrt{n})$, for almost every sequence $Z_1,\dots,Z_n$ conclude that uniformly in t in compacta:

$$\frac{1}{n^{2/3}}\sum_{i=1}^n (W_{ni}-1)f_t^n(Z_i) = \frac{1}{\sqrt{n}}\sum_{i=1}^n \frac{(\varepsilon_i-1)}{\sqrt[6]{n}}f_t^n(Z_i) + o_p(1).$$

Now consider the decomposition:

$$f_t^n(z) = \tilde{f}_t^n(z) + r_t^n(z)$$

where for $\tilde{p} = p - p(0)\mathbf{1}_{[0,\infty)}$:

$$r_t^n(z) := n^{1/3}(\tilde{p}(x + n^{-1/3}t - z) - \tilde{p}(x - z)),$$
$$\tilde{f}_t^n(z) := n^{1/3} p(0)\mathbf{1}_{[x,x+n^{-1/3}t)}(z).$$

We obtain the claim that uniformly in t over compacta:

$$W_n^*(t) = \mathbb{W}_n^*(t) + o_p(1), \tag{4.5.7}$$

as soon as we show that for all compact $K \subset \mathbb{R}$ and $\forall\, \varepsilon > 0$:

$$\mathrm{P}\left(\sup_{t\in K} \left| \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{(\varepsilon_i - 1)}{\sqrt[6]{n}} r_t^n(Z_i) \right| > \varepsilon \mid Z_1, \ldots, Z_n \right) \xrightarrow{G_0} 0. \tag{4.5.8}$$

For $\tilde{p} = p - p(0)\mathbf{1}_{[0,\infty)}$, $\tilde{p}$ is γ-Hölder continuous of degree $\gamma > 1/2$. Define:

$$R_n^*(t) := \frac{1}{n^{1/3}} \sum_{i=1}^n (\varepsilon_i - 1)\left(\tilde{p}(x + n^{-1/3}t - Z_i) - \tilde{p}(x - Z_i) \right).$$

The claim is obtained if we show for any $K \in (0,\infty)$, $\sup_{|t|\le K} |R_n^*(t)| \xrightarrow{G_0} 0$.

Let $0 = t_0 < t_1 < \cdots < t_{K_n} = K$ and $t_{-i} = -t_i$, $i = 1, \ldots, K_n$. Now note:

$$\begin{aligned} \sup_{|t|\le K} |R_n^*(t)| &= \max_{-K_n+1\le i\le K_n} \sup_{t\in[t_{i-1},t_i]} |R_n^*(t)| \\ &\le \max_{-K_n+1\le i\le K_n} \left(|R_n^*(t_i)| + \sup_{t\in[t_{i-1},t_i]} |R_n^*(t) - R_n^*(t_i)| \right). \end{aligned}$$

Using Markov's inequality, we have:

$$\begin{aligned} \varepsilon\, \mathrm{P}\left(\sup_{|t|\le K} |R_n^*(t)| > \varepsilon \right) &\le \mathrm{E}\left(\sup_{|t|\le K} |R_n^*(t)| \right) \\ &\le \mathrm{E}\left(\max_{-K_n+1\le i\le K_n} |R_n^*(t_i)| \right) + \mathrm{E}\left(\max_{-K_n+1\le i\le K_n} \sup_{t\in[t_{i-1},t_i]} |R_n^*(t) - R_n^*(t_i)| \right). \end{aligned}$$

For each $t \in [t_{i-1}, t_i]$ and $\mathbb{P}_n = \frac{1}{n}\sum_{i=1}^n \delta_{\varepsilon_i}$

$$\begin{aligned} &|R_n^*(t) - R_n^*(t_i)| \\ &= n^{2/3} \left| \int_0^\infty \int_0^\infty \left(\tilde{p}(x + n^{-1/3}t - z) - \tilde{p}(x + n^{-1/3}t_i - z) \right)(\varepsilon - 1)\, d(\mathbb{G}_n, \mathbb{P}_n)(z, \varepsilon) \right| \\ &\le 2n^{2/3} L n^{-\gamma/3} |t - t_i|^\gamma \left(1 + \frac{1}{n}\sum_{j=1}^n \varepsilon_j \right) = 2n^{(2-\gamma)/3} L |t - t_i|^\gamma \left(1 + \frac{1}{n}\sum_{j=1}^n \varepsilon_j \right). \end{aligned}$$

Take a grid of t_i's such that $|t_i - t_{i-1}|^\gamma = \delta n^{-(2-\gamma)/3}$ then

$$\mathrm{E}\left(\sup_{-K_n+1\le i\le K_n} \sup_{t\in[t_{i-1},t_i]} |R_n^*(t) - R_n^*(t_i)| \right) \le 4L\delta,$$

(which we can make small by taking δ small enough; note also that $K_n = O(n^{(2-\gamma)/3\gamma})$). We can rewrite $R_n^*(t) = \sum_{i=1}^n Y_i$ where:

$$Y_i := n^{-1/3}(\varepsilon_i - 1)\left(\tilde{p}(x + n^{-1/3}t - Z_i) - \tilde{p}(x - Z_i)\right).$$

Note that Y_i has expectation 0 and:

$$\begin{aligned}|Y_i| &\leq n^{-1/3}(Ln^{-\gamma/3}|t|^\gamma + Cn^{-1/3}|t|)|\varepsilon_i - 1| \\ &\leq (C'n^{-(1+\gamma)/3} + CKn^{-2/3})|\varepsilon_i - 1|\end{aligned}$$

and because the exponential random variables have finite Orlicz norm $\|\varepsilon_i\|_{\psi_1} < T$, $0 < T < \infty$ (where the Orlicz norm is defined as: $\|X\|_{\psi_1} := \inf\{c > 0 : \mathrm{E}[\exp(|X|/c)] \leq 2\}$), then for some finite constant K':

$$\|Y_i\|_{\psi_1} \leq K'(C'n^{-(1+\gamma)/3} + CKn^{-2/3}).$$

Therefore,

$$\|Y_i\|_{\psi_1}^2 \leq 2(K'C')^2 n^{-(2+2\gamma)/3} + 2(KCK'C')^2 n^{-4/3} \leq 2(K'C')^2(1 + (KC)^2)n^{-(2+2\gamma)/3}.$$

By Example 2.2.12 and Lemma 2.2.10 in [103], for some constants $C_{1,2,3}$:

$$\begin{aligned}\mathrm{P}(|R_n^*(t)| > x) &= \mathrm{P}\left(\left|\sum_{i=1}^n Y_i\right| > x\right) \\ &\leq 2\exp\left\{-\frac{1}{2}\frac{x^2}{C_1 n^{(1-2\gamma)/3} + (C_2 n^{-(1+\gamma)/3} + C_3 n^{-2/3})x}\right\}.\end{aligned}$$

By applying Lemma 2.2.13 in [103] to $R_n^*(t_{-K_n+1}), \ldots, R_n^*(t_{K_n})$ and $\|\cdot\|_1 \lesssim \|\cdot\|_{\psi_1}$:

$$\begin{aligned}\mathrm{E}\left(\max_{-K_n+1\leq i\leq K_n} |R_n^*(t_i)|\right) \lesssim C\Big((C_2 n^{-(1+\gamma)/3} + C_3 n^{-2/3})\log(1 + 2K_n) &\qquad (4.5.9)\\ + C_1^{1/2} n^{(1-2\gamma)/6}\sqrt{\log(1 + 2K_n)}\Big). &\qquad (4.5.10)\end{aligned}$$

Because $K_n = O(n^{(2-\gamma)/3\gamma})$, the result follows. □

Lemma 4.5.4. *For every compact $K \subset \mathbb{R}$ as $n \to \infty$:*

$$\mathbb{W}_n^* \mid Z_1, \ldots, Z_n \rightsquigarrow \sqrt{\frac{g_0(x)}{k(0)}}\mathbb{W}_2 \quad \text{in} \quad \ell^\infty(K). \tag{4.5.11}$$

where $\mathbb{W}_n^*$ is as in (4.5.6) *and $\mathbb{W}_2$ is a two sided Brownian Motion originating in* 0*.*

Proof. First, for $t \in \mathbb{R}$, we give the proof of the marginal convergence

$$\mathbb{W}_n^*(t) \mid Z_1, \ldots, Z_n \rightsquigarrow \mathbb{W}_2(t) \tag{4.5.12}$$

where $\mathbb{W}_2$ is a two sided Brownian Motion originating in 0. After that, we will show conditional asymptotic equicontinuity of the process $\mathbb{W}_n^*$, proving the claim (4.5.11).

Without loss of generality, let $t \neq 0$ in $\mathbb{R}$. We apply Theorem 6.16 in [63] with ξ_{ni} equal to:

$$\xi_{ni} := \frac{(\varepsilon_i - 1)\tilde{f}_t^n(Z_i)}{n^{2/3}} = \frac{(\varepsilon_i - 1)}{n^{1/3}} p(0)\mathbf{1}_{[x,x+n^{-1/3}t]}(Z_i).$$

Therefore we need to show $\forall\, \delta > 0$:

$$\sum_{i=1}^{n} \mathrm{P}(|\varepsilon_i - 1||\tilde{f}_t^n(Z_i)| > \delta n^{2/3} \mid Z_1, \ldots, Z_n) \xrightarrow{G_0} 0, \tag{4.5.13}$$

$$\frac{1}{n^{4/3}} \sum_{i=1}^{n} \mathrm{E}\left((\varepsilon_i - 1)\tilde{f}_t^n(Z_i)\mathbf{1}_{\{|\varepsilon_i-1||\tilde{f}_t^n(Z_i)|\geq\delta n^{2/3}\}} \mid Z_1, \ldots, Z_n\right) \xrightarrow{G_0} 0, \quad \text{and} \tag{4.5.14}$$

$$\frac{1}{n^{4/3}} \sum_{i=1}^{n} \mathrm{Var}\left((\varepsilon_i - 1)\tilde{f}_t^n(Z_i)\mathbf{1}_{\{|\varepsilon_i-1||\tilde{f}_t^n(Z_i)|<\delta n^{2/3}\}} \mid Z_1, \ldots, Z_n\right) \xrightarrow{G_0} \frac{g_0(x)}{k(0)^2} t. \tag{4.5.15}$$

For (4.5.13), use that $|\tilde{f}_t^n(Z_i)| \leq p(0)n^{1/3}$, for all n big enough, unconditionally:

$$\begin{aligned}\sum_{i=1}^{n} \mathrm{P}(|\varepsilon_i - 1||\tilde{f}_t^n(Z_i)| > \delta n^{2/3}) &\leq n\,\mathrm{P}(|\varepsilon_i - 1| > \delta n^{1/3}/p(0)) \\ &= n \int_{\lambda > 1+|t|^{-1}\delta n^{1/3}} e^{-\lambda} d\lambda = n e^{-1-|t|^{-1}\delta n^{1/3}} \to 0.\end{aligned}$$

Now consider (4.5.14) and recall $Z \sim g_0$ and $\varepsilon \sim \mathrm{Exp}(1)$ in what follows. Because $\mathrm{E}\tilde{f}_t^n(Z) = O(1)$ since, as $n \to \infty$:

$$\begin{aligned}\mathrm{E}\tilde{f}_t^n(Z) &= n^{1/3} \int p(0)\mathbf{1}_{[x,x+n^{-1/3}t]}(z) g_0(z)\, dz \\ &= p(0)n^{1/3}\,\mathrm{P}(Z \in [x, x + n^{-1/3}t]) \to t p(0) g_0(x),\end{aligned}$$

and $\varepsilon \perp\!\!\!\perp Z$, we conclude that:

$$\begin{aligned}&\mathrm{E}\left|\frac{1}{n^{2/3}} \sum_{i=1}^{n} \mathrm{E}\left((\varepsilon_i - 1)\tilde{f}_t^n(Z_i)\mathbf{1}_{\{|\varepsilon_i-1||\tilde{f}_t^n(Z_i)|\geq\delta n^{2/3}\}} \mid Z_1, \ldots, Z_n\right)\right| \\ &\lesssim n^{1/3}\mathrm{E}(|\varepsilon - 1|\mathbf{1}_{\{|\varepsilon-1|\geq\delta n^{1/3}/p(0)\}})\mathrm{E}\tilde{f}_t^n(Z) \leq n^{1/3}(1 + \delta n^{1/3})e^{-1-\delta n^{1/3}} \to 0.\end{aligned}$$

For (4.5.15), some more work is needed. Note that:

$$\begin{aligned}(4.5.15) = \frac{1}{n^{4/3}} \sum_{i=1}^{n} \Big\{ &\mathrm{E}\left((\tilde{f}_t^n(Z_i))^2(\varepsilon_i - 1)^2\mathbf{1}_{\{|\varepsilon_i-1|\tilde{f}_t^n(Z_i)<\delta n^{2/3}\}} \mid Z_1, \ldots, Z_n\right) \\ &- \left(\mathrm{E}\left(\tilde{f}_t^n(Z_i)(\varepsilon_i - 1)\mathbf{1}_{\{|\varepsilon_i-1|\tilde{f}_t^n(Z_i)<\delta n^{2/3}\}} \mid Z_1, \ldots, Z_n\right)\right)^2 \Big\}.\end{aligned}$$

First notice, because $\mathrm{E}(\varepsilon - 1) = 0$ and $\varepsilon \perp\!\!\!\perp Z$:

$$\begin{aligned}&\mathrm{E}\left(\frac{1}{n^{4/3}} \sum_{i=1}^{n} \left(\mathrm{E}\left(\tilde{f}_t^n(Z_i)(\varepsilon_i - 1)\mathbf{1}_{\{|\varepsilon_i-1|\tilde{f}_t^n(Z_i)>\delta n^{2/3}\}} \mid Z_1, \ldots, Z_n\right)\right)^2\right) \\ &\leq \frac{1}{n^{1/3}}\mathrm{E}\left((\tilde{f}_t^n(Z))^2(\varepsilon - 1)^2\mathbf{1}_{\{|\varepsilon-1|\tilde{f}_t^n(Z)>\delta n^{2/3}\}}\right) \\ &\leq \frac{1}{n^{1/3}}\mathrm{E}\left((\tilde{f}_t^n(Z))^2(\varepsilon - 1)^2\mathbf{1}_{\{|\varepsilon-1|>\delta n^{1/3}/p(0)\}}\right) \to 0,\end{aligned}$$

because as $n \to \infty$

$$\frac{1}{n^{1/3}}\mathrm{E}\left((\tilde{f}_t^n(Z))^2\right) = n^{1/3} \int p^2(0)\mathbf{1}_{[x,x+n^{-1/3}t]}(z) g_0(z) g_0(z) dz \to p^2(0) g_0(x) t,$$

and for all n big enough:

$$\begin{aligned}\mathrm{E}\left((\varepsilon-1)^2\mathbf{1}_{\{|\varepsilon-1|>\delta n^{1/3}\}}\right) &\lesssim \int_{1+\delta n^{1/3}}^{\infty}(\lambda-1)^2e^{-\lambda}d\lambda \\ &= (\delta n^{1/3}(\delta n^{1/3}+2)+2)e^{-1-\delta n^{1/3}} \to 0. \end{aligned} \tag{4.5.16}$$

Now using that $\mathrm{E}(\varepsilon-1)^2=1$ and that $\varepsilon \perp\!\!\!\perp Z$, we get for the domination term of (4.5.15):

$$\begin{aligned}&\frac{1}{n^{4/3}}\sum_{i=1}^{n}\mathrm{E}\left((\tilde{f}_t^n(Z_i))^2(\varepsilon_i-1)^2\mathbf{1}_{\{|\varepsilon_i-1|\tilde{f}_t^n(Z_i)<\delta n^{2/3}\}}\mid Z_1,\dots,Z_n\right)\\ &=\frac{1}{n^{4/3}}\sum_{i=1}^{n}\mathrm{E}\left((\tilde{f}_t^n(Z_i))^2(\varepsilon_i-1)^2(1-\mathbf{1}_{\{|\varepsilon_i-1|\tilde{f}_t^n(Z_i)\geq\delta n^{2/3}\}})\mid Z_1,\dots,Z_n\right)\\ &=\frac{1}{n^{4/3}}\sum_{i=1}^{n}(\tilde{f}_t^n(Z_i))^2-\frac{1}{n^{4/3}}\sum_{i=1}^{n}\mathrm{E}\left((\tilde{f}_t^n(Z_i))^2(\varepsilon_i-1)^2\mathbf{1}_{\{|\varepsilon_i-1|\tilde{f}_t^n(Z_i)\geq\delta n^{2/3}\}}\mid Z_1,\dots,Z_n\right).\end{aligned}$$

Now we show $\frac{1}{n^{4/3}}\sum_{i=1}^{n}(\tilde{f}_t^n(Z_i))^2 \xrightarrow{G_0} t\frac{g_0(x)}{k^2(0)}$. We apply Lemma 2.3 from the supplement of [47] with $\xi_{ni}:=(\tilde{f}_t^n(Z_i))^2/n^{4/3}$. For all $\varepsilon>0$ and all n big enough:

$$\begin{aligned}\sum_{i=1}^{n}\mathrm{E}|\xi_{ni}|\mathbf{1}_{\{|\xi_{ni}|>\varepsilon\}} &= \frac{1}{n^{4/3}}\sum_{i=1}^{n}\mathrm{E}\left((\tilde{f}_t^n(Z_i))^2\mathbf{1}_{\{(\tilde{f}_t^n(Z_i))^2>\varepsilon n^{4/3}\}}\right)\\ &\leq \frac{1}{n^{4/3}}\sum_{i=1}^{n}\mathrm{E}\left((\tilde{f}_t^n(Z_i))^2\mathbf{1}_{\{p^2(0)>\varepsilon n^{2/3}\}}\right)=0.\end{aligned}$$

Finally note, because $p^2(0)=\frac{1}{k^2(0)}$:

$$\sum_{i=1}^{n}\mathrm{E}\xi_{ni}=n^{1/3}p^2(0)\int\mathbf{1}_{[x,x+n^{-1/3}t]}(z)g_0(z)\,dz\to t\frac{g_0(x)}{k^2(0)}.$$

The fact that the ξ_{ni} are nonnegative and $\sum_{i=1}^{n}\mathrm{E}\xi_{ni}\to t\frac{g_0(x)}{k^2(0)}$ imply $\sum_{i=1}^{n}\mathrm{E}|\xi_{ni}|=O(1)$. Using (4.5.16) we conclude:

$$\begin{aligned}&\frac{1}{n^{4/3}}\sum_{i=1}^{n}\mathrm{E}\left((\varepsilon_i-1)^2(\tilde{f}_t^n(Z_i))^2\mathbf{1}_{\{|\varepsilon_i-1|\tilde{f}_t^n(Z_i)>\delta n^{2/3}\}}\right)\\ &\leq\frac{1}{n^{4/3}}\sum_{i=1}^{n}(\tilde{f}_t^n(Z_i))^2\mathrm{E}\left((\varepsilon-1)^2\mathbf{1}_{\{|\varepsilon-1|>\delta n^{1/3}\}}\right)\to 0.\end{aligned}$$

because $\frac{1}{n^{4/3}}\sum_{i=1}^{n}(\tilde{f}_t^n(Z_i))^2=O_{G_0}(1)$.

Now we verify the behavior of the covariance structure. Note that $\forall s,t\geq 0$ we have the following convergence in probability as $n\to\infty$: $\mathrm{Cov}(\mathbb{W}_n^*(t),\mathbb{W}_n^*(s)\mid Z_1,\dots,Z_n)$ converges in probability under G_0 as $n\to\infty$ to:

$$\begin{aligned}\mathrm{Cov}(\mathbb{W}_n^*(t),\mathbb{W}_n^*(s)) &= n^{1/3}p^2(0)\int(\mathbf{1}_{[x,x+n^{-1/3}t]}-\mathbf{1}_{[0,x]})(z)(\mathbf{1}_{[x,x+n^{-1/3}s]}-\mathbf{1}_{[0,x]})(z)\,dG_0(z)\\ &= n^{1/3}p^2(0)\,\mathrm{P}(Z\in[x,x+n^{-1/3}\min(s,t)])\to\frac{g_0(x)}{k^2(0)}\min(s,t),\end{aligned}$$

conclude the convergence marginal convergence in (4.5.12).

Now to prove

$$\mathbb{W}_n^* \mid Z_1, \ldots, Z_n \rightsquigarrow \sqrt{\frac{g_0(x)}{k(0)}}\mathbb{W}_2 \quad \text{in } \ell^\infty(K) \tag{4.5.17}$$

take the random class of functions (for $\varepsilon \sim \text{Exp}(1)$), for $T > 0$

$$\tilde{\mathcal{F}}_n^{T,\delta} = \{z \mapsto n^{-1/6}(\varepsilon - 1)(\tilde{f}_t^n - \tilde{f}_s^n)(z) : s, t \in \mathbb{R}, |s - t| < \delta, \max\{|s|, |t|\} < T\},$$

and the class of functions $\mathcal{H}_{n,3}^{T,\delta}$ whose functions for $f \in \tilde{\mathcal{F}}_n^{T,\delta}$ are given by $f/(\varepsilon - 1)$ thus without the $\varepsilon - 1$. These classes have square integrable envelopes $\tilde{F}_{n,\delta}$, $H_{n,\delta}$ that satisfy for all $\delta > 0$ small and all n big enough:

$$\begin{aligned}\|\tilde{F}_{n,\delta}\|_2^2 = \text{E}(\varepsilon - 1)^2\|H_{n,\delta}\|_{2,G_0}^2 &= \frac{1}{n^{1/3}}\int (\tilde{f}_t^n - \tilde{f}_s^n)^2(z)g_0(z)\,dz \\ &= n^{1/3}p^2(0)\int \mathbf{1}_{[x+n^{-1/3}s\wedge t, x+n^{-1/3}s\vee t]}(z)g_0(z)\,dz \lesssim p^2(0)g_0(x)\delta.\end{aligned}$$

For fixed $s, t \in \mathbb{R}$, in view of (4.2.10), we prove conditional asymptotic equicontinuity of $\mathbb{W}_n^*$, i.e. $\forall\, \eta > 0$ as $n \to \infty$, followed by $\delta > 0$ the right hand side of the expression below converges to 0 under G_0:

$$\text{P}\left(\sup_{|s-t|<\delta} |\mathbb{W}_n^*(t) - \mathbb{W}_n^*(s)| > \eta \mid Z_1, \ldots, Z_n\right) \leq \frac{1}{\eta}\text{E}\left[\sup_{|s-t|<\delta} |\mathbb{W}_n^*(t) - \mathbb{W}_n^*(s)| \mid Z_1, \ldots, Z_n\right].$$

By the maximal inequality in Theorem 2.14.1 in [103], as $\delta \to 0$

$$\text{E}\left[\sup_{|s-t|<\delta} |\mathbb{W}_n^*(t) - \mathbb{W}_n^*(s)|\right] \lesssim J(1, \tilde{\mathcal{F}}_n^{T,\delta}, L_2)\|\tilde{F}_{n,\delta}\|_2 \lesssim \sqrt{\delta} \to 0.$$

In the last convergence we use that:

$$J(1, \tilde{\mathcal{F}}_n^{T,\delta}, \mathbb{L}_2) := \sup_R \int_0^1 \sqrt{\log N(\eta\|\tilde{F}_{n,\delta}\|_2, \tilde{\mathcal{F}}_n^{T,\delta}, \mathbb{L}_2(R))}\,d\eta,$$

where the $\mathbb{L}_2(R)$ - square distance is given for s, t and $s', t' \in \mathbb{R}$ by:

$$\frac{1}{n^{1/3}}\int \left((\tilde{f}_s^n - \tilde{f}_t^n)(z) - (\tilde{f}_{s'}^n - \tilde{f}_{t'}^n)(z)\right)^2 dR(z),$$

(note that we did not take $\varepsilon - 1$ into account as $\text{E}(\varepsilon - 1)^2 = 1$). Furthermore we use $J(1, \tilde{\mathcal{F}}_n^{T,\delta}, \mathbb{L}_2) < \infty$ as a consequence of Exercise 20 in [103] (p. 221), which implies that the function classes under consideration are VC of uniform bounded index.

All together this proves (4.5.11).

□

Lemma 4.5.5. *Assume Condition 1 holds and that F_0 is continuous at x. For any $T > 0$ and $x \in [0, T)$, as $n \to \infty$, $\forall\, \varepsilon > 0$:*

$$\Pi_n(|\hat{F}_G(x) - F_0(x)| > \varepsilon \mid Z_1, \ldots, Z_n) \xrightarrow{G_0} 0.$$

Proof. Using Proposition 4.3 in [42], we obtain:

$$\mathrm{E}(H_G(x) \mid Z_1, \ldots, Z_n) = \frac{1}{|\alpha| + n} \int_0^x p(x - z)\, d\left(\alpha + \sum_{i=1}^{n} \delta_{Z_i}\right)(z)$$

$$\mathrm{Var}(H_G(x) \mid Z_1, \ldots, Z_n) = \frac{1}{1 + |\alpha| + n} \frac{1}{|\alpha| + n} \int_0^x p^2(x - z)\, d\left(\alpha + \sum_{i=1}^{n} \delta_{Z_i}\right)(z)$$

Thus, using the conditional Chebyshev inequality, $\forall\, \varepsilon > 0$:

$$\Pi_n(|H_G(x) - H_0(x)| > \varepsilon \mid Z_1, \ldots, Z_n) \xrightarrow{G_0} 0 \quad \text{a.s.}$$

Because of the continuity of H_0, the monotonicity of H_G and H_0 and that $H_G(0) = H_0(0) = 0$, $\forall\, T > 0, \varepsilon > 0$:

$$\Pi_n\left(\sup_{0 \le x \le T} |H_G(x) - H_0(x)| > \varepsilon \mid Z_1, \ldots, Z_n\right) \to 0 \quad \text{a.s.}$$

Now, since $x \in [0, T)$ and F_0 is continuous at x, fix $\varepsilon > 0$ and $h > 0$ such that:

$$\left|\frac{H_0(x) - H_0(x - h)}{h} - F_0(x)\right| < \varepsilon$$

By the convexity of $H_G^\star$ (recall that it is the GCM of H_G) we have:

$$\frac{H_G^\star(x) - H_G^\star(x - h)}{h} \le \hat{F}_G(x) \le \frac{H_G^\star(x + h) - H_G^\star(x)}{h}$$

Thus:

$$\begin{aligned}
&\Pi_n(\hat{F}_G(x) - F_0(x) > 2\varepsilon \mid Z_1, \ldots, Z_n) \\
&\qquad \le \Pi_n\left(\frac{H_G^\star(x) - H_G^\star(x - h)}{h} \ge \frac{H_0(x) - H_0(x - h)}{h} + \varepsilon \mid Z_1, \ldots, Z_n\right) \xrightarrow{G_0} 0.
\end{aligned}$$

Similarly:

$$\Pi_n(\hat{F}_G(x) - F_0(x) < -2\varepsilon \mid Z_1, \ldots, Z_n) \xrightarrow{G_0} 0.$$

□

Lemma 4.5.6. *For Z_G as in* (4.2.5)*, there exists a stochastically bounded sequence $\hat{M}_n$ such that:*

$$\Pi_n\left(\inf\{t \in [-xn^{1/3}, (T - x)n^{1/3}] : Z_G(t) - \textit{at is minimal}\} \le \tilde{M}_n \mid Z_1, \ldots, Z_n\right) \xrightarrow{G_0} 1 \quad (4.5.18)$$

Proof. First, for T_G as in (4.2.4), we need to show consistency of T_G at x. By the switch relation $\forall\, \varepsilon > 0$,

$$\begin{aligned}
&\Pi_n(|T_G(F_0(x)) - x| > \varepsilon \mid Z_1, \ldots, Z_n) \\
&\quad = \Pi_n(T_G(F_0(x)) > x + \varepsilon \mid Z_1, \ldots, Z_n) + \Pi_n(T_G(F_0(x)) < x - \varepsilon \mid Z_1, \ldots, Z_n) \\
&\quad \le \Pi_n(\hat{F}_G(x + \varepsilon) < F_0(x) \mid Z_1, \ldots, Z_n) + \Pi_n(\hat{F}_G(x - \varepsilon) > F_0(x) \mid Z_1, \ldots, Z_n) \xrightarrow{G_0} 0
\end{aligned} \quad (4.5.19)$$

where in (4.5.19) we use that F_0 is strictly increasing at x by assumption and Lemma 4.5.5 in the above which gives for every $\eta > 0$: $\Pi_n(|\hat{F}_G(x \pm \varepsilon) - F_0(x \pm \varepsilon)| > \eta \mid Z_1, \ldots, Z_n) \to 0$ under G_0.

The claim (4.5.18) is implied by the unconditional version of this convergence and this entails proving that the rate for $|T_G(F_0(x)) - x|$ is $n^{1/3}$. Because of Lemma 4.5.5, we can proceed by verifying the rest of the conditions of Theorem 3.2.5 in [103] (or see Theorem 3.4.1 for a more general version).

The argmin process we are interested in is: $\tilde{t}_n := \operatorname{argmin}_t\{Z_G(t) - at\}$. Thus note that we can write it as: $\tilde{t}_n = n^{1/3} \operatorname{argmax}_t\{-\mathbb{M}_n(t)\}$ where

$$\mathbb{M}_n(t) := \int (p(x+t-z) - p(x-z) - F_0(x)t)\, d(G - \mathbb{G}_n + G_0)(z) + \int (p(x+t-z) - p(x-z))\, d(\mathbb{G}_n - G_0)(z) - atn^{-1/3}.$$

Moreover define the deterministic function;

$$\mathbb{M}(t) := \int (p(x+t-z) - p(x-z) - F_0(x)t)\, dG_0(z)$$

Note $\mathbb{M}_n(0) = \mathbb{M}(0) = 0$. Moreover note that as a consequence of Taylor's theorem $\exists\, \delta_0 > 0$: $\forall t : |t| < \delta_0$:

$$-\mathbb{M}(t) + \mathbb{M}(0) \lesssim -|t|^2$$

Following Theorem 3.2.5. in [103], define $d(s,t) := |s-t|$. Consider the class of functions $m_t : \mathbb{R} \to \mathbb{R}$, for $\eta > 0$:

$$\mathcal{M}_\eta = \{m_t : m_t(z) = p(x+t-z) - p(x-z) - F_0(x)t,\ 0 < t < \eta\}$$

We show that the envelope $M_\eta = \sup_{0<t<\eta} |m_t|$ of $\mathcal{M}_\eta$,

$$\int M_\eta^2(z) g_0(z) dz \lesssim \int (p(x-z+\eta) - p(x-z))^2 g_0(z)\, dz + \eta^2 F_0(x)$$

where $\int (p(x-z+\eta) - p(x-z))^2 g_0(z)\, dz = p^2(0) g_0(x) \eta^2 + o(\eta^2)$. Thus for all η small enough:

$$\int M_\eta^2(z) g_0(z)\, dz \lesssim (p^2(0) g_0(x) + F_0(x))\eta^2$$

Similarly if we define the class of functions for $\eta > 0$:

$$\tilde{\mathcal{M}}_\eta = \{\tilde{m}_t : \tilde{m}_t(z) = p(x+t-z) - p(x-z) : 0 < t < \eta\}$$

as aconsequence of the above, for the envelope $\tilde{M}_\eta := \sup_{0<t<\eta} |\tilde{m}_t|$ we have:

$$\int \tilde{M}_\eta^2(z) g_0(z) dz \lesssim p^2(0) g_0(x) \eta^2.$$

By Proposition G.10 in [42], for independent $Q \sim \mathrm{DP}(\alpha)$, $\mathbb{B}_n \sim \mathrm{DP}(n\mathbb{G}_n)$, and $V_n \sim \mathrm{Be}(|\alpha|, n)$, the following representation in distribution of $G \sim \mathrm{DP}(\alpha + n\mathbb{G}_n)$: $G = V_n Q + (1 - V_n)\mathbb{B}_n$, and thus

$$\mathbb{M}_n(t) - \mathbb{M}(t) = (1 - V_n) \int (p(x+t-z) - p(x-z) - F_0(x)t)\ d(\mathbb{B}_n - \mathbb{G}_n)(z) \qquad (4.5.20)$$

$$+ V_n \int (p(x+t-z) - p(x-z) - F_0(x)t)\; d(Q - \mathbb{G}_n)(z) \tag{4.5.21}$$

$$+ \int (p(x+t-z) - p(x-z))\, d(\mathbb{G}_n - G_0)(z) - atn^{-1/3} \tag{4.5.22}$$

By applying the maximal inequality in Theorem 2.14.1 in [103]

$$\begin{aligned} \mathrm{E}\left[\sup_{|t|<\delta} \sqrt{n}|\mathbb{M}_n - \mathbb{M}|(t)\right] &\lesssim J(1, M_\delta, \mathbb{L}_2)\sqrt{\int M_\delta^2\, dG_0} + \sqrt{n}\frac{1}{n}\delta^\gamma + |a|\delta\sqrt{n}n^{-1/3} \\ &\quad + J(1, \tilde{M}_\delta, \mathbb{L}_2)\sqrt{\int \tilde{M}_\delta^2\, dG_0} \lesssim \delta + n^{-1/2}\delta^\gamma + a\delta n^{1/6} \\ &\sim \delta(1 + an^{1/6}) \end{aligned}$$

Because δ must satisfy: $\delta \geq n^{-1/3}$.

All together the above proves that the rate for $\operatorname{argmax}_t\{-\mathbb{M}_n(t)\}$ is δ^{-1}. □

4.5.2 Review of the integral equation

A very important question regarding integral equation (4.1.1) is whether there exist a function p satisfying it. In this section, we discuss conditions on the kernel k that are sufficient for the existence of a solution p to (4.1.1). This appendix is just a review based on the known results in [60] and [108] (no new results are given). We emphasize that the main results of this chapter (Theorem 4.2.2 and Theorem 4.2.1) only require p to satisfy Condition 2, which is a Hölder continuity condition on p itself. The sufficient conditions discussed in this appendix, centering on the special case of *decreasing* densities k, are therefore not necessary for the main results; they serve to illustrate a concrete and well-studied class of kernels for which Condition 2 can be verified. As demonstrated in the simulation study (Section 3), Condition 2 is also met by a wider variety of kernels that are not necessarily decreasing. Suppose:

Condition 3. $k : [0, \infty) \to [0, \infty)$ is a decreasing density and $0 < k(0) < \infty$.

Without loss of generality we may take k to be right continuous. For a kernel satisfying Condition 3, properties of the associated function p solving equation (4.1.1) can be obtained using some results from *renewal theory* (see e.g. [50]). Define the distribution function J on $[0, \infty)$ by

$$J(x) := 1 - \frac{k(x)}{k(0)}.$$

Note that $\mu_J = \int_0^\infty x\, dJ(x) = 1/k(0)$. Renewal theory assures the existence of a renewal function m on $[0, \infty)$ satisfying the renewal equation

$$m(x) = J(x) + \int_0^x m(x-y)\, dJ(y).$$

Integrating this equation with respect to x, we obtain

$$\mathbf{1} * m = (\mathbf{1} + m) * J. \tag{4.5.23}$$

Writing

$$p(x) = \frac{1 + m(x)}{k(0)} = \mu_J(1 + m(x)) \quad \text{for } x \geq 0, \tag{4.5.24}$$

it follows that p solves (4.1.1). Indeed, for $x > 0$,

$$p * k(x) = \frac{1}{k(0)}(\mathbf{1} + m) * k(x) = (\mathbf{1} + m) * (\mathbf{1} - J)(x) = \mathbf{1} * \mathbf{1}(x)$$

by the definition of J and (4.5.23).

Lemma 4.5.7 (Lemma 1 in [108])**.** *Let the density k on $[0, \infty)$ satisfy Condition 3. Then*

a) there is a uniquely determined solution p to (1) which is bounded on bounded intervals and vanishes on $(-\infty, 0)$

b) writing $J^{*1} = J$ and $J^{*n} = J^{*(n-1)} * J$ for $n = 2, 3, \ldots$, this uniquely determined p on $[0, \infty)$ is given by

$$p(x) = \frac{1}{k(0)}\left(1 + \sum_{n=1}^{\infty} J^{*n}(x)\right) = \frac{1}{k(0)}\left(1 + \sum_{n=1}^{\infty} \mathrm{P}\left(\sum_{j=1}^{n} T_j \le x\right)\right), \tag{4.5.25}$$

where $T_1, T_2, \ldots$ is an iid sequence of random variables such that T_1 has distribution function J.

c) the function p is nonnegative and nondecreasing on $[0, \infty)$.

d)

$$\lim_{x \to \infty} \frac{p(x)}{x} = 1.$$

Proof from [108]. All statements follow from relation (5) together with known results from renewal theory. See, for example, [50] Theorem 10.1.11 for a), Lemma 10.1.7 for b) and Theorem 10.2.3 for d). Part c) is obvious from the definition of a renewal function. □

Because of Condition 2, which we use to derive our main result Theorem 4.2.2, one interesting question is to know conditions on k under which p is Lipschitz continuous on intervals $[0, T]$ for any bounded T. Functions p associated with kernels satisfying Condition 3 do not have this property without further assumptions. It follows from (4.5.25) that a jump of k at a point $x > 0$ causes infinitely many jumps in the function p. The condition on k given below in Lemma 4.5.8 is sufficient for k to guarantee that p is Lipschitz continuous on bounded intervals in $[0, \infty)$. It gives thus a sufficient condition for Condition 2 and thus also for Condition 1 to hold.

Lemma 4.5.8 (Lemma 1 in [109])**.** *Let $0 < T < \infty$. Suppose the density k can be written as*

$$k(x) = k(0)\left(1 + \int_0^x l(u)\, du\right), \quad x \in [0, T]$$

for some bounded Borel measurable function $l : [0, T] \to \mathbb{R}$. Then the unique continuous (on $(0, T)$) solution p of (4.1.1) allows the representation

$$p(x) = \frac{1}{k(0)}\left(1 + \int_0^x q(u) du\right), \quad x \in [0, T],$$

where q is a bounded Borel measurable function on $[0, T]$.

Proof from [109]. Consider the type-II Volterra convolution equation

$$q(x) + \int_0^x l(x-u)q(u)du = -l(x),$$

or, equivalently

$$q + l * q = -l.$$

By Theorem 3.5 in Gripenberg, Londen and Staffans (1990), it follows that the solution q of this equation uniquely exists. It is bounded and Borel measurable on $[0, T]$ whenever l is. Now define $p = k(0)^{-1}(\mathbf{1} + \mathbf{1} * q)$ and observe

$$k * p = k(0)(\mathbf{1} + \mathbf{1} * l) * \frac{\mathbf{1} + \mathbf{1} * q}{k(0)} = \mathbf{1} * \mathbf{1} + \mathbf{1} * \mathbf{1} * (q + l + q * l) = \mathbf{1} * \mathbf{1}.$$

□

4.6 Bibliographic Remarks

There is a vast literature on deconvolution problems, especially in the frequentist nonparametric literature, where the nonparametric maximum likelihood estimator (NPMLE) is a central focus. For the special case where the kernel k is a decreasing density on $[0, \infty)$ and $F_0(0) = 0$, [54] study the NPMLE for F_0, demonstrating its consistency and conjecturing its asymptotic distribution. They also introduce the broader problem of one-sided errors in convolution models, deriving NPMLEs for the c.d.f. in this context. While these NPMLEs often lack explicit expressions and require iterative computation, some particular cases yield more direct solutions. For instance, uniform deconvolution is addressed by [104, 106], and more recently by [107] who propose kernel-based estimators and inversion formulas. Exponential deconvolution is tackled by [60, 61]. For situations lacking such explicit forms, [109] circumvent this by proposing an isotonic inverse estimator. Many of these maximum likelihood estimators, and the works surrounding them, share cube root asymptotics similar to the Grenander estimator for decreasing densities and NPMLEs in current status censoring problems (c.f. [53]), with a general focus on deriving the asymptotic distributions of these monotonic estimators. An alternative approach to deconvolution utilizes the convolution structure for inversion via Fourier transform techniques. Kernel estimators based on this methodology are introduced by several authors (c.f. [33] and [28]), using direct inversion formulas for gamma and Laplace deconvolution are found in [105]. While these kernel-based methods can achieve faster convergence rates if the unknown F_0 is smooth, a notable disadvantage is that their resulting estimators for F_0 are not inherently monotone, unlike the NPMLE and isotonic inverse estimators.

Within the broader field of deconvolution, a lot of work is done to address the estimation of a signal distribution f observed with additive, known-distribution noise. This noise generally is conceptualized as centered, differentiating it from one-sided error scenarios. Foundational studies on convergence rates and optimality for kernel-based estimators include those by [12], [97], [98], [33], and [31]. Establishing sharp asymptotic optimality further is done by [9] and [10, 11]. Adaptive methodologies, particularly for bandwidth selection in models with known error distributions, also are extensively developed. Some examples include wavelet-based methods [82], direct bandwidth selection techniques [26], and penalized projection strategies [20], with comprehensive treatments available in works like [74]. The estimation of the cumulative distribution function (c.d.f.) in convolution models also receives attention. Contributions

from [116], [33], [55], [24], [23], and [25] primarily focus on pointwise estimation procedures, a consequence of the c.d.f. not being square-integrable on $\mathbb{R}$. The latter two studies address c.d.f. estimation with unknown error distributions, contingent on specific tail decay properties of the error's characteristic function, and achieve optimal rates for Sobolev-class target functions. It is worth noting that the deconvolution of non-negative variables also emerges in applied contexts such as actuarial and insurance modeling. Finally, some financial applications see contributions from [59] and [86] addressing one-sided errors, focusing on optimal adaptive estimation in non-parametric regression where errors (often exponential) are not assumed to be centered.

A similar methodology to the one proposed in this chapter, using an unconstrained conjugate prior for the distribution function of the observables and projecting the posterior samples into the space of monotone functions was very recently adopted in isotonic regression settings (c.f. [17, 112, 113]), isotonic density estimation (c.f. [18]), ODEs settings (c.f. [5, 6]) and for Wicksell's problem ([44],[47]). [5, 6] develop Bayesian two-step methods for parameter inference in ODE models by embedding them in a nonparametric regression framework with B-spline priors. The parameter posterior is obtained by minimizing a discrepancy between the estimated regression function and the ODE solution, either via derivative matching or Runge–Kutta approximations. [6] further establishes a Bernstein–von Mises theorem for the Runge–Kutta-based method, showing that the resulting Bayes estimator is asymptotically efficient. However, their framework relies on a very different type of prior modelling and the projection method does not entail isotonization. In [17], the authors study a nonparametric Bayesian regression model for a response variable Y with respect to a predictor variable $X \in [0, 1]$, given by $Y = f(X) + \varepsilon$, where f is a monotone increasing function and ε is a mean-zero random error. They propose a "projection-posterior" estimator f^*, constructed using a finite random series of step functions with normal basis coefficients as a prior for f. Their main finding is that when f^* is centered at the maximum likelihood estimator (MLE) and rescaled appropriately, the Bernstein-von Mises (BvM) theorem does not hold. However, when f^* is instead centered at the true function f_0, a BvM result is obtained. This phenomenon resonates with the known inconsistency of the bootstrap for the Grenander estimator in isotonic regression (c.f. [93]) and represents a key similarity with the present work.

One of the key advantages our approach shares with the works in [5, 6, 17, 112, 113] is the computational efficiency, driven by posterior conjugacy, as well as the relative simplicity of the asymptotic analysis.

Chapter 5

Proximal inference based on mixtures of linear models

5.1 Introduction

In causal inference using observational data, a frequent assumption is that there is no unmeasured confounding. This means that the available covariates are informative enough to ensure that individuals within each stratum of the covariates are comparable across different treatment groups. However, this assumption is often met with doubt, as it requires researchers to accurately capture all relevant confounders, which is rarely achievable in practice. Confounding mechanisms are not always fully identifiable from the measured covariates, meaning that these variables may only serve as imperfect representations of the true confounders.



Figure 5.1: A Directed Acyclic Graph (DAG) for the classical setting of proximal inference.

Proximal inference is a method used to uncover treatment effects when standard methods fall short due to unobserved confounders. The key idea of this method is to leverage auxiliary data, the so called *proxy variables*, which carry enough information about the unobserved confounding factors to allow for the estimation of treatment effects. Negative control exposure (NCE) and negative control outcome (NCO) variables are examples of auxiliary data used in this context. An NCO variable Z is a variable known not to be causally affected by an exposure

A and an NCE variable W is a variable known not to cause an outcome Y. See Figure 5.1 for a typical graph that shows the relations of the proxy variables with the other variables (U contains the unobserved confounders).

There has been a surge in research on formal methods in proximal inference (see [70, 71, 75, 94, 95]). These methods typically depend on assumptions like linear models for outcomes and unmeasured confounders (see [36–38, 111]), rank preservation (see [100]), monotonicity (see [96]), or categorical unmeasured confounders (see [94]) to identify treatment effects. In [75] and [21] significant contributions were made by presenting conditions for nonparametric identification of treatment effects using treatment and outcome proxies within point treatment contexts.

The method they present relies on the existence of a *bridge function* that connects the observed data. This function, denoted as $(a, w) \mapsto h_a(w)$, must satisfy the following relationship for each $A = a$:

$$\mathrm{E}(Y \mid Z, A = a) = \mathrm{E}\left(h_a(W) \mid Z, A = a\right) \qquad Z - \text{a.s.} \tag{5.1.1}$$

Under (5.1.1) and the following additional assumptions:

A. $Y(a) = Y$ if $A = a$,

B. $(Y(a), W) \perp\!\!\!\perp (A, Z) \mid U$,

C. For any set $\mathfrak{Z}$ with $\mathrm{P}(Z \in \mathfrak{Z}) = 1$, the set of conditional distributions $\{\mathcal{L}(U \mid Z = z, A = a) : z \in \mathfrak{Z}\}$ is complete,

the average treatment effect of setting A to a is $\mathrm{E}\, Y(a) = \mathrm{E}\, h_a(W)$. Note that condition B. implies $Y(a) \perp\!\!\!\perp A \mid U$, $Y \perp\!\!\!\perp Z \mid A, U$ and $W \perp\!\!\!\perp (A, Z) \mid U$. The average treatment effect $\mathrm{E}\, Y(a)$ can be estimated from observational data, without needing direct information about the unmeasured confounder U, if the function h_a is known. One of the greatest challenges of the approach based on bridge functions lies in finding the function h_a. In the machine learning literature adversarial methods have been developed to estimate h_a from the data (see [64, 68]). However in many cases the existence of such function is not clear. A simple setting in which the existence of a bridge function is guaranteed is under the assumptions of linear models for outcomes and unmeasured confounders (see [36–38, 111]). However this model is too simple to handle most of the applications.

In this chapter, we make two key contributions. First, we introduce a causal model based on mixtures of linear models for the regression of the outcome Y on the unmeasured confounders and on the covariates. This approach offers flexibility in modeling the distribution of the outcome Y given the covariates A and Z. Importantly, we prove that this extension allows the treatment effect to be identifiable, meaning we can derive a closed-form expression for the bridge function h_a. Second, we propose a Bayesian nonparametric estimation method. This method models the distribution of the outcome given the covariates using a probit stick-breaking process, with mixing weights that depend on the covariates. We also prove that our methodology is consistent across a range of true underlying causal models. The key advantage of our approach is that it eliminates concerns about the existence of the bridge function h_a, as we demonstrate that such a function in this context exists and has an explicit form. However, a limitation of our method is that we still rely on the assumptions of a simpler linear model for the distribution of W given the covariates A and Z.

The structure of the chapter is as follows: In Section 5.2, we begin by revisiting the classical linear model (see [36–38, 111]) to provide a solid foundation for understanding

our extension to mixtures of linear models. We then introduce our proposed model, along with a submodel that utilizes simplified mixing weights. In Section 5.3, we present the Bayesian nonparametric estimation procedure based on a dependent stick-breaking process and demonstrate that it achieves consistency for a range of underlying causal models.

5.2 Mixture of linear models

The classical linear model.

One simple case where the form of h_a is known is if we use the restrictive assumption of linear models for the regressions of Y and W on the variables A, U, Z and of U on A and Z:

$$\mathrm{E}(Y \mid A, U, Z) = \beta_0^y + \beta_1^y A + \beta_2^y U \tag{5.2.1}$$

$$\mathrm{E}(W \mid A, U, Z) = \beta_0^w + \beta_1^w U \tag{5.2.2}$$

$$\mathrm{E}(U \mid A, Z) = \beta_0^u + \beta_1^u A + \beta_2^u Z \tag{5.2.3}$$

By taking conditional expectations given A, Z across the two equations and after some algebra (provided that $\beta_1^w \neq 0$), we can obtain the bridge function h_a as:

$$\mathrm{E}(Y \mid A, Z) = \mathrm{E}(\beta_0^y - \beta_2^y \beta_0^w / \beta_1^w + \beta_1^y A + \beta_2^y / \beta_1^w W \mid A, Z).$$

In the display above we can read out the form of h_a and $\mathrm{E}\, Y(a+1) - \mathrm{E}\, Y(a) = \beta_1^y$.

Since (5.2.1)-(5.2.3) ensure that the regressions of Y on A, Z and of W on A, Z are linear, there exist coefficients θ_i^y, θ_i^w for $i = 1, 2, 3$:

$$\mathrm{E}(Y \mid A, Z) = \theta_0^y + \theta_1^y A + \theta_2^y Z, \tag{5.2.4}$$

$$\mathrm{E}(W \mid A, Z) = \theta_0^w + \theta_1^w A + \theta_2^w Z, \tag{5.2.5}$$

and we can obtain after some algebra (if $\theta_2^w \neq 0$, thus $\beta_2^u \neq 0$) that the average treatment effect is given by:

$$\beta_1^y = \theta_1^y - \theta_2^y \theta_1^w / \theta_2^w.$$

Note that within this model, the assumptions (5.2.1)-(5.2.3) remove the need of the completeness assumption of the law of $\mathcal{L}(U \mid Z, A = a)$. It is easy to show that if instead of (5.2.1)-(5.2.3) we assume the completeness of the law of $\mathcal{L}(U \mid Z, A = a)$ (which intuitively tells that Z must be sufficiently informative about U) with A.-C. and that the regressions of Y and W on A, Z are linear as in (5.2.4)-(5.2.5), with $\theta_2^w \neq 0$, then we can also obtain h_a and thus $\mathrm{E}\, Y(a)$.

Although this model is mathematically intriguing because it offers a closed-form example of a bridge function, its reliance on linearity assumptions renders it unsuitable for many scenarios. In the following subsection we will introduce a generalization of this model based on mixture of normals and we will show that it is suitable for a much larger class of underlying models. This model will also provide a closed form expression for the bridge function h_a.

Extension 1 of the linear model.

The underlying model. We retain the classical proximal DAG of Figure 5.1 and the linear regressions (5.2.2)-(5.2.3) of the proxy W and of the confounder U, but we relax the linearity of the outcome regression (5.2.1) to

$$\mathrm{E}(Y \mid A, U, Z) = f_0(A) + f_2(A)\, U,$$

for arbitrary measurable functions f_0, f_2: the dependence on the exposure is now unrestricted, and only linearity in the unobserved confounder is kept (the latter cannot be relaxed; see Remark 5.2.2).

The working model. The underlying model leaves f_0, f_2 in $\mathrm{E}(Y \mid A, U, Z) = f_0(A) + f_2(A)U$ entirely free as functions of A. We approximate this regression function by letting the coefficients of the classical linear model vary across a discrete latent *stratum*: a latent variable S, drawn given A alone, indexes a family of classical models for the outcome,

$$\mathrm{E}(Y \mid A, U, S) = \beta^y_{0,S} + \beta^y_{1,S} A + \beta^y_{2,S} U, \tag{5.2.6}$$

while W and U keep the unstratified regressions (5.2.2)–(5.2.3), i.e. S enters the outcome mechanism only. The DAG of Figure 5.2a displays this construction.



(a) DAG with latent node S. (b) Marginal DAG after integrating out S.

Figure 5.2: Extension 1. a: the DAG that motivates the shape of the working model, with the latent stratum variable S entering the outcome mechanism; b: its marginalization over S, which coincides with the classical proximal DAG of Figure 5.1 on which the underlying model is based. The DAG a is motivation only and is nowhere assumed.

Marginalising out S returns the DAG of Figure 5.2b, which coincides with the classical proximal DAG of Figure 5.1 on which the underlying model is based: the DAG implies $S \perp\!\!\!\perp (W, U, Z) \mid A$ and $Y \perp\!\!\!\perp Z \mid A, U, S$, whence

$$\mathrm{E}(Y \mid A, U, Z) = \mathrm{E}(\beta^y_{0,S} \mid A) + \mathrm{E}(\beta^y_{1,S} \mid A)\, A + \mathrm{E}(\beta^y_{2,S} \mid A)\, U,$$

which is of the underlying form $f_0(A) + f_2(A)U$: as S ranges over strata with weights depending on A, the stratum average $A \mapsto \mathrm{E}(\beta^y_{2,S} \mid A)$ can approximate an arbitrary function of A, which is exactly the extra flexibility that the underlying model allows over the classical one. The same independences turn (5.2.6) into stratum-wise linear regression of the observables,

$$\mathrm{E}(Y \mid A, S, Z) = \theta^y_{0,S} + \theta^y_{1,S} A + \theta^y_{2,S} Z, \qquad \mathrm{E}(W \mid A, Z) = \theta^w_0 + \theta^w_1 A + \theta^w_2 Z,$$

with $\theta^y_{i,S}$ an explicit affine function of $\beta^y_{0,S}, \beta^y_{1,S}, \beta^y_{2,S}$ and θ^w_i as in the classical model (the exact substitution is carried out in the remark following the theorem). Averaging over $S \mid A$ turns this into a mixture representation of the observables,

$$S \mid A \sim (w_k(A))_{k \in \mathbb{N}} \tag{5.2.7}$$

$$Y \mid A, Z \sim \sum_{k=1}^{\infty} w_k(A)\, N(\theta^y_{0,k} + \theta^y_{1,k} A + \theta^y_{2,k} Z, \sigma^2_{y,k}) \tag{5.2.8}$$

$$W \mid A, Z \sim N(\theta^w_0 + \theta^w_1 A + \theta^w_2 Z, \sigma^2_w), \tag{5.2.9}$$

and it is a model of this form — with $(\theta^y_{i,k})_{i\in\{0,1,2\},k\in\mathbb{N}}$, $(\theta^w_i)_{i\in\{0,1,2\}}$ and $(w_k(A))_{k\in\mathbb{N}}$ now treated as free parameters, rather than fixed at the values determined by the betas above — that we fit to a sample (Z, A, Y, W) via the Bayesian nonparametric procedure of Section 5.3.

The DAG, the latent S, and the betas above were introduced only to motivate this particular family of mixtures as working model; nothing about them is assumed of the mechanism that generates the data. What matters instead is a property of the family (5.2.7)–(5.2.9) that we call *stratum representation* of the observables. This concept is formalized in part (b) of the theorem below. In essence a *stratum representation* of the observables is an enrichment of the law of (Z, A, Y, W) with a latent variable S that allows us to express the conditional expectation of $Y \mid A, Z$ as a mixture of linear regressions within each strata. Part (a) in the theorem below identifies the ATE of the underlying model by an explicit functional of the laws of the observables, and part (b) shows that every stratum representation collapses this functional into the closed form computed by the fitted working model.

Theorem 5.2.1 (Extension 1)**.** *Suppose (U, Z, A, Y, W) follows the underlying model: the DAG of Figure 5.1 (hence the consistency A. and the conditional independence B.) and*

1. $\mathrm{E}(Y \mid A, U, Z) = f_0(A) + f_2(A)U$*, for some measurable functions* f_0, f_2*,*
2. $\mathrm{E}(W \mid A, U, Z) = \beta^w_0 + \beta^w_1 U$*, with* $\beta^w_1 \neq 0$*,*
3. $\mathrm{E}(U \mid A, Z) = \beta^u_0 + \beta^u_1 A + \beta^u_2 Z$*, with* $\beta^u_2 \neq 0$*,*

and fix $a \in \mathcal{A}$ with $\mathrm{Var}(Z \mid A = a) \in (0, \infty)$*. Then:*

(a) *(Identification) The regressions of the observables are,*

$$\mathrm{E}(Y \mid A = a, Z) = g_0(a) + g_2(a)Z, \qquad g_2(a) = f_2(a)\beta^u_2, \tag{5.2.10}$$

$$\mathrm{E}(W \mid A, Z) = \theta^w_0 + \theta^w_1 A + \theta^w_2 Z, \qquad \theta^w_2 = \beta^w_1 \beta^u_2 \neq 0, \tag{5.2.11}$$

and the ATE is given by

$$\mathrm{E}\, Y(a) = \mathrm{E}[\mathrm{E}(Y \mid A = a, Z)] + \gamma(a)\, \mathrm{E}\big[\mathrm{E}(W \mid A, Z) - \mathrm{E}(W \mid A = a, Z)\big], \tag{5.2.12}$$

where $\gamma(a)$ is identified from the laws of the observables:

$$\gamma(a) = \frac{\mathrm{Cov}(Z, \mathrm{E}(Y \mid A = a, Z) \mid A = a)}{\mathrm{Cov}(Z, \mathrm{E}(W \mid A = a, Z) \mid A = a)} = \frac{g_2(a)}{\theta^w_2}.$$

(b) *(Closed form under a stratum representation) Let (Z, A, Y, W, S) be any stratum representation of the observables: a random vector, on some probability space, whose first four coordinates are distributed as the observables and such that*

$$S \perp\!\!\!\perp Z \mid A \qquad \text{and} \qquad \mathrm{E}(Y \mid A, S, Z) = \theta^y_{0,S} + \theta^y_{1,S} A + \theta^y_{2,S} Z, \tag{5.2.13}$$

for some coefficients $(\theta^y_{i,s})_{i\in\{0,1,2\},s\in\mathbb{N}}$. Then, for $\boldsymbol{\theta}_i(a) = \mathrm{E}(\theta^y_{i,S} \mid A = a)$,

$$\frac{\boldsymbol{\theta}_2(a)}{\theta^w_2} = \gamma(a),$$

and the ATE takes the closed form

$$\begin{aligned}\mathrm{E}\,Y(a) &= \mathrm{E}\,\mathrm{E}(Y \mid A=a, Z) + \frac{\boldsymbol{\theta}_2(a)}{\theta_2^w}\,\mathrm{E}\big[\mathrm{E}(W \mid A, Z) - \mathrm{E}(W \mid A=a, Z)\big] \\ &= \boldsymbol{\theta}_0(a) + \boldsymbol{\theta}_1(a)\,a + \frac{\boldsymbol{\theta}_2(a)}{\theta_2^w}\Big(\mathrm{E}W - \theta_0^w - \theta_1^w a\Big). \end{aligned} \tag{5.2.14}$$

Remark 5.2.1 (A stronger but easier to interpret sufficient condition for part (b)). The stratum representation condition (5.2.13) can be replaced by the stronger, more directly interpretable assumption that $S \perp\!\!\!\perp (U, Z) \mid A$ and

$$\mathrm{E}(Y \mid A, U, S) = \beta^y_{0,S} + \beta^y_{1,S} A + \beta^y_{2,S} U,$$

i.e. that (5.2.6) holds within every stratum. Together with assumption 3 of the theorem ($\mathrm{E}(U \mid A, Z) = \beta^u_0 + \beta^u_1 A + \beta^u_2 Z$), iterated expectation gives

$$\mathrm{E}(Y \mid A, S, Z) = (\beta^y_{0,S} + \beta^y_{2,S}\beta^u_0) + (\beta^y_{1,S} + \beta^y_{2,S}\beta^u_1)\,A + \beta^y_{2,S}\beta^u_2\,Z,$$

which is exactly (5.2.13) with $\theta^y_{0,S} = \beta^y_{0,S} + \beta^y_{2,S}\beta^u_0$, $\theta^y_{1,S} = \beta^y_{1,S} + \beta^y_{2,S}\beta^u_1$, and $\theta^y_{2,S} = \beta^y_{2,S}\beta^u_2$. This is strictly stronger than (5.2.13) because it constrains the stratum structure at the level of the unobserved confounder U, not merely at the level of the observables. The theorem is stated in the weaker form (5.2.13) because that is all that is needed.

Proof. Part (a). Write $m_Y(a, z) = \mathrm{E}(Y \mid A = a, Z = z)$ and $m_W(a, z) = \mathrm{E}(W \mid A = a, Z = z)$. By the tower property of conditional expectation,

$$\begin{aligned}\mathrm{E}(Y \mid A, Z) &= f_0(A) + f_2(A)\mathrm{E}(U \mid A, Z) \\ &= f_0(A) + f_2(A)(\beta^u_0 + \beta^u_1 A + \beta^u_2 Z) =: g_0(A) + g_2(A)Z,\end{aligned}$$

where $g_2(A) = f_2(A)\beta^u_2$, and g_0 collects the remaining terms; this is (5.2.10). Similarly, by iterated expectation,

$$\mathrm{E}(W \mid A, Z) = \beta^w_0 + \beta^w_1 \mathrm{E}(U \mid A, Z) = \theta^w_0 + \theta^w_1 A + \theta^w_2 Z,$$

with $\theta^w_2 = \beta^w_1\beta^u_2 \neq 0$, which is (5.2.11).

The slope $g_2(a)$ is identified from the observable regression $m_Y(a, \cdot)$: letting Q_a denote the law of $Z \mid A = a$ and using $\mathrm{Var}(Z \mid A = a) > 0$, it admits the explicit form

$$g_2(a) = \frac{\mathrm{E}[(Z - \mathrm{E}(Z \mid A = a))\, m_Y(a, Z) \mid A = a]}{\mathrm{Var}(Z \mid A = a)}. \tag{5.2.15}$$

To verify (5.2.15), write $m_Y(a, Z) = g_0(a) + g_2(a)Z$ and note that

$$\begin{aligned}\mathrm{E}[(Z - \mu_a) m_Y(a, Z) \mid A = a] &= \mathrm{E}[(Z - \mu_a)(g_0(a) + g_2(a)Z) \mid A = a] \\ &= g_2(a)\mathrm{E}[(Z - \mu_a)Z \mid A = a] = g_2(a)\mathrm{Var}(Z \mid A = a),\end{aligned}$$

where $\mu_a = \mathrm{E}(Z \mid A = a)$; dividing by $\mathrm{Var}(Z \mid A = a)$ yields (5.2.15). (If $Z \in \{0, 1\}$, this reduces to $g_2(a) = m_Y(a, 1) - m_Y(a, 0)$.) In the same way, since $m_W(a, z) = h_0(a) + \theta^w_2 z$ for fixed a,

$$\theta^w_2 = \frac{\mathrm{E}\big[(Z - \mu_a)\, m_W(a, Z) \mid A = a\big]}{\mathrm{Var}(Z \mid A = a)}. \tag{5.2.16}$$

Combining (5.2.15)–(5.2.16) yields

$$\frac{g_2(a)}{\theta_2^w} = \frac{\mathrm{E}[(Z-\mu_a)m_Y(a,Z) \mid A=a]}{\mathrm{E}[(Z-\mu_a)m_W(a,Z) \mid A=a]} = \frac{\mathrm{Cov}(Z, m_Y(a,Z) \mid A=a)}{\mathrm{Cov}(Z, m_W(a,Z) \mid A=a)} = \gamma(a),$$

the denominator being nonzero by $\theta_2^w \neq 0$ and $\mathrm{Var}(Z \mid A=a) > 0$. Finally, using the consistency A. and $Y(a) \perp\!\!\!\perp A \mid (U,Z)$, which is implied by B., together with the tower property of conditional expectation, we obtain

$$\begin{aligned}\mathrm{E}\, Y(a) &= \mathrm{E}[\mathrm{E}(Y(a) \mid A=a, U, Z)] = \mathrm{E}[f_0(a) + f_2(a)U] = f_0(a) + f_2(a)\mathrm{E}(U) \\ &= \mathrm{E}[\mathrm{E}(Y \mid A=a, Z)] + f_2(a)\mathrm{E}(\mathrm{E}(U \mid Z) - \mathrm{E}(U \mid A=a, Z)) \\ &= \mathrm{E}[\mathrm{E}(Y \mid A=a, Z)] + \frac{g_2(a)}{\theta_2^w}\mathrm{E}(\mathrm{E}(W \mid A, Z) - \mathrm{E}(W \mid A=a, Z)),\end{aligned}$$

which is (5.2.12).

Part (b). A stratum representation leaves the law of the observables unchanged, so every conditional expectation below may be computed on the space carrying it. By $S \perp\!\!\!\perp Z \mid A$ and (5.2.13),

$$\mathrm{E}(Y \mid A=a, Z) = \mathrm{E}(\theta_{0,S}^y + \theta_{1,S}^y\, a + \theta_{2,S}^y\, Z \mid A=a, Z) = \boldsymbol{\theta}_0(a) + \boldsymbol{\theta}_1(a)\, a + \boldsymbol{\theta}_2(a)\, Z,$$

while part (a) gives $\mathrm{E}(Y \mid A=a, Z) = g_0(a) + g_2(a)Z$. These are two affine representations of the same regression $z \mapsto \mathrm{E}(Y \mid A=a, Z=z)$: their difference, an affine function of z, vanishes for Q_a-almost every z, and since $\mathrm{Var}(Z \mid A=a) > 0$ the law Q_a charges at least two points, so both its coefficients are zero. Whence $\boldsymbol{\theta}_2(a) = g_2(a)$ and $\boldsymbol{\theta}_2(a)/\theta_2^w = g_2(a)/\theta_2^w = \gamma(a)$ by part (a). Substituting into the ATE formula (5.2.12) of part (a) and using (5.2.11),

$$\begin{aligned}\mathrm{E}\, Y(a) &= \mathrm{E}\,\mathrm{E}(Y \mid A=a, Z) + \frac{\boldsymbol{\theta}_2(a)}{\theta_2^w}\,\mathrm{E}[\mathrm{E}(W \mid A, Z) - \mathrm{E}(W \mid A=a, Z)] \\ &= \boldsymbol{\theta}_0(a) + \boldsymbol{\theta}_1(a)\, a + \boldsymbol{\theta}_2(a)\,\mathrm{E}Z + \frac{\boldsymbol{\theta}_2(a)}{\theta_2^w}\,\theta_1^w(\mathrm{E}A - a),\end{aligned}$$

and since $\mathrm{E}W = \theta_0^w + \theta_1^w \mathrm{E}A + \theta_2^w \mathrm{E}Z$ by (5.2.11), the right-hand side equals $\boldsymbol{\theta}_0(a) + \boldsymbol{\theta}_1(a)a + \frac{\boldsymbol{\theta}_2(a)}{\theta_2^w}(\mathrm{E}W - \theta_0^w - \theta_1^w a)$, which is (5.2.14). □

After choosing a prior for $((\theta_{i,k}^y)_{k\in\mathbb{N}, i\in\{0,1,2\}}, (\theta_i^w)_{i\in\{0,1,2\}}, (w_k(A))_{k\in\mathbb{N}})$, any posterior draw of the parameters of the working model (5.2.7)-(5.2.9) carries a stratum representation of the fitted law of the observables: the mixing weights depend on the exposure only, so $S \perp\!\!\!\perp Z \mid A$, and the mixture components are linear, which is (5.2.13). With $\boldsymbol{\theta}_i(a) = \sum_k w_k(a)\, \theta_{i,k}^y$, the closed form (5.2.14) of Theorem 5.2.1(b) reads

$$\mathrm{E}\, Y(a) = \sum_{k\geq 1} w_k(a)\left(\theta_{0,k}^y + \theta_{1,k}^y a - \frac{\theta_{2,k}^y}{\theta_2^w}\left(\theta_0^w + \theta_1^w a\right)\right) + \sum_{k=1} \frac{\theta_{2,k}^y}{\theta_2^w} w_k(a)\mathrm{E}\,(W), \tag{5.2.17}$$

so the posterior on the parameters induces, through (5.2.17), a posterior on the ATE. The scope of this procedure is exactly the content of Theorem 5.2.1: the quantity estimated by (5.2.17) equals $\mathrm{E}\, Y(a)$ under every underlying model satisfying the assumptions of the theorem, whichever stratum representation the fit selects; in particular, whether or not the working model is correctly specified with respect to the U. Moreover, since by part (a) the ATE is a continuous functional of the regressions of the observables, consistent estimation of these regressions translates into consistent estimation of the ATE; this is made precise in Section 5.3.

Remark 5.2.2 (Linearity in U cannot be dropped). One may wonder whether it is possible to generalize assumption 1 of the underlying model class to an expression nonlinear in U. We show it is not possible. Consider an underlying model in which, for some f_0:

$$\begin{aligned}\mathrm{E}(Y \mid A, U) &= f_0(A, U)\\ \mathrm{E}(W \mid A, U) &= \beta_0^w + \beta_1^w U\\ \mathrm{E}(U \mid A, Z) &= \beta_0^u + \beta_1^u A + \beta_2^u Z,\end{aligned}$$

therefore we can find θ_i^w such that $\mathrm{E}(W \mid A, Z) = \theta_0^w + \theta_1^w A + \theta_2^w Z$. Then $\mathrm{E}\, Y(a) = \mathrm{E} f_0(a, U)$ and define $z_a := \frac{1}{\theta_2^w}(\theta_1^w(\mathrm{E}A - a) + \theta_2^w \mathrm{E}Z)$. Then:

$$\begin{aligned}&\mathrm{E}(Y \mid A = a, Z) = \mathrm{E}(f_0(a, U) \mid A = a, Z)\\ &\mathrm{E}(W \mid A = a, Z = z_a) = \mathrm{E}W\end{aligned}$$

on the other hand:

$$\mathrm{E}(W \mid A, Z) = \beta_0^w + \beta_1^w \mathrm{E}(U \mid A, Z) \implies \mathrm{E}W = \beta_0^w + \beta_1^w \mathrm{E}(U)$$

therefore:

$$\beta_0^w + \beta_1^w \mathrm{E}(U) = \mathrm{E}W = \mathrm{E}(W \mid A = a, Z = z_a) = \beta_0^w + \beta_1^w \mathrm{E}(U \mid A = a, Z = z_a)$$

which implies: $\mathrm{E}(U) = \mathrm{E}(U \mid A = a, Z = z_a)$. Thus in this case the following formula is true only if we can exchange f_0 with the expectation, i.e. f_0 is linear in U ($f_0(a, U) = f_0(a) + f_2(a)U$):

$$\begin{aligned}&\mathrm{E}(Y \mid A = a, Z = z_a) = \mathrm{E}(f_0(a, U) \mid A = a, Z = z_a)\\ &= f_0(a) + f_2(a)\mathrm{E}(U \mid A = a, Z = z_a) = f_0(a) + f_2(a)\mathrm{E}(U) = \mathrm{E} f_0(a, U) = \mathrm{E}\, Y(a)\end{aligned}$$

Extension 2 of the linear model.

The underlying model. In the second extension we enlarge the class of underlying models by allowing the proxy Z to affect the outcome directly. The underlying model now follows the DAG of Figure 5.3b, which augments the classical proximal DAG with the edge $Z \to Y$, and the outcome regression may depend on Z and interact it with the confounder,

$$\mathrm{E}(Y \mid A, U, Z) = f_0(A, Z) + \{\alpha_0(A) + \alpha_1(A)Z\}\, U,$$

while the regressions of W and U keep the same form as before. The DAG encodes the consistency assumption and the conditional independence read from its SWIG,

1. $Y(a) = Y$ if $A = a$
2. $Y(a) \perp\!\!\!\perp A \mid U, Z$

Although the DAG of Figure 5.3b is not identical to the canonical proximal DAG, it remains proximal in the sense relevant for identification, since proximal assumptions are structural constraints on proxy roles rather than exact graphical equalities: W is still a valid negative-control outcome proxy (it is not causally affected by A, and $W \perp\!\!\!\perp (A, Z) \mid U$), Z is still a valid pre-exposure proxy (it does not causally affect Y), and both proxies remain informative about the latent confounder U, which is the key requirement enabling proximal identification.

The working model. As in Extension 1, an outcome regression that interacts A, Z and U through the arbitrary functions f_0, α_0, α_1 cannot be fitted directly. We extend the Extension 1 construction by letting the stratum label also depend on the proxy Z: a label S, drawn given (A, Z), indexes a family of classical models for the outcome as in (5.2.6),

$$\mathrm{E}(Y \mid A, U, S, Z) = \beta^y_{0,S} + \beta^y_{1,S} A + \beta^y_{2,S} U,$$

while W and U keep the unstratified regressions (5.2.2)–(5.2.3). The DAG of Figure 5.3a displays this construction: it extends that of Extension 1 by adding the edge $Z \to S$, so that marginalising out S returns the DAG of Figure 5.3b on which the underlying model is based (the edge $Z \to Y$ is inherited from the path $Z \to S \to Y$). Unlike in Extension 1, this marginalisation does not automatically land on the underlying form $f_0(A, Z) + \{\alpha_0(A) + \alpha_1(A)Z\}U$: it requires the constraint on the betas made precise in part (b) of the theorem below.



(a) DAG with latent node S. (b) Marginal DAG after integrating out S.

Figure 5.3: Extension 2. a: the DAG that motivates the shape of the working model, with the latent stratum variable S now influenced by Z as well; b: its marginalization over S, which augments the classical proximal DAG of Figure 5.1 with the edge $Z \to Y$ and on which the underlying model is based. The DAG a is motivation only and is nowhere assumed.

The same tower-property computation as in Extension 1 turns (5.2.6) into a stratum-wise linear regression of the observables, $\mathrm{E}(Y \mid A, S, Z) = \theta^y_{0,S} + \theta^y_{1,S} A + \theta^y_{2,S} Z$; averaging over $S \mid A, Z$, the observables therefore follow the mixture

$$S \mid A, Z \sim (w_k(A, Z))_{k \in \mathbb{N}} \tag{5.2.18}$$

$$Y \mid A, Z \sim \sum_{k=1}^{\infty} w_k(A, Z)\, N(\theta^y_{0,k} + \theta^y_{1,k} A + \theta^y_{2,k} Z, \sigma^2_{y,k}) \tag{5.2.19}$$

$$W \mid A, Z \sim N(\theta^w_0 + \theta^w_1 A + \theta^w_2 Z, \sigma^2_w) \tag{5.2.20}$$

with $(\theta^y_{i,k})_{i \in \{0,1,2\}, k \in \mathbb{N}}$, $(\theta^w_i)_{i \in \{0,1,2\}}$ and $(w_k(A, Z))_{k \in \mathbb{N}}$ now treated as free parameters, rather than fixed at the values determined by the betas above, that we fit to a sample (Z, A, Y, W) via the Bayesian nonparametric procedure of Section 5.3.

The following theorem is the analogue of Theorem 5.2.1: part (a) identifies the ATE of the underlying model by an explicit functional of the laws of the observables, and part (b) turns this functional into the closed form computed by the fitted working model. The contrast with Extension 1 lies in the stratum representation required in part (b): a stratum structure on the

observables alone no longer suffices, and the representation must now involve the unobserved confounder U; unlike in Extension 1, the working model does not automatically supply such a representation, since the mixture (5.2.18)–(5.2.20) is fitted to the observables alone and says nothing, by itself, about U. Its existence is therefore a genuine compatibility condition between the underlying model and the stratum structure of the working model.

Theorem 5.2.2 (Extension 2)**.** *Suppose* (U, Z, A, Y, W) *follows the underlying model: the DAG of Figure 5.3b (hence the consistency 1 and the conditional independence 2) and*

1. *For some measurable* f_0 *and functions* α_0, α_1*,*

$$\mathrm{E}(Y \mid A, U, Z) = f_0(A, Z) + \{\alpha_0(A) + \alpha_1(A)Z\}U;$$

2. $\mathrm{E}(W \mid A, U, Z) = \beta_0^w + \beta_1^w U$ *with* $\beta_1^w \neq 0$*;*

3. $\mathrm{E}(U \mid A, Z) = \beta_1^u A + \beta_2^u Z$ *with* $\beta_2^u \neq 0$*.*

Fix $a \in \mathcal{A}$*, let* Q_a *denote the conditional law of* $Z \mid A = a$*, and assume moreover:*

(i) (Orthogonality) $\mathrm{E}[f_0(a, Z)Z^j \mid A = a] = 0$ *for* $j = 0, 1, 2$*;*

(ii) (Invertibility) $G_a = \mathrm{E}[\phi(Z)\phi(Z)^\top \mid A = a]$ *with* $\phi(Z) = (1, Z, Z^2)^\top$ *is invertible.*

Then:

(a) *(Identification)* $\mathrm{E}(W \mid A, Z) = \theta_0^w + \theta_1^w A + \theta_2^w Z$ *with* $\theta_2^w = \beta_1^w \beta_2^u \neq 0$*, and the ATE admits the representation*

$$\mathrm{E}\,Y(a) = \mathrm{E}[\mathrm{E}(Y \mid A = a, Z)] + \mathrm{E}\big[\gamma(a, Z)\,\{\mathrm{E}(W \mid A, Z) - \mathrm{E}(W \mid A = a, Z)\}\big], \quad (5.2.21)$$

where $\gamma(a, z)$ *is identified from the laws of the observables:*

$$\gamma(a, z) = \frac{1}{\theta_2^w}\Big\{b_1(a) + b_2(a)\Big(z - a\frac{\theta_1^w}{\theta_2^w}\Big)\Big\},$$

with $(b_0(a), b_1(a), b_2(a))$ *the* $L_2(Q_a)$*-projection coefficients of* $\mathrm{E}(Y \mid A, Z)$ *onto the* $\mathrm{span}\{1, z, z^2\}$*.*

(b) *(Closed form under a stratum representation) Let* (U, Z, A, Y, W, S) *be any stratum representation of the* full *vector, including the unobserved confounder: a random vector, on some probability space, whose first five coordinates are distributed as* (U, Z, A, Y, W) *and such that*

$$S \perp\!\!\!\perp U \mid A, Z \qquad \textit{and} \qquad \mathrm{E}(Y \mid A, U, S, Z) = \beta_{0,S}^y + \beta_{1,S}^y A + \beta_{2,S}^y U, \quad (5.2.22)$$

for some coefficients $(\beta_{i,s}^y)_{i\in\{0,1,2\}, s\in\mathbb{N}}$*, and suppose* $\mathrm{Var}(U \mid A = a, Z) > 0$ Q_a*-a.s., i.e. the confounder is not a deterministic function of the observed covariates. Then there exist coefficients* $(\theta_{i,s}^y)_{i\in\{0,1,2\}, s\in\mathbb{N}}$ *such that*

$$\mathrm{E}(Y \mid A, S, Z) = \theta_{0,S}^y + \theta_{1,S}^y A + \theta_{2,S}^y Z, \quad (5.2.23)$$

and, for $\boldsymbol{\theta}_i(a,Z) := \mathrm{E}(\theta^y_{i,S} \mid A=a,Z)$, $i=0,1,2$,

$$\frac{\boldsymbol{\theta}_2(a,Z)}{\theta^w_2} = \gamma(a,Z) \quad Q_a\text{-a.s.},$$

so the ATE takes the closed form

$$\begin{aligned}\mathrm{E}\,Y(a) &= \mathrm{E}\,\mathrm{E}(Y \mid A=a,Z) + \mathrm{E}\left[\frac{\boldsymbol{\theta}_2(a,Z)}{\theta^w_2}(\mathrm{E}(W \mid A,Z) - \mathrm{E}(W \mid A=a,Z))\right] \\ &= \mathrm{E}\left[\boldsymbol{\theta}_0(a,Z) + \boldsymbol{\theta}_1(a,Z)\,a - \frac{\boldsymbol{\theta}_2(a,Z)}{\theta^w_2}(\theta^w_0 + \theta^w_1 a)\right] + \mathrm{E}\left[\frac{\boldsymbol{\theta}_2(a,Z)}{\theta^w_2}\,W\right]. \quad (5.2.24)\end{aligned}$$

The proof of Theorem 5.2.2 is provided in the Appendix.

Remark 5.2.3. By the tower property, condition (5.2.22) combined with $S \perp\!\!\!\perp U \mid A,Z$ implies $\mathrm{E}[\beta^y_{2,S} \mid A,Z] = \alpha_0(A) + \alpha_1(A)Z$. This represents a genuine constraint for the augmented joint of (U,Z,A,Y,W,S).

After choosing a prior for $((\theta^y_{i,k})_{k\in\mathbb{N},i\in\{0,1,2\}}, (\theta^w_i)_{i\in\{0,1,2\}}, (w_k(A,Z))_{k\in\mathbb{N}})$, any posterior draw of the parameters of the working model (5.2.18)-(5.2.20) defines a stratum structure on the fitted law of the observables, with $\boldsymbol{\theta}_i(a,Z) = \sum_k w_k(a,Z)\,\theta^y_{i,k}$, and the closed form (5.2.24) of Theorem 5.2.2(b) reads

$$\mathrm{E}\,Y(a) = \sum_{k\geq 1} \mathrm{E}\left[w_k(a,Z)\left(\theta^y_{0,k} + \theta^y_{1,k}a - \frac{\theta^y_{2,k}}{\theta^w_2}(\theta^w_0 + \theta^w_1 a)\right)\right] + \sum_{k\geq 1}\frac{\theta^y_{2,k}}{\theta^w_2}\mathrm{E}[w_k(a,Z)\,W], \quad (5.2.25)$$

so the posterior on the parameters induces, through (5.2.25), a posterior on the ATE. The scope of this procedure is again given by the theorem, with one difference from Extension 1: the fitted mixture provides strata for the observables only, whereas part (b) requires a stratum representation involving the unobserved confounder, as in (5.2.22). The quantity estimated by (5.2.25) therefore equals $\mathrm{E}\,Y(a)$ under every underlying model for which the fitted stratum structure extends to a representation compatible with the confounder. By part (a), consistent estimation of the regressions of the observables still translates into consistent estimation of the ATE. Corollary 5.3.1 in Section 5.3 makes this precise for the Bayesian procedure that we propose.

The representation (5.2.21) also clarifies the role of the proximal correction in this extended setting: $\mathrm{E}\,Y(a)$ separates into a baseline outcome regression term, $\mathrm{E}[\mathrm{E}(Y \mid A=a,Z)]$, and a proxy-based bias-correction term driven by $\mathrm{E}(W \mid A,Z) - \mathrm{E}(W \mid A=a,Z)$. Hence, the departure from standard outcome-regression identification is entirely encoded in the second term, whose weight $\gamma(a,Z)$ is itself identified from observable regressions. This decomposition is useful both conceptually and practically. Conceptually, it shows that allowing the edge $Z \to Y$ does not invalidate proximal identification; rather, it changes the form of the bridge through $\gamma(a,Z)$ while preserving identification from observables. Practically, the theorem gives transparent sufficient conditions for consistency: accurate recovery of $\mathrm{E}[Y \mid A=a,Z=z]$ and $\mathrm{E}[W \mid A=a,Z=z]$, together with stable estimation of (θ^w_1,θ^w_2) and a non-degenerate denominator $|\theta^w_2| > 0$. In particular, weak signal in θ^w_2 is the main source of instability, so inference should be accompanied by diagnostics on the strength of the Z-coefficient in the W-model.

Remark 5.2.4 (Possible extension). A natural extension of Proposition 5.2.2 is to replace the linear interaction in Z, $\{\alpha_0(A) + \alpha_1(A)Z\}U$, by a richer dictionary interaction $\{\alpha_0(A) + \sum_{k=1}^{K} \alpha_k(A)\varphi_k(Z)\}U$.

Under the same proximal assumptions on W and U, together with the analogous orthogonality and invertibility conditions for the enlarged basis

$$\psi(Z) = (1, \varphi_1(Z), \ldots, \varphi_K(Z), Z, Z\varphi_1(Z), \ldots, Z\varphi_K(Z)),$$

one expects the same bridge form

$$\mathrm{E}\, Y(a) = \mathrm{E}[\mathrm{E}(Y \mid A = a, Z)] + \mathrm{E}\big[\gamma(a, Z)\{\mathrm{E}(W \mid A, Z) - \mathrm{E}(W \mid A = a, Z)\}\big],$$

with $\gamma(a, Z)$ identified through $L_2(Q_a)$-projection coefficients on $\mathrm{span}\{\psi\}$. In that extension, writing

$$b(a) = G_a^{-1}\mathrm{E}[\mathrm{E}(Y \mid A = a, Z)\psi(Z) \mid A = a],$$

with components

$$b(a) = (b_0(a),\, b_{\varphi_1}(a), \ldots, b_{\varphi_K}(a),\, b_Z(a),\, b_{Z\varphi_1}(a), \ldots, b_{Z\varphi_K}(a)),$$

the corresponding bridge coefficient is

$$\gamma(a, z) = \frac{1}{\theta_2^w}\left\{b_Z(a) + \sum_{k=1}^{K} b_{Z\varphi_k}(a)\varphi_k(z) - a\frac{\theta_1^w}{\theta_2^w}\sum_{k=1}^{K} b_{\varphi_k}(a)\varphi_k(z)\right\}.$$

5.3 Contraction

In this section, we present the key theorems that provide sufficient conditions for posterior consistency of the ATE within a class of underlying causal models. We state the results for the second extension of the linear model (DAG 5.3a). The analogous results for the first extension (DAG 5.2a) do not follow automatically: in that simpler model the mixture weights $w_k(A)$ depend only on A, so the prior has strictly smaller support than the covariate-dependent prior used here. The KL property and the sieve entropy bound (Theorems 5.3.1 and 5.3.2) were established for the covariate-dependent weights (5.3.5); extending them to A-only weights requires verifying that the true density $f_0(\cdot \mid a, z)$ lies in the support of the restricted prior, which holds when f_0 itself depends on (a, z) only through a. Under that restriction the proofs carry over verbatim, and we leave the straightforward adaptation to the reader.

5.3.1 *Priors*

For the law $\mathcal{L}(W \mid A, Z)$ we fit a parametric model normal as follows:

$$W \mid A, Z \sim N(\theta_0^w + \theta_1^w A + \theta_2^w Z, \sigma_w^2) \tag{5.3.1}$$

The prior on the weights $\boldsymbol{\theta}^w := (\theta_0^w, \theta_1^w, \theta_2^w)$ is a normal with mean μ_{θ^w} and diagonal covariance matrix Σ_w. We put an Inverse Gamma prior on the standard deviation $\sigma_w \sim \mathrm{InvGamma}(\gamma_1, \gamma_2)$.

On the other hand, for the law of the observables $\mathcal{L}(Y \mid A, Z)$ we fit a nonparametric predictor dependent countable mixture of Gaussian linear regressions:

$$Y \mid A, Z \sim \sum_{k=1}^{\infty} w_k^n(A, Z)\, N(\theta_{0,k}^y + \theta_{1,k}^y A + \theta_{2,k}^y Z, \sigma_{y,k}^2) \tag{5.3.2}$$

The nonparametric aspect of this prior consists in the construction of the weights $(a, z) \mapsto w_k^n(a, z)$ for all $k, n \in \mathbb{N}$. For $k \in \mathbb{N}$ consider three independent discrete-time Gaussian sequences $\{\eta_{i,m}^k\}_{m=1}^{\infty}$, $i = 0, 1, 2$, supported on $\mathbb{N}$ such that, for integer $l \geq 0$:

$$\mathrm{E}\eta_{i,m}^k = \mu_{i,m}^k \quad \text{and} \quad \mathrm{Cov}(\eta_{i,m}^k, \eta_{i,m+l}^k) = {}_k\sigma_{i,m}^2 \delta_m(m+l), \qquad i = 0, 1, 2, \ \forall\, k \in \mathbb{N}. \tag{5.3.3}$$

Thus $\forall\, m, k \in \mathbb{N}$, $\boldsymbol{\mu}_m^k = (\mu_{0,m}^k, \mu_{1,m}^k, \mu_{2,m}^k)$ and $\Sigma_m^k = \mathrm{diag}({}_k\sigma_{0,m}^2, {}_k\sigma_{1,m}^2, {}_k\sigma_{2,m}^2)$, $\boldsymbol{\eta}_m^k \sim N(\boldsymbol{\mu}_m^k, \Sigma_m^k)$ and $\boldsymbol{\eta}_m^k \perp\!\!\!\perp \boldsymbol{\eta}_l^k$ for $l \neq m$.

Now for realizations $A = a, Z = z$, define $\boldsymbol{x} = (a, z)$. Then for integers $N_n \to \infty$ as $n \to \infty$, define, for every fixed $n \in \mathbb{N}$ partitions $\cup_{m=1}^{N_n} T_{m,n}$ of the domain $\mathcal{X} = \mathcal{A} \times \mathcal{Z}$ of the covariate space:

$$\alpha_k^n(\boldsymbol{x}) = \sum_{m=1}^{N_n} (\eta_{0,m}^k + \eta_{1,m}^k a + \eta_{2,m}^k z) \mathbf{1}_{(a,z) \in T_{m,n}} \tag{5.3.4}$$

$$w_k^n(\boldsymbol{x}) = \Phi(\alpha_k^n(\boldsymbol{x})) \prod_{l<k} \left\{1 - \Phi(\alpha_l^n(\boldsymbol{x}))\right\} \tag{5.3.5}$$

where $\Phi(\cdot)$ denotes the cumulative distribution function for the standard normal distribution. We thus conclude that the prior is given by the dependent probit-stick breaking mixtures of normals, for $\bar{\boldsymbol{x}} = (1, a, z)$ and for a base measure G_0:

$$G_{\boldsymbol{x}}^n = \sum_{k=1}^{\infty} w_k^n(\boldsymbol{x}) \delta_{(\boldsymbol{\theta}^y, \sigma_y)}, \qquad (\boldsymbol{\theta}^y, \sigma_y) \sim G_0 \tag{5.3.6}$$

$$f(y \mid \boldsymbol{x}) = \int \frac{1}{\sigma_y} \phi\left(\frac{y - \bar{\boldsymbol{x}}' \boldsymbol{\theta}^y}{\sigma_y}\right) dG_{\boldsymbol{x}}^n(\boldsymbol{\theta}^y, \sigma_y). \tag{5.3.7}$$

Note that $w_k^n(\boldsymbol{x}) \geq 0$ are random functions of $\boldsymbol{x}$ such that $\sum_{k=1}^{\infty} w_k^n(\boldsymbol{x}) = 1$ a.s. for each fixed $\boldsymbol{x} \in \mathcal{X}$ and $G_0 = G_{0,\boldsymbol{\theta}^y} \times G_{0,\sigma_y}$ is a probability distribution on $\mathbb{R}^3 \times \mathbb{R}^+$where $G_{0,\boldsymbol{\theta}^y}$ and G_{0,σ_y} are probability distributions on $\mathbb{R}^3$ and $\mathbb{R}^+$ respectively. We additionally assume that the prior satisfies the following two properties, where we denote the supremum norm on the covariate space by $\|\cdot\|$ and the variance by $\mathbb{V}$: $0 < \min_{1 \leq k \leq m_n} \|\mathbb{V}(\alpha_k^n)\| \leq \max_{1 \leq k \leq m_n} \|\mathbb{V}(\alpha_k^n)\| < \infty, \quad \forall\, n \in \mathbb{N}$.

5.3.2 *Results*

Assume the Y_i is observed independently given the covariates (A_i, Z_i), $i = 1, 2, \ldots$ which are drawn independently from a probability distribution Q on $\mathcal{X}$ which admits density q with respect to the Lebesgue measure and which is known. Similarly the W_i is observed independently given the covariates (A_i, Z_i), $i = 1, 2, \ldots$. Denote the observed data of size n by $\mathcal{D}^{(n)} = (Y^{(n)}, W^{(n)}, A^{(n)}, Z^{(n)})$, where $x^{(n)} = (x_1, \ldots, x_n)$. We denote by p_0 the joint density of Y, W, A, Z relative to the probability measure P_0. Given the prior Π we denote a version of the posterior by $\Pi_n\left(\cdot \mid \mathcal{D}^{(n)}\right)$.

We assume q is known, but it could be estimated separately from the data. Note that our model in Section 5.2 does not impose any restriction on q. There are several well known nonparametric estimation methods in the literature that under mild restrictions give consistency in density estimation (e.g. Dirichlet Process Mixtures or exponential transforms of Gaussian Process, see [42]).

Now we present the main result of this section, which confirms consistency of the ATE when using the above constructed priors. Most of the proofs of this section are based on existing work done in [81] and [80].

Corollary 5.3.1. *Consider any underlying causal model that satisfies the assumptions of Proposition 5.2.2 and assume the joint law Q of Z, A as well as the conditional law Q_a of $Z \mid A = a$ with density $z \mapsto q(z \mid a)$ for any fixed $a \in \mathcal{A}$ are known. Consider the prior defined in* (5.3.1)-(5.3.7)*. Suppose furthermore that the assumptions of Theorem 5.3.2 and Proposition 5.3.1 are satisfied, and that:*

A. There exists a constant $C < \infty$ such that

$$\Pi_n\left(\sup_{(a,z)\in\mathcal{X}} \int y^2 f(y \mid a, z)\, dy \leq C \,\middle|\, \mathcal{D}^{(n)}\right) \to 1, \quad P_0^{(n)}\text{-a.s.}$$

Then for every fixed $a \in \mathcal{A}$:

$$\Pi_n(|\mathrm{E}_0 Y(a) - \mathrm{E}\, Y(a)| > \varepsilon \mid \mathcal{D}^{(n)}) \to 0, \qquad P_0^{(n)} - a.s. \tag{5.3.8}$$

Remark 5.3.1. Assumption A is a mild posterior regularity condition. It is implied, for instance, if the sieve $\mathcal{F}_n^a$ in the proof of Theorem 5.3.2 is further restricted to have component parameters $\|\boldsymbol{\theta}_k^y\| \leq R$ and $\sigma_{y,k} \leq R$ for a fixed constant $R < \infty$ independent of n, since in that case $\int y^2 f(y \mid a, z)dy \leq R^2(1 + |a| + |z|)^2 + R^2 \leq 4R^2(1 + C_{\mathcal{X}})^2$ uniformly over $(a, z) \in \mathcal{X}$ and over the truncated sieve. Moreover, under the underlying model the true second moment $\int y^2 f_0(y \mid a, z)dy = \mathrm{Var}(Y \mid A = a, Z = z) + m_Y^0(a, z)^2$ is bounded uniformly over compact $\mathcal{X}$ by the assumptions of Proposition 5.2.2, so the true density automatically satisfies the same bound.

The proof of this corollary is a consequence of Theorem 5.3.2, Proposition 5.3.1 (provided below) and the triangle inequality and it is provided in Appendix B. We now outline the main results involved in the proofs that lead to the conclusion of Corollary 5.3.1.

Because we are concerned with consistency of the distributions of $Y \mid A, Z \sim f_0(\cdot \mid A, Z)$ and of $W \mid A, Z \sim g_0(\cdot \mid A, Z)$, we have that the marginals of interest are:

$$h_0^y(y, a, z) = f_0(y \mid a, z)q(a, z), \qquad h_0^w(w, a, z) = g_0(w \mid a, z)q(a, z).$$

Recall $\boldsymbol{x} = (a, z)$. For $f_0 \in \mathcal{F}_d$ such that $h_0^y(\boldsymbol{x}, y) = q(\boldsymbol{x})f_0(y \mid \boldsymbol{x})$ is the true joint data-generating density, where $\mathcal{F}_d$ is the space of conditional densities which are continuous in its arguments (see Def. 5.5.1), we define an ε-sized KL neighborhood around f_0. For $\mathrm{KL}\left(h_0^y, h^y\right) = \int h_0^y \log\left(h_0^y / h^y\right)$:

$$K_\varepsilon(f_0) = \left\{f : f \in \mathcal{F}_d, \mathrm{KL}\left(h_0^y, h^y\right) < \varepsilon, h^y(\boldsymbol{x}, y) = q(\boldsymbol{x})f(y \mid \boldsymbol{x}),\ \forall\, y \in \mathcal{Y}, \boldsymbol{x} \in \mathcal{X}\right\} \tag{5.3.9}$$

Then, Π_n is said to have KL property at $f_0 \in \mathcal{F}_d$, denoted $f_0 \in \mathrm{KL}(\Pi_n)$, if, $\forall\, \varepsilon > 0$:

$$\liminf_{n\to\infty} \Pi_n\{K_\varepsilon(f_0)\} > 0.$$

Remark 5.3.2. Note that $f_0 \in \mathrm{KL}(\Pi_n)$ implies $\forall\, \varepsilon > 0$, as $n \to \infty$:

$$\Pi_n\left(f \,:\, d_w(f, f_0) > \varepsilon \mid \mathcal{D}^{(n)}\right) \to 0, \qquad \text{in } P_0^{(n)} - \text{probability.} \tag{5.3.10}$$

d_w is the metric relative to the weak neighborhoods, whose sub-base is defined in (5.5.3).

Theorem 5.3.1. *Consider the prior defined in* (5.3.2)-(5.3.7)*. Assume $\mathcal{X}$ is a compact subset of $\mathbb{R}^3$. If:*

A0. the base measure G_0 is absolutely continuous with respect to $\lambda\left(\mathbb{R}^3 \times \mathbb{R}^+\right)$.

A1. f_0 is nowhere zero and bounded by $M < \infty$.

A2. $\left|\int_{\mathcal{X}} \int_{\mathcal{Y}} f_0(y \mid \boldsymbol{x}) \log f_0(y \mid \boldsymbol{x})\, dy\, q(\boldsymbol{x})\, d\boldsymbol{x}\right| < \infty$.

A3. $\left|\int_{\mathcal{X}} \int_{\mathcal{Y}} f_0(y \mid \boldsymbol{x}) \log \frac{f_0(y|\boldsymbol{x})}{\psi_{\boldsymbol{x}}(y)}\, dy\, q(\boldsymbol{x})\, d\boldsymbol{x}\right| < \infty$, *where* $\psi_{\boldsymbol{x}}(y) = \inf_{t\in[y-1,y+1]} f_0(t \mid \boldsymbol{x})$.

A4. $\exists\, \eta > 0$ *such that* $\int_{\mathcal{X}} \int_{\mathcal{Y}} |y|^{2(1+\eta)} f_0(y \mid \boldsymbol{x})\, dy\, q(\boldsymbol{x})\, d\boldsymbol{x} < \infty$.

A5. $(\boldsymbol{x}, y) \mapsto f_0(y \mid \boldsymbol{x})$ is jointly continuous.

Then $f_0 \in \mathrm{KL}(\Pi_n)$.

The proof of Theorem 5.3.1 is given in Appendix B. Because we adopted a parametric model for the law $\mathcal{L}(W \mid A, Z)$ consistency of the functional $\mathrm{E}[W \mid Z]$ and $\mathrm{E}[W \mid A = a, Z]$ in $\mathbb{L}_1$ is easily granted. The proof of this proposition, provided in Appendix B, is a straightforward application of the standard theory of consistency for parametric models.

Proposition 5.3.1. *Under the assumption that the underlying law of $W \mid A, Z$ is as in Proposition 5.2.2, that the density $(a, z) \mapsto q(z \mid a)$ is bounded, that the covariate space $\mathcal{W}$ is a bounded subset of $\mathbb{R}$, and using the prior* (5.3.1)*, for fixed $a \in \mathcal{A}$ and $\forall\, \varepsilon > 0$, as $n \to \infty$:*

$$\Pi(\,\mathrm{E}_0|\mathrm{E}_0(W \mid A = a, Z) - \mathrm{E}(W \mid A = a, Z)| > \varepsilon \mid \mathcal{D}^{(n)}) \to 0, \qquad P_0^{(n)} - a.s.$$
$$\Pi(\,\mathrm{E}_0|\mathrm{E}_0(W \mid A, Z) - \mathrm{E}(W \mid A, Z)| > \varepsilon \mid \mathcal{D}^{(n)}) \to 0, \qquad P_0^{(n)} - a.s.$$

where $\mathrm{E}_0|\mathrm{E}_0(W \mid A, Z) - \mathrm{E}(W \mid A, Z)| = \iint |\int w(g_0(w \mid a, z) - g(w \mid a, z))\, dw|\, q(a, z)\, da\, dz$ *and* $\mathrm{E}_0|\mathrm{E}_0(W \mid A = a, Z) - \mathrm{E}(W \mid A = a, Z)| = \iint |\int w(g_0(w \mid a, z) - g(w \mid a, z))\, dw|\, q(z \mid a)\, dz$.

We now prove consistency in $\mathbb{L}_1$ integrated with respect to the density $q(z \mid a)$ (see the metric defined by (5.5.5) in Appendix A). For fixed $a \in \mathcal{A}$ let $(a, z) \mapsto q(z \mid a)$ be the density of the underlying probability measure associated to the law of $Z \mid A$. Then we denote the joints:

$$f_0^a(y, z \mid a) = f_0(y \mid a, z)\, q(z \mid a) \tag{5.3.11}$$

Note that the metric of interest (c.f. (5.5.5)) can be conveniently rewritten as follows using Fubini's theorem and the nonnegativity of $z \mapsto q(z \mid a)$ for fixed $a \in \mathcal{A}$:

$$\begin{aligned}\int \|f(\cdot \mid a, z) - f_0(\cdot \mid a, z)\|_1\, q(z \mid a)\, dz &= \int \int |f(y \mid a, z) - f_0(y \mid a, z)|\, q(z \mid a)\, dz\, dy \\ &= \|f_0^a - f^a\|_1.\end{aligned}$$

Theorem 5.3.2. *Fix $a \in \mathcal{A}$. Consider the prior defined in* (5.3.2)-(5.3.7)*. Let the assumptions of Theorem 5.3.1 hold true. Assume $\mathcal{X}$ is a compact subset of $\mathbb{R}^3$ and that, for fixed $a \in \mathcal{A}$, the assumptions of Theorem 5.3.1 are satisfied with respect to the density $z \mapsto q(z \mid a)$. Moreover if:*

B0. the base measure G_0 *satisfies* $G_0\left(([-a_n, a_n]^3)^c \times [l_n, h_n]^c\right) \leq m_n^{-1} e^{-m_n \log m_n}$, *where* $a_n = \sqrt{n}$, $h_n = \sqrt{n}$, $l_n = 1/\sqrt{n}$, $m_n = n/\log n$.

B1. for any fixed $a \in \mathcal{A}$, $(z, y) \mapsto f_0(y \mid a, z)$ *is jointly continuous.*

Then $\forall\, \varepsilon > 0$, *as* $n \to \infty$ *:*

$$\Pi_n\left(f \,:\, \|f^a - f_0^a\|_1 > \varepsilon \mid \mathcal{D}^{(n)}\right) \to 0, \qquad P_0^{(n)} - a.s.$$

The proof of Theorem 5.3.2 is provided in Appendix B. This result is the last fundamental piece that leads to consistency of the ATE, since it gives consistency in $\mathbb{L}_1$ metric for the density of $Y \mid A, Z$ when modelled with the Bayesian nonparametric procedure we proposed.

5.4 Conclusion

In this chapter, we introduced a more versatile approach to causal inference when unmeasured confounding is present. By building upon the classical linear model in the setting of proximal inference and transitioning to mixtures of linear models, we created a more flexible framework for modeling the distribution of outcomes given the covariates. We proved that in this framework the average treatment effect remains identifiable and, by deriving it, we obtained a closed-form expression for the bridge function. This extension improves on the limitations of simpler linear models, which often fail to capture the intricacies of real-world data.

Additionally, we presented a Bayesian nonparametric estimation method that employs a probit stick-breaking process with mixing weights that depend on the covariates. Our findings indicate that this method yields consistency across various causal models. One significant benefit of our approach is its guarantee of the bridge function's existence, which simplifies the estimation process within this framework. Although our method enhances flexibility, it still relies on assumptions regarding the simpler linear model for the distribution of W given the covariates. Future studies could explore relaxing these assumptions to further extend the model's applicability. Nevertheless, our work represents a substantial advancement in proximal inference, offering a practical solution for estimating the average causal effect in presence of unmeasured confounding.

5.5 Appendix

5.5.1 Definitions of function spaces and topologies used in proofs

First we define all the relevant function spaces for our consistency study.

Definition 5.5.1. Let $\mathcal{X} = \mathcal{A} \times \mathcal{Z}$ be the covariate space which is a compact subset of $\mathbb{R}^2$ and $\mathcal{Y} = \mathbb{R}$ be the response space. Let:

$$\mathcal{F}_d = \Big\{ f : \mathcal{X} \times \mathcal{Y} \to (0, \infty),\ \int f(y \mid \boldsymbol{x})\, dy = 1,\ \forall\, \boldsymbol{x} \in \mathcal{X} \text{ and the map from } \boldsymbol{x} \mapsto \mathbb{L}_1(\lambda, \mathcal{Y}),\ \boldsymbol{x} \mapsto f(\cdot \mid \boldsymbol{x}) \text{ is continuous} \Big\}.$$

Similarly, we define the space $\mathcal{G}_d$ for the conditional densities $g : \mathcal{X} \times \mathcal{W} \to (0, \infty)$.

Definition 5.5.2. Define collection of conditional probability measures on $(\mathbb{R}^3\times\mathbb{R}^+, \mathcal{B}(\mathbb{R}^3\times\mathbb{R}^+))$ denoted by $\mathscr{G}_x^*$ such that $\forall\, B \in \mathcal{B}(\mathbb{R}^3 \times \mathbb{R}^+)$,

- $\boldsymbol{x} \mapsto F_{\boldsymbol{x}}(B)$ is continuous,
- for any sequence of sets $A_n \subset \mathbb{R}^3 \times \mathbb{R}^+ \downarrow \emptyset$, $\sup_{\boldsymbol{x}\in\mathcal{X}} F_{\boldsymbol{x}}(A_n) \downarrow 0$.

We now define all the relevant neighborhoods for our consistency study. For the mixing measures $G_{\boldsymbol{x}} \in \mathscr{G}_x^*$ we consider two types of neighborhoods.

Definition 5.5.3. For a bounded continuous function $g : \mathbb{R}^3 \times \mathbb{R}^+ \times \mathcal{X} \to \mathbb{R}$ and $\varepsilon > 0$, a sub-base of the weak neighborhood of a conditional probability measure $\{F_{\boldsymbol{x}}, \boldsymbol{x} \in \mathcal{X}\}$ is defined as

$$S_\varepsilon^w(F_{\boldsymbol{x}}, q) := \left\{\{G_{\boldsymbol{x}} : \boldsymbol{x} \in X\} : \left|\int_{\mathcal{X}\times\mathbb{R}^3\times\mathbb{R}^+} g(\boldsymbol{\theta}, \sigma, \boldsymbol{x}) d\left(G_{\boldsymbol{x}}(\boldsymbol{\theta}, \sigma) - F_{\boldsymbol{x}}(\boldsymbol{\theta}, \sigma)\right) q(\boldsymbol{x})\, d\boldsymbol{x}\right| < \varepsilon\right\}. \tag{5.5.1}$$

A conditional probability measure $\{G_{\boldsymbol{x}}, \boldsymbol{x} \in \mathcal{X}\}$ lies in the weak support of Π_n if it assigns positive probability to every basic neighborhood generated by the sub-base of the type (5.5.1).

Definition 5.5.4. A weak sup-neighborhood of a $F_{\boldsymbol{x}} \in \mathscr{G}_x^*$ is defined, for $\varepsilon > 0$, by:

$$S_\varepsilon^1(F_{\boldsymbol{x}}, q) := \left\{\{G_{\boldsymbol{x}} : \boldsymbol{x} \in X\} : \sup_{\boldsymbol{x}\in\mathcal{X}}\left|\int_{\mathbb{R}^3\times\mathbb{R}^+} g(\boldsymbol{\theta}, \sigma) d\left(G_{\boldsymbol{x}}(\boldsymbol{\theta}, \sigma) - F_{\boldsymbol{x}}(\boldsymbol{\theta}, \sigma)\right)\right| < \varepsilon\right\} \tag{5.5.2}$$

g bounded continuous function.

We introduce the weak and q-integrated neighborhoods for a set of conditional densities $\{f_0(\cdot \mid \boldsymbol{x}), \boldsymbol{x} \in \mathcal{X}\}$ as follows.

Definition 5.5.5. A sub-basis for a weak neighborhood for a set of conditional densities $\{f_0(\cdot \mid \boldsymbol{x}), \boldsymbol{x} \in \mathcal{X}\}$ is given, for $\varepsilon > 0$, by

$$W_{\varepsilon,k}(f_0) = \left\{f : f \in \mathcal{F}_d, \left|\int_{\mathcal{X}\times\mathcal{Y}} kh^y - \int_{\mathcal{X}\times\mathcal{Y}} kh_0^y\right| < \varepsilon\right\}, \tag{5.5.3}$$

where $k : \mathcal{X} \times \mathcal{Y} \to \mathbb{R}$ is a bounded, continuous function and $h^y(y, \boldsymbol{x}) = f(y \mid \boldsymbol{x})q(\boldsymbol{x})$. We define the same types of neighborhoods also for $g \in \mathcal{G}_d$.

A weak neighborhood basis is constructed from finite intersections of sets of the form (5.5.3).

Definition 5.5.6. The q-integrated $\mathbb{L}_1$ neighborhood is given, for $\varepsilon > 0$, by

$$W_\varepsilon(f_0; q) = \left\{f : f \in \mathcal{F}_d, \int \|f(\cdot \mid \boldsymbol{x}) - f_0(\cdot \mid \boldsymbol{x})\|_1\, q(\boldsymbol{x})\, d\boldsymbol{x} < \varepsilon\right\}. \tag{5.5.4}$$

Also in this case we define the same types of neighborhoods for $g \in \mathcal{G}_d$.

In the topology defined by (5.5.4), the space $\mathcal{F}_d$ is regarded as a closed subset of $\mathbb{L}_1(\lambda \times q, \mathcal{Y} \times \mathrm{supp}(q))$, a complete and separable metric space. As a result, measurability concerns do not emerge under these topologies. It is clear that to obtain consistency for the ATE we need to define new neighborhoods for fixed $a \in \mathcal{A}$.

Definition 5.5.7. For any fixed $a \in \mathcal{A}$, the q^a-integrated $\mathbb{L}_1$ neighborhood is given, for $\varepsilon > 0$, by

$$W_\varepsilon^a(f_0; q) = \left\{f : f \in \mathcal{F}_d, \int \|f(\cdot \mid a, z) - f_0(\cdot \mid a, z)\|_1\, q(z \mid a)\, dz < \varepsilon\right\}. \tag{5.5.5}$$

Similarly, we define the same neighborhoods for $g \in \mathcal{G}_d$.

5.5.2 Proofs

***Proof of Theorem** 5.2.2. Part (a).* Write $m_Y(a,z) = \mathrm{E}(Y \mid A=a, Z=z)$ and $m_W(a,z) = \mathrm{E}(W \mid A=a, Z=z)$. By iterated expectation and Assumption 3,

$$\begin{aligned}\mathrm{E}(Y \mid A,Z) &= \mathrm{E}\{\mathrm{E}(Y \mid A,U,Z) \mid A,Z\}\\ &= f_0(A,Z) + \{\alpha_0(A) + \alpha_1(A)Z\}\mathrm{E}(U \mid A,Z)\\ &= f_0(A,Z) + \{\alpha_0(A) + \alpha_1(A)Z\}\{\beta_1^u A + \beta_2^u Z\}.\end{aligned}$$

Fix a and set $c_0(a) = \beta_1^u a$. Then

$$m_Y(a,z) = f_0(a,z) + b_0(a) + b_1(a)z + b_2(a)z^2, \tag{5.5.6}$$

with

$$b_0(a) = \alpha_0(a)c_0(a), \qquad b_1(a) = \alpha_0(a)\beta_2^u + \alpha_1(a)c_0(a), \qquad b_2(a) = \alpha_1(a)\beta_2^u. \tag{5.5.7}$$

Consider $L_2(Q_a)$ with inner product $\langle r,s\rangle_a = \mathrm{E}[r(Z)s(Z) \mid A=a]$. Let $\mathcal{S}_a = \mathrm{span}\{1, Z, Z^2\}$ and $\phi(Z) = (1, Z, Z^2)^\top$. By (i), $f_0(a,\cdot) \perp \mathcal{S}_a$, hence the polynomial term $p_a(Z) = b_0(a) + b_1(a)Z + b_2(a)Z^2$ is the $L_2(Q_a)$-projection of $m_Y(a,\cdot)$ onto $\mathcal{S}_a$. By (ii), the projection coefficients are uniquely identified via the normal equations,

$$(b_0(a), b_1(a), b_2(a))^\top = G_a^{-1}\mathrm{E}[m_Y(a,Z)\phi(Z) \mid A=a], \qquad G_a = \mathrm{E}[\phi(Z)\phi(Z)^\top \mid A=a].$$

Define $g_2(a,z) := \beta_2^u\{\alpha_0(a) + \alpha_1(a)z\}$. Using (5.5.7), $\beta_2^u\alpha_1(a) = b_2(a)$ and, since $c_0(a) = \beta_1^u a$,

$$b_1(a) = \beta_2^u\alpha_0(a) + \alpha_1(a)c_0(a) \quad \Rightarrow \quad \beta_2^u\alpha_0(a) = b_1(a) - \alpha_1(a)c_0(a) = b_1(a) - \frac{b_2(a)}{\beta_2^u}\beta_1^u a.$$

Therefore

$$g_2(a,z) = b_1(a) + b_2(a)\Big(z - a\frac{\beta_1^u}{\beta_2^u}\Big). \tag{5.5.8}$$

Next, by iterated expectation and Assumption 2,

$$\mathrm{E}(W \mid A,Z) = \beta_0^w + \beta_1^w\mathrm{E}(U \mid A,Z) = \beta_0^w + \beta_1^w(\beta_1^u A + \beta_2^u Z) =: \theta_0^w + \theta_1^w A + \theta_2^w Z,$$

where $\theta_1^w = \beta_1^w\beta_1^u$ and $\theta_2^w = \beta_1^w\beta_2^u \neq 0$. Hence

$$\frac{\beta_1^u}{\beta_2^u} = \frac{\theta_1^w}{\theta_2^w}. \tag{5.5.9}$$

Combining (5.5.8)–(5.5.9) yields the identified form

$$\frac{g_2(a,z)}{\theta_2^w} = \frac{1}{\theta_2^w}\Big\{b_1(a) + b_2(a)\Big(z - a\frac{\theta_1^w}{\theta_2^w}\Big)\Big\} =: \gamma(a,z). \tag{5.5.10}$$

We now derive the ATE representation. Using the consistency 1 and the conditional independence 2,

$$\mathrm{E}\{Y(a)\} = \mathrm{E}[\mathrm{E}\{Y \mid A=a, U, Z\}] = \mathrm{E}[f_0(a,Z) + \{\alpha_0(a) + \alpha_1(a)Z\}U].$$

Also,

$$\mathrm{E}[\mathrm{E}(Y \mid A = a, Z)] = \mathrm{E}[f_0(a, Z) + \{\alpha_0(a) + \alpha_1(a)Z\}\mathrm{E}(U \mid A = a, Z)].$$

Subtracting gives

$$\mathrm{E}\{Y(a)\} = \mathrm{E}[\mathrm{E}(Y \mid A = a, Z)] + \mathrm{E}[\{\alpha_0(a) + \alpha_1(a)Z\}\{\mathrm{E}(U \mid A, Z) - \mathrm{E}(U \mid A = a, Z)\}]\,.$$

Since $\mathrm{E}(W \mid A, Z) = \beta_0^w + \beta_1^w \mathrm{E}(U \mid A, Z)$, we have

$$\begin{aligned}\mathrm{E}(U \mid A, Z) - \mathrm{E}(U \mid A = a, Z) &= \frac{1}{\beta_1^w}\{\mathrm{E}(W \mid A, Z) - \mathrm{E}(W \mid A = a, Z)\} \\ &= \frac{\beta_2^u}{\theta_2^w}\{\mathrm{E}(W \mid A, Z) - \mathrm{E}(W \mid A = a, Z)\}.\end{aligned}$$

Substitute and recall $g_2(a, Z) = \beta_2^u\{\alpha_0(a) + \alpha_1(a)Z\}$ to obtain

$$\mathrm{E}[Y(a)] = \mathrm{E}[\mathrm{E}(Y \mid A = a, Z)] + \mathrm{E}\left[\frac{g_2(a, Z)}{\theta_2^w}\{\mathrm{E}(W \mid A, Z) - \mathrm{E}(W \mid A = a, Z)\}\right].$$

Finally, replace g_2/θ_2^w by $\gamma(a, Z)$ from (5.5.10), yielding (5.2.21).

Part (b). Since a stratum representation preserves the joint law of (U, Z, A, Y, W), all conditional expectations below may be computed on the space carrying it, where S is available. First, by (5.2.22) and Assumption 3 of the theorem,

$$\begin{aligned}\mathrm{E}(Y \mid A, Z, S) &= \mathrm{E}(\mathrm{E}(Y \mid A, U, Z, S) \mid A, Z, S) = \mathrm{E}\left(\beta_{0,S}^y + \beta_{1,S}^y A + \beta_{2,S}^y U \mid A, Z, S\right) \\ &= \beta_{0,S}^y + \beta_{1,S}^y A + \beta_{2,S}^y \mathrm{E}(U \mid A, Z) = \beta_{0,S}^y + \left(\beta_{1,S}^y + \beta_{2,S}^y \beta_1^u\right) A + \beta_{2,S}^y \beta_2^u Z,\end{aligned}$$

where the third equality uses $S \perp\!\!\!\perp U \mid A, Z$. This is (5.2.23) with

$$\begin{cases} \theta_{0,S}^y = \beta_{0,S}^y \\ \theta_{1,S}^y = \beta_{1,S}^y + \beta_{2,S}^y \beta_1^u \\ \theta_{2,S}^y = \beta_{2,S}^y \cdot \beta_2^u \end{cases} \tag{5.5.11}$$

Next we show that marginalising out S recovers the underlying specification. By (5.2.22), i.e. the stratum-wise outcome regression together with $S \perp\!\!\!\perp U \mid A, Z$,

$$\begin{aligned}\mathrm{E}(Y \mid A, U, Z) &= \mathrm{E}(\mathrm{E}(Y \mid A, U, S, Z) \mid A, U, Z) \\ &= \mathrm{E}(\beta_{0,S}^y \mid A, Z) + \mathrm{E}(\beta_{1,S}^y \mid A, Z)\, A + \mathrm{E}(\beta_{2,S}^y \mid A, Z)\, U,\end{aligned}$$

while Assumption 1 of the theorem gives $\mathrm{E}(Y \mid A, U, Z) = f_0(A, Z) + \{\alpha_0(A) + \alpha_1(A)Z\}U$. These are two affine representations of the same regression as a function of U: their difference, an affine function of U, vanishes on the support of $U \mid A = a, Z = z$, which, for Q_a-almost every z, contains at least two points because $\mathrm{Var}(U \mid A = a, Z = z) > 0$; so both its coefficients are zero (the exact analogue of the argument in the proof of Theorem 5.2.1(b), with U in place of Z). Hence

$$\mathrm{E}\left(\beta_{2,S}^y \mid A = a, Z\right) = \alpha_0(a) + \alpha_1(a)Z, \quad Q_a\text{-a.s.} \tag{5.5.12}$$

Averaging (5.5.11) over $S \mid A = a, Z$ and using (5.5.12):

$$\boldsymbol{\theta}_2(a, Z) = \mathrm{E}\left(\theta_{2,S}^y \mid A = a, Z\right) = \beta_2^u\, \mathrm{E}\left(\beta_{2,S}^y \mid A = a, Z\right) = \beta_2^u\{\alpha_0(a) + \alpha_1(a)Z\}.$$

From equations (5.5.7)–(5.5.10) in the proof of part (a), with $\theta_2^w = \beta_1^w \beta_2^u \neq 0$:

$$\gamma(a,Z) = \frac{1}{\theta_2^w}\{b_1(a) + b_2(a)(Z - a\tfrac{\theta_1^w}{\theta_2^w})\} = \frac{\alpha_0(a) + \alpha_1(a)Z}{\beta_1^w},$$

so that

$$\frac{\boldsymbol{\theta}_2(a,Z)}{\theta_2^w} = \frac{\beta_2^u\{\alpha_0(a) + \alpha_1(a)Z\}}{\beta_1^w\beta_2^u} = \gamma(a,Z) \quad Q_a\text{-a.s.}$$

Finally we derive the closed form. Substituting $\gamma(a,Z) = \boldsymbol{\theta}_2(a,Z)/\theta_2^w$ into the ATE representation (5.2.21) of part (a):

$$\mathrm{E}\, Y(a) = \mathrm{E}\,\mathrm{E}(Y \mid A = a, Z) + \mathrm{E}\left[\frac{\boldsymbol{\theta}_2(a,Z)}{\theta_2^w}(\mathrm{E}(W \mid A, Z) - \mathrm{E}(W \mid A = a, Z))\right].$$

By (5.2.23) and the tower property, $\mathrm{E}(Y \mid A = a, Z) = \boldsymbol{\theta}_0(a,Z) + \boldsymbol{\theta}_1(a,Z)\, a + \boldsymbol{\theta}_2(a,Z)\, Z$; by part (a), $\mathrm{E}(W \mid A, Z) - \mathrm{E}(W \mid A = a, Z) = \theta_1^w(A - a)$; and, again by the tower property,

$$\mathrm{E}\left[\frac{\boldsymbol{\theta}_2(a,Z)}{\theta_2^w}\, W\right] = \mathrm{E}\left[\frac{\boldsymbol{\theta}_2(a,Z)}{\theta_2^w}(\theta_0^w + \theta_1^w A + \theta_2^w Z)\right].$$

Combining the three identities,

$$\begin{aligned}
&\mathrm{E}\left[\boldsymbol{\theta}_0(a,Z) + \boldsymbol{\theta}_1(a,Z)\, a - \frac{\boldsymbol{\theta}_2(a,Z)}{\theta_2^w}(\theta_0^w + \theta_1^w a)\right] + \mathrm{E}\left[\frac{\boldsymbol{\theta}_2(a,Z)}{\theta_2^w}\, W\right] \\
&\quad = \mathrm{E}[\boldsymbol{\theta}_0(a,Z) + \boldsymbol{\theta}_1(a,Z)\, a + \boldsymbol{\theta}_2(a,Z)\, Z] + \mathrm{E}\left[\frac{\boldsymbol{\theta}_2(a,Z)}{\theta_2^w}\,\theta_1^w(A - a)\right] \\
&\quad = \mathrm{E}\,\mathrm{E}(Y \mid A = a, Z) + \mathrm{E}\left[\frac{\boldsymbol{\theta}_2(a,Z)}{\theta_2^w}(\mathrm{E}(W \mid A, Z) - \mathrm{E}(W \mid A = a, Z))\right] = \mathrm{E}\, Y(a),
\end{aligned}$$

which is the closed form (5.2.24), completing the proof. □

Lemma 5.5.1. *Fix* $\{F_{\boldsymbol{x}} : \boldsymbol{x} \in \mathcal{X}\} \in \mathscr{G}_x$ *and a measurable partition* $\{A_k, k = 1, \ldots, m\}$ *of* $\mathbb{R}^3 \times \mathbb{R}^+$. $\forall\, \varepsilon_m > 0$*:*

$$\liminf_{n\to\infty} \Pi\left(\sup_{\boldsymbol{x}\in\mathcal{X}} \left|w_k^n(\boldsymbol{x}) - F_{\boldsymbol{x}}(A_k)\right| < \varepsilon_m,\ k = 1, \ldots, m\right) > 0.$$

Proof. Without loss of generality, let $0 < F_{\boldsymbol{x}}(A_k) < 1$, $k = 1, \ldots, m$, $\forall\, \boldsymbol{x} \in \mathcal{X}$ and for, λ Lebesgue measure on $\mathbb{R}^3 \times \mathbb{R}^+$, $\lambda(\mathcal{X}) = 1$. Define $\forall\, \boldsymbol{x} \in \mathcal{X}$:

$$g_1(\boldsymbol{x}) = F_{\boldsymbol{x}}(A_1), \quad g_k(\boldsymbol{x}) \textstyle\prod_{l<k}(1 - g_l(\boldsymbol{x})) = F_{\boldsymbol{x}}(A_k), \quad 2 \leqslant k \leqslant m-1, \quad g_m(\boldsymbol{x}) = 1,$$

because the requirement $\sum_{k=1}^m F_{\boldsymbol{x}}(A_k) = 1$ implies $g_m(\boldsymbol{x}) = 1$. For $k < m$ it is easy to solve the above equations and get by recursion:

$$\begin{aligned}
&g_1(\boldsymbol{x}) = F_{\boldsymbol{x}}(A_1), \quad g_2(\boldsymbol{x}) = \frac{F_{\boldsymbol{x}}(A_2)}{1 - F_{\boldsymbol{x}}(A_1)}, \quad g_3(\boldsymbol{x}) = \frac{F_{\boldsymbol{x}}(A_3)}{1 - F_{\boldsymbol{x}}(A_1) - F_{\boldsymbol{x}}(A_2)}, \\
&g_4(\boldsymbol{x}) = \frac{F_{\boldsymbol{x}}(A_4)}{1 - F_{\boldsymbol{x}}(A_1) - F_{\boldsymbol{x}}(A_2) - F_{\boldsymbol{x}}(A_3)} \quad \cdots \quad g_k(\boldsymbol{x}) = \frac{F_{\boldsymbol{x}}(A_k)}{1 - \sum_{l<k} F_{\boldsymbol{x}}(A_l)}.
\end{aligned}$$

Recall that the functions α_k^n in the prior are of the form

$$\alpha_k^n(\boldsymbol{x}) = \sum_{m=1}^{N_n} \left(\eta_{0,m}^k + \eta_{1,m}^k a + \eta_{2,m}^k z\right) \mathbf{1}_{(a,z)\in T_{m,n}} \tag{5.5.13}$$

for a sequence of partitions $\{T_{m,n}\}_{m=1}^{N_n}$ of $\mathcal{X}$ indexed by $n \in \mathbb{N}$ ($N_n \to \infty$, as $n \to \infty$) and $\left(\eta_{0,m}^k, \eta_{1,m}^k, \eta_{2,m}^k\right)$ is normally distributed, with diagonal covariance matrix.

Now notice that $\forall\, \varepsilon_m > 0$, since the functions g_k, $k = 1, \ldots, m$ are continuous on a compact $\mathcal{X}$ with compact codomain contained in [0, 1], Φ is Lipschitz continuous and $\Phi(\Phi^{-1}(g_k(\boldsymbol{x}))) = g_k(\boldsymbol{x})$, there exist coefficients $\left\{\eta_{0,m}^k, \eta_{1,m}^k, \eta_{2,m}^k\right\}_{m=1}^{\infty}$ and a $N_\varepsilon \in \mathbb{N}$ such that $\forall\, N_n \geq N_\varepsilon$:

$$\sup_{\boldsymbol{x}\in\mathcal{X}} \left| \Phi\left(\sum_{m=1}^{N_n} \left(\eta_{0,m}^k + \eta_{1,m}^k a + \eta_{2,m}^k z\right) \mathbf{1}_{(a,z)\in T_{m,n}} \right) - g_k(\boldsymbol{x}) \right| \tag{5.5.14}$$

$$\lesssim \sup_{\boldsymbol{x}\in\mathcal{X}} \left| \sum_{m=1}^{N_n} \left(\eta_{0,m}^k + \eta_{1,m}^k a + \eta_{2,m}^k z\right) \mathbf{1}_{(a,z)\in T_{m,n}} - \Phi^{-1}\left(g_k(\boldsymbol{x})\right) \right| < \frac{\varepsilon_m}{2}. \tag{5.5.15}$$

This follows from the Arzelà-Ascoli theorem, as the approximating functions form a uniformly bounded sequence of real-valued differentiable functions on a compact domain, with uniformly bounded derivatives. This proves that the functions $\Phi^{-1}(g_k)$ are in the linear span of the piecewise linear functions in $\boldsymbol{x} = (a, z)$.

Because the coefficients $\left\{\eta_{0,m}^k, \eta_{1,m}^k, \eta_{2,m}^k\right\}_{m=1}^{\infty}$ are in the support of the prior with positive probability, as the prior on the weights has full support (c.f. proof of Lemma 2.2 in [42]), it follows thus from (5.5.14) and (5.5.13) that:

$$\liminf_{n\to\infty} \Pi\left(\sup_{\boldsymbol{x}\in\mathcal{X}} \left|\Phi(\alpha_k^n(\boldsymbol{x})) - g_k(\boldsymbol{x})\right| < \varepsilon_m \right) > 0$$

and for $k = m$:

$$\Pi\left(\sup_{x\in\mathcal{X}} \left|\Phi\left(\alpha_m^n(\boldsymbol{x})\right) - 1\right| < \varepsilon_m \right) = \Pi\left(\inf_{\boldsymbol{x}\in\mathcal{X}} \alpha_m^n(\boldsymbol{x}) > \Phi^{-1}\left(1 - \varepsilon_m\right) \right)$$

choose $M > \Phi^{-1}\left(1 - \varepsilon_m\right) + \varepsilon_m$. We have $0 < M < 1$ and

$$\left\{ \sup_{x\in\mathcal{X}} \left|\alpha_m^n(\boldsymbol{x}) - M\right| < \varepsilon_m \right\} \subset \left\{ \inf_{\boldsymbol{x}\in\mathcal{X}} \alpha_m^n(\boldsymbol{x}) > \Phi^{-1}\left(1 - \varepsilon_m\right) \right\}$$

So:

$$\liminf_{n\to\infty} \Pi\left(\sup_{\boldsymbol{x}\in\mathcal{X}} \left|\Phi\left(\alpha_m^n(\boldsymbol{x})\right) - 1\right| < \varepsilon_m \right) > \liminf_{n\to\infty} \Pi\left(\sup_{\boldsymbol{x}\in\mathcal{X}} \left|\alpha_m^n(\boldsymbol{x}) - M\right| < \varepsilon_m \right) > 0$$

Now let S_{m-1} be the m-dimensional simplex. Define $\bar{\Phi}_n = \left(\Phi\left(\alpha_1^n(\boldsymbol{x})\right), \ldots, \Phi\left(\alpha_m^n(\boldsymbol{x})\right)\right)$, $\bar{g} = (g_1(\boldsymbol{x}), \ldots, g_m(\boldsymbol{x}))$. Let $z = (z_1, \ldots, z_m)^\top$, $f_k : S_{m-1} \to \mathbb{R}$ the function that maps $z \mapsto z_k \prod_{l<k}(1 - z_l)$, $k = 1, \ldots, m$. Note: $\left\| f_k\left(\bar{\Phi}_n\right) - f_k(\bar{g}) \right\|_\infty = \sup_{\boldsymbol{x}\in\mathcal{X}} \left| w_k^n(\boldsymbol{x}) - F_{\boldsymbol{x}}\left(A_k\right) \right|$

$$\left\| f_k\left(\bar{\Phi}_n\right) - f_k(\bar{g}) \right\|_\infty = \left\| \Phi\left(\alpha_k^n(\boldsymbol{x})\right) \left\{ 1 - \sum_{l<k} f_l\left(\bar{\Phi}_n\right) \right\} - g_k(\boldsymbol{x}) \left\{ 1 - \sum_{l<k} f_l(\bar{g}) \right\} \right\|_\infty$$

$$\leqslant (k-1) \sup_{\boldsymbol{x}\in\mathcal{X}} \left| \Phi\left(\alpha_k^n(\boldsymbol{x})\right) - g_k(\boldsymbol{x}) \right| + \sum_{l<k} \left\| f_l\left(\bar{\Phi}_n\right) - f_l(\bar{g}) \right\|_\infty$$

Thus one can find $\varepsilon_m > 0$ such that:

$$\begin{aligned}&\left\{\sup_{\boldsymbol{x}\in\mathcal{X}}\left|\Phi\left(\alpha_k^n(\boldsymbol{x})\right) - g_k(\boldsymbol{x})\right| < \varepsilon_m,\ k = 1,\ldots,m\right\}\\ &\subset \left\{\left\|f_1\left(\bar{\Phi}_n\right) - f_1(\bar{g})\right\|_\infty < \varepsilon_m, \ldots, \left\|f_m\left(\bar{\Phi}_n\right) - f_m(\bar{g})\right\|_\infty < \varepsilon_m\right\}\end{aligned}$$

Conclude by noticing:

$$\begin{aligned}&\liminf_{n\to\infty}\Pi\left(\left\|f_1\left(\bar{\Phi}_n\right) - f_1(\bar{g})\right\|_\infty < \varepsilon_m, \ldots, \left\|f_m\left(\bar{\Phi}_n\right) - f_m(\bar{g})\right\|_\infty < \varepsilon_m\right)\\ &> \liminf_{n\to\infty}\Pi\left\{\sup_{\boldsymbol{x}\in\mathcal{X}}\left|\Phi\left(\alpha_k^n(\boldsymbol{x})\right) - g_k(\boldsymbol{x})\right| < \varepsilon_m,\ k = 1,\ldots,m\right\} =\\ &= \prod_{k=1}^{m}\liminf_{n\to\infty}\Pi\left\{\sup_{\boldsymbol{x}\in\mathcal{X}}\left|\Phi\left(\alpha_k^n(\boldsymbol{x})\right) - g_k(\boldsymbol{x})\right| < \varepsilon_m\right\} > 0\end{aligned}$$

□

***Proof of Theorem** 5.3.1.* The technique we will use to show $f_0 \in \mathrm{KL}(\Pi_n)$ is to use the tail conditions on $f_0(y \mid \boldsymbol{x})$ to approximate it only on a compact interval by $\tilde{f}(y \mid \boldsymbol{x}) = \int \frac{1}{\sigma}\phi(\frac{y-\bar{\boldsymbol{x}}'\boldsymbol{\theta}^y}{\sigma})\, d\tilde{G}_{\boldsymbol{x}}(\boldsymbol{\theta}^y, \sigma)$, where $\{\tilde{G}_{\boldsymbol{x}}, \boldsymbol{x} \in \mathcal{X}\}$ is compactly supported. Since:

$$\mathrm{KL}\left(h_0^y, h^y\right) = \underbrace{\int_{\mathcal{X}}\int_{\mathcal{Y}} f_0(y \mid \boldsymbol{x}) \log\frac{f_0(y \mid \boldsymbol{x})}{\tilde{f}(y \mid \boldsymbol{x})}\, dy\, q(\boldsymbol{x})\, d\boldsymbol{x}}_{(1)} + \underbrace{\int_{\mathcal{X}}\int_{\mathcal{Y}} f_0(y \mid \boldsymbol{x}) \log\frac{\tilde{f}(y \mid \boldsymbol{x})}{f(y \mid \boldsymbol{x})}\, dy\, q(\boldsymbol{x})\, d\boldsymbol{x}}_{(2)}$$

we construct $\tilde{f}$ which makes (1) sufficiently small. We next show that (2) is also sufficiently small if $\{G_{\boldsymbol{x}}, \boldsymbol{x} \in \mathcal{X}\}$ lies inside a finite intersection of neighborhoods of $\{\tilde{G}_{\boldsymbol{x}}, \boldsymbol{x} \in \mathcal{X}\}$ of the type (5.5.2).

For brevity, in the following proofs we will denote $\boldsymbol{\theta}^y$ simply by $\boldsymbol{\theta}$ and σ_y by σ.

Proof of $(1) < \varepsilon/2$.

Define a sequence of conditional densities $f_k(y \mid \boldsymbol{x})$ for $k \geq 1$:

$$f_k(y \mid \boldsymbol{x}) = \int \frac{1}{\sigma}\phi\left(\frac{y - \bar{\boldsymbol{x}}'\boldsymbol{\theta}}{\sigma}\right) d\tilde{G}_{\boldsymbol{x}}^k(\boldsymbol{\theta}, \sigma) = \frac{\int_{-kx_1}^{kx_1} \frac{1}{\sigma_k}\phi\left(\frac{y-t}{\sigma_k}\right) f_0(t \mid \boldsymbol{x}) dt}{\int_{-kx_1}^{kx_1} f_0(t \mid \boldsymbol{x}) dt}$$

where for $\eta > 0$, $\sigma_k = k^{-\eta}$. The above given f_k, which is a mollification of f_0, can be obtained using as measure $\tilde{G}_{\boldsymbol{x}}^k$:

$$d\tilde{G}_{\boldsymbol{x}}^k(\boldsymbol{\theta}, \sigma) = \frac{\mathbf{1}_{\theta_1\in[-k,k]} f_0\left(\boldsymbol{x}'\boldsymbol{\theta} \mid \boldsymbol{x}\right) \prod_{j\neq 1}\delta_0\left(\theta_j\right)\delta_{\sigma_n}(\sigma)}{\int_{-k}^{k} f_0\left(x_1\theta_1 \mid \boldsymbol{x}\right) d\theta_1}$$

An application of DCT using the conditions A1-A5 will yield

$$\int_{\mathcal{X}}\int_{\mathcal{Y}} f_0(y \mid \boldsymbol{x}) \log\frac{f_0(y \mid \boldsymbol{x})}{f_k(y \mid \boldsymbol{x})}\, dy\, q(\boldsymbol{x})\, d\boldsymbol{x} \to 0 \quad \text{as } n \to \infty. \tag{5.5.16}$$

Therefore choose $\tilde{f} = f_{k_0}$ for sufficiently large k_0. We now show formally (5.5.16). We work with:

$$f_k(y \mid \boldsymbol{x}) = t_k \int_{-k}^{k} \frac{1}{\sigma_k} \phi(y-t) f_0(t \mid \boldsymbol{x})\, dt, \qquad t_k = \left(\int_{-k}^{k} f_0(t \mid \boldsymbol{x}) dt \right)^{-1} \tag{5.5.17}$$

Find positive constant ξ s.t. $\int_{-\xi}^{\xi} \phi(t)dt > 1 - \varepsilon$. For sufficiently large $k \in \mathbb{N}$ such that $[y - \xi\sigma_k, y + \xi\sigma_k] \subset [-k, k]$, since $t_k \to 1, \sigma_k \to 0 \Rightarrow f_k(y \mid \boldsymbol{x}) \to f_0(y \mid \boldsymbol{x})$ as $k \to \infty$ by the continuity of f_0. Thus:

$$\log \frac{f_0(y \mid \boldsymbol{x})}{f_k(y \mid \boldsymbol{x})} \to 0, \quad \forall\, (y, \boldsymbol{x}) \in \mathcal{Y} \times \mathcal{X}.$$

Since t_k decreases and $\|f_0\|_\infty < M$:

$$f_k(y \mid \boldsymbol{x}) = t_k \int_{-k}^{k} \frac{1}{\sigma_k} \phi\left(\frac{y-t}{\sigma_k} \right) f_0(t \mid \boldsymbol{x})\, dt \le M t_k \le M t_1.$$

Now fix $y \in \mathcal{Y}$. Since $|y - t| \le |y| + k,\ \forall\, t \in [-k, k]$ and $t_k \geqslant 1, \forall\, k \leqslant |y|$:

$$f_k(y \mid \boldsymbol{x}) \ge \frac{1}{\sigma_k} \phi\left(\frac{|y| + k}{\sigma_k} \right) = k^\eta \phi(k^\eta(|y| + k)) \ge |y|^\eta \phi\left(2|y|^{1+\eta}\right)$$

The last inequality follows from the fact that $\tau^\eta \phi(\tau^\eta(|y| + \tau))$ is decreasing in τ for $\tau \ge 1$. Let $\psi^\eta_{\boldsymbol{x}}(y) = \inf_{t \in [y-\sigma_k, y+\sigma_k]} f_0(t \mid \boldsymbol{x})$. Let $A_k = [-k, k] \cap [y - \sigma_k, y + \sigma_k]$ and $c = \int_0^1 \phi(t)\, dt < 1,\ \forall\, k > |y|$:

$$\begin{aligned} f_k(y \mid \boldsymbol{x}) &\geqslant t_k \int_{A_k} \frac{1}{\sigma_k} \phi\left(\frac{y-t}{\sigma_k} \right) f_0(t \mid \boldsymbol{x})\, dt \\ &\geqslant t_k \psi^k_{\boldsymbol{x}}(y) \int_{A_k} \frac{1}{\sigma_k} \phi\left(\frac{y-t}{\sigma_k} \right) dt \end{aligned}$$

since $t_k \ge 1, \psi^k_{\boldsymbol{x}}(y) \geqslant \psi_{\boldsymbol{x}}(y)$ and $\int_{A_k} \frac{1}{\sigma_k} \phi\left(\frac{y-t}{\sigma_k}\right) dt \ge \int_0^1 \phi(t)dt = c,\ \forall\, k \geqslant 1$ and $\boldsymbol{x} \in \mathbb{R}^3$ then

$$f_k(y \mid \boldsymbol{x}) \geqslant c\psi_{\boldsymbol{x}}(y), \quad \forall\, k > |y|$$

So:

$$f_k(y \mid \boldsymbol{x}) \ge \begin{cases} c\psi_{\boldsymbol{x}}(y), & |y| < 1 \\ \min\left\{ |y|^\eta \phi\left(2|y|^{1+\eta}\right), c\psi_{\boldsymbol{x}}(y) \right\}, & |y| \geqslant 1 \end{cases}$$

Therefore:

$$\left| \log \frac{f_0(y \mid \boldsymbol{x})}{f_k(y \mid \boldsymbol{x})} \right| \le \log \frac{M t_1}{f_0(y \mid \boldsymbol{x})} + \log \frac{f_0(y \mid \boldsymbol{x})}{c\psi_{\boldsymbol{x}}(y)} + \mathbf{1}_{|y| \ge 1} \log \frac{f_0(y \mid \boldsymbol{x})}{|y|^\eta \phi\left(2|y|^{1+\eta}\right)}$$

The last expression is integrable with respect to $h_0^y(\boldsymbol{x}, y) = f_0(y \mid \boldsymbol{x}) q(\boldsymbol{x})$ by assumption.

Proof of $(2) < \varepsilon/2$.

We first prove that the prior gives positive prior mass to arbitrarily small neighborhoods of the form (5.5.2) of $\{\tilde{G}^{k_0}_{\boldsymbol{x}} : \boldsymbol{x} \in \mathcal{X}\}$. 1) $\{\tilde{G}^{k_0}_{\boldsymbol{x}} : \boldsymbol{x} \in x\}$ is compactly supported. 2) $\boldsymbol{x} \mapsto \tilde{G}^{k_0}_{\boldsymbol{x}}(A)$ is

continuous for every measurable set A. So $\tilde{G}_{\boldsymbol{x}}^{k_0} \in \mathscr{G}_x$. Because $f_0(y \mid \boldsymbol{x})$ is an arbitrary function of $\boldsymbol{x}$ we must be able to approximate generic measures $\boldsymbol{x} \mapsto F_{\boldsymbol{x}} \in \mathscr{G}_x$.

Fix $\{F_{\boldsymbol{x}} : \boldsymbol{x} \in \mathcal{X}\} \in \mathscr{G}_x$. Without loss of generality, it is enough to show for a bounded uniformly continuous function $g : \mathbb{R}^3 \times \mathbb{R}^+ \mapsto [0,1]$ such that $g(\boldsymbol{\theta}, \sigma) \to 0$, as $\|\boldsymbol{\theta}\| \to \infty, \sigma \to \infty$, that $\varepsilon > 0$;

$$\liminf_{n\to\infty} \Pi_n \left\{ \{G_{\boldsymbol{x}}^n : \boldsymbol{x} \in x\} : \sup_{\boldsymbol{x}\in\mathcal{X}} \left| \int_{\mathbb{R}^3\times\mathbb{R}^+} g(\boldsymbol{\theta}, \sigma)\, d\left(G_{\boldsymbol{x}}^n(\boldsymbol{\theta}, \sigma) - F_{\boldsymbol{x}}(\boldsymbol{\theta}, \sigma)\right) \right| < \varepsilon \right\} > 0 \tag{5.5.18}$$

Fix $\varepsilon > 0$, then $\exists\, a, \bar{\sigma}, \underline{\sigma} > 0$ not depending on $\boldsymbol{x}$ such that $F_{\boldsymbol{x}}([-a,a]^3 \times [\bar{\sigma}, \underline{\sigma}]) \overset{(A)}{>} 1 - \varepsilon,\ \forall\, \boldsymbol{x} \in \mathcal{X}$. Let $C = [-a,a]^3 \times [\bar{\sigma}, \underline{\sigma}]$ and using the definition of the prior

$$\begin{aligned} &\sup_{\boldsymbol{x}\in\mathcal{X}} \left| \int_{\mathbb{R}^3\times\mathbb{R}^+} g(\boldsymbol{\theta}, \sigma)\, d\left(G_{\boldsymbol{x}}^n(\boldsymbol{\theta}, \sigma) - F_{\boldsymbol{x}}(\boldsymbol{\theta}, \sigma)\right) \right| \\ &\overset{(A)}{\leq} \sup_{\boldsymbol{x}\in\mathcal{X}} \left| \sum_{k=1}^{\infty} w_k^n(\boldsymbol{x}) g\left(\boldsymbol{\theta}_k, \sigma_k\right) - \int_C g(\boldsymbol{\theta}, \sigma) dF_{\boldsymbol{x}}(\boldsymbol{\theta}, \sigma) \right| + \varepsilon \end{aligned} \tag{5.5.19}$$

w_k^n are as in the prior and $(\boldsymbol{\theta}_k, \sigma_k) \sim G_0$. Now for each $\boldsymbol{x} \in \mathcal{X}$, construct a Riemann sum approximation of: $\int_C g(\boldsymbol{\theta}, \sigma, \boldsymbol{x})\, dF_{\boldsymbol{x}}(\boldsymbol{\theta}, \sigma)$. Let $\{A_{k,m}, k = 1, \ldots, m\}$ be a sequence of partitions of C with increasing refinement as $m \to \infty$. Assure $\max_{1\leq k\leq m} \operatorname{diam}(A_{k,m}) \to 0$ as $n \to \infty$. Fix $(\tilde{\boldsymbol{\theta}}_{k,m}, \tilde{\sigma}_{k,m}) \in A_{k,m},\ k = 1, \ldots, m$. This is a uniformly continuous sequence of functions on $\mathcal{X}$ which decreases monotonically to its limit, thus as $m \to \infty$

$$\sup_{\boldsymbol{x}\in\mathcal{X}} \left| \sum_{k=1}^{m} g\left(\tilde{\boldsymbol{\theta}}_{k,m}, \tilde{\sigma}_{k,m}\right) F_{\boldsymbol{x}}\left(A_{k,m}\right) - \int_C g(\boldsymbol{\theta}, \sigma) dF_{\boldsymbol{x}}(\boldsymbol{\theta}, \sigma) \right| \to 0.$$

Hence there exists m_1 such that for all $m \geqslant m_1$:

$$(5.5.19) \leqslant \sup_{\boldsymbol{x}\in\mathcal{X}} \left| \sum_{k=1}^{\infty} w_k^n(\boldsymbol{x}) g\left(\boldsymbol{\theta}_k, \sigma_k\right) - \sum_{k=1}^{m} g\left(\tilde{\boldsymbol{\theta}}_{k,m}, \tilde{\sigma}_{k,m}\right) F_{\boldsymbol{x}}\left(A_{k,m}\right) \right| + 2\varepsilon \tag{5.5.20}$$

Define now the set:

$$\Omega_{1,n} := \left\{ \omega \,:\, \sup_{\boldsymbol{x}\in\mathcal{X}} \left| w_k^n(\boldsymbol{x})(\omega) - F_{\boldsymbol{x}}\left(A_k\right)(\omega) \right| < \varepsilon_m,\ k = 1, \ldots, m \right\}.$$

We proved in Lemma 5.5.1: $\liminf_{n\to\infty} \Pi(\Omega_{1,n}) > 0$. As $\sum_{k=1}^{\infty} w_k(\boldsymbol{x}) = 1$ a.s. there exists Ω_{00} with prior probability 1 such that on Ω_{00}, $\sum_{k=1}^{m} w_k(\boldsymbol{x}) \to 1$ as $m \to \infty$. Note that, because of the boundedness in $\boldsymbol{x}$, the convergence is uniform in $\boldsymbol{x}$ by Dini's theorem as these are continuous functions defined on a compact set $\mathcal{X}$, monotonically increasing to a continuous function identically equal to 1. Because Ω_{00} has prior probability 1, it follows that $\liminf_{n\to\infty} \Pi(\Omega_{1,n} \cap \Omega_{00}) > 0$. By Egoroff's theorem, there exists a measurable subset $\Omega_{2,n} \subset \Omega_{1,n} \cap \Omega_{00}$ with $\liminf_{n\to\infty} \Pi(\Omega_{2,n}) > 0$ such that within this subset $\sum_{k=1}^{m} w_k(\boldsymbol{x}) \to 1$ as $m \to \infty$ uniformly in $\boldsymbol{x}$ and in $\omega \in \Omega_{2,n}$. Thus $\forall\, \varepsilon > 0$ there exists a positive integer $m_\varepsilon \in \mathbb{N}$ independent of ω such that on $\Omega_{2,n}$;

$$\sup_{\boldsymbol{x}\in\mathcal{X}} \sum_{k=m_\varepsilon+1}^{\infty} w_k(\boldsymbol{x}) < \varepsilon. \tag{5.5.21}$$

Consider now any fixed $m_\varepsilon \in \mathbb{N}$. For ω on a set Ω_0 of prior probability 1 and $\forall\, k \in \mathbb{N}$, let $\alpha_k(\boldsymbol{x})$ be the limiting functions of the sequence of functions $\{\alpha_k^n(\boldsymbol{x})\}_n$. Because the sequence of these functions is uniformly bounded on a compact domain with uniformly bounded derivatives, by Arzela-Ascoli theorem: $\sup_{\boldsymbol{x}\in\mathcal{X}} |\alpha_k^n(\boldsymbol{x}) - \alpha_k(\boldsymbol{x})| \to 0$. Now note that Φ is Lipschitz-continuous, as well as $(z,\ldots,z_k) \mapsto z_k \prod_{l<k}(1-z_l)$. Any finite sum of m_ε Lipschitz continuous functions is Lipschitz continuous. We thus have that on a set Ω_0 of prior probability 1, for any fixed $m_\varepsilon \in \mathbb{N}$, as $1 - \sum_{k=m_\varepsilon+1}^{\infty} w_k^n(\boldsymbol{x}) = \sum_{k=1}^{m_\varepsilon} w_k^n(\boldsymbol{x})$, as $n \to \infty$:

$$\sup_{\boldsymbol{x}\in\mathcal{X}} \left| \sum_{k=m_\varepsilon+1}^{\infty} \left(w_k^n(\boldsymbol{x}) - w_k(\boldsymbol{x})\right) \right| = \sup_{\boldsymbol{x}\in\mathcal{X}} \left| \sum_{k=1}^{m_\varepsilon} \left(w_k^n(\boldsymbol{x}) - w_k(\boldsymbol{x})\right) \right| \to 0 \tag{5.5.22}$$

We conclude therefore, for $\Omega_{3,n} := \Omega_{2,n} \cap \Omega_0$: $\liminf_{n\to\infty} \Pi(\Omega_{3,n}) > 0$.

By definition of the functions g, $\exists\, k > 0$ independent of $\boldsymbol{x}$, such that $g(\boldsymbol{\theta},\sigma) < \varepsilon$ if $\|\boldsymbol{\theta}\| > k$, $\sigma > k$.

Let

$$\Omega_{4,n} = \Omega_{3,n} \cap \{(\boldsymbol{\theta}_{m+1},\sigma_{m+1}) \in A_1,\ldots,(\boldsymbol{\theta}_{m_\varepsilon-1},\sigma_{m_\varepsilon-1}) \in A_1\}$$

and

$$A_1 = \{(\boldsymbol{\theta},\sigma) : \|\boldsymbol{\theta}\| > k, \sigma > k\}$$

(note that these sets have positive prior probability and their intersection $\Omega_{4,n}$ satisfies $\liminf_{n\to\infty} \Pi(\Omega_{4,n}) > 0$ as they are independent of each others). Furthermore $\exists\, B_k$, $k = 1,\ldots,m$ independent of $\boldsymbol{x}$, such that if $(\boldsymbol{\theta}_k,\sigma_k) \in B_k$

$$|g(\boldsymbol{\theta}_k,\sigma_k) - g(\tilde{\boldsymbol{\theta}}_{k,m},\tilde{\sigma}_{k,m})| < \varepsilon$$

So let $\Omega_{5,n} = \Omega_{4,n} \cap \{(\boldsymbol{\theta}_1,\sigma_1) \in B_1,\ldots,(\boldsymbol{\theta}_m,\sigma_m) \in B_m\}$. Then $\liminf_{n\to\infty} \Pi(\Omega_{5,n}) > 0$ (all the other subsets were chosen independently).

Recall that we denote by $w_k(\boldsymbol{x})$ the pointwise in $\boldsymbol{x}$ almost sure limit of each $w_k^n(\boldsymbol{x})$. Moreover m_ε is chosen as in (5.5.21). Then, using $\|g\|_\infty < \infty$ and the triangle inequality:

$$\begin{aligned}
(5.5.20) \le{}& \sup_{\boldsymbol{x}\in\mathcal{X}} \left| \sum_{k=m_\varepsilon+1}^{\infty} \left(w_k^n(\boldsymbol{x}) - w_k(\boldsymbol{x})\right) \right| + \sup_{\boldsymbol{x}\in\mathcal{X}} \sum_{k=m_\varepsilon+1}^{\infty} w_k(\boldsymbol{x}) \\
&+ \sup_{\boldsymbol{x}\in\mathcal{X}} \sum_{k=m+1}^{m_\varepsilon} w_k^n(\boldsymbol{x}) \left|g\left(\boldsymbol{\theta}_k,\sigma_k\right)\right| + \sup_{\boldsymbol{x}\in\mathcal{X}} \left| \sum_{k=1}^{m} \left(w_k^n(\boldsymbol{x}) - F_{\boldsymbol{x}}\left(A_{k,m}\right)\right) g\left(\boldsymbol{\theta}_k,\sigma_k\right) \right| \\
&+ \sup_{\boldsymbol{x}\in\mathcal{X}} \left| \sum_{k=1}^{m} F_{\boldsymbol{x}}\left(A_{k,m}\right) \left(g(\boldsymbol{\theta}_k,\sigma_k) - g(\tilde{\boldsymbol{\theta}}_{k,m},\tilde{\sigma}_{k,m})\right) \right| + 2\varepsilon < 7\varepsilon.
\end{aligned}$$

This proves that the prior gives positive prior mass to arbitrarily small neighborhoods of the form (5.5.2) of $\{\tilde{G}_{\boldsymbol{x}}^{k_0} : \boldsymbol{x} \in \mathcal{X}\}$.

Note that $\{\tilde{G}_{\boldsymbol{x}}^{k_0}, \boldsymbol{x} \in \mathcal{X}\} \in \mathcal{G}_x$. Let thus $B = \left([-a,a]^3 \times (\underline{\sigma},\bar{\sigma})\right)$ be the support of $\{\tilde{G}_{\boldsymbol{x}}^{k_0}, \boldsymbol{x} \in \mathcal{X}\}$. Choose $k > 3a + \bar{\sigma}$ such that

$$\int_{\mathcal{X}} \int_{|y|>k} f_0(y \mid \boldsymbol{x}) \left\{ \frac{|y| + 3a}{2\underline{\sigma}^2} \right\}^2 dy q(\boldsymbol{x}) d\boldsymbol{x} < \frac{\varepsilon}{2}$$

Take $V = \left\{\{G_{\boldsymbol{x}}^n \mid \boldsymbol{x} \in \mathcal{X}\} : \inf_{\boldsymbol{x}\in\mathcal{X}} G_{\boldsymbol{x}}^n(B) > \frac{\underline{\sigma}}{\bar{\sigma}}\right\}$. By approximating 1_B by a bounded continuous function, we can show for all n big enough that V contains a neighborhood V' of $\{\tilde{G}_{\boldsymbol{x}}^{k_0} \mid \boldsymbol{x} \in \mathcal{X}\}$

of the type (5.5.2). For any density $f \in \mathcal{F}_d, f(y \mid \boldsymbol{x}) = \int \frac{1}{\sigma}\phi(\frac{y-\bar{\boldsymbol{x}}'\boldsymbol{\theta}}{\sigma})dG^n_{\boldsymbol{x}}(\boldsymbol{\theta},\sigma), \boldsymbol{x} \in \mathcal{X}$, with $\{G^n_{\boldsymbol{x}} \mid \boldsymbol{x} \in \mathcal{X}\} \in V'$,

$$\begin{aligned}
&\int_{\mathcal{X}}\int_{|y|>k} f_0(y \mid \boldsymbol{x}) \log\left\{\frac{\tilde{f}(y \mid \boldsymbol{x})}{f(y \mid \boldsymbol{x})}\right\} dy\, q(\boldsymbol{x})\, d\boldsymbol{x} \\
&\leq \int_{\mathcal{X}}\int_{|y|>k} f_0(y \mid \boldsymbol{x}) \log \frac{\int_B \frac{1}{\sigma}\phi\left(\frac{y-\bar{\boldsymbol{x}}'\boldsymbol{\theta}}{\sigma}\right) d\tilde{G}^{k_0}_{\boldsymbol{x}}(\boldsymbol{\theta},\sigma)}{\int_B \frac{1}{\sigma}\phi\left(\frac{y-\bar{\boldsymbol{x}}'\boldsymbol{\theta}}{\sigma}\right) dG^n_{\boldsymbol{x}}(\boldsymbol{\theta},\sigma)}\, dy\, q(\boldsymbol{x})\, d\boldsymbol{x} \\
&\leq \int_{\mathcal{X}}\int_{|y|>k} f_0(y \mid \boldsymbol{x}) \log \frac{\frac{1}{\bar{\sigma}}\phi\left(\frac{|y|-3a}{\bar{\sigma}}\right)}{\frac{1}{\sigma}\phi\left(\frac{|y|+3a}{\sigma}\right) G^n_{\boldsymbol{x}}(B)}\, dy\, q(\boldsymbol{x})\, d\boldsymbol{x} \\
&\leq \int_{\mathcal{X}}\int_{|y|>k} f_0(y \mid \boldsymbol{x}) \left\{\frac{|y|+3a}{2\underline{\sigma}^2}\right\}^2 dy\, q(\boldsymbol{x})\, d\boldsymbol{x} < \frac{\varepsilon}{2}
\end{aligned}$$

Let $\inf_{\{y:|y|\leq k\}\times\mathcal{X}} \inf_{(\boldsymbol{\theta},\sigma)\in B} \frac{1}{\sigma}\phi\left(\frac{y-\bar{\boldsymbol{x}}'\boldsymbol{\theta}}{\sigma}\right) = c$. Consider the uniformly equicontinuous family of functions

$$\left\{g_{y,\boldsymbol{x}} : g_{y,\boldsymbol{x}} : \mathbb{R}^3 \times \mathbb{R}^+ \to \mathbb{R}, (\boldsymbol{\theta},\sigma) \mapsto \frac{1}{\sigma}\phi\left(\frac{y-\bar{\boldsymbol{x}}'\boldsymbol{\theta}}{\sigma}\right), (y,\boldsymbol{x}) \in [-k,k]\times\mathcal{X}\right\}$$

By the Arzela-Ascoli theorem, given $\delta > 0$, there exist finitely many points:

$$\{(y_i, \boldsymbol{x}_i) \in [-k,k]\times\mathcal{X}, i = 1,\ldots,m\}$$

such that for any $(y,\boldsymbol{x}) \in [-k,k]\times\mathcal{X}$, $\exists\ i$ such that

$$\sup_{(\boldsymbol{\theta},\sigma)\in B} \left|g_{y,\boldsymbol{x}}(\boldsymbol{\theta},\sigma) - g_{y_i,\boldsymbol{x}_i}(\boldsymbol{\theta},\sigma)\right| < c\delta$$

Let $g_{y_i,\boldsymbol{x}_i} * G^n_{\boldsymbol{x}} = \int \frac{1}{\sigma}\phi(\frac{y_i-\boldsymbol{x}_i'\boldsymbol{\theta}}{\sigma})\, dG^n_{\boldsymbol{x}}(\boldsymbol{\theta},\sigma)$ and consider:

$$E_n = \left\{\{G^n_{\boldsymbol{x}} \mid \boldsymbol{x} \in \mathcal{X}\} : \sup_{\boldsymbol{x}\in\mathcal{X}} \left|g_{y_i,\boldsymbol{x}_i} * G^n_{\boldsymbol{x}} - g_{y_i,\boldsymbol{x}_i} * \tilde{G}^{k_0}_{\boldsymbol{x}}\right| < c\delta, i = 1,2,\ldots,m\right\}$$

E_n is a neighborhood of $\{\tilde{G}^{k_0}_{\boldsymbol{x}} \mid \boldsymbol{x} \in \mathcal{X}\}$ formed by finite intersections of sets of the type (5.5.2), that has positive prior mass for all n big enough, as proved in (5.5.18). For $\{G^n_{\boldsymbol{x}} \mid \boldsymbol{x} \in \mathcal{X}\} \in E_n$ and $(y,\boldsymbol{x}) \in [-k,k]\times\mathcal{X}$, choosing the appropriate $(y_i,\boldsymbol{x}_i)$ and using a simple triangulation argument:

$$\left|\frac{\int \frac{1}{\sigma}\phi\left(\frac{y-\bar{\boldsymbol{x}}'\boldsymbol{\theta}}{\sigma}\right) dG^n_{\boldsymbol{x}}(\boldsymbol{\theta},\sigma)}{\int \frac{1}{\sigma}\phi\left(\frac{y-\bar{\boldsymbol{x}}'\boldsymbol{\theta}}{\sigma}\right) d\tilde{G}^{k_0}_{\boldsymbol{x}}(\boldsymbol{\theta},\sigma)} - 1\right| < 3\delta$$

and thus:

$$\left|\frac{\int \frac{1}{\sigma}\phi\left(\frac{y-\bar{\boldsymbol{x}}'\boldsymbol{\theta}}{\sigma}\right) d\tilde{G}^{k_0}_{\boldsymbol{x}}(\boldsymbol{\theta},\sigma)}{\int \frac{1}{\sigma}\phi\left(\frac{y-\bar{\boldsymbol{x}}'\boldsymbol{\theta}}{\sigma}\right) dG^n_{\boldsymbol{x}}(\boldsymbol{\theta},\sigma)} - 1\right| < \frac{3\delta}{1-3\delta}$$

for $\delta < \frac{1}{3}$. Thus given any $\varepsilon > 0$, there exist neighborhoods E_n of $\{\tilde{G}^{k_0}_{\boldsymbol{x}} \mid \boldsymbol{x} \in \mathcal{X}\}$, with positive prior mass for all n big enough, such that for $\{G^n_{\boldsymbol{x}} \mid \boldsymbol{x} \in \mathcal{X}\} \in E_n$ with $f(y \mid \boldsymbol{x}) = \int \frac{1}{\sigma}\phi(\frac{y-\boldsymbol{x}'\boldsymbol{\theta}}{\sigma})\, dG^n_{\boldsymbol{x}}(\boldsymbol{\theta},\sigma)$,

$$\int_{\mathcal{X}}\int_{y:|y|\leq k} f_0(y \mid \boldsymbol{x}) \log\left\{\frac{\tilde{f}(y \mid \boldsymbol{x})}{f(y \mid \boldsymbol{x})}\right\} dy\, q(\boldsymbol{x})\, d\boldsymbol{x} < \frac{\varepsilon}{2}$$

Taking $W_n = V' \cap E_n$ and since W_n is a finite intersection of neighborhoods of $\tilde{G}_{\boldsymbol{x}}$ of the type (5.5.2) for all n, the result follows. □

***Proof of Proposition** 5.3.1.* First we prove consistency in the q^a-integrated $\mathbb{L}_1$ metric defined in (5.5.5) of the densities $g(\cdot \mid a, Z)$. In other words we prove $\forall\, \varepsilon > 0$:

$$\Pi\left(W_\varepsilon^a(g_0; q) \mid \mathcal{D}^{(n)}\right) \to 1, \qquad P_0^{(n)} - \text{a.s.} \tag{5.5.23}$$

First note that:

$$\begin{aligned}\int \|g(\cdot \mid a, z) - g_0(\cdot \mid a, z)\|_1 \, q(z \mid a)\, dz &= \int\int |g(w \mid a, z) - g_0(w \mid a, z)| q(z \mid a)\, dz\, dw \\ &= \|g_0^a - g^a\|_1 \end{aligned} \tag{5.5.24}$$

We can therefore use Theorem 5.5.1. First note, that being a parametric model, the KL property, i.e. $g_0^a \in \text{KL}(\Pi)$, is automatically satisfied, as the prior has full support. We define sieves for the parameter space. Consider the prior for $g(\cdot \mid a, Z)$ given in (5.3.1), where $\boldsymbol{\theta}^w := (\theta_0^w, \theta_1^w, \theta_2^w) \sim N(\mu_{\theta^w}, \Sigma_w)$ with diagonal covariance matrix Σ_w and $\sigma_w \sim \text{InvGamma}(\gamma_1, \gamma_2)$. Construct a sieve for $(\boldsymbol{\theta}^w, \sigma_w)$, where $\phi_{\boldsymbol{\theta}^w, \sigma_w} = \frac{1}{\sigma_w}\phi(\frac{y - \tilde{\boldsymbol{x}}'\boldsymbol{\theta}^w}{\sigma_w})$, where $\boldsymbol{x} = (a, z)$:

$$\Theta_{a_n, h_n, l_n} = \{\phi_{\boldsymbol{\theta}^w, \sigma_w} \,:\, \|\boldsymbol{\theta}^w\| \leq a_n, l_n \leq \sigma_w \leq h_n\}. \tag{5.5.25}$$

we choose $a_n = O(\sqrt{n})$, $h_n = e^n$ and $l_n = O(1/\sqrt{n})$. Because from [101] for $\sigma_{2,w} > \sigma_{1,w} > \frac{\sigma_{2,w}}{2}$ and for each $\boldsymbol{x} \in \mathcal{X}$,

$$\int_{\mathcal{W}} \left|\phi_{\boldsymbol{\theta}_1^w, \sigma_{1,w}}(\boldsymbol{x}, w) - \phi_{\boldsymbol{\theta}_2^w, \sigma_{2,w}}(\boldsymbol{x}, w)\right| dw \leq \left(\frac{2}{\pi}\right)^{1/2} \frac{\left\|\boldsymbol{\theta}_2^w - \boldsymbol{\theta}_1^w\right\| \sqrt{3}}{\sigma_{2,w}} + \frac{3\,(\sigma_{2,w} - \sigma_{1,w})}{\sigma_{1,w}}$$

Using the same proof as in Lemma 4.1 in [101] we conclude that there exists constants $d_1, d_2 > 0$ such that: $N(\Theta_{a_n, h_n, l_n}, \varepsilon, \|\cdot\|_1) \leq \frac{d_1}{\varepsilon^2}\frac{a_n}{l_n} + \frac{d_2}{\varepsilon}\log\frac{h_n}{l_n} + 1$. Combined with the given choice of a_n, h_n, l_n satisfies the second condition of Theorem 5.5.1. For the first condition of Theorem 5.5.1, for constants C, C_1, C_2, C_3 and n big enough:

$$\Pi(\Theta^c_{a_n, h_n, l_n}) \leq \Pi(\|\boldsymbol{\theta}^w\| > C_1\sqrt{n})(\Pi(\sigma_w > C_2 e^n) + \Pi(\sigma_w < C_3/\sqrt{n})) \lesssim e^{-Cn}$$

where we used the classical tail bound for Gaussians and for the inverse gamma distribution. All this proves:

$$\Pi(g \,:\, \|g^a - g_0^a\|_1 > \varepsilon \mid \mathcal{D}^{(n)}) \to 0, \qquad P_0^{(n)} - \text{a.s.}$$

Because of the boundedness of $\mathcal{W}$, the above proof gives consistency of $\text{E}[W \mid A = a, Z]$ in $\mathbb{L}_1$, as:

$$\int \left|\int w(g_0(w \mid a, z) - g(w \mid a, z))\, dw\right| q(z \mid a)\, dz \lesssim \|g_0^a - g^a\|_1.$$

Using the same type of proof as above, with the classical Theorem 6.23 in [42], we obtain:

$$\Pi(g \,:\, \|g - g_0\|_1 > \varepsilon \mid \mathcal{D}^{(n)}) \to 0, \qquad P_0^{(n)} - \text{a.s.}$$

(note that these neighborhoods are now integrated with respect to the joint $q(a, z)$). This proves on the other hand consistency of $\mathrm{E}[W \mid Z] = \mathrm{E}[\mathrm{E}[W \mid A, Z] \mid Z]$, as using the boundedness of q:

$$\int \left| \int \int w(g_0(w \mid a, z) - g(w \mid a, z))\, dw\, q(a \mid z)\, da \right| q(z)\, dz \lesssim \|g_0 - g\|_1.$$

Finally, using the same technique used for (5.5.26) it is easy to prove, for fixed $a \in \mathcal{A}$, consistency of $g(w \mid a, z = 1)$ and of $g(w \mid a, z = 0)$ in the $\mathbb{L}_1$ metric, this implying consistency of θ_2^w. All the above combined, with the triangle inequality, imply the claim. □

***Proof of Theorem** 5.3.2.* We have that $f_0 \in \mathrm{KL}(\Pi)$. We can therefore turn to verifying the conditions of Theorem 5.5.1. Define the sieves for fixed $a \in \mathcal{A}$:

$$\mathcal{F}_n^a = \Big\{ f : f(y \mid a, z) = \sum_{k=1}^{\infty} w_k^n(a, z) \frac{1}{\sigma_k} \phi\left(\frac{y - (1, a, z)' \boldsymbol{\theta}_k}{\sigma_k} \right),$$
$$\{\phi_{\boldsymbol{\theta}_k, \sigma_k}\}_{k=1}^{m_n} \in \Theta_{a_n, h_n, l_n},\ \alpha_k^n \in B_{k,n}, k = 1, \ldots, m_n,\ \sup_{\boldsymbol{x} \in \mathcal{X}} \sum_{k \geq m_n + 1} w_k^n(\boldsymbol{x}) \leq \varepsilon \Big\},$$

where:

$$\Theta_{a_n, h_n, l_n} = \{\phi_{\boldsymbol{\theta}, \sigma} \ :\ \|\boldsymbol{\theta}\| \leq a_n, l_n \leq \sigma \leq h_n\}, \tag{5.5.26}$$

and for a sequence of partitions $\{T_{m,n}\}_{m=1}^{\infty}$ indexed by $n \in \mathbb{N}$:

$$B_{k,n} := \Big\{ \alpha_k^n(a, z) = \sum_{m=1}^{N_n} (\eta_{0,m}^k + \eta_{1,m}^k a + \eta_{2,m}^k z) \mathbf{1}_{(a,z) \in T_{m,n}} \ : \tag{5.5.27}$$
$$: \boldsymbol{\eta}_i^k = (\eta_{i,1}^k, \ldots, \eta_{i,N_n}^k),\ i \in \{0, 1, 2\},\ \|\boldsymbol{\eta}_i^k\|_1 \leq r_n \Big\}. \tag{5.5.28}$$

Let $\phi_{\boldsymbol{\theta}_k, \sigma_k}(a, z, y) = \frac{1}{\sigma_k} \phi\left(\frac{y - (1, a, z)' \boldsymbol{\theta}_k}{\sigma_k} \right)$ for $y \in \mathcal{Y}$ and $(a, z) \in \mathcal{A} \times \mathcal{Z}$. Without loss of generality assume $\mathcal{X} = [0, 1]^2$. From [101], we obtain for $\sigma_2 > \sigma_1 > \frac{\sigma_2}{2}$ and for each $(a, z) \in \mathcal{A} \times \mathcal{Z}$, for a constant $K \geq 2$:

$$\int_{\mathcal{Y}} \left| \phi_{\boldsymbol{\theta}_1, \sigma_1}(a, z, y) - \phi_{\boldsymbol{\theta}_2, \sigma_2}(a, z, y) \right| dy \leq \left(\frac{2}{\pi} \right)^{1/2} \frac{\|\boldsymbol{\theta}_2 - \boldsymbol{\theta}_1\|\, \sqrt{K}}{\sigma_2} + \frac{3\,(\sigma_2 - \sigma_1)}{\sigma_1}.$$

For $f_1, f_2 \in \mathcal{F}_n^a$ we have:

$$\begin{aligned}
\left\| f_1^a - f_2^a \right\|_1 &\leq \int_{\mathcal{Z}} \int_{\mathcal{Y}} \sum_{k=1}^{\infty} \left| {}_1 w_k^n(a, z) \phi_{{}_1\boldsymbol{\theta}_k, {}_1\sigma_k}(a, z, y) - {}_2 w_k^n(a, z) \phi_{{}_2\boldsymbol{\theta}_k, {}_2\sigma_k}(a, z, y) \right| dy\, q(z \mid a)\, dz \\
&\leq \int_{\mathcal{Z}} \int_{\mathcal{Y}} \sum_{k=1}^{m_n} {}_1 w_k^n(a, z) \left| \phi_{{}_1\boldsymbol{\theta}_k, {}_1\sigma_k}(a, z, y) - \phi_{{}_2\boldsymbol{\theta}_k, {}_2\sigma_k}(a, z, y) \right| dy\, q(z \mid a)\, dz \\
&+ \int_{\mathcal{Z}} \int_{\mathcal{Y}} \sum_{k=1}^{m_n} \left| {}_1 w_k^n(a, z) - {}_2 w_k^n(a, z) \right| \phi_{{}_2\boldsymbol{\theta}_k, {}_2\sigma_k}(a, z, y)\, dy\, q(z \mid a) dz \\
&+ \int_{\mathcal{Z}} \sum_{k=m_n+1}^{\infty} \left({}_1 w_k^n(a, z) + {}_2 w_k^n(a, z) \right) q(z \mid a)\, dz
\end{aligned}$$

$$\leq \int_{\mathcal{Z}} \sum_{k=1}^{m_n} {}_1w_k^n(a,z) \int_{\mathcal{Y}} \left|\phi_{{}_1\boldsymbol{\theta}_k,{}_1\sigma_k}(a,z,y) - \phi_{{}_2\boldsymbol{\theta}_k,{}_2\sigma_k}(a,z,y)\right| \, dy \, q(z \mid a) \, dz$$
$$+ \sum_{k=1}^{m_n} \left\| {}_1w_k^n - {}_2w_k^n \right\|_\infty + 2\varepsilon$$
$$\leq \sum_{k=1}^{m_n} \left(\frac{2}{\pi}\right)^{1/2} \frac{\|{}_1\boldsymbol{\theta}_k - {}_2\boldsymbol{\theta}_k\| \sqrt{K}}{{}_2\sigma_k} + \frac{3\,({}_2\sigma_k - {}_1\sigma_k)}{{}_1\sigma_k} + \sum_{k=1}^{m_n} \left\| {}_1w_k^n - {}_2w_k^n \right\|_\infty + 2\varepsilon.$$

It is clear that $\Theta_{a_n,h_n,l_n} \subset \{\phi_{\boldsymbol{\theta},\sigma} : \boldsymbol{\theta} \in (-a\sqrt{K}, a\sqrt{K}]^3, l_n \leq \sigma \leq h_n\}$. Let $\kappa < \min(\frac{\varepsilon}{6m_n}, 1)$ and $\sigma_j = l_n(1+\kappa)^j$, $j \geq 0$. Let j_0 be the smallest integer such that $\sigma_{j_0} = l_n(1+\kappa)^{j_0} > h_n$. This implies $j_0 \leq (1+\kappa)^{-1} \log \frac{h_n}{l_n} + 1$. By the choice of σ_j, $j \geq 1$

$$\frac{3\left(\sigma_j - \sigma_{j-1}\right)}{\sigma_{j-1}} < \frac{\varepsilon}{2m_n}.$$

Let $N_j = \left\lceil \left(\frac{32}{\pi}\right)^{1/2} \frac{a_n m_n K \sqrt{K}}{\sigma_{j-1}\varepsilon} \right\rceil$. For $1 \leq i \leq N_j$ and $1 \leq j \leq j_0$, define

$$E_{ij} = \left(-a_n\sqrt{K} + \frac{2a_n\sqrt{K}(i-1)}{N_j}, -a_n\sqrt{K} + \frac{2a_n\sqrt{K}i}{N_j} \right]^3 \times (\sigma_{j-1}, \sigma_j]$$

Clearly for $(\boldsymbol{\theta},\sigma), (\boldsymbol{\theta}',\sigma') \in E_{ij}$,

$$\left\|\phi_{\boldsymbol{\theta},\sigma} - \phi_{\boldsymbol{\theta}',\sigma'}\right\| \leq \left(\frac{2}{\pi}\right)^{1/2} \frac{\|\boldsymbol{\theta} - \boldsymbol{\theta}'\| \sqrt{K}}{\sigma'} + \frac{\varepsilon}{2m_n} \leq \left(\frac{2}{\pi}\right)^{1/2} \frac{2a_n K \sqrt{K}}{\sigma_{j-1} N_j} + \frac{\varepsilon}{2m_n} \leq \frac{\varepsilon}{m_n}$$

Thus, for constants $d_1, d_2 > 0$ to be determined at the end:

$$N\left(\varepsilon/m_n, \Theta_{a_n,h_n,l_n}, \|\cdot\|_1\right) \leq \sum_{j=1}^{j_0} \left\{ \left(\frac{32}{\pi}\right)^{1/2} \frac{a_n m_n K \sqrt{K}}{\sigma_{j-1}\varepsilon} + 1 \right\}^3$$
$$\leq \min\left(1, 2^2\right) \left[\left\{ \left(\frac{32}{\pi}\right)^{1/2} \frac{a_n m_n K \sqrt{K}}{\varepsilon} \right\}^3 \sum_{j=1}^{m_0} \left(\frac{1}{\sigma_{j-1}}\right)^p + j_0 \right]$$
$$\leq \min\left(1, 2^2\right) \left[\left\{ \left(\frac{32}{\pi}\right)^{1/2} \frac{a_n m_n K \sqrt{K}}{\varepsilon l_n} \right\}^3 \frac{1 - (1+\kappa)^{-3j_0}}{1 - (1+\kappa)^{-3}} \right.$$
$$\left. + (1+\kappa)^{-1} \log \frac{h_n}{l_n} + 1 \right] \leq d_1 \left(\frac{a_n m_n}{l_n \varepsilon}\right)^3 + d_2 \log \frac{h_n}{l_n} + 1.$$

Now we look at $\left\| {}_1w_k^n - {}_2w_k^n \right\|_\infty$ for $k = 1, \ldots, m_n$.

Because: $w_k^n(\boldsymbol{x}) = \Phi(\alpha_k^n(\boldsymbol{x})) \prod_{l<k} \left\{1 - \Phi(\alpha_l^n(\boldsymbol{x}))\right\}$ it is easy to see that:

$$\|{}_1w_k^n - {}_2w_k^n\|_\infty \leq \sum_{m=1}^{k} \|\Phi({}_1\alpha_m^n) - \Phi({}_2\alpha_m^n)\|_\infty,$$

by the Lipschitz continuity of Φ there exist a universal constant K_1 such that

$$\|\Phi({}_1\alpha_k^n) - \Phi({}_2\alpha_k^n)\|_\infty \leq K_2 \|{}_1\alpha_k^n - {}_2\alpha_k^n\|_\infty$$

$$= K_2 \sup_{a,z} \left| \sum_{m=1}^{N_n} \left(({}^1\eta^k_{0,m} - {}^2\eta^k_{0,m}) + ({}^1\eta^k_{1,m} - {}^2\eta^k_{1,m})a + ({}^1\eta^k_{2,m} - {}^2\eta^k_{2,m})z \right) \mathbf{1}_{(a,z)\in T_{m,n}} \right|$$

$$\lesssim \sum_{m=1}^{N_n} |{}^1\eta^k_{0,m} - {}^2\eta^k_{0,m}| + |{}^1\eta^k_{1,m} - {}^2\eta^k_{1,m}| + |{}^1\eta^k_{2,m} - {}^2\eta^k_{2,m}|.$$

by taking three $\varepsilon/(3km_n)$ nets on the set $B_{n,k}$ of vectors $\boldsymbol{\eta}^k_i = (\eta^k_{i,1}, \ldots, \eta^k_{i,N_n})$ for $i = 0, 1, 2$ we can bound $\sum_{k=1}^{m_n} \left\| {}_1w^n_k - {}_2w^n_k \right\|_\infty$ by ε. Note that the entropy is by Proposition C.2 in [42]:

$$N\left(\frac{\varepsilon}{3km_n}, \{\boldsymbol{\eta}^k_i = (\eta^k_{i,1}, \ldots, \eta^k_{i,N_n}) \in \mathbb{R}^{N_n},\ \|\boldsymbol{\eta}^k_i\|_1 \le r_n\}, \|\cdot\|_1 \right) \le \left(\frac{6km_nr_n}{\varepsilon} \right)^{N_n}.$$

We therefore conclude that:

$$\log N\left(\varepsilon, \mathcal{F}^a_n, \|\cdot\|_1\right) \le N_n \log\left(\frac{6(m_n+1)!r_n}{\varepsilon} \right) + m_n \log\left(d_1 \left(\frac{a_nm_n}{l_n\varepsilon} \right)^3 + d_2 \log \frac{h_n}{l_n} + 1 \right).$$

Now we verify condition 1 of Theorem 5.5.1. First note:

$$\Pi\left(\mathcal{F}^a \setminus \mathcal{F}^a_n\right)$$
$$\le \Pi\Big(\sup_{\boldsymbol{x}\in\mathcal{X}} \sum_{k\ge m_n+1} w^n_k(\boldsymbol{x}) \ge \varepsilon \Big) + \prod_{k=1}^{m_n} \Pi(\alpha^n_k \notin B_{k,n}) + m_nG_0\left(([-a_n, a_n]^3)^c \times [l_n, h_n]^c \right).$$

The last term is exponentially small by assumption. As $\forall\, k, i,\ \boldsymbol{\eta}^k_i$ is normally distributed, for all n big enough and some constant C:

$$\Pi(\alpha^n_k \notin B_{k,n}) \le \sum_{i=0}^{2} \Pi(\|\boldsymbol{\eta}^k_i\| > r_n) \le \sum_{i=0}^{2} \sum_{m=1}^{N_n} \Pi(|\eta^k_{i,m}| > r_n/N_n)$$
$$\le \sum_{i=0}^{2} \sum_{m=1}^{N_n} \exp\left\{ -\frac{(r_n/N_n - \mathrm{E}|\eta^k_{i,m}|)^2}{\mathrm{Var}(|\eta^k_{i,m}|)} \right\} \lesssim N_n e^{-C\frac{r_n^2}{N_n^2}}.$$

thus we conclude:

$$\prod_{k=1}^{m_n} \Pi(\alpha^n_k \notin B_{k,n}) \lesssim e^{-C^* m_n (\frac{r_n^2}{N_n^2} - \log N_n)}.$$

For $\tilde{\alpha}^n_k = \inf_{\boldsymbol{x}\in\mathcal{X}} \alpha^k_n(\boldsymbol{x})$ and $t_0 = -\log\varepsilon > 0$:

$$\Pi\Big(\sup_{\boldsymbol{x}\in\mathcal{X}} \sum_{k\ge m_n+1} w^n_k(\boldsymbol{x}) \ge \varepsilon \Big) = \Pi\Big(\sup_{\boldsymbol{x}\in\mathcal{X}} \prod_{k=1}^{m_n} (1 - \Phi(\alpha^n_k(\boldsymbol{x}))) > \varepsilon \Big)$$
$$\le \Pi\Big(\prod_{k=1}^{m_n} (1 - \Phi(\tilde{\alpha}^n_k)) > \varepsilon \Big) = \Pi\Big(-\sum_{k=1}^{m_n} \log(1 - \Phi(\tilde{\alpha}^n_k)) < t_0 \Big).$$

In what follows, without loss of generality assume that α^n_k is mean zero. Thus $-\alpha^n_k$ has the same normal distribution as α^n_k. Denote by $\lambda = -\Phi^{-1}\left(1 - e^{-t}\right)$ and for brevity denote the supremum norm by $\|\cdot\|$, $\mathbb{V}$ indicates the variance and C, C', C'', C''' positive constants. $\forall\, t \in (0, t_0)$:

$$\Pi\left(-\log\left(1 - \Phi\left(\tilde{\alpha}^n_k\right)\right) < t \right) = \Pi\left(\left(1 - \Phi\left(\tilde{\alpha}^n_k\right)\right) > e^{-t} \right)$$

$$
\begin{aligned}
&= \Pi\left(\Phi\left(\tilde{\alpha}_k^n\right) < 1 - e^{-t}\right) = \Pi\left(\inf_{\boldsymbol{x}\in\mathcal{X}} \alpha_n^k(\boldsymbol{x}) < \Phi^{-1}\left(1 - e^{-t}\right)\right) \\
&= \Pi\left(-\inf_{x\in\mathcal{X}} \alpha_k^n(x) > -\Phi^{-1}\left(1 - e^{-t}\right)\right) = \Pi\left(\sup_{\boldsymbol{x}\in\mathcal{X}} -\alpha_k^n(\boldsymbol{x}) > -\Phi^{-1}\left(1 - e^{-t}\right)\right) \\
&= \Pi\left(\sup_{x\in\mathcal{X}} \alpha_k^n(x) > -\Phi^{-1}\left(1 - e^{-t}\right)\right) \leqslant \Pi\left(\left\|\alpha_k^n\right\| > \lambda\right) \\
&\overset{(1)}{\leq} C\lambda^2\left(1 - \Phi\left(\frac{\lambda}{\|\mathbb{V}(\alpha_k^n)\|^{\frac{1}{2}}}\right)\right) \leq C\lambda^2 \frac{\|\mathbb{V}(\alpha_k^n)\|^{\frac{1}{2}}}{\sqrt{2\pi}\lambda} \exp\left\{-\frac{1}{2}\frac{\lambda^2}{\|\mathbb{V}(\alpha_k^n)\|}\right\} \\
&\leq C' \frac{\|\mathbb{V}(\alpha_k^n)\|^{\frac{1}{2}}}{\sqrt{2\pi}\lambda} \exp\left\{-\frac{1}{4}\frac{\lambda^2}{\|\mathbb{V}(\alpha_k^n)\|}\right\} \leq C''\left(1 - \Phi\left(\frac{\lambda}{\sqrt{2}\|\mathbb{V}(\alpha_k^n)\|^{\frac{1}{2}}}\right)\right) \\
&= C''\left(\Phi\left(\frac{\Phi^{-1}\left(1 - e^{-t}\right)}{\sqrt{2}\|\mathbb{V}(\alpha_k^n)\|^{\frac{1}{2}}}\right)\right) \leq C'''\left(1 - \exp\left\{-\gamma_0 t\right\}\right).
\end{aligned}
$$

We used in (1) Theorem 5.2 from [1], (2.1) of [42] and for any constant $c > 0$, all $\lambda \geq 0$ and big enough $K \geq 2$: $\frac{\lambda}{c} \leq K e^{\frac{1}{4}\frac{\lambda^2}{c^2}}/(\frac{\lambda}{2c} + \frac{2c}{\lambda})$. Furthermore we used that for all $\lambda \geq 0$, $\phi(\lambda)/(\lambda + \lambda^{-1}) \leq 1 - \Phi(\lambda)$, where ϕ is the density of the standard normal, that Φ is continuous and monotone increasing and that $\frac{1}{c}\Phi^{-1}(1 - e^{-t}) \leq \Phi^{-1}(1 - e^{-\max\{1/c,1\}t})$ for $t > 0$, together with $0 < \min_{1\leq k\leq m_n} \|\mathbb{V}(\alpha_k^n)\| \leq \max_{1\leq k\leq m_n} \|\mathbb{V}(\alpha_k^n)\| < \infty$ by construction of the prior.

We proved that for all $t \in (0, t_0)$ and a constant C''':

$$
\Pi\left(-\log\left(1 - \Phi\left(\tilde{\alpha}_k^n\right)\right) < t\right) \leq C'''\left(1 - \exp\left\{-\gamma_0 t\right\}\right). \tag{5.5.29}
$$

Using Lemma E.1 from [81] which states that for non-negative r.v.s. A_i, B_i, if $\Pi(A_i \leq u) \leq C_i\Pi(B_i \leq u)$ for $u \in (0, t_0), t_0 > 0, i = 1, 2, \Pi(A_1 + A_2 \leq t_0) \leq C_1C_2\Pi(B_1 + B_2 \leq t_0)$. Let γ_0 and C_* be the maximal constants such that (5.5.29) holds for all $1 \leq k \leq m_n$. Thus by induction, for $Z_k \sim \mathrm{Ga}(1, \gamma_0)$:

$$
\begin{aligned}
&\Pi\Big(-\sum_{k=1}^{m_n} \log(1 - \Phi(\tilde{\alpha}_k^n)) < t_0\Big) \leq C_*^{m_n}\Pi\Big(\sum_{k=1}^{m_n} Z_k < t_0\Big) \\
&= C_*^{m_n}\int_0^{t_0} \frac{1}{(m_n - 1)!} r^{m_n - 1}(\gamma_0)^{m_n} e^{-\gamma_0 r} dr \leq \frac{(C_* t_0 \gamma_0)^{m_n}}{m_n!} \leq \left(\frac{C_* e\, t_0 \gamma_0}{m_n}\right)^{m_n}
\end{aligned}
$$

The last inequality follows, since $x^N/N! \leq e^x$, for every $x > 0$, including $x = N$. The last expression can be bounded by $e^{-(m_n/2)\log m_n}$.

By choosing $a_n = \sqrt{n}$, $h_n = \sqrt{n}$, $l_n = 1/\sqrt{n}$, $m_n = n/\log n$, $N_n = \sqrt{\log n}$ and $r_n = m_n^{m_n/N_n}/(m_n + 1)!$ we conclude for constants $C > 0, \xi < 1/2$, and sufficiently large n,

$$
\Pi(\mathcal{F}^a \setminus \mathcal{F}_n^a) \leq e^{-Cn}, \quad \log N(\varepsilon, \mathcal{F}_n^a, \|\cdot\|_1) \leq \xi n\varepsilon^2.
$$

□

Proof of Corollary 5.3.1. Fix $a \in \mathcal{A}$ and $\varepsilon > 0$. Let $m_Y(a, z) := \int y f(y \mid a, z)\, dy$ and $m_Y^0(a, z) := \int y f_0(y \mid a, z)\, dy$ denote the posterior and true conditional means of $Y \mid A = a, Z = z$, respectively. Define $m_W(a, z) := \theta_0^w + \theta_1^w a + \theta_2^w z$ and $m_W^0(a, z) := \theta_0^{w,0} + \theta_1^{w,0} a + \theta_2^{w,0} z$. By

Proposition 5.2.2, the true ATE admits the representation

$$\mathrm{E}_0 Y(a) = \underbrace{\int m_Y^0(a,z)\, q(z \mid a)\, dz}_{T_1^0} + \underbrace{\iint \gamma_0(a,z)\{m_W^0(a',z) - m_W^0(a,z)\}\, q(a',z)\, da'\, dz}_{T_2^0},$$

where $\gamma_0(a,z) = \{b_1^0(a) + b_2^0(a)(z - a\theta_1^{w,0}/\theta_2^{w,0})\}/\theta_2^{w,0}$, with

$$(b_0^0, b_1^0, b_2^0)^\top = G_a^{-1}\mathrm{E}_0[m_Y^0(a,Z)(1,Z,Z^2)^\top \mid A = a].$$

Since $Q = Q_0$ is known, the posterior estimator of $\mathrm{E}_0 Y(a)$ is the same functional evaluated at the posterior parameters, so the error satisfies $|\mathrm{E}_0 Y(a) - \mathrm{E}Y(a)| \le |T_1 - T_1^0| + |T_2 - T_2^0|$.

Step 1: Consistency of T_1. For any $M > 0$ and any f with $C_f := \sup_{(a,z)\in\mathcal{X}} \int y^2 f(y \mid a, z)\, dy \le C$, split the integrand at $|y| \le M$:

$$\int \left| \int y(f - f_0)(y \mid a, z)\, dy \right| q(z \mid a)\, dz \le M \, \|f^a - f_0^a\|_1 + \frac{C + C_0}{M},$$

where $C_0 := \sup_{(a,z)\in\mathcal{X}} \int y^2 f_0(y \mid a, z)\, dy < \infty$ (which is finite under the assumptions of Proposition 5.2.2 and compactness of $\mathcal{X}$). The first inequality uses $|\int_{|y|\le M} y(f - f_0)\, dy| \le M \int |f - f_0|\, dy$; the second uses $\int_{|y|>M} |y| g\, dy \le M^{-1} \int y^2 g\, dy$ for $g \in \{f, f_0\}$. Choosing $M = M(\varepsilon) := (C + C_0)^{1/2}\varepsilon^{-1/2}$, the right-hand side is at most $M\|f^a - f_0^a\|_1 + \varepsilon^{1/2}(C + C_0)^{1/2}$. Hence, on the event $\{\|f^a - f_0^a\|_1 \le \delta\} \cap \{C_f \le C\}$ with $\delta := \varepsilon M^{-1}$:

$$|T_1 - T_1^0| \le 2\varepsilon^{1/2}(C + C_0)^{1/2}.$$

Since both $\Pi_n(\|f^a - f_0^a\|_1 > \delta \mid \mathcal{D}^{(n)}) \to 0$ (Theorem 5.3.2) and $\Pi_n(C_f > C \mid \mathcal{D}^{(n)}) \to 0$ (Assumption C) $P_0^{(n)}$-a.s., we conclude $\Pi_n(|T_1 - T_1^0| > \varepsilon \mid \mathcal{D}^{(n)}) \to 0$ $P_0^{(n)}$-a.s.

Step 2: Consistency of $\gamma(a,\cdot)$. Since Q_a is known and $G_a = \mathrm{E}_0[\phi(Z)\phi(Z)^\top \mid A = a]$ (with $\phi(Z) = (1, Z, Z^2)^\top$) is invertible by Proposition 5.2.2(ii), the projection coefficients satisfy

$$|b_j(a) - b_j^0(a)| \le \|G_a^{-1}\|_F \cdot C_{\mathcal{Z}}^j \cdot \int |m_Y(a,z) - m_Y^0(a,z)|\, q(z \mid a)\, dz, \qquad j = 0, 1, 2,$$

where $C_{\mathcal{Z}} := \sup_{z\in\mathcal{Z}} |z| < \infty$ uses compactness of $\mathcal{Z}$. By Step 1, the right-hand side converges to 0 in posterior probability. Combined with the consistency of (θ_1^w, θ_2^w) and the lower bound $|\theta_2^{w,0}| > 0$ from Proposition 5.3.1, continuous mapping gives

$$\sup_{z\in\mathcal{Z}} |\gamma(a,z) - \gamma_0(a,z)| \to 0 \quad \text{in posterior probability, } P_0^{(n)}\text{-a.s.}$$

Step 3: Consistency of T_2. Decompose $T_2 - T_2^0$ into three terms via $\gamma = \gamma_0 + (\gamma - \gamma_0)$ and $m_W = m_W^0 + (m_W - m_W^0)$:

$$|T_2 - T_2^0| \le \underbrace{\sup_z |\gamma(a,z) - \gamma_0(a,z)|}_{\to 0,\ \text{Step 2}} \cdot \underbrace{\iint |m_W^0(a',z) - m_W^0(a,z)|\, q(a',z)\, da'\, dz}_{\text{bounded, } \mathcal{X} \text{ compact}}$$

$$+ \underbrace{\sup_z |\gamma_0(a,z)|}_{\text{bounded}} \cdot \Big(\underbrace{\iint |m_W(a',z) - m_W^0(a',z)|\, q(a',z)\, da'\, dz}_{\to 0,\ \text{Prop. 5.3.1}}$$

$$+ \underbrace{\int |m_W(a,z) - m_W^0(a,z)|\, q(z \mid a)\, dz}_{\to 0,\ \text{Prop. 5.3.1}} \Big)$$

$$+ \underbrace{\sup_z |\gamma(a,z) - \gamma_0(a,z)|}_{\to 0} \cdot \underbrace{\iint |m_W(a',z) - m_W^0(a,z)|\, q(a',z)\, da'\, dz}_{\text{bounded} \to 0}.$$

Each product converges to 0 in posterior probability $P_0^{(n)}$-a.s.

Conclusion. By the triangle inequality and Steps 1–3, for any $\varepsilon > 0$, $P_0^{(n)}$-a.s.

$$\Pi_n(|\mathrm{E}_0 Y(a) - \mathrm{E} Y(a)| > \varepsilon \mid \mathcal{D}^{(n)}) \le \Pi_n(|T_1 - T_1^0| > \varepsilon/2 \mid \mathcal{D}^{(n)}) + \Pi_n(|T_2 - T_2^0| > \varepsilon/2 \mid \mathcal{D}^{(n)}) \to 0$$

□

5.5.3 Consistency by entropy

In this section we give a straightforward adaptation of Theorem 6.23 in [42]. The only difference with the general result given in [42] is that we are concerned with conditional laws at a fixed value in the conditioning. For fixed $a \in \mathcal{A}$ and $(a,z) \mapsto q(z \mid a)$ the density of the underlying probability measure associated to the law of $Z \mid A$, for any measurable set B:

$$P^a(B) = \int_B p(u \mid a,z) q(z \mid a)\, d(u,z), \tag{5.5.30}$$

It follows that $p^a(u,z \mid a) = p(u \mid a,z) q(z \mid a)$. Define the *conditional for fixed* $a \in \mathcal{A}$ Hellinger distance:

$$h(r^a, p^a) = \int (\sqrt{r(u \mid a,z)} - \sqrt{p(u \mid a,z)})^2\, q(z \mid a)\, d(u,z). \tag{5.5.31}$$

The *conditional for fixed* $a \in \mathcal{A}$ Hellinger affinity:

$$\rho_{1/2}(r^a, p^a) = \int \sqrt{r(u \mid a,z) p(u \mid a,z)}\, q(z \mid a)\, d(u,z) \tag{5.5.32}$$

then: $h^2(r^a, p^a) = 2 - 2\rho_{1/2}(r^a, p^a)$ and also: $\|p^a - r^a\|_1 \le 2h(p^a, q^a)$. The minimax risk for testing a probability P^a versus a set $\mathcal{R}^a$ of probability measures is defined by

$$\pi(P^a, \mathcal{R}^a) = \inf_\phi (P^a \phi + \sup_{R^a \in \mathcal{R}^a} R^a (1 - \phi))$$

where the infimum is taken over all tests, i.e. measurable functions $\phi : \mathfrak{X} \to [0,1]$. As a consequence of Proposition D.1 and Lemma D.6 in [42] adapted to the current notation, we obtain for every convex set of measures $\mathcal{R}^a$:

$$\pi((P^a)^n, (\mathcal{R}^a)^n) \le \rho_{1/2}((P^a)^n, \mathrm{conv}((\mathcal{R}^a)^n)) \le \rho_{1/2}(P^a, \mathcal{R}^a)^n \tag{5.5.33}$$

Thus any convex set $\mathcal{R}^a$ with Hellinger affinity to P^a smaller than 1 can be tested with error probabilities that decrease exponentially in the number of observations. For any probability measure P^a and convex, dominated set of probability measures $\mathcal{R}^a$ with $h(p^a, r^a) > \varepsilon$ for every $r^a \in \mathcal{R}^a$ and any $n \in \mathbb{N}$, there exists a test ϕ such that

$$(P^a)^n\phi \leq e^{-n\varepsilon^2/2}, \quad \sup_{R^a \in \mathcal{R}^a} (R^a)^n(1-\phi) \leq e^{-n\varepsilon^2/2} \tag{5.5.34}$$

Indeed we have $\rho_{1/2}(p^a, r^a) = 1 - h^2(p^a, r^a)/2$, which is bounded above by $1 - \varepsilon^2/2$. Combined with (5.5.33) we see that $\pi\left((P^a)^n, (\mathcal{R}^a)^n\right) \leq \left(1 - \varepsilon^2/2\right)^n \leq e^{-n\varepsilon^2/2}$, since $1 - x \leq e^{-x}$, for every x.

Given a semimetric d on a set $\mathcal{R}$ and $\varepsilon > 0$, the covering number $N(\varepsilon, \mathcal{R}, d)$ is defined as the minimal number of balls of radius ε needed to cover $\mathcal{R}$. Then it is possible to extend the result of (5.5.34) to non-convex alternatives (as the ones generated using the Hellinger distance) leveraging the concept of covering numbers. Using the same proof or Proposition D.8 in [42], for every $\varepsilon > 0$ and n there exists a test ϕ such that

$$(P^a)^n\phi \leq N(\varepsilon/4, \mathcal{R}^a, h)e^{-n\varepsilon^2/8}, \quad \sup_{R^a \in \mathcal{R}^a : h(p^a, r^a) > \varepsilon} (R^a)^n(1-\phi) \leq e^{-n\varepsilon^2/8}. \tag{5.5.35}$$

Theorem 5.5.1. *The posterior distribution is strongly consistent relative to the $\mathbb{L}_1$-distance at every $p_0^a \in \mathrm{KL}(\Pi)$ if for every $\varepsilon > 0$ there exist measurable sets $\mathcal{P}_n^a \subset \mathcal{P}^a$ (which may depend on ε) such that, for constants $C > 0, \xi < 1/2$, and sufficiently large n,*

$$\Pi\left(\mathcal{P}^a \backslash \mathcal{P}_n^a\right) \leq e^{-Cn}, \quad \log N\left(\varepsilon, \mathcal{P}_n^a, \|\cdot\|_1\right) \leq \xi n\varepsilon^2$$

Proof. Because the $\mathbb{L}_1$-distance is bounded above by twice the Hellinger distance, we can use (5.5.35) as $N\left(\varepsilon/2, \mathcal{P}_n^a, d\right) \leq N\left(\varepsilon, \mathcal{P}_n^a, \|\cdot\|_1\right) \leq e^{\xi n\varepsilon^2}$. Thus there exists a test ϕ_n with

$$(P_0^a)^n\phi_n \leq e^{\xi n\varepsilon^2}e^{-n\varepsilon^2/2}, \quad \sup_{p^a \in \mathcal{P}_n^a : \left\|p^a - p_0^a\right\|_1 > 4\varepsilon} (P^a)^n\left(1-\phi_n\right) \leq e^{-n\varepsilon^2/2}$$

Thus the conditions of the extension of Schwartz's theorem (Theorem 6.17 in [42]) are satisfied, where we take $\mathcal{U}$ the ball of radius 5ε around p_0^a and set the constant C equal to the minimum of $(1/2 - \xi)\varepsilon^2/2$ and the present constant C. □

Summary

In many scientific problems, the quantity one wants to estimate is not directly observable. Instead, the data is a noisy, indirect image of it, produced by a known forward transformation. Recovering the unknown from the observed data requires inverting that transformation: a statistical inverse problem. A canonical example is Wicksell's stereological problem: spherical particles of unknown size are embedded in an opaque material, and the only observable data are the radii of the circular cross-sections left by a cutting plane, which are a known geometric transform of the true sphere radii. The same mathematical structure appears in astronomy, deconvolution problems in epidemiology, and, in a different guise, in causal inference with unmeasured confounders.

Two difficulties make these problems hard. The first is *ill-posedness*: the forward operator may smooth the data, so inverting it amplifies estimation errors; naive unconstrained estimators are erratic and often have infinite variance. Exploiting *shape constraints* like monotonicity, in some of the problems treated here, is the natural regularisation. The second difficulty is *uncertainty quantification* (UQ). Even after constructing a well-behaved estimator, building valid confidence sets is non-trivial: the limiting distributions depend on unknown local properties of the distribution F_0 of the unknown (the density of the observations $g_0(x)$ and the local Hölder smoothness γ of F_0), which are themselves hard to estimate. This makes naive plug-in confidence intervals unreliable.

The main approach developed in this thesis is to construct Bayesian nonparametric posteriors and then isotonise the posterior draws — projecting each draw onto the set of monotone functions via the least concave majorant. The result, the *Isotonized Inverse Posterior* (IIP), is computationally inexpensive and, as we show, automatically valid for UQ without requiring the estimation of any nuisance parameter.

Chapter 2 studies Wicksell's problem from a frequentist perspective. We prove that the *isotonic inverse estimator* (IIE) — the right-derivative of the least concave majorant of a naive empirical functional — adapts automatically to the local Hölder smoothness $\gamma > 1/2$ of F_0 at the estimation point, without knowing γ. We also prove a matching minimax lower bound, showing that no estimator can do better. The IIE is therefore the unique efficient nonparametric estimator in this problem. When F_0 is locally flat, the IIE even attains the faster $\sqrt{n}$ rate automatically.

Chapter 3 constructs the Bayesian counterpart via a Dirichlet Process prior on the observable distribution and isotonisation of the posterior. We prove semiparametric Bernstein-von Mises theorems for the resulting IIP in both Wicksell's problem and the deconvolution problem on the half-line. The credible sets derived from the IIP have asymptotically correct frequentist coverage at the optimal rate $\sqrt{n}/\sqrt{\log n}$, adapting to γ without estimating it.

Chapter 4 extends the IIP methodology to general half-line deconvolution, where the forward operator is a convolution with a known monotone kernel. We derive both a Bernstein-von

Mises theorem for the IIP and an asymptotic distribution result for the frequentist isotonized estimator, providing the first nuisance-parameter-free UQ results for this class of problems.

Chapter 5 turns to causal inference. We introduce a causal model for proximal inference based on mixtures of linear models for the regression of the outcome on an unmeasured confounder. Under this model the bridge function, which links observed proxies to the causal effect, exists and has a closed-form expression, making the average treatment effect identifiable in a richer class of settings than was previously known. We propose a Bayesian nonparametric estimator based on a probit stick-breaking process and prove it is consistent.

Samenvatting

In veel wetenschappelijke problemen is de grootheid die men wil schatten niet direct waarneembaar. In plaats daarvan zijn de gegevens een ruisig, indirect beeld daarvan, geproduceerd door een bekende voorwaartse transformatie. Het terugwinnen van de onbekende grootheid uit de waargenomen gegevens vereist het omdraaien van die transformatie: een statistisch invers probleem. Een kanoniek voorbeeld is Wicksells stereologische probleem: bolvormige deeltjes van onbekende grootte zijn ingebed in een ondoorzichtig materiaal. De enige waarneembare gegevens zijn de stralen van de cirkelvormige doorsneden die door een snijvlak worden achtergelaten, welke een bekende meetkundige transformatie zijn van de werkelijke bolstralen. Dezelfde wiskundige structuur komt voor in de astronomie, deconvolutieproblemen in de epidemiologie, en, in een andere gedaante, in causale gevolgtrekking met ongemeten verstorende variabelen.

Twee moeilijkheden maken deze problemen lastig. De eerste is *slecht-gesteldheid*: de voorwaartse operator kan de gegevens afvlakken, waardoor het inverteren ervan schattingsfouten versterkt; naïeve onbeperkte schatters zijn grillig en hebben vaak oneindige variantie. Het gebruikmaken van *vormrestricties* zoals monotoniciteit, in sommige van de hier behandelde problemen, is de natuurlijke regularisatie. De tweede moeilijkheid is *onzekerheidskwantificering* (UQ). Zelfs na het construeren van een goed-gedragende schatter is het bouwen van geldige betrouwbaarheidsgebieden niet triviaal: de limietverdelingen hangen af van onbekende lokale eigenschappen van de verdeling F_0 (de dichtheid van de waarnemingen $g_0(x)$ en de lokale Hölder-gladheid γ van F_0), die op zichzelf moeilijk te schatten zijn. Dit maakt naïeve insteek-betrouwbaarheidsintervallen onbetrouwbaar.

De voornaamste aanpak die in dit proefschrift wordt ontwikkeld, is het construeren van Bayesiaanse niet-parametrische posteriorverdelingen en vervolgens het isotoniseren van de posteriortrekkingen — door elke trekking te projecteren op de verzameling van monotone functies via de kleinste concave majorant. Het resultaat, de *Geïsotoniseerde Inverse Posterior* (IIP), is computationeel goedkoop en, zoals wij aantonen, automatisch geldig voor onzekerheidskwantificering zonder dat het schatten van enige parameter vereist is.

Hoofdstuk 2 bestudeert Wicksells probleem vanuit een frequentistisch perspectief. Wij bewijzen dat de *isotonische inverse schatter* (IIE) — de rechter afgeleide van de kleinste concave majorant van een naïeve empirische schatter — zich automatisch aanpast aan de lokale Hölder-gladheid $\gamma > 1/2$ van F_0 in het schattingspunt, zonder γ te kennen. Wij bewijzen ook een overeenkomende minimax-ondergrens, waaruit blijkt dat geen enkele schatter het beter kan doen. De IIE is daarmee de uniek efficiënte niet-parametrische schatter in dit probleem. Wanneer F_0 lokaal vlak is, bereikt de IIE zelfs automatisch de snellere $\sqrt{n}$-snelheid.

Hoofdstuk 3 construeert het Bayesiaanse analogon via een Dirichlet-procesprior op de waarneembare verdeling en isotonisering van de posterior. Wij bewijzen semiparametrische Bernstein-von Mises-stellingen voor de resulterende IIP in zowel Wicksells probleem als het

deconvolutieprobleem op de halve rechte lijn. De geloofwaardigheidsverzamelingen afgeleid uit de IIP hebben asymptotisch correcte frequentistische dekking op de optimale snelheid $\sqrt{n}/\sqrt{\log n}$, met aanpassing aan γ zonder deze te schatten.

Hoofdstuk 4 breidt de IIP-methodologie uit naar algemene deconvolutie op de halve rechte lijn, waarbij de voorwaartse operator een convolutie is met een bekende monotone kern. Wij leiden zowel een Bernstein-von Mises-stelling voor de IIP als een asymptotisch verdelingsresultaat voor de frequentistische geïsotoniseerde schatter af, en leveren daarmee de eerste resultaten voor onzekerheidskwantificering zonder parameter voor deze klasse van problemen.

Hoofdstuk 5 richt zich op causale gevolgtrekking. Wij introduceren een causaal model voor proximale inferentie gebaseerd op mengsels van lineaire modellen voor de regressie van de uitkomst op een ongemeten verstorende variabele. Onder dit model bestaat de brugfunctie, die waargenomen proxies koppelt aan het causale effect, en heeft zij een expliciete uitdrukking, waardoor het gemiddelde behandelingseffect identificeerbaar is in een rijkere klasse van situaties dan voorheen bekend was. Wij stellen een Bayesiaanse niet-parametrische schatter voor gebaseerd op een probit-stokbreekproces en bewijzen dat deze consistent is.

Acknowledgements

The first ones I want to thank are Aad and Geurt. Obviously, I want to thank you for being great coauthors and statisticians. But also, and especially, for the following reasons: Aad, thank you for your incredibly deep knowledge, for your work ethic, for your intuition and attention to detail, for your patience, for your intellectual honesty, and ultimately for your ability to teach by *walking the way yourself* and not just by *showing the way*. Geurt, thank you for your passion, for your astonishing knowledge of isotonic methods, for your genuine interest in everything that happens around you, for your work ethic, for your optimism, for your energy, for your intellectual honesty, for your human side, and for your *Buongiornissimo* in the morning; I will miss it.

During these years I met a lot of extraordinary people at TU Delft, some of whom became very good friends. Ardjen, thank you for all the time we spent together, in person and over video calls, for all the laughs, for your interesting views about the world, for being the linear-regression-king, for being brothers in arms and coming back victorious (*you know what I mean*), and ultimately for being a great friend and the very best cow in the world. Matthias and Francesco, thank you for your friendship. Francesco, thank you for your generosity, for your always-surprising points of view, for your humour, for your openness, for your hard work, and for all the fun we had together; Matthias, thank you for your rationality, for your genuine optimism, for your healthy outlook on the world around you, for your serenity, and obviously for all the fun we had together! I also want to thank Thorben for all the support and for all the climbing sessions, thank you Lasse for your inspiring passion for mathematics and all the laughs we had in the early days of my PhD. Thank you Wieger for being a truly kind and good-hearted person – the world needs more people like you. Thank you Koen for teaching me something about football – I still think I am a lost cause in that sense. Thank you Serena and Matteo for the good times together in Delft, it was fun. Thank you Andreas for your Greek strength, and thank you Andrea for your Italian strength. Thank you Thomas for being a great office mate. Thank you Julia for the vivid, first-hand demonstration that people can disappear from one day to the next.

Outside TU Delft, I also met great people. First of all, thanks to all the great colleagues at WEBB Traders; thanks to Oscar and Armand for the good work and for the fun we had together, thank you to Dennis and Jesus for the planking sessions, thank you to Azar for the good conversations, and thank you to Jop and Mo for the opportunity and everything I learnt. Many thanks to Martine and Pelagia for your kindness. From my time with G-Research in France, I kept one friend: Felix, you are a great and smart guy. From the many conferences I attended, I kept one non-mathematical friendship: thank you so much, Charlotte, for the fun we had together and for your good heart. From my time in Amsterdam during my master's, I kept two good friends, Diogo and Rapha, whom I want to thank for all the good times together. Diogo, you are an extremely smart and kind person; I am thankful to have you in my life, and

happy you are back in the Netherlands. Looking forward to more adventures together. Rapha, thank you for your multifaceted knowledge and for all the conversations ranging from music to literature and mathematics. At Uber, I want to thank Pablo for your kindness, good heart, and for welcoming me into your great team; it has been such a good journey working together these past months. I am also very thankful to my teammates Vicente and Antoine for their incredible knowledge and for all the good times we shared. From the people back in Turin, I especially want to thank Busso for being the goat, Edo for his passion for the mountains, and Franchino (Lollo) for all the good times we spent together back in the day. Many thanks to Renzo too, for teaching me philosophy, the importance of pondering every thought, and the beauty of classical art and thinking. My love for maths, logic, and understanding the underlying truth is rooted in our conversations. Outside work and study circles, I want to thank Simone. That short conversation in a bouldering gym back in 2023 gave birth to one of the best climbing teams the Netherlands has ever seen, haha! We spent so much time together climbing, and sharing thoughts and views on all sorts of topics. I am truly grateful for your friendship and your always surprising and unconventional way of thinking about the world. I regret that I have recently had little time to spend with you since you have been travelling, but because you are Goldmund (and I suppose I am Narcissus?), I will always be there for you whenever you come back.

Finally, I would like to thank the most important people in my life. First of all, I want to thank Stefano. We have known each other for around 20 years now, and we have walked very similar paths in life up until this point; I would not be the person I am today without you. Thank you for all the good and bad times we went through together, for all your wise words and all your stupid jokes, and for the universe of references that we share, which make every sentence either a punch line or a philosophical dispute. Thank you for inspiring me to always strive to be a better version of who I am. You are the best. I hope these 20 years are just the beginning. Thank you, Matias, for being the greatest alpinist I know and for teaching me the beauty of literature and art and the pleasure of knowledge, for showing me how to look at the world around me with different eyes, ready to capture the beauty of the unexpected and the deeper meaning of things. I am looking forward to seeing you, Amparo, Diego, and Matias again. Evy, no words in any language I know are enough to express all the gratitude and happiness I feel when I think of you. It is not an exaggeration to say that you made my life in the Netherlands worth living. We have been through so many things together over the last 3 years, and we have been to so many places too (3 continents!). I want to thank you for your love, for your trust, for your serene way of looking at things, for your optimism, for your patience, for your perseverance, for your rationality, for all the laughs: in short, for being exactly what I needed and need. I also want to thank Vera and Tim for all the festivals and trips together, Iris and David for all the fun times, and Mirjam and Wil for your good hearts and patience with my choice of discussion topics. Finally, I want to thank my family, because I owe them everything I have accomplished and will accomplish in my life. Thank you to Marco and Annalisa for all the great childhood memories, thank you zia Nucci and zio Daniele for being so caring and always encouraging me, thank you zio Robi for all the laughs and Dave Gahan's songs. Thank you nonna for teaching me what it means to be Piemontese and for all the times you took care of me and my sister when we were kids. Thank you to my parents for all their sacrifices, for all the hard work they put into raising me and my sister, for teaching us the value of hard work, for always being there when we needed them most, for teaching us strong moral values without the need for a religious pill, for showing us the value of well-crafted objects and ideas, for teaching us that good stuff takes time, for teaching us the value of loyalty and protecting the ones we love, for bringing us to the mountains and showing

us that pain and sweat are not a reason to give up and thus the value of things that are hard to get, for showing us that taking care of the body is as important as taking care of the mind (Anima Sana In Corpore Sano), for giving us the intellectual tools to always be critical of the world around us, for teaching us the value of humility, for giving us, early on, the chance to be fully independent (unlike many other Italian fellows...), and ultimately for all the love they gave us.

Curriculum Vitæ

Francesco Gili f.gili@tudelft.nl | www.francescogili.nl

Education

2022–2026	PhD in Mathematical Statistics, Delft University of Technology
2020–2022	MSc Stochastics and Financial Mathematics, University of Amsterdam, *cum laude* (GPA 9.03/10)
2017–2020	BSc Mathematics for Finance and Insurance, University of Turin, *cum laude* (GPA 29/30)

Professional experience

2026 Sep – present	Data Scientist I, Ops, Uber
2026 Feb – Jul	Quant Analyst (Trainee), WEBB Traders, Amsterdam
2025 Apr	Spring Quant Program (SiQF25), G-Research, Nice
2022 Jan – Aug	Machine Learning Group Intern, CWI, Amsterdam
2021 Jul – Sep	Data Scientist Intern, Alliander, Amsterdam

Teaching

Lecturer and teaching assistant at TU Delft (MSc/BSc level) in Advanced Methods in Statistics, Causality, and Probability. Teaching assistant at the University of Amsterdam in Stochastic Processes for Finance and Stochastic Integration. Teacher at Digital Math Training Project (Maple) at University of Turin.

Languages

Italian (native), English (C1), Dutch (B2), Spanish (C1).

List of Publications

1. **F. Gili**, G. Jongbloed, A. van der Vaart. *Adaptive and efficient isotonic estimation in Wicksell's problem. Journal of Nonparametric Statistics* **37**(3), 491–531, 2025.

2. **F. Gili**, G. Jongbloed, A. van der Vaart. *Semiparametric Bernstein–von Mises phenomenon via Isotonized Posterior in Wicksell's problem. The Annals of Statistics* **54**(1), 383–407, 2026.

3. **F. Gili**, G. Jongbloed, A. van der Vaart. *Asymptotically efficient estimation under local constraint in Wicksell's problem. Journal of Statistical Planning and Inference* **240**, 106299, 2026.

4. **F. Gili**, G. Jongbloed. *Semiparametric Uncertainty Quantification via Isotonized Posterior for Deconvolutions.* Preprint, arXiv:2602.18210. *Submitted.*

5. **F. Gili**, K. Hu, A. van der Vaart. *Proximal inference based on mixtures of linear models. In preparation.*

Bibliography

[1] Adler, R.J. *An Introduction to Continuity, Extrema, and Related Topics for General Gaussian Processes*, volume 12 of *Lecture Notes-Monographs, Institute of Mathematical Statistics*. Institute of Mathematical Statistics, 1990. URL https://www.jstor.org/stable/pdf/4355563.pdf. https://www.jstor.org/stable/pdf/4355563.pdf.

[2] Alonso, J., Matio, M., and Sen, B. Uncloaking Globular Clusters of the Inner Galaxy. *Proceedings of the International Astronomical Union*, 3:359–360, 2007. doi: 10.1088/0004-6256/143/3/70. URL https://doi.org/10.1088/0004-6256/143/3/70. https://doi.org/10.1088/0004-6256/143/3/70.

[3] Antoniak, C. Mixtures of Dirichlet Processes with Applications to Bayesian Nonparametric Problems. *The Annals of Statistic*, 2(6):1152–1174, 1974. doi: 10.1214/aos/1176342871. URL https://doi.org/10.1214/aos/1176342871. https://doi.org/10.1214/aos/1176342871.

[4] Ayer, M., Brunk, H. D., Ewing, G. M., Reid, W. T., and Silverman, E. An Empirical Distribution Function for Sampling with Incomplete Information. *The Annals of Mathematical Statistics*, 26(4):641–647, 1955. URL http://www.jstor.org/stable/2236377. http://www.jstor.org/stable/2236377.

[5] Bhaumik, P. and Ghosal, S. Bayesian Two-Step Estimation in Differential Equation Models. *Electronic Journal of Statistics*, 9(2):3124–3154, 2015. doi: 10.1214/15-EJS1099. URL https://doi.org/10.1214/15-EJS1099. https://doi.org/10.1214/15-EJS1099.

[6] Bhaumik, P. and Ghosal, S. Efficient Bayesian estimation and uncertainty quantification in ordinary differential equation models. *Bernoulli*, 23:3537–3570, 2015. doi: 10.3150/16-BEJ856. URL https://doi.org/10.3150/16-BEJ856. https://doi.org/10.3150/16-BEJ856.

[7] Bolthausen, E., van der Vaart, A., and Perkins, E. *Lectures on Probability Theory and Statistics, Ecole d'Eté de Probabilités de Saint-Flour XXIX - Part III. Aad van der Vaart*. Springer, New York, 1999.

[8] Bracewell, R. N. and Roberts, J. A. Aerial Smoothing in Radio Astronomy. *Australian Journal of Physics*, 7(4):615–640, 1954. doi: 10.1071/PH540615. URL https://doi.org/10.1071/PH540615. https://doi.org/10.1071/PH540615.

[9] Butucea, C. Deconvolution of supersmooth densities with smooth noise. *Can. J. Stat.*, 32:181–192, 2004. doi: 10.2307/3315941. URL https://doi.org/10.2307/3315941. https://doi.org/10.2307/3315941.

[10] Butucea, C. and Tsybakov, A. B. Sharp optimality in density deconvolution with dominating bias I. *Theory Probab. Appl.*, 52:24–39, 2008. doi: 10.1137/S0040585X97982840. URL https://doi.org/10.1137/S0040585X97982840.

https://doi.org/10.1137/S0040585X97982840.

[11] Butucea, C. and Tsybakov, A. B. Sharp optimality in density deconvolution with dominating bias II. *Theory Probab. Appl.*, 52:237–249, 2008. doi: 10.1137/S0040585X97982992. URL https://doi.org/10.1137/S0040585X97982992. https://doi.org/10.1137/S0040585X97982992.

[12] Carroll, R. J. and Hall, P. Optimal rates of convergence for deconvolving a density. *J. Am. Stat. Assoc.*, 83:1184–1186, 1988. doi: 10.2307/2290153. URL https://doi.org/10.2307/2290153. https://doi.org/10.2307/2290153.

[13] Castillo, I. A semiparametric Bernstein–von Mises theorem for Gaussian process priors. *Probability Theory and Related Fields*, 152:53–99, 2012. doi: 10.1007/s00440-010-0316-5. URL https://doi.org/10.1007/s00440-010-0316-5. https://doi.org/10.1007/s00440-010-0316-5.

[14] Castillo, I. and Nickl, R. Nonparametric Bernstein–von Mises Theorems in Gaussian White Noise. *The Annals of Statistics*, 41(4):1999–2028, 2013. doi: 10.1214/13-AOS1133. URL https://doi.org/10.1214/13-AOS1133. https://doi.org/10.1214/13-AOS1133.

[15] Castillo, I. and Nickl, R. On the Bernstein-Von Mises Phenomenon for Nonparametric Bayes Procedures. *The Annals of Statistics*, 42(5):1941–1969, 2014. URL http://www.jstor.org/stable/43556349. http://www.jstor.org/stable/43556349.

[16] Castillo, I. and Rousseau, J. A Bernstein–von Mises Theorem for Smooth Functionals in Semiparametric Models. *The Annals of Statistics*, 43(6):2353 – 2383, 2015. doi: 10.1214/15-AOS1336. URL https://doi.org/10.1214/15-AOS1336. https://doi.org/10.1214/15-AOS1336.

[17] Chakraborty, M. and Ghosal, S. Coverage of Credible Intervals in Nonparametric Monotone Regression. *Annals of Statistics*, 49(2):1011–1028, 2021. doi: 10.1214/20-AOS1989. URL https://doi.org/10.1214/20-AOS1989. https://doi.org/10.1214/20-AOS1989.

[18] Chakraborty, M. and Ghosal, S. Rates and coverage for monotone densities using projection-posterior. *Bernoulli*, 28(2):1093 – 1119, 2022. doi: 10.3150/21-BEJ1379. URL https://doi.org/10.3150/21-BEJ1379. https://doi.org/10.3150/21-BEJ1379.

[19] Chan, K. C. G. and Qin, J. Nonparametric Maximum Likelihood Estimation for the Multisample Wicksell Corpuscle Problem. *Biometrika*, 103:273–286, 2016. doi: 10.1093/biomet/asw011. URL https://doi.org/10.1093/biomet/asw011. https://doi.org/10.1093/biomet/asw011.

[20] Comte, F., Rozenholc, Y., and Taupin, M.-L. Penalized contrast estimator for adaptive density deconvolution. *Can. J. Stat.*, 34:431–452, 2006. URL https://www.jstor.org/stable/20445213. https://www.jstor.org/stable/20445213.

[21] Cui, Y., Pu, H., Shi, X., Miao, W., and Tchetgen Tchetgen, E. Semiparametric Proximal Causal Inference. *Journal of the American Statistical Association*, 119(546):1348–1359, 2023. doi: 10.1080/01621459.2023.2191817. URL https://doi.org/10.1080/01621459.2023.2191817. https://doi.org/10.1080/01621459.2023.2191817.

[22] Cuzzi, J. and Olson, D. Recovering 3D Particle Size Distributions from 2D Sections. *Meteoritics & Planetary Science*, 52(3):532–545, 2017. doi: 10.1111/maps.12812. URL https://doi.org/10.1111/maps.12812. https://doi.org/10.1111/maps.12812.

[23] Dattner, I. and Reiser, B. Estimation of distribution functions in measurement error

models. *J. Stat. Plann. Inference*, 143:479–493, 2013. doi: 10.1016/j.jspi.2012.09.004. URL https://doi.org/10.1016/j.jspi.2012.09.004. https://doi.org/10.1016/j.jspi.2012.09.004.

[24] Dattner, I., Goldenshluger, A., and Juditsky, A. On deconvolution of distribution functions. *Ann. Statist.*, 39:2477–2501, 2011. doi: 10.1214/11-AOS907. URL https://doi.org/10.1214/11-AOS907. https://doi.org/10.1214/11-AOS907.

[25] Dattner, I., Reiss, M., and Trabs, M. Adaptive quantile estimation in deconvolution with unknown error distribution. *Bernoulli*, 22:143–192, 2016. doi: 10.3150/14-BEJ626. URL https://doi.org/10.3150/14-BEJ626. https://doi.org/10.3150/14-BEJ626.

[26] Delaigle, A. and Gijbels, I. Bootstrap bandwidth selection in kernel density estimation from a contaminated sample. *Ann. Inst. Statist. Math.*, 56:19–47, 2004. doi: 10.1007/BF02506823. URL https://doi.org/10.1007/BF02530523. https://doi.org/10.1007/BF02530523.

[27] Deng, H., Han, Q., and Zhang, C. Confidence Intervals for Multiple Isotonic Regression and other Monotone Models. *The Annals of Statistics*, 49(4):2021–2052, 2021. doi: 10.1214/20-AOS2025. URL https://doi.org/10.1214/20-AOS2025. https://doi.org/10.1214/20-AOS2025.

[28] Diggle, P. J. and Hall, P. A Fourier Approach to Nonparametric Deconvolution of a Density Estimate. *Journal of the Royal Statistical Society. Series B (Methodological)*, 55 (2):523–531, 1993. doi: 10.2307/2346211. URL http://www.jstor.org/stable/2346211. http://www.jstor.org/stable/2346211.

[29] Durot, C. and Tocquet, A.-S. On the Distance between the Empirical Process and its Concave Majorant in a Monotone Regression Framework. *Annales de l'Institut Henri Poincaré (B) Probability and Statistics*, 39(2):217–240, 2003. URL https://www.sciencedirect.com/science/article/pii/S0246020302000134. https://doi.org/10.1016/S0246-0203(02)00013-4.

[30] Durrett, R. *Probability: Theory and Examples*. Cambridge University Press, 2010.

[31] Efromovich, S. Density estimation for the case of supersmooth measurement errors. *J. Am. Stat. Assoc.*, 92:526–535, 1997. doi: 10.1080/01621459.1997.10474005. URL https://doi.org/10.1080/01621459.1997.10474005. https://doi.org/10.1080/01621459.1997.10474005.

[32] Efromovich, S. A sharp minimax lower bound for the nonparametric estimation of Sobolev densities of order $\frac{1}{2}$. *Statistics and Probability Letters*, 80:77–81, 2010. doi: 10.1016/j.spl.2009.08.011. URL https://doi.org/10.1016/j.spl.2009.08.011. https://doi.org/10.1016/j.spl.2009.08.011.

[33] Fan, J. On the optimal rates of convergence for nonparametric deconvolution problems. *The Annals of Statistics*, 19:1257–1272, 1991. doi: 10.1214/aos/1176348248. URL https://doi.org/10.1214/aos/1176348248. https://doi.org/10.1214/aos/1176348248.

[34] Ferguson, T. S. Prior Distributions on Spaces of Probability Measures. *The Annals of Statistics*, 2(4):615 – 629, 1974. doi: 10.1214/aos/1176342752. URL https://doi.org/10.1214/aos/1176342752. https://doi.org/10.1214/aos/1176342752.

[35] Ferguson, T.S. *Mathematical Statistics: A Decision Theoretic Approach*. Academic Press, New York, 1967.

[36] Flanders, W.D., Klein, M., Darrow, L.A., Strickland, M.J., Sarnat, S.E., Sarnat, J.A., Waller, L.A., Winquist, A., and Tolbert, P.E. A method for detection of residual confounding in time-series and other observational studies. *Epidemiology (Cambridge,*

Mass.), 22:59–67, 2011. doi: 10.1097/EDE.0b013e3181fdcabe. URL https://doi.org/10.1097/EDE.0b013e3181fdcabe. https://doi.org/10.1097/EDE.0b013e3181fdcabe.

[37] Flanders, W.D., Strickland, M.J., and Klein, M. A New Method for Partial Correction of Residual Confounding in Time-Series and Other Observational Studies. *American Journal of Epidemiology*, 185(10):941–949, 2015. doi: 10.1093/aje/kwx013. URL https://doi.org/10.1093/aje/kwx013. https://doi.org/10.1093/aje/kwx013.

[38] Gagnon-Bartsch, J.A. and Speed, T.P. Using control genes to correct for unwanted variation in microarray data. *Biostatistics*, 13(3):539–552, 2012. doi: 10.1093/biostatistics/kxr034. URL https://doi.org/10.1093/biostatistics/kxr034. https://doi.org/10.1093/biostatistics/kxr034.

[39] Gao, F. and van der Vaart, A. Posterior Contraction Rates for Deconvolution of Dirichlet-Laplace Mixtures. *Electronic Journal of Statistics*, 10:608–627, 2016. doi: 10.1214/16-EJS1119. URL https://doi.org/10.1214/16-EJS1119. https://doi.org/10.1214/16-EJS1119.

[40] Ghosal, S. and van der Vaart, A. Entropies and Rates of Convergence for Maximum Likelihood and Bayes Estimation for Mixtures of Normal Densities. *The Annals of Statistics*, 29(5):1233–1263, 2001. doi: 10.1214/aos/1013203452. URL https://doi.org/10.1214/aos/1013203452. https://doi.org/10.1214/aos/1013203452.

[41] Ghosal, S. and van der Vaart, A. Posterior Convergence Rates of Dirichlet Mixtures at Smooth Densities. *The Annals of Statistics*, 35(2):697–723, 2007. URL http://www.jstor.org/stable/25463573. http://www.jstor.org/stable/25463573.

[42] Ghosal, S. and van der Vaart, A. *Fundamentals of Nonparametric Bayesian Inference*. Cambridge University Press, 2017.

[43] Gili, F., Jongbloed, G., and van der Vaart, A. Adaptive and Efficient Isotonic Estimation in Wicksell's Problem. *Journal of Nonparametric Statistics*, 37(3):491–531, 2025. doi: 10.1080/10485252.2024.2397680. URL https://doi.org/10.1080/10485252.2024.2397680. https://doi.org/10.1080/10485252.2024.2397680.

[44] Gili, F., Jongbloed, G., and van der Vaart, A. Asymptotically Efficient Estimation under Local Constraint in Wicksell's Problem. *Journal of Statistical Planning and Inference*, 240:106299, 2026. doi: 10.1016/j.jspi.2025.106299. URL https://doi.org/10.1016/j.jspi.2025.106299. https://doi.org/10.1016/j.jspi.2025.106299.

[45] Gili, F., Jongbloed, G., and van der Vaart, A. Semiparametric Bernstein-von Mises Phenomenon via Isotonized Posterior in Wicksell's problem. *The Annals of Statistics*, 54 (1):383–407, 2026. doi: 10.1214/25-AOS2571. URL https://doi.org/10.1214/25-AOS2571. https://doi.org/10.1214/25-AOS2571.

[46] Gili, F., Jongbloed, G., and van der Vaart, A. Supplement to "Semiparametric Bernstein-von Mises Phenomenon via Isotonized Posterior in Wicksell's problem". *The Annals of Statistics*, 54(1), 2026. doi: 10.1214/25-AOS2571. URL https://doi.org/10.1214/25-AOS2571. Online supplement to [45].

[47] Gili, F., Jongbloed, G., and van der Vaart, A. Semiparametric Bernstein-von Mises Phenomenon via Isotonized Posterior in Wicksell's problem. *The Annals of Statistics*, 54 (1):383–407, 2026. doi: 10.1214/25-AOS2571. URL https://doi.org/10.1214/25-AOS2571. https://doi.org/10.1214/25-AOS2571.

[48] Golubev, G. K. and Enikeeva, F. N. Asymptotically Efficient Smoothing in the Wicksell Problem under Squared Losses. *Problems of Information Transmission*, 37(1):12–

21, 2001. doi: 10.1023/A:1010495609901. URL https://doi.org/10.1023/A:1010495609901. https://doi.org/10.1023/A:1010495609901.

[49] Golubev, G. K. and Levit, B. Y. Asymptotically Efficient Estimation in the Wicksell Problem. *The Annals of Statistics*, 26(6):2407–2419, 1998. doi: 10.1214/aos/1024691477. URL https://doi.org/10.1214/aos/1024691477. https://doi.org/10.1214/aos/1176324310.

[50] Grimmett, G. and Stirzaker, D. *Probability and Random Processes*. Oxford University Press, 1987.

[51] Londen, S. O. Gripenberg, G. and Staffans, O. *Volterra Integral and Functional Equations*. Cambridge University Press, 1990. URL https://doi.org/10.1017/CBO9780511662805.

[52] Groeneboom, P. and Jongbloed, G. Isotonic Estimation and Rates of Convergence in Wicksell's Problem. *The Annals of Statistic*, 23(5):1528–1542, 1995. doi: 10.1214/aos/1176324310. URL https://doi.org/10.1214/aos/1176324310. https://doi.org/10.1214/aos/1176324310.

[53] Groeneboom, P. and Jongbloed, G. *Nonparametric Estimation Under Shape Constraints*. Cambridge University Press, 2014.

[54] Groeneboom, P. and Wellner, J. A. Information bounds and nonparametric maximum likelihood estimation. *DMV Seminar, Birkhäuser Verlag, Basel*, 1992. Book.

[55] Hall, P. and Lahiri, S. N. Estimation of distributions, moments and quantiles in deconvolution problems. *Ann. Statist.*, 36:2110–2134, 2008. URL https://www.jstor.org/stable/25464705. https://www.jstor.org/stable/25464705.

[56] Hall, P. and Smith, R. L. The Kernel Method for Unfolding Sphere Size Distributions. *J. Comput. Phys.*, 74:409–421, 1988. doi: 10.1016/0021-9991(88)90085-X. URL https://doi.org/10.1016/0021-9991(88)90085-X. https://doi.org/10.1016/0021-9991(88)90085-X.

[57] Hjort, N. and Petrone, S. Nonparametric Quantile Inference using Dirichlet Processes, 2007. URL https://doi.org/10.1142/9789812708298_0023. https://doi.org/10.1142/9789812708298_0023.

[58] Hu, J. K., Zorzetto, D., and Dominici, F. Estimate an Exposure Response Function with Negative Controls: A Bayesian Nonparametric Approach. *American Journal of Epidemiology*, page kwag073, 2026. doi: 10.1093/aje/kwag073. URL https://doi.org/10.1093/aje/kwag073. https://doi.org/10.1093/aje/kwag073.

[59] Jirak, M., Meister, A., and Reiss, M. Adaptive function estimation in nonparametric regression with one-sided errors. *Ann. Statist.*, 42:1970–2002, 2014. doi: 10.1214/14-AOS1248. URL https://doi.org/10.1214/14-AOS1248. https://doi.org/10.1214/14-AOS1248.

[60] Jongbloed, G. *Three Statistical Inverse Problems, Estimators-algorithms-asymptotics*. PhD Thesis, TU Delft, 1995. URL https://repository.tudelft.nl/record/uuid:1412cd89-70ff-41ae-8ba2-04abcadc92a9.

[61] Jongbloed, G. Exponential deconvolution: Two asymptotically equivalent estimators. *Stat. Neerl.*, 52:6–17, 1998. doi: 10.1111/1467-9574.00065. URL https://doi.org/10.1111/1467-9574.00065. https://doi.org/10.1111/1467-9574.00065.

[62] Jongbloed, G. Sieved Maximum Likelihood Estimation in Wicksell's Problem and Related Deconvolution Problems. *Scandinavian Journal of Statistics*, 2(6):1152–1174, 2001. doi: 10.1111/1467-9469.00230. URL https://doi.org/10.1111/1467-9469.00230. https://doi.org/10.1111/1467-9469.00230.

[63] Kallenberg, O. *Foundations of Modern Probability*. Probability Theory and Stochastic

Modelling. Springer, third edition, 2021. URL https://link.springer.com/book/10.1007/978-3-030-61871-1. https://link.springer.com/book/10.1007/978-3-030-61871-1.

[64] Kallus, N., Mao, X., and Uehara, M. Causal inference under unmeasured confounding with negative controls: A minimax learning approach, 2021. URL https://doi.org/10.48550/arXiv.2103.14029. Preprint at https://doi.org/10.48550/arXiv.2103.14029.

[65] Kim, J. and Pollard, D. Cube Root Asymptotics. *The Annals of Statistics*, 18(1): 191–219, 1990. doi: 10.1214/aos/1176347498. URL https://doi.org/10.1214/aos/1176347498. https://doi.org/10.1214/aos/1176347498.

[66] Kim, Y. The Bernstein–von Mises Theorem for the Proportional Hazard Model. *The Annals of Statistics*, 34(4):1678 – 1700, 2006. doi: 10.1214/009053606000000533. URL https://doi.org/10.1214/009053606000000533. https://doi.org/10.1214/009053606000000533.

[67] Kim, Y. and Lee, J. A Bernstein–von Mises Theorem in the Nonparametric Right-Censoring Model. *Annals of Statistics*, 32:1492–1512, 2004. URL https://api.semanticscholar.org/CorpusID:88519037. https://api.semanticscholar.org/CorpusID:88519037.

[68] Kompa, B., Bellamy, D.R., Kolokotrones, T., Robins, J.M., and Beam, A.L. Deep learning methods for proximal inference via maximum moment restriction. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 35, pages 11189–11201, 2022. URL https://papers.neurips.cc/paper_files/paper/2022/file/487c9d6ef55e73aa9dfd4b48fe3713a6-Paper-Conference.pdf. https://papers.neurips.cc/paper_files/paper/2022/file/487c9d6ef55e73aa9dfd4b48fe3713a6-Paper-Conference.pdf.

[69] Koshevnik, Yu. A. and Levit, B. Ya. On a Non-Parametric Analogue of the Information Matrix. *Theory of Probability and its Applications*, 21, 2023. URL https://doi.org/10.1137/1121087. https://doi.org/10.1137/1121087.

[70] Kuroki, M. and Pearl, J. Measurement bias and effect restoration in causal inference. *Biometrika*, 101:423–437, 2014. doi: 10.1093/biomet/ast066. URL https://doi.org/10.1093/biomet/ast066. https://doi.org/10.1093/biomet/ast066.

[71] Lipsitch, M., Tchetgen, E.T., and Cohen, T. Negative controls: a tool for detecting confounding and bias in observational studies. *Epidemiology (Cambridge, Mass.)*, 21:383–388, 2010. doi: 10.1097/EDE.0b013e3181d61eeb. URL https://doi.org/10.1097/EDE.0b013e3181d61eeb. https://doi.org/10.1097/EDE.0b013e3181d61eeb.

[72] Lopez-Sanchez, M. A. and LLana-Fúnez, s. An Extension of the Saltykov Method to Quantify 3D Grain Size Distributions in Mylonites. *Journal of Structural Geology*, 93: 149–161, 2016. doi: 10.1016/j.jsg.2016.10.008. URL https://doi.org/10.1016/j.jsg.2016.10.008. https://doi.org/10.1016/j.jsg.2016.10.008.

[73] Lucy, L. B. An iterative technique for the rectification of observed distributions. *The Astronomical Journal*, 79:745–754, 1974. doi: 10.1086/111605. URL https://doi.org/10.1086/111605. https://doi.org/10.1086/111605.

[74] Meister, A. Deconvolution problems in nonparametric statistics. *Lecture Notes in Statistics, Springer-Verlag, Berlin*, 2009. doi: 10.1007/978-3-540-87557-4. URL https://doi.org/10.1007/978-3-540-87557-4. Book, DOI: 10.1007/978-3-540-87557-4.

[75] Miao, W., Geng, Z., and Tchetgen Tchetgen, E.J. Identifying treatment effects with

proxy variables of an unmeasured confounder. *Biometrika*, 105:987–993, 2018. doi: 10.1093/biomet/asy038. URL https://doi.org/10.1093/biomet/asy038. https://doi.org/10.1093/biomet/asy038.

[76] Monard, F., Nickl, R., and Paternain, G.P. Efficient Nonparametric Bayesian Inference for X-ray Transforms. *The Annals of Statistics*, 47(2):1113 – 1147, 2019. doi: 10.1214/18-AOS1708. URL https://doi.org/10.1214/18-AOS1708. https://doi.org/10.1214/18-AOS1708.

[77] Nguyen, X. Convergence of Latent Mixing Measures in Finite and Infinite Mixture Models. *The Annals of Statistics*, 41(1):370–400, 2013. doi: 10.1214/12-AOS1065. URL https://doi.org/10.1214/12-AOS1065. https://doi.org/10.1214/AOS1065.

[78] Nickl, R. Bernstein-von Mises Theorems for Statistical Inverse Problems I: Schrödinger equation. *Journal of the European Mathematical Society*, 22, 2017. doi: 10.4171/JEMS/975. URL https://doi.org/10.4171/JEMS/975. https://doi.org/10.4171/JEMS/975.

[79] Nickl, R. and Söhl, J. Bernstein-von Mises Theorems for Statistical Inverse Problems II: compound Poisson processes. *Electronic Journal of Statistics*, 13(2):3513–3571, 2019. doi: 10.1214/19-ejs1609. URL https://doi.org/10.1214/19-ejs1609. https://doi.org/10.1214/19-ejs1609.

[80] Norets, A. and Pelenis, J. Posterior Consistency in Conditional Density Estimation by Covariate Dependent Mixtures. *Econometric Theory*, 30:606–646, 2014. doi: 10.1017/S026646661300042X. URL https://doi.org/10.1017/S026646661300042X. https://doi.org/10.1017/S026646661300042X.

[81] Pati, D., Dunson, D. B., and Tokdar, S. Posterior consistency in conditional distribution estimation. *Journal of Multivariate Analysis*, 116:456–472, 2013. doi: 10.1016/j.jmva.2013.01.011. URL https://doi.org/10.1016/j.jmva.2013.01.011. https://doi.org/10.1016/j.jmva.2013.01.011.

[82] Pensky, M. and Vidakovic, B. Adaptive wavelet estimator for nonparametric density deconvolution. *Ann. Statist.*, 27:2033–2053, 1999. doi: 10.1214/aos/1017939249. URL https://doi.org/10.1214/aos/1017939249. https://doi.org/10.1214/aos/1017939249.

[83] Pollard, D. *Convergence of Stochastic Processes*. Springer, New York, 1984.

[84] Ray, K. Adaptive Bernstein-von Mises Theorems in Gaussian White Models. *The Annals of Statistics*, 45(6):2511–2536, 2017. URL http://www.jstor.org/stable/26362949. http://www.jstor.org/stable/26362949.

[85] Ray, K. and van der Vaart, A. Semiparametric Bayesian Causal Inference. *The Annals of Statistics*, 48:2999–3020, 2020. doi: 10.1214/19-AOS1919. URL https://doi.org/10.1214/19-AOS1919. https://doi.org/10.1214/19-AOS1919.

[86] Reiss, M. and Selk, L. Estimating nonparametric functionals efficiently under one-sided errors. *Bernoulli*, 23:1022–1055, 2017. URL https://doi.org/10.48550/arXiv.1407.4229. https://doi.org/10.48550/arXiv.1407.4229.

[87] Richardson, W. H. Bayesian-Based Iterative Method of Image Restoration. *Journal of the Optical Society of America*, 62(1):55–59, 1972. doi: 10.1364/JOSA.62.000055. URL https://doi.org/10.1364/JOSA.62.000055. https://doi.org/10.1364/JOSA.62.000055.

[88] Rivoirard, V. and Rousseau, J. Bernstein-von Mises Theorem for Linear Functionals of the density. *The Annals of Statistics*, 40(3):1489–1523, 2012. URL http://www.jstor.org/stable/41713683. http://www.jstor.org/stable/41713683.

[89] Robinson, E. A. Predictive Decomposition of Seismic Traces. *Geophysics*, 22(4):767–778, 1957. doi: 10.1190/1.1438419. URL https://doi.org/10.1190/1.1438419. https://doi.org/10.1190/1.1438419.

[90] Rudin, W. *Real and Complex Analysis*. McGraw-Hill, New York, 3rd edition, 1987.

[91] Sen, B. *A Gentle Introduction to Empirical Process Theory and Applications*. Lecture notes, Columbia University, 2022.

[92] Sen, B. and Woodroofe, F. N. Bootstrap Confidence Intervals for Isotonic Estimators in a Stereological Problem. *Bernoulli*, 18(4):1249–1266, 2012. doi: 10.3150/12-BEJ378. URL https://doi.org/10.3150/12-BEJ378. https://doi.org/10.3150/12-BEJ378.

[93] Sen, B., Banerjee, M., and Woodroofe, M. Inconsistency of Bootstrap: The Grenander Estimator. *The Annals of Statistics*, 38(4):1953–1977, 2010. doi: 10.1214/09-AOS777. URL https://doi.org/10.1214/09-AOS777. https://doi.org/10.1214/09-AOS777.

[94] Shi, X., Miao, W., Nelson, J.C., and Tchetgen Tchetgen, E.J. Multiply Robust Causal Inference with Double Negative Control Adjustment for Categorical Unmeasured Confounding. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 82(2):521–540, 2019. doi: 10.1111/rssb.12361. URL https://doi.org/10.1111/rssb.12361. https://doi.org/10.1111/rssb.12361.

[95] Shi, X., Miao, W., and Tchetgen Tchetgen, E. A Selective Review of Negative Control Methods in Epidemiology. *Current Epidemiology Reports*, 7(4):190–202, 2020. doi: 10.1007/s40471-020-00243-4. URL https://doi.org/10.1007/s40471-020-00243-4. https://doi.org/10.1007/s40471-020-00243-4.

[96] Sofer, T., Richardson, D.B., Colicino, E., Schwartz, J., and Tchetgen Tchetgen, E.J. On negative outcome control of unobserved confounding as a generalization of difference-in-differences. *Statistical Science*, 31(3):348–361, 2016. doi: 10.1214/16-STS558. URL https://doi.org/10.1214/16-STS558. https://doi.org/10.1214/16-STS558.

[97] Stefanski, L. A. Rates of convergence of some estimators in a class of deconvolution problems. *Statist. Probab. Lett.*, 9:229–235, 1990. doi: 10.1016/0167-7152(90)90061-B. URL https://doi.org/10.1016/0167-7152(90)90061-B. https://doi.org/10.1016/0167-7152(90)90061-B.

[98] Stefanski, L. A. and Carroll, R. J. Deconvoluting kernel density estimators. *Statistics*, 21:169–184, 1990. doi: 10.1080/02331889008802238. URL https://doi.org/10.1080/02331889008802238. https://doi.org/10.1080/02331889008802238.

[99] Stoyan, D. and Kendall, W. S. *Stochastic Geometry and Its Applications*. Wiley, New York, 1987.

[100] Tchetgen Tchetgen, E. The Control Outcome Calibration Approach for Causal Inference With Unobserved Confounding. *American Journal of Epidemiology*, 179(5):633–640, 2014. doi: 10.1093/aje/kwt303. URL https://doi.org/10.1093/aje/kwt303. https://doi.org/10.1093/aje/kwt303.

[101] Tokdar, S. Posterior Consistency of Dirichlet Location-Scale Mixture of Normals in Density Estimation and Regression. *The Indian Journal of Statistics*, 67:90–110, 2006. URL https://www.jstor.org/stable/25053478. https://www.jstor.org/stable/25053478.

[102] van der Vaart, A. *Asymptotic Statistics*. Cambridge University Press, 1998.

[103] van der Vaart, A. and Wellner, J. *Weak Convergence and Empirical Processes*. Springer, New York, 2023.

[104] van Es, A. J. Uniform Deconvolution: Nonparametric Maximum Likelihood and Inverse Estimation. In Roussas, G., editor, *Nonparametric Functional Estimation and Related Topics*, volume 335. Springer, 1991. doi: 10.1007/978-94-011-3222-0_14. URL https://doi.org/10.1007/978-94-011-3222-0_14. https://doi.org/10.1007/978-94-011-3222-0_14.

[105] van Es, A. J. and Kok, A. R. Simple kernel estimators for certain nonparametric deconvolution problems. *Statistics and Probability Letters*, 39(2):151–160, 1998. doi: 10.1016/S0167-7152(98)00054-6. URL https://doi.org/10.1016/S0167-7152(98)00054-6. https://doi.org/10.1016/S0167-7152(98)00054-6.

[106] van Es, A. J. and van Zuijlen, M. C. A. Convex Minorant Estimators of Distributions in Non-Parametric Deconvolution Problems. *Scandinavian Journal of Statistics*, 23(1):85–104, 1996. doi: 10.2307/4616388. http://www.jstor.org/stable/4616388.

[107] van Es, B. Combining kernel estimators in the uniform deconvolution problem. *Stat. Neerl.*, 65:275–296, 2011. doi: 10.1111/j.1467-9574.2011.00485.x. URL https://doi.org/10.1111/j.1467-9574.2011.00485.x. https://doi.org/10.1111/j.1467-9574.2011.00485.x.

[108] van Es, B., Jongbloed, G., and van Zuijlen, M. Nonparametric deconvolution for decreasing kernels. *Delft University of Technology, Faculty of Technical Mathematics and Informatics*, 1995. URL https://www.researchgate.net/publication/2362108_Nonparametric_Deconvolution_for_Decreasing_Kernels.

[109] van Es, B., Jongbloed, G., and van Zuijlen, M. Isotonic inverse estimators for nonparametric deconvolution. *The Annals of Statistics*, 26:2395–2406, 1998. doi: 10.1214/aos/1024691476. URL https://doi.org/10.1214/aos/1024691476. https://doi.org/10.1214/aos/1024691476.

[110] Villani, C. *Optimal Transport: Old and New*. Springer Berlin Heidelberg, Berlin, Heidelberg, 2009.

[111] Wang, J., Zhao, Q., Hastie, T., and Owen, A.B. Confounder adjustment in multiple hypothesis testing. *Ann. Statist.*, 45(5):1863–1894, 2017. doi: 10.1214/16-AOS1511. URL https://doi.org/10.1214/16-AOS1511. https://doi.org/10.1214/16-AOS1511.

[112] Wang, K. and Ghosal, S. Coverage of Credible Intervals in Bayesian Multivariate Isotonic Regression. *Annals of Statistics*, 51(3):1376–1400, 2023. doi: 10.1214/23-AOS2298. URL https://doi.org/10.1214/23-AOS2298. https://doi.org/10.1214/23-AOS2298.

[113] Wang, K. and Ghosal, S. Posterior Contraction and Testing for Multivariate Isotonic Regression. *Electronic Journal of Statistics*, 17(1):798–822, 2023. doi: 10.1214/23-EJS2115. URL https://doi.org/10.1214/23-EJS2115. https://doi.org/10.1214/23-EJS2115.

[114] Watson, G. S. Estimating Functionals of Particle Size Distributions. *Biometrika*, 58:483–490, 1971. doi: 10.1093/biomet/58.3.483. URL https://doi.org/10.1093/biomet/58.3.483. https://doi.org/10.1093/biomet/58.3.483.

[115] Wicksell, S. D. The Corpuscle Problem. *Biometrika*, 17:84–99, 1925. doi: 10.1093/biomet/17.1-2.84. URL https://doi.org/10.1093/biomet/17.1-2.84. https://doi.org/10.1093/biomet/17.1-2.84.

[116] Zhang, C.-H. Fourier methods for estimating mixing densities and distributions. *Ann. Statist.*, 18:806–831, 1990. doi: 10.1214/aos/1176347627. URL https://doi.org/10.1214/aos/1176347627. https://doi.org/10.1214/aos/1176347627.