

ICFP M2 - STATISTICAL PHYSICS 2
Homework n° 6
Random Matrices and the Wigner surmise

Grégory Schehr, Francesco Zamponi

This exercise is a preparation to the lectures and TDs that will focus on Random Matrices and Random Hamiltonians.

Consider a two by two symmetric real random matrix M such that the matrix elements M_{11} , M_{12} and M_{22} are independent Gaussian random variables with zero mean and variances :

$$\mathbb{E}[M_{11}^2] = 1 , \quad \mathbb{E}[M_{22}^2] = 1 , \quad \mathbb{E}[M_{12}^2] = \frac{1}{2} ;$$

by symmetry $M_{21} = M_{12}$. We denote λ_1 and λ_2 the eigenvalues of M , and $\Delta = |\lambda_1 - \lambda_2|$ their spacing.

Find the probability density of Δ , its average value $\mathbb{E}[\Delta]$, and deduce that the normalized spacing $s = \Delta/\mathbb{E}[\Delta]$ has the probability density

$$P(s) = \frac{\pi}{2} s e^{-\frac{\pi}{4} s^2} ,$$

known as the Wigner surmise.

It is what Wigner proposed as an approximation for the probability density function of the normalised mean-level spacing of very complex nuclei, see Figure 1. The connection between random matrices and the Hamiltonian of a very complex non-random Hamiltonian will be discussed in the next lectures and TDs.




FIGURE 1 – Nearest neighbor spacing distribution for the “Nuclear Data Ensemble” comprising 1726 spacings (histogram) versus $s = S/D$ with D the mean level spacing and S the actual spacing. For comparison, the Wigner surmise labelled GOE is shown (don’t mind about the curve labelled Poisson).