

Prequel to Hawkes Processes: An Overview of Spatial, Temporal and Spatio-Temporal Point Processes and Some Simulations

Frances Lin

June 2021

Major Professor: James Molyneux **1. Help with reference 2. Check definitions especially HPP**

Committee Members: Lisa Madsen & Charlotte Wickham

Abstract

Events such as earthquake epicenters, crime patterns, forest wildfires, financial transactions, etc. often exhibit triggering and clustering behavior. The ability to capture events with such behavior gives Hawkes or self-exciting point processes (SEPP) the potential to become a powerful predictive tool for a wide variety of applications. In this project, we give brief introductions, review definitions, discuss properties and applications of selected spatial and temporal point processes leading up to spatio-temporal SEPP, and simulate some of the processes in 1D and 2D in hope that interested readers have the background knowledge to read and comprehend existing SEPP literature as well as explore the field further.

1 Introduction

Real-world data are often spatial, temporal, or spatio-temporal in nature. Spatial data (e.g. soil properties, housing prices) often involve locations such as points and areas. Temporal data (e.g. sensor readings, stock prices) often involve times such as moments and intervals. Spatio-temporal data are data that relates to both locations and times. Examples of spatio-temporal data include earthquake epicenters, forest inventories, remotely sensed images, disease cases, map services and travel times, to name a few.

Various statistical methods have been developed to model spatial, temporal, or spatio-temporal data, and which models to use depend on questions of interest. For this project, we focus primarily on the point process models. These models are useful for describing phenomena that occurs at random locations, times,

or locations and times, and the questions of interests typically are: Does the rate for the occurrence of events vary with space, time, or space-time? Do events appear clustered? Do events trigger subsequent events?

Recall that spatial data can be broadly categorized into three types: geostatistical data, areal data, and point pattern data (Cressie, 2015), we are in the third category in which point pattern data are realizations of spatial point processes. Typical questions are: Is there clustering of events? Can we define a point process to capture the events? Examples of such models include Cox and cluster processes. On the other hand, when dealing with temporal data, *marked* point processes are sometimes used interchangeable with time series, and vice versa (Schoenberg, 2010). One major distinction is that in point processes, time intervals are treated as continuous, whereas in time series, they are treated as discrete. Examples of such models include (temporal) Poisson and Hawkes processes.

Hawkes processes are also known as self-exciting point processes (SEPP). The original Hawkes processes are temporal, whereas the more recently developed SEPP have been extended to account for both the spatial and temporal aspects of the data. The defining characteristic of SEPP is that it ‘self-excites’, i.e., the occurrence of an event increases the occurrence of future events nearby in space and/or time, although the events don’t self-excite in perpetuity. Given the history of events, more recent events also exert more influence on the rate at which events occur. In seismology, an event can be an earthquake that causes aftershocks. In criminology, an event can be a gang rivalry that triggers retaliations following the gang crime. In both cases, the initial event can continue to spawn ‘offspring’ events and the ‘offspring’ events can spawn ‘offspring’ events of their own, but the spawns fade out eventually.

In addition to modeling earthquake epicenters (Ogata, 1988, 1998) and crime patterns (Mohler et al., 2011; Reinhart & Greenhouse, 2018), Hawkes processes have also been used in modeling events such as forest wildfires (Peng, et al., 2005), insurance claims (Stabile et al., 2010), financial transactions (Bauwens and Hautsch, 2009; Embrechts et al., 2011; Bacry et al., 2015), social network events (Zhao et al., 2015; Rizoïu et al., 2017), neuron activities (Johnson, 1996; Gerhard et al., 2017), and disease spread or transmission (Meyer et al., 2012; Meyer & Held, 2014), thus allowing the models find applications in a wide variety of fields such as risk management, finance, social network, etc. Recent work has extended the use of SEPP to novel applications such as mass shootings (Boyd & Molyneux, 2021), COVID-19 transmission (Chiang et al., 2020), and gang violence (Park et al., 2021). However, there is still much work to be done which include computational advances to ease the burden of applying the models to bigger data sets (Holbrook et al., 2021), residual and model diagnostics, methods that make the models more flexible and applicable, etc. Given the flexibility and applicability of SEPP, it is surprising that SEPP have not gained enough attention from the machine learning communities, which would find their predictive capabilities beneficial.

Understanding SEPP also benefits from knowing some of the relevant point processes (e.g. nonhomogeneous Poisson, Cox and cluster processes), which are often left out from graduate-level, introductory spatial statistics and stochastic processes courses. The objective of this project is then to give an overview of various types of point processes so that readers of interest have the background knowledge to read and comprehend existing SEPP literature as well as explore the field further.

The outline of this project is as follows: In *Section 2*, we introduce, define and discuss properties and applications of counting processes, homogeneous and nonhomogeneous Poisson processes, Cox processes, cluster processes, Hawkes and spatio-temporal SEPP. In *Section 3*, we wrap up the aforementioned point processes and mention future work of Hawkes and SEPP. In *Appendix*, we discuss in particular the thinning algorithm (acceptance-rejection method) and **spatstat** package of **R** that are used to simulate selected processes in 1D and 2D, respectively.

2 Introductions, Definitions and Properties

2.1 Counting Process

A counting process counts the occurrence (or number) of events over time, space, space-time or, in the most general sense, any metric space in which events occur and can be counted (Daley & Vere-Jones, 2003). If we were to denote the time of arrival for customers at a super market, then we would have a set of points in time in which we could count the number customers over some interval of time. If we consider the location of trees to occur at a point in space, then in some bounded region of the space, we could count the number of trees. An earthquake's epicenter would be a point in space but we can also capture when the point appears in time.

Let us restrict ourselves to the temporal domain so that interpretations are easier to follow. The counting process requires that 1) the number of events $N(t)$ up to some time t to be greater than zero, 2) the number of events must be an integer, 3) the counts always increase, and 4) the number of events in specific time interval can be obtained by subtracting the number of events in previous interval from that in current interval. In addition, counting processes are independent, stationary, and homogeneous. In other words, the number of events $N(t)$ occurring in disjoint interval t are independent, the distribution of the number of events depends only on the length of the interval t , and the transition probability (i.e. the probability moving from one state to another state) between any two states at two times depends only on the difference between the states.

Before formally defining a counting process, first we need to define stochastic processes and point processes.

A stochastic process is a collection of random variables indexed by time, t , space, s or space-time, (s, t) , but we restrict ourselves to the time domain again and follow with Obral (2016) to define a stochastic process and a point process

Definition 2.1.1 (Stochastic Process) A stochastic process is a family of random variables indexed by time t and is defined as

$$\{X_t\}_{t \in T}.$$

Definition 2.1.2 (Point Process) Let $\{T_i\}_{i \in \mathbb{N}}$ be a sequence of non-negative random variables such that $T_i < T_{i+1} \forall i \in \mathbb{N}$, a point process on R^+ is defined as

$$\{T_i\}_{i \in \mathbb{N}}.$$

A point process relies on the occurrence of an event occurring at a specific point in time. Stochastic processes are more general. It can be related to a time interval, a waiting time, a state (e.g. blue or red) that changes over time, etc.

A counting process is then defined as follows

Definition 2.1.3 (Counting Process) Let $N(t)$ be the number of events up to some time t such that the values are nonnegative, integer valued, and nondecreasing, a stochastic process is said to be a counting process and is defined as

$$\{N(t), t \geq 0\}.$$

Let us look at a more explicit example, which we show in Figure 1. Suppose that $N(t)$ counts the number of events up to some time t and events occur at times $t = 0.1, 1, 1.5, 3, 5$, etc. then $N(2) = 3$ since events occurring at 0.01, 1, and 1.5 all occur in the time interval $(0, 2]$. Similarly, $N(4) = 4$ since 4 events occur in the time interval $(0, 4]$.

The equivalent definition of a counting process, which may be more beneficial as we get into other point processes in the later sections since it is easier to see that for $i \in \mathbb{N}$, if $T_i \leq t$, then the indicator function $I_{\{T_i \leq t\}}$ is equal to 1 and then we sum up all the 1s for events which have occurred, is

Definition 2.1.4 (Counting Process) Let $\{T_i\}_{i \in \mathbb{N}}$ be a point process, a counting process associated with $\{T_i\}_{i \in \mathbb{N}}$ is defined as

$$N(t) = \sum_{i \in \mathbb{N}} I_{\{T_i \leq t\}}.$$



Figure 1: Counting Process

A useful corollary of counting process, which describes more formally some of the properties we stated above and will be helpful to understand as we move into Poisson processes and then beyond, is

Corollary 2.1.1 A counting process satisfies that

1. $N(t) \geq 0$
2. $N(t)$ is an integer
3. If $t \leq t + h$, then $N(t) \leq N(t + h)$
4. If $t < t + h$, then $N(t + h) - N(t)$ is the number of events occur in the interval $(t, t + h]$.

In other words, 1. An event has to occur for it to be counted. 2. We either count an event or we don't. There is no event that results in decimal value. 3. Counts always increase. Once we observe an event and count it, it remains in the counts. 4. The number of events in specific time interval can be obtained by subtracting the number of events in previous interval from that in the current interval.

2.2 Poisson Process

The homogeneous Poisson process (HPP) is one of the simplest yet most-widely used point processes (Baddeley et al., 2015). HPPs can be used to model the number of events such as bus arrivals at a bus stop, car accidents at a site, or the document requests on a web server over time. As we alluded to previously with counting processes, HPPs can also be considered over a space which is often taken to be a two-dimensional plane, such as the surface of Earth, or a three-dimensional volume, such as the interior of Earth.

Like counting processes, HPPs are also independent, stationary, and homogeneous. We also assume that the numbers of events, $N(t)$, follow a Poisson distribution with a constant rate λ and the interarrival times between events, W , are exponentially distributed. We follow with Obral (2016) and Chen (2016) to formally define a Poisson process

Definition 2.2.1 (Poisson Process) If the following conditions hold, a counting process $\{N(t), t \geq 0\}$ is said to be a Poisson Process with constant rate (or intensity) $\lambda > 0$

1. $N(0) = 0$
2. $N(t)$ has independent increments
3. $P(N(t+h) - N(t) = 1) = \lambda h + o(h)$
4. $P(N(t+h) - N(t) > 1) = o(h)$,

where the function of little o, $o(h)$, is given as

$$\lim_{h \rightarrow 0^+} \frac{o(h)}{h} = 0.$$

In other words, 1. An event has to occur for it to be counted. 2. For any disjoint time intervals, the occurrence of an event does not affect the probability of the occurrence of one another event. 3. λ is the rate (i.e. events over time) at which points occur and is constant. 4. No more than 1 event can occur at the same location.

An alternative way to think of HPP is that it is a uniformly random process. If we were to take a realization of a HPP over some time interval $(0, T]$ and ‘bin’ the number of events occurring in some set of equal intervals, then the histogram for the realization would resemble a realization of a uniform distribution over time 0 to T.

Let us look at a realization of a HPP in time, which we show in Figure 2. First, we note that the cumulative number of points is growing at a constant linear rate. We can also see that the histogram of rate appears roughly uniform, i.e., the rates are roughly constant at $\lambda = 10$. The algorithm used for simulating this HPP can be found in the Appendix section.

For an HPP, the numbers of events in any time interval $N(t)$ are Poisson distributed. More formally, we can say that the number of events in any time interval $(t, t+h]$, $N((t, t+h])$, $\sim Pos(\lambda \cdot h)$. That is, for all $t, h \geq 0$ and $n = 0, 1, \dots$,

$$P(N(t+h) - N(t) = n) = \frac{(\lambda h)^n e^{-\lambda h}}{n!}.$$



Figure 2: Homogeneous Poisson process (rate = 10)

Let us also denote the interarrival times between events as W . For example, let T_0 be the starting time of the process while T_1 the time of the first occurrence of event and T_2 the time of the second occurrence of event, then the elapsed time between the start of the process and first event is W_1 and the elapsed time between the first event and second event is W_2 . The interarrival times W are exponentially distributed. More formally, the interarrival times $W, \sim \exp(\frac{1}{\lambda})$. That is, for rate $\lambda > 0$, the interarrival time W_i $i = 1, 2, \dots$,

$$P(W_1 > h) = P(N(h) = 0) = e^{-\lambda h}.$$

This is because the probability of the first point arriving after time h can be thought of as the probability that the first point does *not* arrive in the time interval $(0, h]$. Similarly, W_2 is also $\sim \exp(\frac{1}{\lambda})$ since

$$P(W_2 > h | W_1 = t) = P(N(t+h) - N(t) | N(t) - N(t^-) = 1) = P(N(t+h) - N(t) = 0) = P(N(h) = 0) = e^{-\lambda h}.$$

2.3 Nonhomogeneous Poisson Process

Assuming that the rate in which points occur is constant is often not realistic in practice. We may want a model that allows for more flexibility. The nonhomogeneous Poisson processes (NPPs) are a generalization of homogeneous Poisson processes that allow for the rate (or intensity) λ to vary as function of time t or space s . We previously assumed that for the HPP the intensity λ is constant. If we have reasons to believe that the intensity is not constant, then we should model as a NPP instead. This would be the case if, as in the supermarket example, we have reasons to believe that the arrival rate of customers is higher during

lunch time as compared to say, 2am, or as in the trees in a forest example, we speculate that environmental factors such as temperature and rainfall affect the spatial distribution of the trees.

NPPs are independent but not stationary nor homogeneous. Recall that HPP has stationary increments since the distribution of the numbers of events $N(t)$ that occur in any interval of time t depends only on the length of the interval t but not the location of the interval t , NPP, in contrast, does not have stationary increments since the distribution of $N(t)$ can change when shifted in t . Since stationary implies homogeneity, NPP is nonhomogeneous.

Like HPP, we assume for NPP that $N(t)$ follow a Poisson distribution too but with an intensity function $\lambda(t)$ such that the intensity now varies with a function of time. This then leads to the following definition of a NPP (Obral, 2016; Chen, 2016)

Definition 2.3.1 (Nonhomogeneous Poisson Process) If the following conditions hold, a counting process $\{N(t), t \geq 0\}$ is said to be a nonhomogeneous Poisson Process with intensity function of time $\lambda(t), t > 0$

1. $N(0) = 0$
2. $N(t)$ has independent increments
3. $P(N(t+h) - N(t) = 1) = \lambda(t)h + o(h)$
4. $P(N(t+h) - N(t) > 1) = o(h)$.

NPPs have additional properties such as if the number of events in any time interval $(t, t+h]$, denoted as $N((t, t+h])$, $\sim \text{Pos}(\Lambda(t) = \int_t^{t+h} \lambda(v)dv)$. That is, for all $v, t, h \geq 0$ and $n = 0, 1, \dots$,

$$P(N(t+h) - N(t) = n) = \frac{(\int_t^{t+h} \lambda(v)dv)^n e^{-\int_t^{t+h} \lambda(v)dv}}{n!},$$

where $\lambda(v)$ again denotes a non-constant rate function.

Further, occurrence of the next point can be determined by utilizing the exponential distribution with

$$P(N(t, t+h] = 0) = e^{-\int_t^{t+h} \lambda(v)dv}.$$

A Motivating Example

Before we delve further into other point processes, let us look at a motivating example, which we show in Figure 3. We demonstrate HPP and NPP in space so that visualizations are easier to look at and comprehend.




Figure 3: Left: HPP (rate = 100) Right: NPP (intensity = $400xy$)

HPP in space is also called complete spatial randomness (CSR) (Baddeley et al., 2015). For HPP in space, the number of events in u with area $|u|$, denoted as $N(u)$, $\sim \text{Pos}(\lambda|u|)$. The left figure of Figure 3 is a realization of a HPP with constant $rate = 100$. HPP points appear uniformly distributed in u .

For NPP in space, the number of events in u , denoted as $N(u)$, $\sim \text{Pos}(\int_u \lambda(v)dv)$. The right figure of Figure 3 is a realization of a NPP with *intensity function* = $400xy$. NPP points are not uniformly distributed; they are distributed according to the intensity function of the process. In this example, the points appear to concentrate at the upper-right corner.

2.4 Cox and Cluster Process

Even more flexible models than NPP are Cox and cluster processes. Whereas, previously assumed independence between events which occur at a constant rate λ for the HPP or were independent but depend on an intensity function $\lambda(t)$ for the NPP, we now discuss models that allow for the relaxation of this independence assumption. Additionally, Cox and cluster processes are mainly spatial processes so we are in the space domain.

Examples that are potentially better modelled as Cox processes than HPP include locations of emergent bramble cane (blackberry) plants and *Beilschmiedia* trees (Baddeley et al., 2015). In these examples, there appears to be some observable or unobservable spatial covariate (e.g. light, humidity, soil quality) or external factor that makes it more likely to observe the plants or trees preferentially in some areas as opposed to others.

Cox process (or doubly stochastic Poisson processes) can be defined as a Poisson process with a random intensity function; if the intensity surface $\Lambda(u)$ is known, then it becomes a Poisson process with intensity function $\Lambda(u)$ (Baddeley et al., 2015). This spatially varying intensity function of the Cox process is random since it relies on a set of random variables called random field to capture the covariates or external factors

(Baddeley et al., 2015). When the intensity is relatively high in some area of space, points occur more frequently. When the intensity is low, fewer points occur. Since the intensity is random and changes based on a set of observed or potentially unobserved set of random variables, this accumulation of points sometimes appears to be clustered, and this is one way to observe the dependence structure of the Cox process. Within the general class of Cox processes, mixed Poisson process, the simplest example of all, involves generating a random variable Λ and, given the value of Λ , generating a Poisson process with intensity function Λ .

On the other hand, example that can be modelled as cluster processes includes seedlings and saplings of California redwood (Baddeley et al., 2015). In this example, unobserved ‘parent’ trees give rise to clusters of observed ‘offspring’ trees. It is also not hard to notice significant overlap between Cox processes and cluster processes since these processes tend to share a considerable amount of information in their constructions. While Cox processes can sometimes appear to exhibit clustering of points due to the underlying random field, cluster processes tend to make the clustering of points more explicit. For a cluster process, we first have some set of unobserved ‘parent’ points \mathbf{Y} generated by the process. Next, each ‘parent’ point $y_i \in \mathbf{Y}$ gives rise to a random number of ‘offspring’ points z_{ij} . These ‘offspring’ points Z_{ij} then form a cluster process \mathbf{X} around the set of parents \mathbf{Y} and only the offspring points are observed. Within the general class of cluster processes, there are numerous models which can be utilized depending on the choice of assumptions. Matern cluster processes, for one example, involves generating homogeneous Poisson parents and each parent gives rise to Poisson number of offspring uniformly distributed in a disc of radius r centered around the parent.

Alternatively, we can think of Cox process as a hierarchical model with two levels and cluster process as a hierarchical model with three levels (Geyer, 2020). For the Cox process, 1) there is some set of random variables that influences the intensity function and 2) based on this intensity function, we observe the set of points. For the cluster process, 1) there is some random or non-random intensity function, 2) based on the intensity function, some set of ‘parent’ (or ‘center’) points are laid down which we often don’t observe, but 3) given the location of the ‘parent’ points, some set of ‘offspring’ points are generated around the ‘parent’ points which we observe.

The left and right figure of Figure 4 is a realization of a mixed Poisson process with *intensity function* = $\exp(1, 1/100)$ and a Matern cluster process with $kappa = 20, r = 0.05, mu = 5$, respectively. Both Cox process and Matern cluster process points appear clustered, but the way the points cluster differ. Points in Cox process cluster accordingly to some specified distribution (i.e. a exponential distribution), whereas points in cluster process cluster in some defined area (i.e. a disc). The **spatstat** package of R used for simulating this HPP can be found in the Appendix section (Baddeley & Turner, 2005).

Having given some background on both the Cox and cluster processes, we follow with Coeurjolly (2015) and




Figure 4: Left: Cox (intensity = $\exp(n = 1, \text{rate} = 1/100)$) Right: Matern ($\kappa = 20, r = 0.05, \mu = 5$)

Daley & Vere-Jones (2003) now with their more formal definitions

Definition 2.4.1 (Cox Process) Let $\Lambda = (\Lambda(u))_{u \in S \subseteq \mathbb{R}^d}$ be a non-negative random field such that the values of $\Lambda(u)$ is non-negative and $\Lambda(u)$ is a locally integrable function, where Λ is a random field means that $\Lambda(u)$ is a random variable $\forall u \in S$ and $\Lambda(u)$ is a locally integrable function means that $E(\Lambda(u))$ exists and is locally integrable with probability 1, if $X \mid \Lambda \sim \text{Pos}(\Lambda)$, then X is said to be a Cox process driven by Λ with intensity function $\lambda(u) = E(\Lambda(u))$. That is,

$$P(N(u) = n) = \frac{(\lambda(u))^n e^{-\lambda(u)}}{n!} = \frac{(E(\Lambda(u)))^n e^{-E(\Lambda(u))}}{n!},$$

where

$$\Lambda(u) \stackrel{a.s.}{=} \int_u^\infty \lambda(x) dx = \int_0^\infty \frac{x^n e^{-x} F_u(dx)}{n!}.$$

Definition 2.4.2 (Cluster Process) Let x ('parent' points) be points in a point process N and replacing every x with a cluster of points N_x ('offspring' point) centered at x and assume also that each N_x is independent of one another, then the union of all the clusters forms a cluster process N_c . That is,

$$N_c = \bigcup_{x \in N} N_x.$$

Next, we follow with Baddeley et al. (2015) to state the following model assumptions for cluster processes

1. 'Parent' points follow a Poisson distribution.
2. Clusters are independent of one another.
3. Clusters are identically distributed, which means that clusters, when shifted, have the same distributions.

4. The locations of ‘offspring’ points of each parent point are independently and identically distributed.
5. The number of ‘offspring’ points of each parent point follows a Poisson distribution.
6. Clusters are isotropic, which means that the distribution of ‘offspring’ points for each parent point depends only on the distance between the ‘parent’ and the ‘offspring’.

Under assumption 1 - 4, we have a Neyman-Scott process. Under assumption 1 - 5, the cluster process is a Cox process. And finally, under assumption 1 - 6, we have a Matern or Thomas cluster process.

2.5 Hawkes Process

The final class of point process models we present are the Hawkes processes. Hawkes processes are also commonly known as self-exciting point processes (SEPP). Like Cox and cluster process, the Hawkes process model allows for dependence between events; however, their dependence differs. In Hawkes processes, the occurrence rate of the events depends not only on time t but also past events \mathcal{H}_t up to some time t . This is the distinguishing feature of the Hawkes process as neither Cox nor cluster processes depends on the past history of events, which is also what makes Hawkes processes ‘self-excite’. Hawkes processes can be powerful predictive models as they naturally capture triggering and clustering behavior (Reinhart, 2018).

As previously mentioned, examples of applications for Hawkes processes include locations of earthquake epicenters, locations of crimes, and locations of patients with a communicable disease. The major sharing feature in each of these examples is that an occurrence of an event leads to an increase in the occurrence of subsequent events (i.e. ‘self-exciting’) in nearby time and/or space. In seismology, a main quake is often followed by aftershocks. In gang crimes, one act of violence often features an act of retaliatory violence. In communicable diseases, one patient may infect others.

Since the intensity of the process is now as a function of past history, we refer to the expected rate as a conditional intensity function. Hawkes processes self-excite because of this conditional intensity function since conditioning on the past set of events, the expected rates at which points occur are expected to be higher when points have occurred in nearby time and/or space and then they gradually decline as the events get further away. This ‘self-exciting’ characteristic is captured by the summing part of the conditional intensity function, which we show in Definition 2.5.2.

Another characteristic is that more recent events exert more influence on the intensity but the intensity will fade out or decay until the next event. This ‘decaying’ characteristic is captured by the triggering part of the conditional intensity function. The triggering function is also what makes the Hawkes process

a self-exciting as well as cluster process as it allows for additional points to occur in nearby time and/or space before ultimately decaying back to the background rate. For example, an earthquake happens today, as supposed to more than a year ago, may lead us believe that there is a higher chance of another quake that will occur tomorrow.

A Hawkes process can be uniquely identified through its conditional intensity function (Daley & Vere-Jones, 2003) and the formal definition of which is

Definition 2.5.1 (Conditional Intensity Function) Let $N(t)$ be the numbers of events $N(t)$ that occur in any interval of time t , the conditional intensity function $\lambda(t)$ with respect to the history of the process up to time t , \mathcal{H}_t , is defined as

$$\lambda(t|\mathcal{H}_t) = \lim_{h \rightarrow 0^+} \frac{E(N(t, t+h)|\mathcal{H}_t)}{h},$$

where \mathcal{H}_t is the history prior to time t .

Having defined the conditional intensity function, we next more formally define the Hawkes process (Rizoiu, 2017; Reinhart, 2018)

Definition 2.5.2 (Hawkes Process) A counting process $\{N(t), t \geq 0\}$ associated with past events $\{\mathcal{H}_t, t > 0\}$ is said to be a Hawkes process with conditional intensity function $\lambda(t|\mathcal{H}_t), t > 0$ and takes the form

$$\lambda(t|\mathcal{H}_t) = \lambda_0(t) + \sum_{i: T_i < t} \phi(t - T_i),$$

where $\lambda_0(t)$ (or μ) is the the constant background rate, $T_i < t$ are the events time occur before current time t , $\phi(\cdot)$ is the kernel function (or $g(\cdot)$ the triggering function) through which intensity function depends on past events, and \mathcal{H}_t is the past history which represents the internal history of N up to time t .

Typical choices of $\phi(\cdot)$ include, for example, exponentially decaying function and power-law kernel (Rizoiu, 2017) and they take the forms of

$$\phi(x) = \alpha e^{-\beta x}$$

and

$$\phi(x) = \frac{\alpha}{(x + \beta)^{\eta+1}}.$$

Exponentially decaying function, which decays faster, has been applied to financial data (Embrechts et al., 2011). On the other hand, power-law kernel, which has a flatter tail, has been used in seismology (Ozaki, 1979) and social media (Rizoiu et al., 2017).




Figure 5: Hawkes Process

In Figure 5, we show a realization of a Hawkes process with the exponentially decaying triggering function ($\mu = 0.5, \alpha = 0.7, \beta = 0.5$). We can see that each time the event arrives, the intensity (conditional intensity function) increases and then it declines back to the background rate. When the next event arrives, the intensity jumps again and then declines. The intensity for this realization gradually increases over time too since the points of this simulation occur fairly regularly. The algorithm used for simulating this Hawkes process can be found in the Appendix section.

2.6 Spatio-Temporal Self-Exciting Point Process

Spatio-temporal SEPP are an extension of temporal Hawkes processes. Recall that the conditional intensity function for Hawkes processes take the form of

$$\lambda(t|\mathcal{H}_t) = \mu + \sum_{i:T_i < t} g(t - t_i),$$

the conditional intensity function for SEPP take the form of

$$\lambda(t|\mathcal{H}_t) = \mu(s) + \sum_{i:T_i < t} g(s - s_i, t - t_i),$$

where $s_i, i = 1, 2, \dots$ are the sequence of locations of events and $t_i, i = 1, 2, \dots$ are the times of events. The triggering function is defined to be separable in space and time for simplicity. In addition, the background rate here can be taken to be a function of space, which is often a NPP or constant. Further, the triggering function here is now a function of space and time and, as a result, events that are closer to some given spatial

and temporal location are more heavily influenced by the points that occur nearby in both space and time.

3 Conclusions and Discussion

In this project, we first introduce the counting process, which can be used to count the number of events over space, time, etc. Next, we introduce the HPP, which is one of the simplest yet most widely used point processes for modeling the occurrence of events with constant arrival rate. Then, we relax the stationarity and homogeneity assumption to talk about the NPP. Allowing the intensity to vary with space or time makes the model more realistic in practice. We follow up with the HPP and NPP in space for the purpose of comparison. We further relax the independence assumption and introduce the Cox and cluster processes. Allowing the events to be dependent makes the models even more flexible, as a lot of the environmental and ecological applications rely on the dependence on covariates and/or external factors. Finally, we get to the Hawkes and SEPP. With the intensity now depending on past history, the models become a suitable choice for capturing events that exhibit both triggering and clustering behavior while having the potential to become a powerful predictive tool for a wide variety of applications. At this point, interested readers should have the background knowledge to comprehend Hawkes and SEPP through additional, existing literature and explore areas for future work such as applications for larger dataset, residual and model diagnostics, etc. On my end, I really enjoy this month-long journey of learning the various point processes as well as witnessing their many practical and important uses for real-world data.

Acknowledgments

I want to thank Dr. James Molyneux for his constant support and encouragement while advising this Master's project, Dr. Lisa Madsen & Dr. Charlotte Wickham for their willingness to co-advise as I press on for the PhD program here at OSU, Dr. Sarah Emerson for both academic and nonacademic support that played major roles in my decision to pursue the PhD here at OSU, my cohort for summer comp (comprehensive exam) studies, theories HW and various other group projects, our previous cohort and other PhD students for reassuring us that we will make it too despite the ongoing challenges, my family and friends for keeping me sane while being remote, and my 92-year-old grandma for hanging in there while battling brain tumor since last year. But no thanks, COVID-19 pandemic, 2020 Oregon wildfires and the social unrest that sparked the Black Lives Matter & Stop Asian Hate movement.

Reference

- Bacry, E., Mastromatteo, I., & Muzy, J. F. (2015). Hawkes processes in finance. *Market Microstructure and Liquidity*, 1(01), 1550005.
- Baddeley, A., Rubak, E., & Turner, R. (2015). *Spatial point patterns: methodology and applications with R*. CRC press.
- Bauwens, L., & Hautsch, N. (2009). Modelling financial high frequency data using point processes. In *Handbook of financial time series* (pp. 953-979). Springer, Berlin, Heidelberg. Boyd, P., & Molyneux, J. (2021). Assessing the contagiousness of mass shootings with nonparametric Hawkes processes. *PLoS one*, 16(3), e0248437.
- Chen, Y. (2016). *Thinning algorithms for simulating point processes*. Florida State University, Tallahassee, FL.
- Chiang, W. H., Liu, X., & Mohler, G. (2020). Hawkes process modeling of COVID-19 with mobility leading indicators and spatial covariates. *medRxiv*.
- Coeurjolly, Jean-François. (2015). Lecture 4: Models for point processes. Retrieved from https://www-ljk.imag.fr/membres/Jean-Francois.Coeurjolly/documents/M2R/models_cours.pdf
- Cressie, N. (2015). *Statistics for spatial data*. John Wiley & Sons.
- Daley, D. J., & Vere-Jones, D. (2003). *An introduction to the theory of point processes: volume I: elementary theory and methods*. Springer New York.
- Embrechts, P., Liniger, T., & Lin, L. (2011). Multivariate Hawkes processes: an application to financial data. *Journal of Applied Probability*, 48(A), 367-378.
- Gerhard, F., Deger, M., & Truccolo, W. (2017). On the stability and dynamics of stochastic spiking neuron models: Nonlinear Hawkes process and point process GLMs. *PLoS computational biology*, 13(2), e1005390.
- Geyer, C. (2020). *Spatial Point Processes*. Retrieved from <https://www.stat.umn.edu/geyer/8501/>
- Holbrook, A. J., Loeffler, C. E., Flaxman, S. R., & Suchard, M. A. (2021). Scalable Bayesian inference for self-excitatory stochastic processes applied to big American gunfire data. *Statistics and Computing*, 31(1), 1-15.
- Johnson, D. H. (1996). Point process models of single-neuron discharges. *Journal of computational neuroscience*, 3(4), 275-299.

- Krishna, R. (2015). Simulation of Non-Homogeneous Poisson Processes. Retrieved from <https://radhakrishna.typepad.com/simulating-nonhomogeneous-poisson-process.pdf>
- Meyer, S., Elias, J., & Höhle, M. (2012). A space–time conditional intensity model for invasive meningococcal disease occurrence. *Biometrics*, 68(2), 607-616.
- Meyer, S., & Held, L. (2014). Power-law models for infectious disease spread. *Annals of Applied Statistics*, 8(3), 1612-1639.
- Mohler, G. O., Short, M. B., Brantingham, P. J., Schoenberg, F. P., & Tita, G. E. (2011). Self-exciting point process modeling of crime. *Journal of the American Statistical Association*, 106(493), 100-108.
- Obral, K. (2016). Simulation, estimation and applications of hawkes processes. (Master’s thesis, University of Minnesota).
- Ogata, Y. (1988). Statistical models for earthquake occurrences and residual analysis for point processes. *Journal of the American Statistical association*, 83(401), 9-27.
- Ogata, Y. (1998). Space-time point-process models for earthquake occurrences. *Annals of the Institute of Statistical Mathematics*, 50(2), 379-402.
- Ozaki, T. (1979). Maximum likelihood estimation of Hawkes’ self-exciting point processes. *Annals of the Institute of Statistical Mathematics*, 31(1), 145-155.
- Park, J., Schoenberg, F. P., Bertozzi, A. L., & Brantingham, P. J. (2021). Investigating Clustering and Violence Interruption in Gang-Related Violent Crime Data Using Spatial–Temporal Point Processes With Covariates. *Journal of the American Statistical Association*, 1-14.
- Peng, R. D., Schoenberg, F. P., & Woods, J. A. (2005). A space–time conditional intensity model for evaluating a wildfire hazard index. *Journal of the American Statistical Association*, 100(469), 26-35.
- Reinhart, A. (2018). A review of self-exciting spatio-temporal point processes and their applications. *Statistical Science*, 33(3), 299-318.
- Reinhart, A., & Greenhouse, J. (2018). Self-exciting point processes with spatial covariates: modeling the dynamics of crime. *arXiv preprint arXiv:1708.03579*.
- Rizoiu, M. A., Lee, Y., Mishra, S., & Xie, L. (2017). A tutorial on hawkes processes for events in social media. *arXiv preprint arXiv:1708.06401*.
- Rubak, E. (2011a). Spatial point processes: Theory and practice illustrated with R. Retrieved from <https://www-ljk.imag.fr/membres/Jean-Francois.Coeurjolly/documents/lecture1.pdf>

- Rubak, E. (2011b). Spatial point processes: Theory and practice illustrated with R. Retrieved from <https://www-ljk.imag.fr/membres/Jean-Francois.Coeurjolly/documents/lecture2.pdf>
- Rubak, E. (2011c). Spatial point processes: Theory and practice illustrated with R. Retrieved from <https://www-ljk.imag.fr/membres/Jean-Francois.Coeurjolly/documents/lecture3.pdf>
- Rizoiu, M. A., Lee, Y., Mishra, S., & Xie, L. (2017). A tutorial on hawkes processes for events in social media. arXiv preprint arXiv:1708.06401.
- Pasupathy, R. (2010). Generating homogeneous poisson processes. Wiley encyclopedia of operations research and management science.
- Schoenberg, F. P. (2010). Introduction to point processes. Wiley Encyclopedia of Operations Research and Management Science.
- Stabile, G., & Torrisi, G. L. (2010). Risk processes with non-stationary Hawkes claims arrivals. *Methodology and Computing in Applied Probability*, 12(3), 415-429.
- Zhao, Q., Erdogdu, M. A., He, H. Y., Rajaraman, A., & Leskovec, J. (2015, August). Seismic: A self-exciting point process model for predicting tweet popularity. In *Proceedings of the 21th ACM SIGKDD international conference on knowledge discovery and data mining* (pp. 1513-1522).