

Reparameterized SGLMM (Spatial Generalized Linear Mixed Model): A Review

Frances Lin

June 2022

Background and Introduction

The SGLMM (spatial generalized linear mixed model)

- ▶ is a hierarchical model that introduces spatial dependence through a GMRF (Gaussian Markov random field) (Besag et al., 1991).
- ▶ was initially proposed for prediction for count data, but
- ▶ has later been applied to estimation and prediction for other types of data (e.g. binary) and
- ▶ found applications in many fields (e.g. ecology, geology, epidemiology, image analysis, and forestry).

Background and Introduction

The SGLMM has been the dominant model for areal data because of

- ▶ its flexible hierarchical specification,
- ▶ the availability of the software WinBUGS for (Bayesian) data analysis (Lunn et al., 2000) and
- ▶ various theoretical and computational advantages over the competitive model named automodel.

However, SGLMMs suffer from two major shortcomings:

- i) variance inflation due to spatial confounding and
- ii) computational challenges posed by high dimensional latent variables (“random effects”).

Background and Introduction

On the other hand, while another model named RHS model

- ▶ seeks to alleviate confounding (Reich et al., 2006),
- ▶ can result in random-effects structure with negative spatial dependence (i.e. repulsion) that is not typically applied in practice.

A reparameterized model is proposed, and it is able to

- ▶ alleviate confounding, and, at the same time,
- ▶ include patterns of positive spatial dependence (i.e. attraction) while excluding patterns of repulsion.

The proposed model is also one of the first dimension reduction techniques for spatial areal models.

Outline

- ▶ Traditional SGLMM (Spatial Generalized Linear Mixed Model)
- ▶ Spatial confounding (via the automodel)
- ▶ RHZ model
- ▶ Sparse reparameterization of the areal SGLMM
- ▶

Comparison of various SGLMMs

Table 1: Comparison of various SGLMMs

Model	Confounded	Account_G
Traditional SGLMM	Yes	No
Uncentred automodel	Yes	Yes
Centred automodel	No	Yes
RHZ SGLMM	No	No
Sparse SGLMM	No	Yes

Traditional SGLMM

Let

$$G = (V, E)$$

be an undirected, labelled graph, where $V = \{1, 2, \dots, n\}$ is a set of vertices (nodes) and $E = \{i, j\}$ is a set of edges, where $i, j \in V$, $i \neq j$.

- ▶ Each vertex represents an area of interest and each edge represents the proximity of areas i and j .
- ▶ The graph G is represented using an adjacency matrix A , which is a $n \times n$ matrix with $\text{diag}(A) = 0$ and entries $A_{ij} = 1\{(i, j) \in E, i \neq j\}$, where $1(\cdot)$ is an indicator function (i.e. entry $a_{ij} = 1$ if vertex v_i and v_j are adjacent. $a_{ij} = 0$, otherwise).

Traditional SGLMM

Further let $Z = (Z_1, \dots, Z_n)^T$ be the random field of interest, where Z_i is the random variable associated with vertex i . Then, the first stage of the model is given by

$$g(E(Z_i|\beta, W_i)) = X_i\beta + W_i, \quad (1)$$

where

- ▶ g is a link function,
- ▶ X_i is the i th row of the design matrix X ,
- ▶ β is a p -vector of regression parameters and
- ▶ W_i is a spatial random effect associated with vertex i .

Different types of data require different canonical choices of the link function g (e.g. the logit function for spatial binary data and the logarithm function for spatial count data).

GMRF prior for the random effects

The field of random effects $W = (W_1, \dots, W_n)^T$, through which spatial dependence is incorporated, is assumed to follow the intrinsic conditionally autoregressive or improper GMRF prior

$$p(W|\tau) \propto \tau^{\text{rank}(Q)/2} \exp\left(-\frac{\tau}{2} W^T Q W\right), \quad (2)$$

where τ is a smoothing (“precision”) parameter and $Q = \text{diag}(A1) - A$ is a precision matrix. The precision matrix Q incorporates both dependence and prior uncertainty.

Note. A SGLMM

1. can be reformatted by replacing GMRF with GP (Gaussian process) for point-referenced data (or geostatistical) data.
2. is restricted to a Bayesian or restricted maximum likelihood (REML) analysis since the prior (2) is improper.

Spatial confounding (via the automodel)

The automodel, SGLMM's closest competitor, is a Markov random-field model that incorporates dependence directly and is defined as

$$g(E(Z_i|\beta, \eta, Z_{-i})) = X_i\beta + \eta \sum_{(i,j) \in E} Z_j^*, \quad (3)$$

where

- ▶ g is a (canonical) link function,
- ▶ η is the dependence parameter ($\eta > 0$ implies attraction; $\eta < 0$ implies repulsion), and
- ▶ Z_{-i} is the field excluding the i th observation.

Spatial confounding (via the automodel)

- ▶ For the centred automodel, $Z_j^* = Z_j - \mu_j$, where μ_j is the independence expectation of Z_j (i.e. $\mu_j = E(Z_j|\beta, \eta = 0) = g^{-1}(X_j\beta)$).
- ▶ For the uncentred automodel, $Z_j^* = Z_j$.

The sum term $\sum_{(i,j) \in E} Z_j^*$ is called the autocovariate, and it is considered as a synthetic predictor.

- ▶ The centred autocovariate makes the dependence parameter easily interpretable (η captures the relativity of an observation to its neighbours, conditional on the hypothesized regression component).
- ▶ The uncentred autocovariate not only poses conceptual challenge but also shows spatial confounding.

The RHS model

The traditional SGLMM can also be shown to be confounded.

Consider P be a projection onto $C(X)$

$$P = X(X^T X)^{-1} X^T$$

and let P^\perp be the projection onto $C(X)$'s complement such that $P^\perp = I - P$, then equation (1) can be rewritten as

$$g(E(Z_i|\beta, W_i)) = X_i\beta + K_i\gamma + L_i\delta,$$

where K and L are orthogonal bases for $C(X)$ and $C(X)^\perp$ respectively and γ and δ are random coefficients.

K and X now share the same column space, and this is the source of the spatial confounding.

The RHS model

Since K has no practical meaning, setting $\gamma = 0$, the first stage of the RHS model is given by

$$g(E(Z_i|\beta, \delta)) = X_i\beta + L_i\delta,$$

and, compared to equation (2), the prior for the random effects δ becomes

$$p(\delta|\tau) \propto \tau^{(n-p)/2} \exp\left(-\frac{\tau}{2} \delta^T Q_R \delta\right),$$

where $Q_R = L^T Q L$.

The sparse, reparameterized areal SGLMM

However, the RHZ model does not allow parsimonious fitting of the residual clustering.

- ▶ The geometry corresponding to the projection P^\perp fails to account for the underlying graph G , thus permitting structure of negative spatial dependence (i.e. repulsion) in the random effects.
- ▶ This is not useful in practice since neighboring observation tends to be similar, rather than dissimilar.

The proposed reparameterized model

- i) considers an alternative projection that captures the geometry of the models, thus allowing *only* patterns of positive spatial dependence (i.e. attraction).
- ii) utilizes the geometry of the models, which leads to dimension reduction of the random effects naturally.

The sparse, reparameterized areal SGLMM

Consider the operator $(I - 11^T/n)A(I - 11^T/n)$ that appears in the numerator (top) of Moran's I -statistic (a commonly used for nonparametric method for spatial dependence)

$$I(A) = \frac{n}{1^T A 1} \frac{Z^T (I - 11^T/n) A (I - 11^T/n) Z}{Z^T (I - 11^T/n) Z},$$

where I is a $n \times n$ identity matrix and 1 is a n -vector of 1s.

Next replace $I - 11^T/n$ with P^\perp , then the resulting operator called the Moran operator for X with respect to G , $P^\perp A P^\perp$, appears in the numerator of the generalized Moran's I -statistic

$$I_x(A) = \frac{n}{1^T A 1} \frac{Z^T P^\perp A P^\perp Z}{Z^T P^\perp Z}.$$

The sparse, reparameterized areal SGLMM

The sparse, reparameterized areal SGLMM

Replace L with M in the RHS model, then the first stage of the reparameterized model is given by

$$g(E(Z_i|\beta, \delta_S)) = X_i\beta + M_i\delta_S,$$

and the prior for the random effects δ_S becomes

$$p(\delta_S|\tau) \propto \tau^{q/2} \exp(-\frac{\tau}{2} \delta_S^T Q_S \delta_S),$$

where $Q_S = M^T Q M$.

Dimension reduction for spatial models