# Feasibility signals compared with what happened in the case-study trial

| Planning question | Feasibility assessment | What subsequently occurred |
|---|---|---|
| **Is recruitment demanding?** | Planned 454/year = 92nd percentile of comparable trials; 8% chance of finishing in 10 months | 377 randomised in 10.2 months (about 446/year): upper-tail performance was achieved |
| **Are the variability assumptions realistic?** | Comparable trials: CV 37% (AUC), 40% (Ctrough) Protocol assumed 50% and 84% | Observed CV 36% and 44% |
| **What safety burden should sites expect?** | Historical arm-level estimate 39.5%; simulated cohort incidence 39.7% (150 of 378) | 39.0% / 40.5% (subcutaneous / intravenous) |
| **Is the population composition fixed?** | Simulated eligible pool: 40% women, 15% Asian, 5% Hispanic (one global mix) | 29% women, 29% Asian, 31% Hispanic: recruitment geography shaped the mix |
| **Will the required PK numbers be retained?** | P(AUC 0-6 weeks ≥ 318) 100.0%; P(Ctrough ≥ 240) 96.6% at 378 enrolled | 371 / 303 evaluable (AUC / Ctrough) |

## Synthetic-control exploration: control arm (95% interval)

Feasible to generate; too uncertain to replace the control arm

