Feasibility signals compared with what happened in the case-study trial

Planning question	Feasibility assessment	What subsequently occurred
Is recruitment demanding?	Planned 454/year = 92nd percentile of comparable trials; 8% chance of finishing in 10 months	377 randomised in 10.2 months (about 446/year): upper-tail performance was achieved
Are the variability assumptions realistic?	Comparable trials: CV 37% (AUC), 40% (Ctrough) Implied by the protocol's power statement: CV 50% (AUC), 84% (Ctrough)	Observed CV 36% and 44%
What safety burden should sites expect?	Historical arm-level estimate 39.5%; simulated cohort incidence 39.7% (150 of 378)	39.0% / 40.5% (subcutaneous / intravenous)
Is the population composition fixed?	Simulated eligible pool: 40% women, 15% Asian, 5% Hispanic (one global mix)	29% women, 29% Asian, 31% Hispanic: recruited population differed from the modelled global mix
Will the required PK numbers be retained?	P(AUC 0-6 weeks ≥ 318) 100.0%; P(Ctrough ≥ 240) 96.6% at 378 enrolled	371 / 303 evaluable (AUC / Ctrough)

Synthetic-control exploration: control arm (95% interval)

Feasible to generate; too uncertain to replace the control arm

