Desenvolvimento de módulo de Análise de Dados para identificação de situações de fraude e corrupção em processos de contratação pública

Francisco Valente Pereira


Master's in Computational Statistics and Data Analysis Departament of Mathematics Faculty of Sciences of the University of Porto 2024

Supervisor

Prof. Dr. Sílvio Gama Faculty of Sciences of the University of Porto

Co-supervisor

Prof. Dr. Margarida Brito Faculty of Sciences of the University of Porto





Todas as correções determinadas pelo júri, e só essas, foram efetuadas.

O Presidente do Júri,



Francisco Valente Pereira

Desenvolvimento de módulo de Análise de Dados para identificação de situações de fraude e corrupção em processos de contratação pública



Supervisor: Prof. Dr. Sílvio Gama Co-Supervisor: Prof. Dr. Margarida Brito

Department of Mathematics
Faculty of Sciences of the University of Porto
2024

Acknowledgements

Resumo

Abstract

Motivation

Table of Contents

A	cknov	vledgements	vii
Re	esumo		ix
Al	bstrac	et e e e e e e e e e e e e e e e e e e	xi
M	otivat	cion Company of the C	xiii
Ta	ble of	f Contents	xv
Li	st of l	Figures	xvii
Li	st of [Tables	xix
Li	st of A	Abbreviations	xxi
1	Intr	oduction	1
	1.1	Intership	1
		1.1.1 Something else	1
2	Con	clusions and Future Work	3
	2.1	Conclusions	3
	2.2	Future Work	4
A			5
Bi	bliogi	raphy	5
A 1	DDEN	DICES	_

List of Figures

List of Tables

List of Abbreviations

CPV Common Procurement Vocabulary

OCDS Open Contracting Data Standard

FCUP Faculdade Ciências da Universidade do Porto

CHAPTER 1

Introduction

1.1 Intership

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetuer id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

1.1.1 Something else

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

CHAPTER 2

Conclusions and Future Work

2.1 Conclusions

In this work, we synthesized $Ca_3Mn_{2-x}Ti_xO_7$ n=2 Ruddlesden-Popper samples with various compositions (x=0.1,0.25,0.3) via a common solid-state reaction method and studied its structural phase transitions and properties through different methods: X-Ray Powder Diffraction (XRPD), Raman Spectroscopy, Perturbed Angular Correlation (PAC), and Density Functional Theory (DFT) simulations, with special focus on the x=0.25 (C_2MTO) system.

After the last annealing of the synthesis process, Rietveld refinements on the XRPD data were performed to check the purity of the formed n=2 phases. We observed an approximate 12% presence of Ca₂Mn_{0.875}Ti_{0.125}O₄ (n=1 C₁MTO) in C₂MTO, however with a- and b-latticeparameters considerably lower (by $\approx 0.6 \text{Å}$) of what can be found in the literature for the undoped Ca₂MnO₄ system.

Raman and PAC spectroscopies, respectively in the 93-653K and 74-1224K temperature ranges, revealed a first order structural phase transition from what appears to be a the ferroelectric $A2_1am$ (s.g. 36) phase to an intermediate and paraelectric Acaa (s.g. 68) symmetry within a 300-550K temperature range, where both structural phases coexist. Thus, when comparing CMTO to the undoped $Ca_3Mn_2O_7$ (C_2MO), we report an increase in the percentage of the ferroelectric phase at room temperature by introducing the x=0.25 Ti⁴⁺ doping. However, our results point to the possibility of the ground state not corresponding to the $A2_1am$ symmetry, but to one with inequivalent A-sites at the rock-salt of the RP system. PAC spectroscopy also revealed a second order structural phase transition from the intermediate orthorhombic Acaa (s.g. 68) to the tetragonal I4/mmm (s.g. 139) at 1150K.

We came across an unexpected result at temperatures below 300K, where we clearly detected the probing of a second local environment, albeit without a clear physical origin. Interestingly, PAC measurements did not found evidence of the C_1MTO phase observed from XRPD refinements, and we believe this suggests that the $A2_1am$ space group might not correspond to the ground state symmetry for every composition of $Ca_3Mn_{2-x}Ti_xO_7$ mixed B-site systems. Phonon calculations on these doped systems in the $A2_1am$ symmetry could reveal if there are any lattice



instabilities, shedding a light on whether the $A2_1am$ still corresponds to the ground state when Ti is added.

The C₂MO system was studied with DFT in it's ground state $A2_1am$ symmetry. We determined a Hubbard-U correction by first principles ($U=6\,\mathrm{eV}$) to properly describe the Mn-3d states, and then performed a series of structural and electronic properties calculations with our ab-initio value and with U=0 and $U=3.9\,\mathrm{eV}$, a value commonly found in the literature. We determined the cell parameters and bulk moduli for the mentioned U values as well as multiple electronic properties, such as the projected density-of-states (PDOS), Bader charges, and the Electric Field Gradient at multiple sites. When we manage to define and adjust the most adequate parameters to compute the EFGs for this system, we intend to probe the Ti^{4+} doped $\mathrm{Ca_3Mn_{2-x}Ti_xO_7}$ structures, and analyse how the EFGs vary as a function of doping concentration.

2.2 Future Work

In the future, we intend to deepen the discussion at several stages of this system's characterization. Regarding the observation of a C₁MTO phase impurity in XRPD experiments, and the underestimated lattice parameters, we intend to pursue a new refinement considering also the intermediate *Acaa* phase. To complement this measurements, XAS and XPS spectroscopies would help us to confirm the oxidation states of Mn and Ti in the sample, to account for some deviations in the XRPD patterns. Additionally, neutron diffraction would confirm the structural and magnetic structural properties of these samples.

More complex Raman Spectroscopy setups could be pursued to study the evolution of the softer modes that drive the $A2_1am - Acaa$ structural phase transition, and DFT calculations could assign the peak positions for each mode in the doped C_2MTO structure.

The interpretation PAC results below 300K is still open to discussion, and we'll be pursing other methods to clarify them. Doped CMTO DFT calculations of the EFG parameters is a work in progress and the study of other doped systems is also scheduled, to later compare with undoped systems. In fact, $n = 1 \text{ Ca}_2(\text{Mn,Ti})\text{O}_4$ RP compounds have already been measured by PAC and will be analysed in the future. Future phonon calculations on these B-site mixed systems may become a key point in understanding these systems.

APPENDIX A

Add appendices.