

Journal notes on:

# Introduction to topological superconductivity

by Francisco Lobo



## CONTENTS

Acknowledgments	4
Preface	5
<b>I Overview of superconductivity theories</b>	<b>6</b>
I. London theory	6
A. London equations	6
B. London penetration depth and Meissner effect	6
C. London coherence length	7
II. Ginzburg-Landau theory	8
A. Superconductive order parameter	8
B. Ginzburg-Landau equations	10
C. Flux quantization	11
D. Type I and type II superconductors	12
E. Little-Parks experiment	13
F. Josephson effects	14
1. DC Josephson effect	15
2. AC Josephson effect	15
3. Inverse AC Josephson effect*	16
4. SQUIDs*	16
III. Bardeen–Cooper–Schrieffer theory*	17
A. Effective Hamiltonian	18
B. The BCS ground state	19
C. The gap equation	19
1. Cooper pair insight	20
2. Critical temperature	21
D. Spin-triplet $p$ -wave superconductivity	22
<b>II Introduction to topological superconductivity</b>	<b>22</b>
IV. Concepts of symmetry and topology	22
A. Introduction to topological invariants in 0D models	24
1. Time-reversal symmetry	24
2. Sublattice symmetry	25
3. Particle-hole symmetry	26
4. Combining symmetries	27
B. Introduction to topological invariant in higher dimensions	27
1. Aharonov-Bohm effect	29
2. Quantum Thouless pump	30

V.	Topological superconductivity in 1D models	30
A.	Kitaev model	30
1.	Majorana modes at a domain wall	34
2.	Kitaev ring	35
B.	SSH model	35
C.	Oreg-Lutchyn models	35
VI.	Topological superconductivity in 2D models	38
A.	Quantum hall effect	38
B.	Anomalous quantum Hall effect in the Haldane model	38
C.	Quantum spin Hall effect in the Kane-Mele model	39
D.	Integer quantum Hall effect	39
E.	Fractional quantum Hall effect	39
<b>III</b>	<b>Topics on topological superconductor theory</b>	39
VII.	Dirty superconductor theory	39
A.	Green's function formulation	39
1.	Abrikosov-Gorkov Green's Functions	39
2.	Eilenberger-Larkin-Ovchinnikov equations	39
3.	Usadel diffusion equation	39
VIII.	Self-consistent topological superconductivity	39
A.	Self-consistent Hartree-Fock-Bogoliubov mean-field theory*	39
1.	Mean-field self-energy in Nambu space	41
2.	Orbital-dependent interactions	45
3.	Hubbard model	47
4.	The gap equation accounting for Zeeman effect	48
B.	Self-consistent Majorana nanowire	48
C.	Self-consistent Bernal graphene	48
<b>IV</b>	<b>Appendices</b>	48
D.	Peierls substitution	48
E.	Overview of simpler systems	49
1.	Linear lattice	49
2.	Square lattice	49
F.	Overview of graphene systems	49
1.	Monolayer graphene	49
2.	Bilayer Bernal graphene	52
3.	Twisted bilayer graphene	54
4.	Kekulé modulation	54
<b>V</b>	<b>Bibliography</b>	55

## ACKNOWLEDGMENTS

My personal website is at <https://franciscolobo1880.github.io/>.

As supplementary material, there is a GitHub repository at <https://github.com/franciscolobo1880/topoSC> where you can check the code that generate the figures of the various models. This is done in *Julia* using the *Quantica.jl* package by Pablo San-Jose, my PhD advisor. Check *Quantica.jl*'s repository and it's tutorial at <https://github.com/pablosanjose/Quantica.jl>.

I thank Pablo San-Jose and Elsa Prada for their tutoring. I thank César Robles, Carlos Paya and Tiago Antão for their help and useful discussions.

By the very nature of being personal journal notes, the sections are never truly finished and being worked one passively. However, really unfinished sections that are not mostly blank are denoted with an asterisk \*. I would refrain to reading those.

**PREFACE**

## Part I

# Overview of superconductivity theories

As a precursor to topological superconductivity theory, we make a brief recap of the main superconductivity theories. Our intentions is not to make a complete mathematical description of said theories but act more as a memory refreshment of the core ideas and concepts. Also, since this section will serve more as consultation, we highlight the famous, useful, equations, be omitting any derivations (although somewhat explaining it in text so one can follow). I will try to keep a linear storytelling of the various theories with some exception of qualitatively nodding to context further ahead in the text to strengthen intuition.

### I. LONDON THEORY

The first theoretical explanation for the occurrence of superconductivity in metallic superconductors was proposed by the London brothers, Fritz London and Heinz London, in 1935. They began with the premise that if electrons in a superconductor do not encounter resistance, they will continue to accelerate under the influence of an applied electric field. Under this notion, they formulated the London equations, which serve as constitutive relations for a superconductor, describing the relationship between its superconducting current and the surrounding electromagnetic fields. While Ohm's law represents the simplest constitutive relation for an ordinary conductor, the London equations provide the most fundamental and meaningful description of superconducting phenomena.

#### A. London equations

Let us then start from the base concept of electrons accelerating with no resistance under the influence of an applied electric field  $\mathbf{E}$ . The equation of motion of these electrons in the superconducting state will then read  $m(d\mathbf{v}_s/dt) = -e\mathbf{E}$  with  $m$ ,  $\mathbf{v}_s$ ,  $e$  and  $n_s$  their mass, velocity, charge and density respectively. On the other hand, the superconducting current density is given by  $\mathbf{J}_s = -en_s\mathbf{v}_s$ . Differentiating it with respect to time and substituting  $d\mathbf{v}_s/dt$  yields the first London equation

$$\frac{d\mathbf{J}_s}{dt} = \frac{n_s e^2}{m} \mathbf{E} \quad (1)$$

Furthermore, taking the curl on both sides, making use of Faraday's law  $\nabla \times \mathbf{E} = -\partial_t \mathbf{B}$ , and integrating both sides of the equation on obtains the second London equation

$$\nabla \times \mathbf{J}_s = -\frac{n_s e^2}{m} \mathbf{B} \quad (2)$$

where the constant of integration is set zero to account for the fact that there is no resistivity in superconductors.

#### B. London penetration depth and Meissner effect

Consider Ampere's law  $\nabla \times \mathbf{B} = -\mu_0 \mathbf{J}$ , with  $\mu_0$  the vacuum magnetic permeability, which relates the magnetic field along a closed path to the total current following through any surface bounded by the path. If one takes its curl from both sides and makes use the no magnetic monopole law  $\nabla \cdot \mathbf{B} = 0$

one obtains  $\nabla^2 \mathbf{B} = -\mu_0 \nabla \times \mathbf{J}$ . Substituting the curl of the generic current  $\mathbf{J}$  for our superconducting current  $\mathbf{J}_s$  as given by London's 2nd equation one obtains the equation that describes the Meissner effect, reading

$$\nabla^2 \mathbf{B} = \left( \mu_0 \frac{n_s e^2}{m} \right) \mathbf{B} \equiv \frac{1}{\lambda_0^2} \mathbf{B} \quad (3)$$

where  $\lambda_0^2$  has dimension of length and is known as London's penetration depth. This equation tells us that that the magnetic field is exponentially suppressed as it penetrates inward a bulk superconductor. For example, see that a magnetic field  $\mathbf{B} = B \hat{\mathbf{z}}$  that penetrates a superconductor within the semi-infinite plane  $xOz$  is damped as  $\mathbf{B}(x) = B_0 \exp(-x/\lambda_0) \hat{\mathbf{z}}$  while inside the superconductor.

This exclusion of magnetic field is a manifestation of the superdiamagnetism emerged during the phase transition from conductor to superconductor, for example by reducing the temperature below critical temperature. In the presence of a weak external magnetic field—one that is below the critical threshold for the breakdown of superconductivity—a superconductor nearly completely expels the magnetic flux by generating electric currents in a thin layer near its surface. Specifically, the magnetic field induces a magnetization within the London penetration depth, which in turn establishes screening currents. These currents serve to protect the superconductor's internal bulk from the external field. Moreover, because the flux expulsion remains invariant over time, the so-called persistent (or screening) currents sustaining this effect do not decay.

See that this penetration depth is inversely proportional to the square root of the electron density in the superconductive state  $n_s$ , which in turn should depend on temperature. Concretely, one expects that as the temperature rises,  $n_s$  decreases and, consequently, the extent of flux penetration increases. At some critical temperature  $T_c$ ,  $n_s$  drops to zero, allowing the magnetic field to fully penetrate the material and causing the superconductor to revert to its normal state. The London brothers did not find exactly what this temperature dependence law should look and mostly miscalculated  $\lambda_0$  of different materials just because  $n_s$  could not be merely treated as a free electron density as it is done on metals; rather, the electrons in these superconductive phase were latter found to interact coherently. The actual temperature-dependent London penetration depth will be described in the next section.

### C. London coherence length

In addition to the London penetration depth  $\lambda_L$ , there is another fundamental length scale that governs superconducting behavior. Together, these two length scales play a crucial role in defining the properties of a superconductor.

While  $\lambda_L$  characterizes the extent to which an external magnetic field can penetrate a superconductor,  $\xi$  defines the spatial region over which the superconducting electron density remains relatively uniform, preventing abrupt variations in the presence of a non-uniform magnetic field. This distinction is particularly relevant in the context of the London equation, which establishes a *local* relationship between the supercurrent density  $\mathbf{J}_s(\mathbf{r})$  and the vector potential  $\mathbf{A}(\mathbf{r})$ , requiring the  $\mathbf{J}_s(\mathbf{r})$  to follow *exactly* any spatial variations in  $\mathbf{A}(\mathbf{r})$ . The coherence length sets a natural limit to this locality, representing the characteristic distance over which the vector potential must be averaged to determine the corresponding supercurrent density.

Any deviation from spatial uniformity incurs an additional kinetic energy cost, in other words, that any modulation of the superconducting wavefunction  $\psi_s(\mathbf{k}, \mathbf{r})$  identified by it's momentum state  $\mathbf{k}$  cost the system energy. Concretely, the increase of energy required for a modulation  $\psi_s(\mathbf{k}, \mathbf{r}) \rightarrow \psi_s(\mathbf{k} + \mathbf{q}, \mathbf{r})$  with  $|\mathbf{q}| \ll |\mathbf{k}|$  corresponds to  $\delta E = \hbar^2 |\mathbf{k}| |\mathbf{q}| / 2m$ . However, if  $\delta E$  exceeds the superconductive energy gap  $E_g$ , superconductivity will be destroyed. The critical value  $\mathbf{q}_0$  at which this happens is given  $E_g = \hbar^2 |\mathbf{k}_F| |\mathbf{q}_0| / 2m$  with  $k_F$  the momentum at the Fermi surface. We can then define an intrinsic coherence length  $\xi_0$  related to this critical modulation as  $\xi_0 = 1/q_0$  reading

$$\xi_0 = \frac{\hbar^2 k_F}{2mE_g} \quad (4)$$

As an additional complication, understand that both the coherence length  $\xi$  and the penetration depth  $\lambda$  of superconductors must be influenced by the mean free path of electrons  $\ell_e$  in the normal state. For now we do not know their specific dependence on  $\ell_e$  but we can at least guess for it qualitatively by considering the nature of the electron's wavefunctions in disordered systems. In a so-called dirty superconductors, one that has a smaller mean free path of electrons, the wavefunction exhibits inherent spatial fluctuations due to disorder. This means that a localized variation in current density can be constructed with lower energy using these pre-existing wiggled wavefunctions, as opposed to the smoother wavefunctions found in a pure superconductor, where greater energy would be required to introduce similar variations. Hence, one can expect that  $\xi < \xi_0$  for smaller  $\ell_e$ . On the other hand, since the ability to screen an external magnetic field depends on how effectively the supercurrent can be set up across the sample. In the dirty limit the superconducting electrons will not be able to coordinate over long distances resulting in an overall weaker screening currents. Weaker screening means that the magnetic field penetrates deeper into the material, and thus one can also expect  $\lambda > \lambda_0$  for smaller  $\ell_e$ .

## II. GINZBURG-LANDAU THEORY

Historically, the Ginzburg-Landau (GL) framework was introduced before the microscopic BCS theory of superconductivity. Although it was initially developed on largely phenomenological grounds, later work showed that it can be derived from the microscopic theory in certain limits. As a result, Ginzburg-Landau theory remains a cornerstone for describing superconductors near their critical temperature, providing both qualitative insights and quantitative tools for analyzing a wide range of superconducting phenomena.

Ginzburg-Landau theory offers a phenomenological way to understand how systems transition into the superconducting state building on the broader concept of second-order phase transitions at a given critical temperature. In this sense, one introduces an order parameter that captures how the system reorganizes itself at the threshold of the transition. This is analogous to how a ferromagnet spontaneously picks a magnetization  $\mathbf{M}$  direction. When the system is in its non-magnetic state the magnetization is effectively zero, but as the temperature cools below a given critical temperature  $T_c$  (dubbed Curie temperature for the case of ferromagnets) it acquires a nonzero value.

Ginzburg-Landau theory clarifies the relationship between the two London's characteristic length scales—the penetration depth  $\lambda$  which quantifies how far magnetic fields can penetrate into the superconductor, and the coherence length  $\xi$  which quantifies how quickly the order parameter can change in space. The balance between these scales determines whether a material expels magnetic fields completely, dubbed type I superconductors, or admits them in quantized flux tubes, dubbed type II superconductors.

### A. Superconductive order parameter

For superconductors, Ginzburg and Landau proposed that this order parameter is not just a simple number but a complex quantity that can vary in space, namely

$$\Psi(\mathbf{r}) = |\Psi(\mathbf{r})|e^{i\phi(\mathbf{r})} \quad (5)$$

whose magnitude  $|\Psi(\mathbf{r})|$  and phase  $\phi(\mathbf{r})$  convey key features of superconductivity. The effective number density of electrons  $n_s$  on the superconductive state is related to this magnitude, concretely  $n_s =$

$|\Psi(\mathbf{r})|^2$ , and the current flowing locally at a given point  $\mathbf{r}$  is related the gradient of the phase, concretely  $|\nabla\phi(\mathbf{r})|^2$ . Intuitively you can think of the magnitude as how “strong” the superconductivity is while the phase is instead related to collective quantum behavior that underlies phenomena such as persistent currents and flux quantization. Moreover, since this order parameter is smoothly varying in space he needs not be uniform near boundaries or in the presence of impurities.

The Ginzburg-Landau theory is formulated by employing a minimization of the Helmholtz free energy density  $f_s$  (thermodynamic potential that measures the useful work that a system held at constant temperature can perform) in terms of  $|\Psi(\mathbf{r})|^2$  and  $|\nabla\Psi(\mathbf{r})|^4$  under constraints imposed by external parameters such as temperature  $T$  and magnetic field  $\mathbf{H}$  with respect to variations in the order parameter  $\Psi$  and the vector potential  $\mathbf{A}$ . Understand that you cannot have powers of  $\Psi(\mathbf{r})$  in  $f_s$  because it must be real; nor can you just expand it terms of  $\text{Re}\{\Psi(\mathbf{r})\}$  since  $f_s$  must not depend on the absolute phase of  $\Psi(\mathbf{r})$ . Moreover, odd powers of  $|\Psi(\mathbf{r})|^2$  are also excluded because they are not analytic at  $\Psi(\mathbf{r}) = 0$ .

As we will see, this procedure results in a set of coupled differential equations governing the behavior of the order parameter  $\Psi(\mathbf{r})$ , dubbed the 1st GL equation, and the electromagnetic vector potential  $\mathbf{A}(\mathbf{r})$ , dubbed the 2nd GL equation. This interplay between the spatially varying superconducting order parameter and the electromagnetic field lies at the heart of the Ginzburg-Landau description.

The fundamental GL postulate asserts that if the magnitude of order parameter is small and varies gradually in space (local electrodynamic approximation) then the Helmholtz free energy density  $f_s$  near the transition temperature  $T_c$  can be expanded into the power series expansion

$$\begin{aligned} f_s(T) &= f_{\text{normal}} + f_{\text{condensate}} + f_{\text{kinetics}} + f_{\text{magnetic}} \\ &= f_n + \left[ \alpha(T)|\Psi(\mathbf{r})|^2 + \frac{\beta(T)}{2}|\Psi(\mathbf{r})|^4 \right] + \frac{\hbar^2}{2m^*} \left| \left( \nabla - \frac{ie^*}{\hbar c} \mathbf{A}(\mathbf{r}) \right) \Psi(\mathbf{r}) \right|^2 + \frac{H^2}{8\pi} \end{aligned} \quad (6)$$

with  $f_n$  the Helmholtz free energy density in the normal state,  $\alpha$  and  $\beta$  some phenomenological parameters to be determined experimentally (in conventional BCS superconductors these parameters be derived from microscopic theory),  $e^*$  and  $m^*$  the effective charge and mass of the superconducting carriers respectively,  $\mathbf{A}(\mathbf{r})$  the electromagnetic vector potential, and  $\mathbf{B} = \nabla \times \mathbf{A}$  the external magnetic field magnitude. The 2nd and 3rd terms correspond to the condensation free energy density, allures to the fact that the superconducting state is to be more ordered than the normal state, the 4th term corresponds to the kinetic energy density of the charged superconducting carriers in the presence of a magnetic field leading to supercurrents (the 2nd of its term to be precise), and the 5th term to the energy density associated with the magnetic field itself.

#### Bulk solutions (absence of field and currents)

Deep inside the bulk of the superconductor, several London penetration length's in, if the system is at the critical temperature  $T = T_c$  then the Helmholtz free energy density at the phase transition must be continuous, i.e that  $f_s(T_c) - f_n(T_c) = \alpha(T_c)|\Psi_\infty|^2 + \beta(T_c)/2|\Psi_\infty|^4 \stackrel{!}{=} 0$ , with  $\Psi_\infty$  the order parameter in the deep bulk regime notation. One the hand, minimizing  $f_s$  with respect to  $|\Psi(\mathbf{r})|$ , one obtains that

$$|\Psi_\infty|^2 \stackrel{!}{=} n_s^* = -\frac{\alpha}{\beta} \quad (7)$$

Substituting back into the previous condition one finds that

$$f_s(T_c) - f_n(T_c) = -\frac{\alpha(T_c)^2}{2\beta(T_c)} \equiv -\frac{H_c}{8\pi} \quad (8)$$

with  $H_c$  the critical magnetic field. See that the  $\beta(T)$  parameter must always be positive, even if  $\alpha(T) > 0$ , because otherwise there would be a *finite* potential barrier that, if crossed, would result

in infinite free energy. Oppositely, the  $\alpha(T)$  parameter can take whatever value. If  $\alpha(T) \geq 0$  the minimum free energy occurs at  $|\Psi(\mathbf{r})| = 0$ , corresponding to the normal state since  $n_s = |\Psi(\mathbf{r})|^2$  states no density of electrons on the superconductive state. One the other hand, if  $\alpha(T) < 0$  then the minimum free energy occurs at  $|\Psi(\mathbf{r})| > 0$ , corresponding to the superconductor state since it gives a lower free energy state. [see Fig.(1)].




Figure 1. Ginzburg-Landau theory Helmholtz free energy density  $f_s$

#### *Temperature dependency*

Since the  $\alpha(T)$  must change from positive to negative at  $T = T_c$  let us make a Taylor's series expansion around  $T_c$  but keeping only the linear term, reading  $\alpha(t_s) = \alpha_s(1 - t_s)$  with  $t_s = T/T_c$  and  $\alpha_s < 0$ , such that in the normal phase  $T > T_c \Rightarrow t_s < 1 \Rightarrow \alpha(t_s) \propto \alpha_s < 0$  and in the superconducting phase  $T < T_c \Rightarrow t_s > 1 \Rightarrow \alpha(t_s) \propto -\alpha_s > 0$ . Inserting the empirical notations that  $B_c \propto (1 - t_s^2)$  one can then infer deep that inside the bulk the temperature dependent behavior of the London's penetration length goes like  $\lambda(t_s) \propto |\Psi_\infty(\alpha(t_s), H_c(t_s))|^2 \propto (1 - t_s^4)^{-1/2}$ .

## B. Ginzburg-Landau equations

Minimizing the total Helmholtz free energy density  $F_s = \int_{\mathcal{V}} d^3\mathbf{r} f_s(\mathbf{r})$  over the volume  $\mathcal{V}$  of the superconductive system with respect to the variation of the order parameter  $\Psi^*(\mathbf{r})$  **why the complex conjugate though?** gives us the 1st Ginzburg-Landau equation

$$\alpha\Psi(\mathbf{r}) + \beta|\Psi(\mathbf{r})|^2\Psi(\mathbf{r}) + \frac{1}{2m^*} \left( -i\hbar\nabla_{\mathbf{r}} - \frac{e^*}{c}\mathbf{A} \right)^2 \Psi(\mathbf{r}) = 0 \quad (9)$$

See that, apart from the nonlinear term, this equation has the form of a Schrodinger equation for particles with energy eigenvalue  $-\alpha$  within the same conditions. The nonlinear term acts like a repulsive potential of  $\Psi(\mathbf{r})$  on itself, tending to favor wavefunctions  $\Psi(\mathbf{r})$  which are spread out as uniformly as possible in space.

One the other hand, the variation of vector potential  $\mathbf{A}$  gives us the 2nd Ginzburg-Landau equation

$$\mathbf{J}_s = \frac{e^*}{m^*}|\Psi(\mathbf{r})|^2 \left( \hbar\nabla_{\mathbf{r}}\phi(\mathbf{r}) - \frac{e^*}{c}\mathbf{A}(\mathbf{r}) \right) \equiv e^*|\Psi(\mathbf{r})|^2\mathbf{v}_s \quad (10)$$

which shows us that also the superconductive current resembles quantum mechanical expressions in the same conditions, concretely the current of probability with the caveat of having an effective number density  $n_s = |\Psi(\mathbf{r})|^2$ , mass  $m^*$  and charge  $e^*$ . In the original formulation of the theory it was assumed without much thought that  $e^*$ ,  $m^*$  and  $n_s^*$  corresponded to their normal electronic values however experimental data surprisingly suggested a better fit for  $e^* = 2e$ ,  $m^* = 2m$  and  $n_s^* = \frac{1}{2}n_s$ . For us time travelers this obviously screams Cooper pairing of electrons as predicted by the microscopic BCS theory. See that the relation  $\lambda = n_s e^2 / m = n_s^* e^{*2} / m^*$  still holds though, ensuring that the London penetration depth remains unchanged due to the pairing mechanism. Notably, see that a decrease in the order parameter results in an increase in the penetration depth.

#### *Boundary conditions*

As an additional and relevant detail, remember that along the variational procedure one must eventually provide a choice of boundary conditions of the superconductive volume. Indeed, in GL theory the boundary condition is that of an insulating surface such that it is ensured that no supercurrent leaks through the superconductor, i.e  $\mathbf{J}_s \cdot \mathbf{n} = 0$  at the interface. Concretely, this means that  $(-i\hbar\nabla_{\mathbf{r}} - e^*/c\mathbf{A}(\mathbf{r}))\Psi(\mathbf{r})]_{\mathbf{n}} = 0$ . From the microscopic theory de Gennes latter shown that the right side, rather than zero, should read instead  $i\hbar/\Psi(\mathbf{r})b$  with  $b$  a real constant. If at the interface  $\mathbf{A} = 0$  then  $b$  corresponds to the extrapolation length to the point outside the boundary at which  $\Psi$  would go to zero if it maintained the slope it had at the surface. The value of  $b$  will depend on the nature of the material to which contact is made, approaching  $b = 0$  for a magnetic material and  $b = +\infty$  for an insulator, with normal metals lying in between.

#### *GL coherence length*

Let us consider a simplified one-dimensional case were no magnetic field are present ( $\mathbf{A} = 0$ ) and analyze GL 1st differential equation in Eq.(9). See that, in this case,  $\Psi(\mathbf{r})$  get to be real since the equation only has real coefficients. Introducing the normalized wavefunction  $\tilde{\Psi} = \sqrt{\beta/|\alpha|}\Psi$  with  $\alpha = -|\alpha|$  the (one-dimensional) equation becomes  $\xi^2 \partial_x^2 \tilde{\Psi} + \tilde{\Psi} - \tilde{\Psi}^3 = 0$  where we identified the characteristic length  $\xi$  of the order parameter variations as  $\xi = \hbar^2 / (2m^{*2}|\alpha(T)|)$ . This is known as the GL coherence length which, as the name implies, plays the same role as the same as London's, describing the distance over which the superconductor can be represented by a wavefunction. Moreover, for my time-travelers fellows, this can also be understood as the distance over which Cooper pairs can be considered to be correlated. Within the deep bulk ( $\mathbf{A} = 0$ ) the order parameter  $\tilde{\psi}$  will not vary in space and thus one can solve the equation by setting the boundary conditions  $\partial_x \tilde{\Psi} = 0$  and  $\tilde{\Psi}^2 = 1$ . One obtains

$$\Psi(x) = \sqrt{\frac{|\alpha|}{\beta}} \tanh\left(\frac{x}{\sqrt{2}\xi}\right) \quad (11)$$

### C. Flux quantization

Consider a superconductor ring with a magnetic flux  $\Phi$  passing through it's perforation inducing a persistent current  $\mathbf{J}_s$  coursing trough it's "inner" "surface" as to counter act the magnetic field in the bulk within a penetration depth  $\lambda$ . Now, consider a circular path  $\mathcal{C}$  within the deep bulk of the ring far away from any persistent currents, such that  $\oint_{\mathcal{C}} \mathbf{J}_s \cdot d\ell = 0$  with  $\mathbf{J}_s$  given by the 2nd GL equation in Eq.(10). Since the system is defined at its minimal energy configuration the order parameter within the deep bulk  $\Psi_\infty$  must have a unique value at every point along the circular path. This leaves us specifically with  $\oint_{\mathcal{C}} \mathbf{v}_s \cdot d\ell = 0$  which is trivial to solve for. For the 1st term, one has that  $\oint_{\mathcal{C}} \nabla_{\mathbf{r}} \phi(\mathbf{r}) \cdot d\ell = 2\pi n$ , since  $\phi(\mathbf{r})$  goes around in a circle and back to here it started acquiring a phase of  $2\pi$  for each  $n \in \mathbb{Z}$  lap, and for the 2nd term one obtain, by definition, the magnetic flux  $\Phi$ , since  $\oint_{\mathcal{C}} \mathbf{A}(\mathbf{r}) \cdot d\ell = \oint_S \nabla \times \mathbf{A}(\mathbf{r}) \cdot dS = \oint_S \mathbf{B} \cdot dS = \Phi$  with  $S$  the surface spanning the over the hole. Note that  $n \neq 0$  requires that the contour cannot be contracted to a single point, meaning that the sample must always contain a hole, as it has in our case. Combining this results one obtains (with the

foresight substitution  $e^* = 2e$ )

$$\Phi = n \frac{hc}{2e} \equiv n\Phi_0, \quad (12)$$

meaning that the flux through the ring is actually quantized in integral multiples of  $\Phi_0$ , the flux quantum, also known as fluxoid. Bear in mind the subtlety that it's the total flux  $\Phi = \Phi_s + \Phi_H$  that is quantized, i.e the sum of the flux from external magnetic field  $\Phi_H$  and the flux from the persistent superconducting currents  $\Phi_s$ . Since there is no quantization condition on the external sources then  $\Phi_s$  itself must adjust appropriately in order that  $\Phi$  assumes a quantized value.

#### *GL coherence length*

As a quick side note, see that putting together Eq.(8) and Eq.(7) along with London's penetration length definition in Eq.(3) and the fluxoid definition in Eq.(12) one can express the GL coherence length as

$$\xi(T) = -\frac{\Phi_0}{2\sqrt{2}\pi H_c(T)\lambda^*} \quad (13)$$

#### D. Type I and type II superconductors

As previously discussed, although currents can flow without any energy dissipation in superconductors, there are certain limitations; the material must operate below a given critical temperature  $T_c$  but also under magnetic field strengths below a critical value  $H_c(T)$ . With regard to their magnetic properties, particularly in the way they expelled magnetic fields superconductors can then be categorized into one of two types, simply named type I and type II.

On one hand, type I superconductors exhibit a sharp normal-superconductive phase transition with all magnetic flux being expelled while in the superconductive phase while type II superconductors exhibit an additional in-between "mixed state", also referred to as "vortex state", where there is partial penetration of flux. This partial penetration occurs as a mechanism to minimize the overall magnetic energy. Surrounding these small localized regions of partial penetration —where the magnetic field is high enough to revert the superconductor into its normal phase—are circulating vortices of quantized screening currents that oppose the magnetic field guaranteeing that the material outside these regions remains in the superconducting state. This process by which superconductivity "kicks off" in small localized pockets is often referred to as nucleation. Understand that, although the sample is not locally superconducting in those regions, it can still have zero electric resistance as a whole since the currents predominantly flows through the superconducting areas. Moreover, understand that to maintain a lossless state these vortices must be pinned in place, for example, by defects within the crystal structure, or else they will move and generate a voltage leading to dissipation.

Another way to qualitatively understand this two types of superconductivity is by examining the interaction energy between superconducting vortices. Rather than performing a full explicit derivation, we can gain insight by considering the broader picture.

The derivation of the vortex interaction energy begins with determining the shape, and consequently the energy, of an individual vortex. This is realized by solving the field equations in cylindrical coordinates for a non-constant  $\Psi(x)$ , as we are dealing with local defects. In this choice of coordinates, the equations take the form of coupled nonlinear differential equations. An important detail in this derivation is that to compute the vortex energy per unit length, one must introduce a cutoff, which reflects the fact that a vortex can only exist within a finite-sized system.

Once the energy of the individual vortices is known, one goes to find the energy of the entire system and then subtract them off to obtain the interaction energy between superconducting vortices. It reads  $E_{\text{int}} \propto d/\lambda - \sqrt{2}d/\xi$  with  $d$  the distance between the vortices. This expression reveals two competing effects: a repulsive interaction caused by vortex currents circulating in opposite directions (analogous

to the force between two parallel wires carrying currents in opposite directions) and an attractive interaction caused by the fact that a superconductor energetically favors a defect-free state, it tends to restore order by merging vortices whenever possible. The balance between these opposing forces determines whether vortices attract or repel. Quantitatively, what governs the nature of this interaction is the ratio between the GL coherence length  $\xi(T)$  and London's penetration depth  $\lambda(T)$ , known as the GL parameter  $\kappa = \lambda(T)/\xi(T)$ . See that apart from being a dimensionless quantity, since both  $\lambda(T)$  and  $\xi(T)$  diverge as  $\sqrt{1 - T/T_c}$  with temperature,  $\kappa$  is also temperature independent. If  $\kappa > 1/\sqrt{2}$  the repulsive interaction dominates and thus the vortices repel from each other, arranging themselves into regular periodic structures, typically a triangular lattice. Since each vortex carries a quanta of flux  $\Phi_0$  this results in partial penetration of the magnetic field, a hallmark of type II superconductors. Conversely, if  $\kappa < 1/\sqrt{2}$ , the attractive interaction prevails, leading to the agglomerate and collapse of all vortices into a single entity. In this case, the superconductor has no mechanism to sustain flux penetration and instead exhibits the Meissner effect, a hallmark of type I superconductor.




Figure 2. The nomenclature type I and type II will be made clear after Ginzburg-Landau theory, for now think of it as simply pure and dirty, respectively

### E. Little-Parks experiment

Instead of the ring, consider now a superconducting cylindrical thin-film shell of radius  $R$  and thickness  $\ell$  with a magnetic flux  $\phi$  passing through its perforation. Specifically, we consider the shell to be so thin so that  $\ell \ll \xi(T)$  with  $\xi(T)$  the London's coherence length. In this case, any small deviation of  $|\Psi(\mathbf{r})|$  would mean an excessively large  $|\Psi(\mathbf{r})|^2$  contribution to the free energy which is not physically realistic. To correct this problem one then needs to approximate the magnitude to a uniform value, i.e  $\Psi(\mathbf{r}) = \Psi_0$ . In this conditions the Helmholtz free energy density  $f_s$  would

approximately read  $f_s^{\text{thin}} \approx f_n + (\alpha + \kappa)|\Psi_0|^2 + \beta/2|\Psi_0|^4 + H^2/8\pi$  with  $\kappa = 1/2m^*v_s^2$  the kinetic energy of the superconducting current. Moreover, we further neglect the free energy term associated with the external magnetic field because it is smaller than the kinetic energy by a ratio of  $\pi R^2$  to  $1/\lambda^2$ . The optimal value of  $|\Psi_0|^2$  is then found by minimizing  $f_s^{\text{thin}}$ , for a given  $v_s$ , reading

$$|\Psi_0|^2 = \Psi_\infty \left[ 1 - \left( \frac{\xi(T)m^*v_s}{\hbar} \right)^2 \right] \quad (14)$$

From the previous quantization condition  $\Phi_s = n\Phi_0 - \Phi_H$  we already know what the supercurrent velocity  $v_s$  should be, it reads as the  $\Phi_H/\Phi_0$ -periodic function  $v_s = \hbar/(m^*R)(n - \Phi_H/\Phi_0)$ .

Let us analyze what happens at the normal-superconductor phase transitions. Substituting directly into Eq.(14) the supercurrent velocity  $v_s$  in its  $\Phi_H/\Phi_0$ -periodic form and setting  $|\Psi_0|^2 = 0$  one finds that, through the temperature dependence of the coherence length  $\xi(T)$  in Eq.(??), there will be a periodic variation  $\delta T_c$  of the critical temperature  $T_c$ , concretely

$$\frac{\delta T_c(H)}{T_c} \propto \begin{cases} \frac{\xi_0^2}{R^2} \left( n - \frac{\Phi_H}{\Phi_0} \right)^2 & \text{for a pure SC} \\ \frac{\xi_0 \ell}{R^2} \left( n - \frac{\Phi_H}{\Phi_0} \right)^2 & \text{for a dirty SC} \end{cases} \quad (15)$$

This is known as the Little-Parks effect. See that the maximum of the depression of  $T_c$  occurs when  $n - \Phi_H/\Phi_0 = 1/2$ .




Figure 3. (left) fluxoid (right) Little-Parks effect

#### F. Josephson effects

This section needs a lot more research and work. . I want a clear, basic, linear, intuitive introduction, I don't want it to be heavy on the math and have to much tangents. I'm working on the figures and plots for this. At the end I want to start with hints to topology and Majoranas with the  $4\pi$  Josephson effect.

The Josephson effect occurs when two superconductors are weakly coupled through a very thin barrier, which may be an insulator, a metallic weak link, or any physical constriction that weakens superconductivity. Within the framework of BCS theory, the weak coupling allows for a probability of Cooper pairs tunneling from one superconductor to the other. Due to the phase coherence embodied

in the superconducting order parameter, a supercurrent will flow even in the absence of an applied voltage. This phenomenon arises from the spontaneous symmetry breaking in the superconducting state, leading to both steady (DC) and oscillatory (AC) tunneling effects.

In the absence of an applied voltage, the relative phase difference between the superconductors remains constant over time, resulting in a steady tunneling supercurrent. The free energy of the system minimized when the order parameters of the two superconductors are optimally aligned. Any deviation from this alignment (i.e., any phase difference) introduces a coupling energy that drives a current. The system reaches a stable state as long as the phase difference is maintained and the supercurrent does not exceed a critical value. Conversely, when a constant voltage is applied across the junction, the phase difference no longer remains static. Instead, it evolves linearly over time in an oscillatory manner, leading to an alternating current. Despite the applied voltage being constant, the tunneling supercurrent alternates with a frequency (known as the Josephson frequency) that is directly proportional to the voltage. This remarkable voltage-to-frequency conversion is the basis for many precision voltage standards.

### 1. DC Josephson effect

Consider a very thin weak link between two superconductor pieces as shown in Fig. ???. Let the phase of order parameter of the left-side superconductor be  $\phi_L$  for  $x < \xi_L$  and  $\phi_R$  for  $x > \xi_R$  for the right-side superconductor such that at the weak link, set at  $x = 0$ , its phase may change rapidly by a very small perturbation (besides the amplitude  $|\Psi|$  being exponentially small). Also, let  $|\Psi| = |\Psi_L| = |\Psi_R|$  and  $\xi = \xi_L = \xi_R$ .

As introduced in the Ginzburg-Landau equations subsections, the boundary conditions  $(-i\hbar\partial_x - e^*/cA_x)\Psi]_{\xi_L} = b\Psi_R$  where, in this case,  $b$  is a small number depending on the properties of the weak link. Moreover, time inversion symmetry demands this boundary conditions to remain valid both  $\Psi \rightarrow \Psi^*$  and  $\mathbf{A} \rightarrow -\mathbf{A}$ , inferring that  $b$  must be real as long as the phase  $\phi$  does not depend on  $\mathbf{A}$ . Hence, for the moment, let us pick appropriate gauge in which  $A_x = 0$  such that  $\partial_x\Psi]_{\xi_L} = i/\hbar b\Psi_R$ . The supercurrent density at  $x = \xi_L$  can then be found from  $J]_{\xi_L} = \hbar e^*/2im^*\left(\Psi_L^* \partial_x\Psi]_{\xi_L} - \Psi_L \partial_x\Psi^*]_{\xi_L}\right)$  which correspond to the supercurrent

$$I = I_c \sin(\varphi) = I_c \sin(\phi_R - \phi_L) \quad (16)$$

with the prefactor being the Josephson junction critical current  $I_c = b\hbar e^*/m^*$  defined at  $\phi_R = \phi_L \pm \pi/2$ , depending upon the junction strength  $b$ . On the other hand, if there is no phase difference, i.e.  $\phi_R = \phi_L$ , no current will flow. A mechanical analog of this equation would be a system of coupled pendulums in the sense that no energy is exchanged if the pendulums are oscillating in phase or out of phase.

### 2. AC Josephson effect

Let us now consider the case when a constant voltage  $V$  is applied across the junction such that the phase difference no longer remains static.

We start from the time-evolution of the *gauge-invariant* phase difference between the two superconductor, also known as Josephson phase  $\varphi_J(t)$ , following directly from the flux quantization condition introduced in the subsection of the same name. It reads

$$\dot{\varphi}_J(t) = \dot{\phi}_R(t) - \dot{\phi}_L(t) - \frac{e^*}{\hbar} \int_{\xi_L}^{\xi_R} dx \dot{A}_x(x, t). \quad (17)$$

Since we are dealing with semi-infinite superconductors at equilibrium one can take the phases  $\phi_R$  and  $\phi_L$  within that bulk to be so slowly varying in time such that any time variation of is negligible

compared to the contribution from the vector potential. As for the second term, we know that the applied voltage in an purely inductive gauge, i.e with no electrostatic difference between the two superconductors, meaning  $\int \nabla \phi_E \cdot d\ell = 0$ , is solely given by the time-variation of the magnetic vector potential  $V = \int \mathbf{A} \cdot d\ell$ . Putting the two expression together one obtains  $V = -e^*/\hbar \dot{\varphi}_J(t)$ , a superconducting analog to electromagnetism Faraday's law of induction with the distinction that the voltage does not come from magnetic flux (there isn't even any in the bulk) but rather from the kinetic energy of the superconducting carriers. This phenomenon is also known as kinetic inductance. The phase difference is then as depending linearly on time given by  $\varphi_J(t) = \omega_J t$  with  $\omega_J = -Ve^*/\hbar$  the Josephson frequency. This is often called the called the second Josephson relation or superconducting phase evolution equation. Finally, the corresponding supercurrent yields

$$I = I_c \sin(\varphi_J(t)) = \sin(\omega_J t), \quad (18)$$

which is known as the first Josephson relation or weak-link current-phase relation.

As a supplementary detail on the electromagnetism comparison, from the two Josephson relations one can then derive the explicit expression for the kinetic inductance  $L(\varphi_J)$  by straightforwardly applying the chain rule to calculate the time derivative of the current and rearrange the result in the form of a current–voltage characteristic of an inductor. One obtains  $L = L_J / \cos(\varphi_J)$  with  $L_J = \Phi_0 / (2\pi I_c)$  the Josephson inductance.

### 3. Inverse AC Josephson effect\*

Let us now consider the case where, instead of applying a DC voltage which has lead to an AC superconducting current, an external (microwave) AC voltage is applied to the junction. As we will see, under the right conditions the junction's phase locks to the external drive leading to the appearance of DC superconducting quantized current plateaus, known as Shapiro steps. We refer to this phenomenon as the inverse AC Josephson effect, as it involves frequency-to-voltage conversion rather than the other way around.




Figure 4. Josephson effects (a) DC (b) AC (c) inverse AC

### 4. SQUIDs\*

Let us now expand on the unidirectional one-dimensional model by connecting the other end point of the superconductors as to form a superconducting loop of two Josephson junctions. Moreover, consider a flux  $\Phi$  flowing thought it's perforation. This device is know as a SQUID, a superconducting quantum interference device, being depicted in Fig.(??).

In this system, each of the two half loops will acquire different phases equal to  $2\pi\Phi/\Phi_0$  and the current will be maximum when the phase difference is an even multiple of  $\pi$  and minimum when the phase difference is an odd multiple of  $\pi$ . One can intuit that the supercurrent, either in the ring will become oscillatory when plotted against the magnetic flux.

In this case, the phase difference will change with the flux in the ring, which is quantized, yielding instead the supercurrent

$$I = I_c \left[ \sin(\phi_R - \phi_L) + 2\pi \frac{\Phi}{\Phi_0} \right] \quad (19)$$

### III. BARDEEN-COOPER-SCHRIEFFER THEORY\*

Bardeen–Cooper–Schrieffer's (BCS) theory of superconductivity was a watershed in modern condensed matter physics. Its key feature is pair condensation, the macroscopic occupation of a bound state of fermion pairs. The binding of fermions into Cooper pairs typically leads to an energy gap in the fermionic excitation spectrum, while condensation of Cooper pairs leads to the breaking of global  $U(1)$  gauge symmetry. This symmetry breaking is linked to the spontaneously choosing of an overall phase  $\varphi$  of the macroscopic wavefunction below the transition temperature  $T_c$  (akin to how a ferromagnet spontaneously picks a magnetization direction) and its generator is the particle number, being related to the fact that  $\varphi$  and  $N$  are canonical conjugate (well, technically only for larger values of  $N$  but this is often the case). In BCS superconductivity considerations,  $\varphi$  is precisely the (conjugate of the) number of Cooper pairs formed. Furthermore, the symmetry breaking of  $U(1)$  implies that the fermionic excitations are no longer charge eigenstates, but each is a coherent superposition of a normal-state particle and hole, e.g.  $\gamma_{\mathbf{k}\sigma} = u_{\mathbf{k}} c_{\mathbf{k}\sigma} + v_{\mathbf{k}}^* c_{-\mathbf{k}\bar{\sigma}}^\dagger$ , with  $c/c^\dagger$  the electronic field operators and where  $u$  and  $v$  are the particle and hole amplitudes (defined by momentum  $\mathbf{k}$  and spin  $\sigma$  [ $\bar{\sigma}$  being the flipped spin]) defining the so called Bogoliubov quasi-particles (or Bogoliubons). Charge conservation is then maintained by an additional channel for charge transport via the coherent motion of the pair condensate. One can then construct the ground state of the superconductor  $|\emptyset\rangle$  (also denoted as  $|\text{GS}\rangle$  or  $|\text{BCS}\rangle$ ) from the condition that it contain no Bogoliubons,  $\gamma|\emptyset\rangle = 0$ , yielding a superposition of states with different number of Cooper pairs  $|\emptyset\rangle = \prod_{\mathbf{k}} (u_{\mathbf{k}} + v_{\mathbf{k}} c_{\mathbf{k}\sigma}^\dagger c_{-\mathbf{k}\bar{\sigma}}^\dagger) |\emptyset\rangle$ , with  $|\emptyset\rangle$  the state containing no electrons.

Expanding beyond conventional BCS, we can distinguish other types of superconductivity by the characteristics of the pair condensation. In conventional BCS superconductors (SCs), the electrons are being Cooper paired with opposite spins, forming a  $S = 0$  spin-singlet state, but it is possible to Cooper pair electrons with parallel spins forming three possible  $S = 1$  spin-triplet states without violating Pauli principle. Concerning with the orbital component we can also distinguish between different angular momentums  $\ell = 0(s), 1(p), 2(d), 3(f)$  and so on. As a first order approximation, one can match the orbital component to the shapes of spherical harmonics, although, of course, with the caveat that the crystal lattice and Fermiology can make the situation more complex in real materials. Because Fermions obey antisymmetric exchange (switching two electrons corresponds to a sign change), if the spin part of the wavefunction is antisymmetric, as is the case for the singlet case, then the orbital part has to be even,  $\ell = 0, 2, \dots$ . Of course, for the same reason, the triplet case must have instead odd orbital part,  $\ell = 1, 3, \dots$ .

### A. Effective Hamiltonian

To investigate the onset of superconductivity, consider the effective Hamiltonian

$$\begin{aligned} H &= H_0 + H_{\text{int}} \\ &= \sum_{\mathbf{k}\sigma} \xi_{\mathbf{k}} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} + \frac{1}{N} \sum_{\mathbf{kk}'} V_{\mathbf{kk}'} c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger c_{-\mathbf{k}'\downarrow} c_{\mathbf{k}'\uparrow} \end{aligned}$$

with  $\xi_{\mathbf{k}} = \varepsilon_{\mathbf{k}} - \mu$  the  $\mathbf{k}$ -state energy apart from the chemical potential shift  $\mu$  and where  $c_{\mathbf{k}\sigma}^\dagger$  ( $c_{\mathbf{k}\sigma}$ ) creates (annihilates) an electron with momentum  $\mathbf{k}$  and spin  $\sigma$ . See that the second term describes the destruction of two electrons with opposite momenta and spin simultaneously and the subsequent creation of another. We call this paring of electrons, with opposite momenta and spin, a Cooper pair.

To proceed, we perform the usual mean-field decoupling of the quartic term, reading

$$\langle c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger c_{-\mathbf{k}'\downarrow} c_{\mathbf{k}'\uparrow} \rangle \approx \langle c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger \rangle c_{-\mathbf{k}'\downarrow} c_{\mathbf{k}'\uparrow} + c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger \langle c_{-\mathbf{k}'\downarrow} c_{\mathbf{k}'\uparrow} \rangle - \langle c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger \rangle \langle c_{-\mathbf{k}'\downarrow} c_{\mathbf{k}'\uparrow} \rangle, \quad (20)$$

In this mean-field scheme the effective Hamiltonian reads

$$H = \sum_{\mathbf{k}\sigma} \xi_{\mathbf{k}} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} - \sum_{\mathbf{k}} (\Delta_{\mathbf{k}} c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger + \Delta_{\mathbf{k}}^* c_{-\mathbf{k}\downarrow} c_{\mathbf{k}\uparrow}) + \sum_{\mathbf{k}} \Delta_{\mathbf{k}} \langle c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger \rangle \quad (21)$$

where we have defined the

$$\Delta_{\mathbf{k}} = -\frac{1}{N} \sum_{\mathbf{k}'} V_{\mathbf{kk}'} \langle c_{-\mathbf{k}'\downarrow} c_{\mathbf{k}'\uparrow} \rangle, \quad (22)$$

known as the gap function. For now there is no reason to call it a gap for now, but we will discuss its meaning very soon.

Furthermore, in order to express this Hamiltonian in its diagonal form we employ the so-called Bogoliubov transformation

$$\begin{aligned} c_{\mathbf{k}\uparrow} &= u_{\mathbf{k}}^* \gamma_{\mathbf{k}\uparrow} + v_{\mathbf{k}} \gamma_{-\mathbf{k}\downarrow}^\dagger \\ c_{-\mathbf{k}\downarrow}^\dagger &= u_{\mathbf{k}} \gamma_{-\mathbf{k}\downarrow}^\dagger - v_{\mathbf{k}}^* \gamma_{\mathbf{k}\uparrow} \end{aligned} \quad (23)$$

where we have defined new fermionic operators  $\gamma_{\mathbf{k}\sigma}$  and coefficients  $u_{\mathbf{k}}, v_{\mathbf{k}}$  whom, in order for the fermionic commutation relations to be satisfied, must satisfy the normalization condition  $|u_{\mathbf{k}}|^2 + |v_{\mathbf{k}}|^2 = 1$ . Substituting directly into the Hamiltonian one finds that additionally the condition

$$2\xi_{\mathbf{k}} u_{\mathbf{k}} v_{\mathbf{k}} - \Delta_{\mathbf{k}} u_{\mathbf{k}}^2 + \Delta_{\mathbf{k}}^* v_{\mathbf{k}}^2 = 0 \Rightarrow \frac{v_{\mathbf{k}}}{u_{\mathbf{k}}} = \frac{\sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2} - \xi_{\mathbf{k}}}{\Delta_{\mathbf{k}}^*} \quad (24)$$

must be held in order to express the Hamiltonian in it's diagonal form. Notice that we picked only the positive root to ensure that the energy of the BCS state is a minimum and not a maximum. Moreover, notice that because the numerator is real, the phase of the complex gap function  $\Delta_{\mathbf{k}}$  must be the same as the relative phase between  $v_{\mathbf{k}}$  and  $u_{\mathbf{k}}$ . Since we can set the phase of  $u_{\mathbf{k}}$  to be zero without loss of generality, it follows that the phases of  $v_{\mathbf{k}}$  and  $\Delta_{\mathbf{k}}$  are the same. This yields the definitions

$$|u_{\mathbf{k}}|^2 = \frac{1}{2} \left( 1 + \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right) \text{ and } |v_{\mathbf{k}}|^2 = \frac{1}{2} \left( 1 - \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \right) \text{ with } E_{\mathbf{k}} = \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2} \quad (25)$$

the excitation energy. The effective Hamiltonian then follows as

$$H = \sum_{\mathbf{k}\sigma} E_{\mathbf{k}} \gamma_{\mathbf{k}\sigma}^\dagger \gamma_{\mathbf{k}\sigma} + E_0 \text{ with } E_0 = \sum_{\mathbf{k}} (\xi_{\mathbf{k}} - E_{\mathbf{k}} + \Delta_{\mathbf{k}}) \langle c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger \rangle \quad (26)$$

the energy of the BCS ground state, denoted by  $|\emptyset\rangle$  (often also as  $|\psi_{\text{BCS}}\rangle$ ). It becomes clear from this equation why we called  $\Delta_{\mathbf{k}}$  the gap function. Even at the Fermi level, where  $\xi_{\mathbf{k}}^2$ , the energy spectrum of the superconductor has a gap of size  $|\Delta_{\mathbf{k}}|$  meaning that we need a minimum energy of  $2|\Delta_{\mathbf{k}}|$  to the system to excite its quasiparticles, usually called Bogoliubons (the ones described by the  $\gamma$  operators). Note from Eq. (23) that a Bogoliubon is a mixture of electrons and holes. From the  $u_{\mathbf{k}}$  and  $v_{\mathbf{k}}$  in Eq.(25), we have that as  $\Delta_{\mathbf{k}} \rightarrow 0$ , the amplitudes behave as  $|u_{\mathbf{k}}|^2 \rightarrow 1$  for  $\xi_{\mathbf{k}} > 0$  and  $|u_{\mathbf{k}}|^2 \rightarrow 0$  for  $\xi_{\mathbf{k}} < 0$  whereas  $|v_{\mathbf{k}}|^2 \rightarrow 1$  for  $\xi_{\mathbf{k}} < 0$  and  $|v_{\mathbf{k}}|^2 \rightarrow 0$  for  $\xi_{\mathbf{k}} > 0$ . Thus, at the normal state, creating a Bogoliubon excitation corresponds to creating an electron for energies above the Fermi level and creating a hole (destroying an electron) of opposite momentum and spin for energies below the Fermi level. At the superconducting state, a Bogoliubon becomes a superposition of both an electron and a hole state. The BCS ground state therefore corresponds to the vacuum of Bogoliubons, i.e  $\gamma|\emptyset\rangle = 0$ .

### B. The BCS ground state

If one desires, one can describe this BCS ground state  $|\emptyset\rangle$  in terms of the original electronic ground state  $|0\rangle$ . We start by expressing the BCS ground state as an arbitrary combination of Cooper pairs, reading

$$|\emptyset\rangle = \mathcal{N} \prod_{\mathbf{q}} e^{\theta_{\mathbf{q}}} |0\rangle \text{ with } \theta_{\mathbf{q}} = \alpha_{\mathbf{q}} c_{\mathbf{q}\sigma}^\dagger c_{-\mathbf{q}\bar{\sigma}}^\dagger, \quad (27)$$

$\mathcal{N}$  a normalization constant and  $\alpha_{\mathbf{q}}$  a function to be determined. See that, if one acts with  $c_{\mathbf{k}\sigma}$  on the ground state above the only term inside the product that does not commute with  $c_{\mathbf{k}\sigma}$  is the one for which  $\mathbf{q} = \mathbf{k}$ . We have  $c_{\mathbf{k}\sigma} e^{\theta_{\mathbf{k}}} |0\rangle = \sum_{n=1}^{+\infty} c_{\mathbf{k}\sigma} \theta_{\mathbf{k}}^n / n! |0\rangle$ . Now, from the vacuum of electrons condition  $c_{\mathbf{k}\sigma} |0\rangle = 0$ , from the vacuum of Bogoliubons conditions  $u_{\mathbf{k}} c_{\mathbf{k}\sigma} |\emptyset\rangle = v_{\mathbf{k}} c_{-\mathbf{k}\bar{\sigma}}^\dagger |\emptyset\rangle$  and from the following commutations relations,  $[c_{\mathbf{k}\sigma}, \theta_{\mathbf{k}}] = \alpha_{\mathbf{k}} c_{-\mathbf{k}\downarrow}^\dagger$  and  $[\theta_{\mathbf{k}}, c_{-\mathbf{k}\bar{\sigma}}^\dagger] = 0$ , one finds that  $\alpha_{\mathbf{k}}$  must correspond to the ratio  $v_{\mathbf{k}}/u_{\mathbf{k}}$ . Finally, from Pauli's exclusion principle  $(c_{\mathbf{k}\sigma}^\dagger c_{-\mathbf{k}\bar{\sigma}}^\dagger)^n = 0$  for  $n > 1$  one finds the normalized BCS ground state as being

$$|\emptyset\rangle = \prod_{\mathbf{k}} \left( u_{\mathbf{k}} + v_{\mathbf{k}} c_{\mathbf{k}\sigma}^\dagger c_{-\mathbf{k}\bar{\sigma}}^\dagger \right) |0\rangle, \quad (28)$$

where the normalization constant is found to be  $\mathcal{N} = \prod_{\mathbf{k}} u_{\mathbf{k}}$  through  $\langle \emptyset | \emptyset \rangle = 1$ . Also, recall that the phase of the Cooper pairs is determined solely by the coefficient  $v_{\mathbf{k}}^*$ , and this phase coincides with the phase of the gap function  $\Delta_{\mathbf{k}}$ .

**Need to finish this paragraph still.** This demonstrates immediately that  $\hat{N}_p = -i\partial/\partial\varphi$  suggesting that particle number and phase are canonically conjugated variables, i.e. there should be a Heisenberg uncertainty relation between both quantities.

### C. The gap equation

Let us now determine an explicit expression for gap function  $\Delta_{\mathbf{k}}$ , given self-consistently by Eq. (22). We start by expressing the electronic operators in terms of the Bogoliubons' using the transformation in Eq.(23). We obtain

$$\Delta_{\mathbf{k}} = -\frac{1}{N} \sum_{\mathbf{k}'} V_{\mathbf{k}\mathbf{k}'} u_{\mathbf{k}'}^* v_{\mathbf{k}'} \left( \langle \gamma_{-\mathbf{k}'\downarrow} \gamma_{-\mathbf{k}'\downarrow}^\dagger \rangle - \langle \gamma_{\mathbf{k}'\uparrow}^\dagger \gamma_{\mathbf{k}'\uparrow} \rangle \right) \quad (29)$$

Now, since the Bogoliubons follow the Fermi-Dirac distribution and have an energy dispersion  $E_{\mathbf{k}}$ , one has that

$$\left\langle \gamma_{-\mathbf{k}'\downarrow}\gamma_{-\mathbf{k}'\downarrow}^\dagger \right\rangle - \left\langle \gamma_{\mathbf{k}'\uparrow}^\dagger\gamma_{\mathbf{k}'\uparrow} \right\rangle = \left(1 - \frac{1}{e^{\beta E_{\mathbf{k}'} + 1}}\right) - \frac{1}{e^{\beta E_{\mathbf{k}'} + 1}} = \tanh\left(\frac{E_{\mathbf{k}'}}{2k_B T}\right). \quad (30)$$

The  $u_{\mathbf{k}'}^* v_{\mathbf{k}'}$  factor can then be explicitly calculated using Eqs.(25) yielding the gap equation

$$\Delta_{\mathbf{k}} = -\frac{1}{N} \sum_{\mathbf{k}'} \frac{V_{\mathbf{k}\mathbf{k}'} \Delta_{\mathbf{k}'}}{2E_{\mathbf{k}'}} \tanh\left(\frac{E_{\mathbf{k}'}}{2k_B T}\right) \quad (31)$$

We can now study for which values of the potential  $V_{\mathbf{k}\mathbf{k}'}$  and of the temperature  $T$  we obtain a non-zero gap, and therefore the BCS solution discussed in the previous section.

To proceed, we need to discuss the form of the potential. From a phonon-mediated electronic interaction standpoint, we consider a constant attractive potential  $V_{\mathbf{k}\mathbf{k}'} = -V_0$  for a shell of thickness  $\hbar\omega_D$  (with  $\omega_D$  the Debye frequency, i.e the "cutoff" frequency where no phonon modes exist in the approximation that atomic vibrations can be treated as phonons confined in the solid's volume) around the Fermi energy, i.e both  $|\xi_{\mathbf{k}}|, |\xi_{\mathbf{k}'}| < \hbar\omega_D$ . Since the potential does not depend on  $\mathbf{k}, \mathbf{k}'$ , we look for a gap function that is also  $\mathbf{k}$  independent and real, meaning that  $\Delta_{\mathbf{k}} = \Delta_{\mathbf{k}'} = \Delta$ . This type of gap function is called an *s*-wave gap. Given this discussion we obtain

$$1 = -\frac{V_0}{N} \sum_{k < k_D} \frac{1}{2E_{\mathbf{k}}} \tanh\left(\frac{E_{\mathbf{k}}}{2k_B T}\right) = V_0 \rho_F \int_{-\hbar\omega_D}^{\hbar\omega_D} \frac{d\varepsilon}{2\sqrt{\varepsilon^2 + \Delta^2}} \tanh\left(\frac{\sqrt{\varepsilon^2 + \Delta^2}}{2k_B T}\right) \quad (32)$$

where we used the fact that  $\hbar\omega_D \ll \mu$  to approximate the density of states *per spin*  $\rho(\varepsilon_{\mathbf{k}})$  by its value at the Fermi level  $\rho_F$ , remembering that  $\xi_{\mathbf{k}} = \varepsilon_{\mathbf{k}} - \mu$ . This self-consistent equation gives the gap function for an arbitrary temperature  $\Delta(T)$ .

Let us now study limiting behaviors of that expression. For example, see that at absolute zero  $T = 0$  the argument of the tanh goes to infinity. Since  $\tanh(x \rightarrow \infty) \rightarrow 1$ , and denoting  $\Delta_0 \equiv \Delta(T = 0)$ , the evaluation of the integral becomes straightforward, giving rise to an  $\arcsin(\hbar\omega_D/\Delta_0)$  term. Moreover, in most cases  $\Delta_0$  is of the order of a few meV, much smaller than  $\hbar\omega_D$ , which is of the order of a few hundreds of meV, allowing us to expand the  $\arcsin(x)$  for large  $x$ . This treatment yields the gap equation at absolute zero

$$\Delta_0 = 2\hbar\omega_D e^{-\frac{1}{V_0 \rho_F}}. \quad (33)$$

This equation tells us that an arbitrarily small attractive interaction  $V_0$  gives rise to a finite gap at zero temperature, showing that the Fermi liquid state is unstable towards the formation of the BCS superconducting state. We also see that superconductivity is a non-perturbative effect, given the dependence on  $\exp(-1/V_0 \rho_F)$ .

### 1. Cooper pair insight

To provide additional insight into the derived gap equation, it is useful to consider the problem from the perspective of single Cooper pair formation (although with some unavoidably repetition of the procedure done above).

Consider two electrons, of total wavefunction  $\Psi(\mathbf{r}_1, \mathbf{r}_1)$ , that interact with each other via an attractive potential  $V(\mathbf{r}_1 - \mathbf{r}_2)$ , with  $\mathbf{r}_1, \mathbf{r}_2$  their position vector. As usual, one proceeds by defining the relative displacement  $\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$ , the position of the center of mass  $\mathbf{R} = (\mathbf{r}_1 + \mathbf{r}_2)/2$ ,  $m^* = 2m$  the total mass and  $\mu = m/2$  the reduced mass. In these point of reference, the potential will not depend on  $\mathbf{R}$  and thus we look for the solution  $\Psi(\mathbf{r}, \mathbf{R}) = \psi(\mathbf{r}) \exp(i\mathbf{K} \cdot \mathbf{R})$ . This yields the Schrodinger equation

$(-\hbar/2\mu\nabla_{\mathbf{r}}^2 + V(\mathbf{r}))\psi(\mathbf{r}) = \tilde{E}\psi(\mathbf{r})$  with  $\tilde{E} = E - \hbar^2 K^2/2m^*$ . For a given eigenvalue  $\tilde{E}$ , the lowest energy  $E$  is the one for which two electrons have opposite momenta because then the momentum of the center of mass  $K$  vanishes, for which  $\tilde{E} = E$ . Depending on the symmetry of the spatial part of the wave-function, even  $\psi(\mathbf{r}) = \psi(-\mathbf{r})$  or odd  $\psi(\mathbf{r}) = -\psi(-\mathbf{r})$ , the spins of the electrons will form either a singlet or a triplet state, respectively, in order to ensure the anti-symmetry of the total wavefunction (switching two fermions corresponds to a sign change). So, for example, if the spin part of the wavefunction is antisymmetric then the orbital part has to be even, meaning  $\ell = 0, 2, \dots$

Fourier transforming the obtained Schrödinger equation and defining an modified wavefunction as  $\Delta(\mathbf{k}) = (E - 2\varepsilon_{\mathbf{k}})\psi(\mathbf{k})$  with  $\varepsilon_{\mathbf{k}} = \hbar^2 k^2/2m$  the free electron energy, one obtains

$$\Delta(\mathbf{k}) = - \int \frac{d\mathbf{k}'}{(2\pi)^3} \frac{V(\mathbf{k} - \mathbf{k}')}{2\varepsilon_{\mathbf{k}} - E} \Delta(\mathbf{k}') \quad (34)$$

See that for the two electrons to form a bound state of binding energy  $E_b = 2\varepsilon_{\mathbf{k}} - E$ , one must have that a total energy that is smaller than the energy of two independent free electrons, i.e  $E < 2\varepsilon_{\mathbf{k}}$ . As done previously, one now considers a potential that is attractive  $V(\mathbf{k} - \mathbf{k}_0) = -V_0$  for  $|\varepsilon_{\mathbf{k}}|, |\varepsilon_{\mathbf{k}'}| < \hbar\omega_D$  and zero otherwise and look for a solution with constant  $\Delta(\mathbf{k}) = \Delta$ . Since this implies an even spatial wave-function, the spins of the two electrons must be anti-parallel, forming a singlet state. The angular dependence of the wave-function will be that of the  $Y_{00}$  spherical harmonic, hence why we refer to the gap function as an *s*-wave gap.

Expressing the gap in terms of the density of states per spin of the two-electron system  $\rho(\varepsilon) \propto \sqrt{\varepsilon}$  yields the equations that determines the value of the bound state energy  $E < 0$  as a function of the attractive potential  $-V_0$ , defining a minimum value  $V_0^{\min}$  of the attractive potential such that  $E \rightarrow 0^-$  remains negative. This would lead us to conclude that a bound state will only form if the attractive interaction is strong enough, however, in this exercise, we overlooked an uttermost important feature—that in the actual many-body system, only the electrons near the Fermi level will be affected by the attractive interaction. To mimic this property, we consider the attractive interaction only for the unoccupied electronic states above the Fermi energy  $\varepsilon_F$  such that  $\varepsilon_{\mathbf{k}'} - \varepsilon_F, \varepsilon_{\mathbf{k}} - \varepsilon_F < \hbar\omega_D$ . Since  $\hbar\omega_D \ll \varepsilon_F$ , we can approximate the density of states for its value at  $\varepsilon_F$ . Moreover, in the limit of small  $V_0\rho_F \ll 1$ ,  $E$  is close to  $2\varepsilon_F$ , and we can approximate  $2\varepsilon_F - E + 2\hbar\omega_D \approx 2\hbar\omega_D$ . The binding energy  $E_b$  then follows exactly as in Eq.(33), apart from additional factors of two as a consequence of the fact that as  $\Delta \rightarrow 0$  one obtains  $E \rightarrow |\xi|$  in absolute value, meaning that it contains two branches of particle-hole excitations, doubling the density of states. This demonstrates that a bound state will ultimately form regardless of how weak the attractive interaction is. Such a bound state is called a Cooper pair. See that this is fundamentally different from the free electron case we considered before, where the attractive interaction has to overcome a threshold to create a bound state. The key property responsible for this different behavior is the existence of a well-defined Fermi surface, separating states that are occupied from states that are unoccupied.

## 2. Critical temperature

Furthermore, to determine at what critical temperature  $T_c$  a non-zero gap first appears we go back to Eq. (32), send  $\Delta \rightarrow 0$ , and use the fact that  $\hbar\omega_D \gg k_B T_c$ . The superconducting transition temperature is then found to be

$$T_c = \frac{2e^{\gamma_E}}{\pi} \frac{\hbar\omega_D}{k_B} e^{-\frac{1}{V_0\rho_F}} \quad (35)$$

with  $\gamma_E \approx 0.577$  the Euler constant. See that  $T_c$  depends on  $\exp(-1/V_0\rho_F)$ , same as the gap  $\Delta_0$ , being non-zero for any arbitrarily small  $V_0$ . This allows us to define the universal ratio  $\Delta_0/k_B T_c \approx 1.76$ . One of the early successes of BCS theory was the verification that this relationship is approximately satisfied

in most of the known superconductors at the time. Furthermore, see that BCS theory also addresses the isotope effect since  $T_c$  depends linearly on the Debye frequency  $\omega_D$ , which in turn varies as the inverse square root of the ionic mass  $M$ , i.e.  $T_c \propto \omega_D \propto M^{-1/2}$ , in agreement with the experimental observations.

#### D. Spin-triplet $p$ -wave superconductivity

## Part II

# Introduction to topological superconductivity

## IV. CONCEPTS OF SYMMETRY AND TOPOLOGY

Topology studies whether objects can be transformed continuously into each other. In condensed matter physics we can ask whether the Hamiltonians of two different systems can be continuously transformed into each other. If that is the case, then we can say that two systems are ‘topologically equivalent’.

In order to understand the concept of topology in condensed matter in the simplest way possible let us consider the transformation of a system described by the Hamiltonian  $\mathcal{H}$  by the tuning of some external parameter  $\alpha$  such that at  $\mathcal{H}_i \equiv \mathcal{H}(\alpha = 0)$  is the initial state Hamiltonian and  $\mathcal{H}_f \equiv \mathcal{H}(\alpha = 1)$  the final. Understand that the transformation of  $\mathcal{H}$  must be physical, meaning that it should be just a matter of point of view. Because of this, not only must  $\mathcal{H}$  be an hermitian matrix, i.e  $\mathcal{H} = \mathcal{H}^\dagger$  (such that it has real eigenenergies), but also that any transformation must be isometric (aka norm-preserving) isomorphisms (aka one-to-one mapping). Due to Wigner’s theorem these transformations can either be unitary  $\mathcal{U}$  or anti-unitary  $\bar{\mathcal{U}}$ . A unitary transformation between two inner product spaces reads as  $\langle \mathcal{U}\varphi | \mathcal{U}\psi \rangle = \langle \varphi | \psi \rangle$  while an anti-unitary transformation reads instead as  $\langle \bar{\mathcal{U}}\varphi | \bar{\mathcal{U}}\psi \rangle = \langle \varphi | \psi \rangle^* = \langle \psi | \varphi \rangle$ . Of course, any anti-unitary operator can be written as the product of a unitary operator and the complex conjugation operator  $\mathcal{K}$ .

### *Unitary transformations*

Unitary transformations do not have particularly interesting consequences for topological classification. Consider an Hamiltonian  $\mathcal{H}$  with the symmetry constraint  $\mathcal{U}^\dagger \mathcal{H} \mathcal{U} = \mathcal{H}$ . See that  $\mathcal{H}$  commutes with  $\mathcal{U}$  meaning that the system has a conservation law, and that the Hamiltonian can be brought to a block-diagonal form

$$\mathcal{H} = \left( \begin{array}{c|c} \mathcal{H}^{(1)} & \\ \hline & \mathcal{H}^{(2)} \end{array} \right) , \text{with } \mathcal{H}^{(n)} = \left( \begin{array}{c|c} h_{11} & h_{12} \\ \hline h_{12}^* & h_{22} \end{array} \right). \quad (36)$$

This procedure can be repeated until one runs out of unitary symmetries and is left with an irreducible block of the Hamiltonian, i.e. one which cannot be block diagonalized. In this case, every one of those  $\mathcal{H}_i^{(n)}$  Hamiltonians at the  $n$  block-diagonal could be continuously deformed into  $\mathcal{H}_f^{(n)}$ , meaning that they are always topologically equivalent.

### *Introduction to CPT symmetries*

One the other hand, anti-unitary transformations do impose constraints on an irreducible Hamiltonian, for example, by forcing it to maintain a (physically) finite energy gap, or to be a real matrix, or to be block off-diagonal. In this case, telling if  $\mathcal{H}_i$  and  $\mathcal{H}_f$  are topologically equivalent is not trivial. There are three fundamental discrete symmetries: chiral symmetry (CS)  $\mathcal{C}$ , parity symmetry  $\mathcal{P}$ , time-reversal symmetry (TRS)  $\mathcal{T}$ , known collectively as CPT symmetry. In a condensed matter picture, we often refer to the chiral symmetry as being a sublattice lattice and the parity symmetry as a particle-hole symmetry (PHS). Sublattice symmetry means that our system can be naturally split into

two interpenetrating sublattices. The Hamiltonian connects only sites from these different sublattices and, as a result, it anticommutes with an operator that distinguishes between them. Particle-hole symmetry means that for every electronic state with energy  $\varepsilon$  there is a corresponding electron-hole (as in absence of an electron) state, at  $-\varepsilon$ . Hence, mirroring the electron's occupancy along the Fermi level, meaning that occupied becomes unoccupied and vice versa, the spectrum remains unchanged. Finally, time reversal symmetry means that our system would have behave the same if time flown backwards. In this backward time frame momentums change sign and spins flip.

There is, however, an important detail: both  $\mathcal{T}$  and  $\mathcal{P}$  are indeed anti-unitary transformations but  $\mathcal{C}$  is not. This is because whenever a system has both TRS and PHS there is also a chiral symmetry  $\mathcal{C} = \mathcal{PT}$ . This also means that if a system only has either but not both, it cannot have a chiral symmetry. In other words, the presence of any two out of the three symmetries implies that the third is also present. Since the product of two anti-unitary operators is a unitary operator then  $\mathcal{C}$  must be unitary. Also, see that if both TRS and PHS are absent, then CS may or may not be present. In these two situations, formally known as *classes*, there are no anti-unitary symmetries, furthering their classification to *complex classes*.

Another important detail is that for TRS we have that  $[\mathcal{H}, \mathcal{T}] = 0$  while for PHS we have that  $\{\mathcal{H}, \mathcal{P}\} = 0$ . By implication of what we just talked, also  $\{\mathcal{H}, \mathcal{C}\} = 0$ .

Furthermore, as the next and final note about this symmetries, know that TRS and PHS may come in two separate flavors, depending on whether they square to plus or minus one. So, for example, a system can behave in three ways concerning TRS: (1) it does not have TRS, (2) it has it and  $\mathcal{T}^2 = +1$  (3) it has it and  $\mathcal{T}^2 = -1$ . On the other hand, the chiral symmetry only comes in one flavor,  $\mathcal{C}^2 = +1$ . Due to flavor combinations we find a total of 10 distinct symmetry classes displayed in Fig.(5). The classifications  $\mathbb{Z}$ ,  $2\mathbb{Z}$  and  $\mathbb{Z}_2$  on the left are to be introduced in the following examples.

class	$\mathcal{C}$	$\mathcal{P}$	$\mathcal{T}$	$d = 0$	1	2	3
A				$\mathbb{Z}$		$\mathbb{Z}$	
AI			1	$\mathbb{Z}$			
AII			-1	$2\mathbb{Z}$		$\mathbb{Z}_2$	$\mathbb{Z}_2$
AIII	1				$\mathbb{Z}$		$\mathbb{Z}$
BDI	1	1	1	$\mathbb{Z}_2$	$\mathbb{Z}$		
C		-1				$2\mathbb{Z}$	
CI	1	-1	1			$2\mathbb{Z}$	
CII	1	-1	-1		$2\mathbb{Z}$	$\mathbb{Z}_2$	
D		1		$\mathbb{Z}_2$	$\mathbb{Z}_2$	$\mathbb{Z}$	
DIII	1	1	-1	$\mathbb{Z}_2$	$\mathbb{Z}_2$	$\mathbb{Z}_2$	$\mathbb{Z}$

Figure 5. Symmetry periodic table with Altland-Zirnbauer classification. For more details on this table, for example, on how to go from  $d = 0$  to  $d > 0$  by adding and removing symmetries and it's Boot clock patterns see Akhmerov's "10 symmetry classes and the periodic table of topological insulators" at [https://topocondmat.org/w8\\_general/classification.html](https://topocondmat.org/w8_general/classification.html).

It is important to have in mind that CPT symmetries may not be the only symmetries at play. Although these are the fundamental symmetries, if one works within a condensed matter framework, the underlying lattice will provide additional, often spatial, symmetries. These include, for example point group symmetries—involution, mirror, and rotational symmetries—, and space group symmetries—

translation, glide, or screw symmetries of the entire crystal lattice. Point group symmetries protect additional degeneracies or enforce selection rules that are not captured by the non-spatial discrete symmetries alone, for example, a mirror symmetry in a crystal that protects gapless modes on certain surfaces or edges that are invariant under reflection. One the other hand space group symmetries constrain the electronic band structure and can lead to phenomena like Dirac or Weyl points that interact with the superconducting pairing.

### A. Introduction to topological invariants in 0D models

In order to study the effects of these symmetries, let us imagine a panoply of different systems and their energy spectrums as a function of  $\alpha$ . Moreover, let us count the number of levels below zero energy (defined at the Fermi level  $\varepsilon_F$ ) at each different  $\alpha$ , denoting it with  $Q$ . This will be our topological invariant prototype. If  $Q$  is the same in the initial and final system and did not change along the tuning of  $\alpha$  then there must be a continuous transformation Hamiltonian which does not close the gap. One the other hand, if  $Q$  changes then the system are *not* topologically equivalent as it would be needed to close the (physically real) gap. Hence, such a crossing changes the topological invariant, dubbed topological phase transition.

For all the examples that follow we assume a zero-dimensional ( $d = 0$ ) system. In a condensed matter realization this could be, for instance, a quantum dot interaction with all kinds of external systems. This will become




Figure 6. Kitaev chain Majorana modes pairing possibilities tbh I still don't fully understand why I can't just change the Fermi energy? I mean, for the spinful TRS I could only get 0 or 2 still. Does the spectrum of the CS and PHS just translate along with  $\varepsilon_F$  while the unitary does not?

#### 1. Time-reversal symmetry

Time-reversal symmetry is represented by an anti-unitary operator, and as such it can always be written as the product  $\mathcal{T} = U\mathcal{K}$  with  $U$  an unitary matrix and  $\mathcal{K}$  complex conjugation. A real Hamiltonian is a manifestation of time-reversal symmetry.

##### *Spinless case*

For example, for a spinless system we have  $\mathcal{T} = \mathcal{K}$  and thus  $\mathcal{T}\mathcal{H}\mathcal{T}^{-1} = \mathcal{H}^* = \mathcal{H}$  is a real matrix. In this case the TRS flavor is positive i.e  $\mathcal{T}^2 = +\mathbb{1}$ . Still, this case is also not interesting because is not different from the previous one, the different energy levels move and the topological invariant

changes by one when one of them crosses zero. In this trivial case the topological invariant is an integer number,  $Q = 0, \pm 1, \pm 2, \dots \in \mathbb{Z}$ . I mean, it should be  $\mathbb{N}$  no? How would the negative numbers appear?

### Spinful case

There is, however, a very important case where time-reversal symmetry makes a real difference. For a 1/2-spin system we the time-reversal operator reads  $\mathcal{T} = i\sigma_y \mathcal{K}$  with  $\sigma_y = [ 0 \ -i ; +i \ 0 ]$  the 2nd Pauli matrix (we reserve  $\sigma$  for Pauli matrices in spin orbital space). In this case the flavor is negative, i.e  $\mathcal{T}^2 = -\mathbb{1}$ , and  $\mathcal{T}\mathcal{H}\mathcal{T}^{-1} = \sigma_y \mathcal{H}^* \sigma_y = \mathcal{H}$  meaning that every energy eigenvalue  $\varepsilon$  is doubly degenerate. This happens because both the electrons with spin up or down have the same eigenenergy. This doubly degeneracy is often refer to as Kramers' degeneracy. Such a Hamiltonian would read in matrix form as

$$\mathcal{H} = \left( \begin{array}{c|c} \varepsilon_1 \mathbb{1} & M \\ \hline M^\dagger & \varepsilon_2 \mathbb{1} \end{array} \right) = \left( \begin{array}{cc|cc} \varepsilon_1 & 0 & m_{11} & m_{12} \\ 0 & \varepsilon_1 & -m_{12}^* & m_{11}^* \\ \hline m_{11}^* & -m_{12} & \varepsilon_2 & 0 \\ m_{12}^* & m_{11} & 0 & \varepsilon_2 \end{array} \right). \quad (37)$$

with  $\varepsilon_1, \varepsilon_2$  real numbers.

We can see the consequences of Kramers' degeneracy on the band spectrum versus  $\alpha$  in Fig.(6). While the spectrum looks quite similar to the previous ones, whenever a line crosses zero energy, our topological invariant makes a jump of two, and not one! In this case, time-reversal symmetry constrains the topological invariant to only take even values,  $Q = 0, \pm 1, \pm 2, \dots \in 2\mathbb{Z}$ . This is an example of how topological properties can be influenced by discrete symmetries.

### 2. Sublattice symmetry

We just saw that time-reversal symmetry can forbid the topological invariant to take a certain set of values. We now study another case where a symmetry changes the topological properties dramatically.

Let's now take a system where we can split all the degrees of freedom into two groups—group  $A$  and group  $B$ —, such that the Hamiltonian only has nonzero matrix elements between two groups, and not inside each group. This situation arises naturally when the a given lattice has two sublattices. For example, for hexagonal boron nitrate (hBN) we can distinguish these sublattices as the boron and nitrogen sites respectively. The matrix of such an Hamiltonian would read

$$\mathcal{H} = \left( \begin{array}{c|c} & M \\ \hline M^\dagger & \end{array} \right) = \left( \begin{array}{cc|cc} 0 & 0 & m_{11} & m_{12} \\ 0 & 0 & m_{21} & m_{22} \\ \hline m_{11}^* & m_{21}^* & 0 & 0 \\ m_{12}^* & m_{22}^* & 0 & 0 \end{array} \right). \quad (38)$$

See that  $\eta_z \mathcal{H} \eta_z = -\mathcal{H}$  where  $\eta_z = [ +1 \ 0 ; 0 \ -1 ]$  is the 3rd Pauli matrix (we reserve  $\eta$  for Pauli matrices in sublattice orbital space). This immediately means that if  $\Psi = [ \psi_A ; \psi_B ]^T$  is an eigenvector of the Hamiltonian with energy  $\varepsilon$ , then  $[ \psi_A ; -\psi_B ]^T$  is an eigenvector with energy  $-\varepsilon$ . A symmetric spectrum is the consequence of sublattice symmetry as seen in Fig.(6). This means that  $Q$  always stays constant and that we can always deform Hamiltonians with sublattice symmetry without closing the gap. This indicates that an extra symmetry, such as this one, may render the topology of a system as trivial.

### 3. Particle-hole symmetry

Another symmetry that has a strong influence on topology is the particle-hole symmetry, showing up in superconducting systems. As we saw in BCS theory, a superconductor will create(annihilate) pairs of electrons by breaking(forming) Cooper pairs costing a pairing energy of  $\Delta$  to the system.

Let us consider that the dynamics of the electrons is described by the an hermitian  $H$  matrix while the pair creation and annihilation is described by an antisymmetric  $\Delta$  matrix. Understand that  $\Delta$  must antisymmetric just because because the fermion operators anticommute. The Hamiltonian describing the full system reads

$$\mathcal{H} = \left( \begin{array}{c|c} H & \Delta \\ \hline -\Delta^* & -H^* \end{array} \right) = \left( \begin{array}{cc|cc} h_{11} & h_{12} & 0 & \Delta \\ h_{12}^* & h_{22} & -\Delta & 0 \\ \hline 0 & -\Delta^* & -h_{11}^* & -h_{12}^* \\ \Delta^* & 0 & -h_{12} & -h_{22}^* \end{array} \right) \quad (39)$$

and is know as the Bogoliubov-de Gennes (BdG) Hamiltonian. Moreover, we now double the amount of degrees of freedom in the system by defining a Nambu spinors

$$\check{c}_i^\dagger = \begin{pmatrix} c_i^\dagger & c_i \end{pmatrix} \quad \text{and} \quad \check{c}_i = \begin{pmatrix} c_i \\ c_i^\dagger \end{pmatrix} \quad (40)$$

such that we can write

$$\check{\mathcal{H}} = \frac{1}{2} \check{c}^\dagger \mathcal{H} \check{c}.$$

This definitions indicates that the Bogoliubov-de Gennes Hamiltonian acts not only on electrons but also on an extra mirror set comprised of eletron-holes. Since holes are related to the electrons,  $\mathcal{H}$  automatically inherits that extra symmetry. This symmetry exchanges electrons with holes, and has an anti-unitary operator  $\mathcal{P} = \tau_x \mathcal{K}$  with  $\tau_x = [0 \ 1; 1 \ 0]$  the 1st Pauli matrix (we reserve  $\sigma$  for Pauli matrices in spin orbital space) and (as before)  $\mathcal{K}$  complex conjugation. Hence we have that  $\mathcal{P}\mathcal{H}\mathcal{P}^{-1} = -\mathcal{H}$ . For this specific case it's flavor is positive, i.e  $\mathcal{P}^2 = +1$ . Indeed, for every eigenvector  $\Psi = [u; v]^T$  with energy  $\varepsilon$ , there will be a particle-hole symmetric eigenvector  $\mathcal{P}\Psi = [v^*; u^*]^T$  with energy  $-\varepsilon$ . As clearly seen in Fig.(6), because of the minus sign in the particle-hole symmetry, the spectrum of  $\mathcal{H}$  must be mirrored around zero energy, that is, the Fermi level).

#### Fermionic parity switches

See that this spectrum mirroring was also the case for sublattice symmetry however, in this case, energy levels do not repel around zero energy, so that crossings at zero energy appear. Unlike in the case of sublattice symmetry, a pair of  $\pm\varepsilon$  energy levels does not corresponds to two distinct quantum states, but to a single quantum state. This quantum state is a coherent superposition of electrons and holes, a so called Bogoliubov quasiparticle. It has an excitation energy  $\varepsilon$ , and it is created by an operator  $\gamma^\dagger = uc^\dagger + vc$ . Populating the partner state at energy  $\varepsilon$  is the same as emptying the positive energy state.

In general a crossing between energy levels happens in the presence of a conserved quantity. While the mean-field Hamiltonian of a superconductor does not conserve the number of particles, it conserves the parity of this number. In other words, forming and breaking Cooper pairs does not affect whether the superconducting contains an even or odd number of electrons so fermion parity is a conserved quantity (provided that isolated electrons do not enter or leave the system). Fermion parity, however, is a many-body quantity, which cannot be directly described in terms of the single particle picture of the BdG Hamiltonian. This is why we had to double the number of degrees of freedom by hand. When a pair of levels crosses zero energy, the excitation energy  $\varepsilon$  of the Bogoliubov quasiparticle changes sign and it becomes favorable to add(remove) a Bogoliubov quasiparticle. In other words, at each crossing the fermion parity in the ground state changes from even to odd (or vice versa), meaning that these crossings are fermion parity switches.

### The Pfaffian invariant

Since the ground state fermion parity is preserved by the superconducting Hamiltonian if there are no Bogoliubov quasiparticles crossing zero energy, the ground state fermion parity is the topological invariant of this system. It is clear however that this invariant is of a different nature than the one of the non-superconducting systems, which is given by the number  $Q$  of negative eigenvalues of the Hamiltonian. The latter cannot change for a BdG Hamiltonian, which has a symmetric energy spectrum, and hence it is not suitable to describe changes in fermion parity. For this kind of systems the actual topological invariant is called the *Pfaffian* and will either take the value  $Q = \pm 1 \in \mathbb{Z}_2$  at every zero-energy crossing. Its rigorous definition is not really that important for our sake so we take a simpler approach.

In order to introduce the Pfaffian invariant, we start by making a basis transformation  $\mathcal{H}'_{\text{BdG}} = \mathcal{U}\mathcal{H}_{\text{BdG}}\mathcal{U}^\dagger$  that makes the Hamiltonian an skew-symmetric matrix, i.e  $\mathcal{H}^T = -\mathcal{H}$ . We do this because the eigenvalues of antisymmetric matrices always come in pairs, i.e  $\pm \varepsilon_n$ . Further reasoning will become apparent as we go. Such a transformation is

$$\mathcal{H}'_{\text{BdG}} = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix} \mathcal{H}_{\text{BdG}} \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix} = \frac{1}{2} \begin{pmatrix} H - H^* + \Delta - \Delta^* & |i(-H - H^* + \Delta + \Delta^*) \\ i(+H + H^* + \Delta + \Delta^*) & H - H^* - \Delta + \Delta^* \end{pmatrix}$$

Indeed, because  $H$  is Hermitian then  $H - H^*$  is antisymmetric and  $H + H^*$  is symmetric, i.e  $\mathcal{H}^T = \mathcal{H}$ ; since  $\Delta$  is antisymmetric then  $\mathcal{H}'_{\text{BdG}}$  is also antisymmetric. In its diagonalized form the determinant of this matrix is just the product of the pairs of eigenenergies, i.e  $\det(\mathcal{H}) = \prod_n (-\varepsilon_n^2)$ . The key feature of the Pfaffian is revealed when taking now the square root of the determinant  $\text{Pf}(\mathcal{H}) = \sqrt{\det(\mathcal{H})} = \pm \prod_n i\varepsilon_n$ . See that it is defined in such a way that the sign of the product is uniquely defined. At a fermion parity switch a single  $\varepsilon_n$  changes sign, so the Pfaffian changes sign as well while the determinant stays the same. We then define the actual topological invariant as

$$Q_{\text{BdG}} = \text{sign} [\text{Pf}(i\mathcal{H})],$$

where we have included a factor of  $i$  just that the Pfaffian is a real number, such that at  $Q_{\text{BdG}}$  changes its value from  $+1$  to  $-1$  at every zero-energy crossing. This means that it is the correct expression for the ground state fermion parity and for the topological invariant. As some sort of intuition, you can think of it as if the number of holeonic levels below zero energy counts negatively to the overall positive electronic levels.

#### 4. Combining symmetries

##### Particle-hole and spinful time-reversal symmetry

Take a system that has both particle-hole symmetry (PHS) and spinful time-reversal symmetry (TRS) described by the Hamiltonian  $\mathcal{H}$ . Let us take an intuitive approach to the band spectrum analysis. By PHS we know that an electronic band is equivalent to a negative holeonic band. On the other hand, by spinful TRS we know that there is Kramer degeneracy. Hence, since a PHS holeonic band counts as negative to the number of bands below zero energy we will always end up with  $Q$  being even and changing sign at a crossing. **This is wrong but can't see the flaw in logic. I mean, looking at the table I can see that  $P^2 = 1$  and  $T^2 = -1$  gives me no constrain on  $Q$  and thus trivial topology.**

## B. Introduction to topological invariant in higher dimensions

In higher dimensional system the discrete energy levels of a  $d = 0$  system are replaced by continuous energy bands defined along the Brillouin zone. In these higher dimensions the topological invariant

cannot be defined merely as counting levels below the Fermi energy or by tracking sign changes of the Pfaffian in superconducting systems. Instead, the central theme of  $d > 0$  dimensional band topology lies in the concept of geometric phases.

As an illustrative example of the concepts to come, consider a vector placed at the earth's north pole, always pointing in the tangent direction to the surface. If one translates the vector to the equator along a meridian, then along the equator for some distance, and back to the north pole, the vector's orientation will have changed relative to how it started by some angle. This angle is called the holonomy. The origin of non-trivial band topological properties is not so different from this example, with the crucial replacement of the vector by a Hamiltonian  $\mathcal{H}(\boldsymbol{\alpha})$  depending on a set of parameters  $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_N)$ , and the earth a manifold (topological space that locally resembles Euclidean space near each point) spanned by those parameters. In the context of Hamiltonians, holonomy manifests as the acquisition of additional geometric phases by the eigenstate of  $\mathcal{H}(\boldsymbol{\alpha})$  as the parameter space manifold is traversed. More concretely, in band topology,  $\boldsymbol{\alpha}$  is taken to be the momenta  $\mathbf{k} = (k_1, \dots, k_d)$ , with  $d$  the space dimensions, together with a set of additional tunable parameters (chemical potential, electric field, Zeeman, integrated out pairing, etc, etc...), and the eigenstate's additional geometric phase is the so-called Berry phases (we will further explore this later concept in just a moment). In this context, the restriction that evolution is adiabatic simply means that the system must remain in a situation where energy bands do not cross, i.e. the system must be gapped.

The idea is that if the parameters  $\boldsymbol{\alpha}$  are varied adiabatically, then at each subsequent value of  $\boldsymbol{\alpha}$ , eigenstates of one set of parameters are smoothly deformed into another set. This is the content of the adiabatic theorem, which states that in the case of adiabatic evolution of the parameters along a curve  $\boldsymbol{\alpha}(t)$ , the Schrodinger equation

$$-i\hbar\partial_t |\psi_n(\boldsymbol{\alpha}(t))\rangle = \varepsilon_n(\boldsymbol{\alpha}(t)) |\psi_n(\boldsymbol{\alpha}(t))\rangle \quad (41)$$

is obeyed instantaneously. Here  $|\psi_n(\boldsymbol{\alpha}(t))\rangle$  represents the eigenstate of the Hamiltonian  $\mathcal{H}(\boldsymbol{\alpha}(t))$  in the  $n$ th band with energy  $\varepsilon_n(\boldsymbol{\alpha}(t))$ . Now, generically, due to the structure of the Schrodinger equation, and the normalization of states, a single degree of freedom exists, which can change the eigenstate as it is moved along the parameter space  $\boldsymbol{\alpha}(t)$ . This corresponds to a phase denoted by  $\theta(t)$ , such that the state can be decomposed as

$$|\psi(\boldsymbol{\alpha}(t))\rangle = e^{i\theta(t)/\hbar} |\phi(\boldsymbol{\alpha}(t))\rangle. \quad (42)$$

A short calculation performed by plugging this form of the state into the Schrodinger equation on both sides, and acting with  $\langle\psi(\boldsymbol{\alpha}(t))|$  on the left is enough to solve for the phase  $\theta(t)$ . One obtains

$$\theta(t) = \int_0^{t'} dt \left[ \varepsilon(\boldsymbol{\alpha}(t)) + \frac{i}{\hbar} \langle\phi(\boldsymbol{\alpha}(t))| \partial_t |\phi(\boldsymbol{\alpha}(t))\rangle \right] \quad (43)$$

There are two contributions to the phase acquired by the eigenstate under adiabatic evolution. The first term is the familiar dynamical phase  $\theta_D(t)$ , which is acquired from evolving in time in the Hilbert space. However, a second term appears, namely

$$\gamma(t) = \frac{i}{\hbar} \int_0^{t'} dt \langle\phi(\boldsymbol{\alpha}(t))| \partial_t |\phi(\boldsymbol{\alpha}(t))\rangle$$

which is called the geometrical phase or Berry phase. This phase can be calculated via the aforementioned time integral, or equivalently by integrating over the curve  $\mathcal{C}$  spanned in the parameter space  $\boldsymbol{\alpha}$  during the adiabatic evolution, reading

$$\gamma_{\mathcal{C}} = \int_{\mathcal{C}} d\boldsymbol{\alpha} \frac{i}{\hbar} \langle\phi(\boldsymbol{\alpha})| \nabla_{\boldsymbol{\alpha}} |\phi(\boldsymbol{\alpha})\rangle \equiv \int_{\mathcal{C}} d\boldsymbol{\alpha} \mathbf{A}(\boldsymbol{\alpha}) \quad (44)$$

with  $\mathbf{A}(\boldsymbol{\alpha})$  the so called Berry connection.

The Berry connection plays the same role in adiabatic evolution as the vector potential in electromagnetism, and indeed, much like in the latter theory, this connection can be used to construct a curvature tensor. In the context of electromagnetism, the curvature tensor is nothing but the electromagnetic tensor  $F_{\mu\nu}$ , while in the context of the adiabatic evolution of quantum systems, it is given the special name of Berry curvature  $\Omega_{\mu\nu}$ . Explicitly, this Berry curvature reads

$$\Omega_{\mu\nu}(\alpha) = \frac{\partial}{\partial\alpha^\mu}A_\nu(\alpha) - \frac{\partial}{\partial\alpha^\nu}A_\mu(\alpha)$$

One often considers the dual pseudo-vector to this tensor, this is  $\Omega_{\mu\nu} = \varepsilon_{\mu\nu\xi}\Omega^\xi$  with  $\varepsilon_{\mu\nu\xi}$  the Levi-Civita symbol, and calls that the Berry curvature instead. This quantity is analogous to the magnetic field  $\mathbf{B} = \nabla \times \mathbf{A}$ . Another similarity between electromagnetism and these concepts is that the Berry connection, like the magnetic vector potential, is defined only up to a gauge choice. This makes it so the Berry phase is only well defined if closed curves  $\mathcal{C}$  in the parameter space are considered.

As a final summary, information about the topology of the target space of  $\mathcal{H}(\alpha)$  is acquired by integrating the Berry connection or curvature over the entire Brillouin zone, or in other words, the holonomy of the Hamiltonian as the Brillouin zone is traversed is sensitive to the band-topology. The integration of this Berry curvature yields quantities called topological invariants which are analogous to the Winding number of the Aharonov-Bohm effect (see the example below). In this  $d > 0$  context, one can also refer to the topological invariants as Chern number.

A fundamental consequence of having a well-defined topological invariant in  $d > 0$  is the so-called bulk-boundary correspondence. This principle asserts that nontrivial topological properties in the bulk of a material inevitably give rise to robust, gapless modes at its boundaries, whether along edges in 2D or surfaces in 3D. One can intuitively see why this should be the case by noting that at the boundary of a topological non-trivial system there is only vacuum, a topological trivial system. This means that, at this boundary, the topological invariant must change from something non-zero to zero which is only possible if the gap closes. This gives rise to emergent gapless edge state which are protected against perturbations that do not close the bulk gap.

### 1. Aharonov-Bohm effect

As a predecessor to the topological band theory, we now introduce the reader to an electromagnetism examples known as the Aharonov-Bohm (A-B) effect as a starting point to understanding the Berry phase, connection, curvature in more detail.

Consider an electron whose movement is restricted to the  $xOy$  place where an infinitely thin and long solenoid pierces through it at its center. Inside the solenoid an electric current flow inducing a magnetic field  $\mathbf{B} = B\hat{z}$  such that a magnetic flux  $\phi$  flows penetrates the the plane of motion of the electron. Although there exists no field or flux outside the solenoid, a magnetic vector potential  $\mathbf{A}$  permeates all space. Now, note that the electron wandering the plane will actually be affected by the vector potential, in that the Hamiltonian describing it will have the form

$$\mathcal{H}(\mathbf{r}) = \frac{\hbar^2}{2m} (\nabla_{\mathbf{r}} - e\nabla_{\mathbf{r}} \cdot \mathbf{A}(\mathbf{r}))^2 \quad (45)$$

with  $\mathbf{p} = -i\hbar\nabla_{\mathbf{r}}$  the momentum operator,  $e$  the electron charge and  $m$  its mass. In this case, the parameters  $\alpha$  can be identified with the actual position of the electron  $\mathbf{r}$ . Since the electron cannot enter the solenoid, which is assumed to be placed at  $\mathbf{r} = 0$ , its movement is restricted to everywhere except there.

As explain in the previous section, as the electron moves following a curve  $\mathcal{C}$  it will acquire a Berry phase, or rather, is in this context, the A-B phase, given by Eq.(44) as

$$\gamma_{\text{A-B}} = \frac{e}{\hbar} \oint_{\mathcal{C}} d\mathbf{r} \mathbf{A}(\mathbf{r}), \quad (46)$$

with  $\mathbf{A}$  being not the general Berry connection but the actual physical vector potential. The A-B phase, in some sense, measures the inability of making a continuous gauge choice for the magnetic vector potential in a punctured plane. The presence of the puncture hole makes it so a discontinuity along a branch cut is a mathematical necessity, and as a result, if an electron loops around the hole, it will acquire a non-trivial phase, dependent only on the number of times it goes around the hole (see Fig. 7). For this reason, it is said that the A-B phase is a topological quantity, depending only on the topology of the electron's trajectory, namely on a quantity called the winding number.

Alternatively, through the usage of Stoke's theorem, it is simple enough to compute the A-B phase as being proportional to the magnetic flux, this is

$$\gamma_{\text{A-B}} = \frac{e}{\hbar} \oint dS \cdot \mathbf{B} = \frac{e}{\hbar} W\phi$$

enclosed by the trajectory's area  $S$ , where  $W \in \mathbb{Z}$  counts the number of loops the electron makes around the solenoid. It's precisely this quantity that corresponds precisely to the winding number.




Figure 7. (a) Holonomy, (b) Aharonov-Bohm effect

## 2. Quantum Thouless pump

## V. TOPOLOGICAL SUPERCONDUCTIVITY IN 1D MODELS

### A. Kitaev model

The *Kitaev chain* or *Kitaev-Majorana chain* is a toy model for a topological superconductor using a 1D hybrid (semiconductor+superconductor) nanowires featuring Majorana bound states. It consists of a 1D linear lattice of  $N$  sites and spinless fermions at zero temperature, subjected to nearest neighbor hoping interactions. The real-space tight-binding Hamiltonian describing such model reads

$$H = \mu \sum_{i=1}^N \left( c_i^\dagger c_i - \frac{1}{2} \right) - t \sum_{i=1}^{N-1} \left( c_{i+1}^\dagger c_i + h.c \right) + \Delta \sum_{i=1}^{N-1} \left( c_{i+1}^\dagger c_i^\dagger + h.c \right) \quad (47)$$

with  $c_i^\dagger$  ( $c_i$ ) fermionic creation (annihilation) operators,  $\mu$  the chemical potential,  $t$  the hopping energy and  $\Delta$  an proximity induced superconducting  $p$ -wave pairing.

The objective of this model definition is to be able to have a Majorana bound states on the edges mode. For this, let us engineering the Hamiltonian in such a special way that it is actually possible to separate two Majoranas. Foremost, we define each site  $n$  as if it has two sublattices,  $s = A$  and  $s = B$ .

We then define Majorana operators relating to the fermionic operators as

$$\gamma_i^A = c_i^\dagger + c_i \quad \text{and} \quad \gamma_i^B = i(c_i^\dagger - c_i) \quad (48)$$

or rather, in the opposite way, as

$$c_i^\dagger = \frac{1}{2}(\gamma_i^A - i\gamma_i^B) \quad \text{and} \quad c_i = \frac{1}{2}(\gamma_i^A + i\gamma_i^B) \quad (49)$$

Indeed, each site can host a fermion or, equivalently, each site hosting two Majorana modes. These Majorana operators are Hermitian  $\gamma_i^s = (\gamma_i^s)^\dagger$ , unitary  $(\gamma_i^s)^2 = 1$  and anticommute as  $\{\gamma_i^s, \gamma_j^{s'}\} = 2\delta_{ij}\delta_{ss'}$ .

Substituting directly into the Hamiltonian of Eq.(47) the fermionic operators as given by Eqs.(49) we obtain

$$H = -i\mu \frac{1}{2} \sum_{i=1}^N \gamma_i^B \gamma_i^A + i \frac{1}{2} \sum_{i=1}^{N-1} (\omega_+ \gamma_i^B \gamma_{i+1}^A + \omega_- \gamma_{i+1}^B \gamma_i^A), \quad \text{with } \omega_\pm = \Delta \pm t \quad (50)$$

From it we can distinguish between two phases—trivial and topological—, corresponding, respectively, to two different ways of pairing these Majoranas states—no unpaired modes or one isolated mode on both edges. These pairing configuration are depicted in Fig.8 in blue and red respectively. This phases can be easily identified, respectively, in their limiting regimes where one sets  $\Delta = t = 0$  and  $\mu = 0$  with  $\Delta = t \neq 0$ .




Figure 8. Kitaev chain Majorana modes pairing possibilities

Indeed, see that by setting  $\Delta = t = 0$  within the Hamiltonian of Eq.(50) we obtain

$$H_{\text{trivial}} = -i\mu \frac{1}{2} \sum_{i=1}^N \gamma_i^B \gamma_i^A, \quad (51)$$

which corresponds to the limiting case of "no unpaired Majorana modes" configuration. The energy cost for each fermion to be occupied is  $\mu$ , with all excitations having an energy of either  $\pm\mu/2$ . The band structure will then have a gapped bulk and no zero energy edge states. Furthermore, see that the wavefunctions of the first three energy states shown in Fig.(9).(middle) in this trivial phase simply resemble the harmonic modes of a string states.

On the other hand, see that by setting  $\mu = 0$  with  $\Delta = t \neq 0$  we obtain

$$H_{\text{topological}} = it \sum_{n=1}^{N-1} \gamma_n^B \gamma_{n+1}^A \quad (52)$$

which corresponds to the "unpaired edge Majorana mode" configuration where every Majorana operator is coupled to a Majorana operator of a different kind in the next site. Note that the summation only goes up to  $n = N - 1$ . Moreover, see that by assigning a new fermion operator  $\tilde{c}_i = 1/2(\gamma_i^B + i\gamma_{i+1}^A)$ , the Hamiltonian can be otherwise expressed as

$$H_{\text{topological}} = 2t \sum_{n=1}^{N-1} \left( \tilde{c}_n^\dagger \tilde{c}_n + \frac{1}{2} \right) \quad (53)$$

which describes a new set of  $N - 1$  Bogoliubov quasiparticles with energy  $t$ . For every Majorana pair we assign an energy difference  $2t$  between the empty and filled state. All states which are not at the ends of the chain have an energy of  $\pm t$  and thus the bands structure has a gapped bulk. However, see that the missing mode  $\tilde{c}_N = 1/2 (\gamma_N^B + i\gamma_1^A)$ , which couples the Majorana operators from the two endpoints of the chain, does not appear in the Hamiltonian and thus it most have zero energy. As the presence of this mode does not change the total energy, the ground state is two-fold degenerate. This condition is a topological superconducting non-trivial phase. This mode is called a Majorana zero mode and is highly delocalized at the edges, as it can be seen in red in Fig(9).(middle). As one tunes  $\mu$  in the direction of the trivial phase, the topological gap, protected by particle-hole symmetry (PHS), gets smaller and smaller and the Majoranas wavefunctions stay less and less localized at the edges. At the transition between the trivial and topological, when the chemical potential takes it's critical value of  $|\mu| = 2t$ , the first energy states stays evenly distributed along the chain.




Figure 9. Kitaev chain (top) band structure (middle) I will eventually plot the 1st, 2nd and 3rd state wavefunction here at each regime, and (bottom) band spectrum for a chain length of  $L = 50$  with lattice spacing  $a_0 = 1$  fixing  $\Delta = t = 1.0$ . The critical  $\mu$  shifts forward to infinity as  $L \rightarrow 0$ .

*Bogoliubov-de Gennes Hamiltonian* Let us now define the Hamiltonian in E.(47) in its Bogoliubov-de Gennes (BdG) form

$$H = \frac{1}{2} \check{c}^\dagger H_{\text{BdG}} \check{c}.$$

where we have defined the Nambu spinor as

$$\check{c}_i^\dagger = \begin{pmatrix} c_i^\dagger & c_i \end{pmatrix} \quad \text{and} \quad \check{c}_i = \begin{pmatrix} c_i \\ c_i^\dagger \end{pmatrix} \quad (54)$$

This proves not only useful to the study of the system's symmetries, but it also a necessary step for the numerical implementation in *Quantical.jl*. Defining  $\tau_x, \tau_y, \tau_z$  as Pauli matrices in the particle-hole

subspace and using we the fermionic anti-commutation properties  $\{c_i, c_j^\dagger\} = \delta_{ij}$  and  $\{c_i, c_j\} = 0$ , one can check that

$$\mu : \quad \check{c}_i^\dagger \tau_z \check{c}_i = \begin{pmatrix} c_i^\dagger & c_i \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} c_i \\ c_i^\dagger \end{pmatrix} = c_i^\dagger c_i - c_i c_i^\dagger = 2c_i^\dagger c_i - 1 \quad (55)$$

$$t : \quad \check{c}_j^\dagger \tau_z \check{c}_i = \begin{pmatrix} c_j^\dagger & c_j \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} c_i \\ c_i^\dagger \end{pmatrix} = c_j^\dagger c_i - c_j c_i^\dagger = c_j^\dagger c_i + h.c \quad (56)$$

$$\Delta : \quad \check{c}_j^\dagger i \tau_y \check{c}_i = \begin{pmatrix} c_j^\dagger & c_j \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} c_i \\ c_i^\dagger \end{pmatrix} = c_j^\dagger c_i^\dagger - c_j c_i = c_j^\dagger c_i^\dagger + h.c \quad (57)$$

Hence the Hamiltonian in Eq.(47) in its BdG form reads as

$$H = \mu \frac{1}{2} \sum_i \check{c}_i^\dagger \tau_z \check{c}_i - t \sum_{i=1}^{N-1} \check{c}_{i+1}^\dagger \tau_z \check{c}_i + \Delta \sum_{i=1}^{N-1} \check{c}_{i+1}^\dagger i \tau_y \check{c}_i \quad (58)$$

See that the Hamiltonian has particle-hole symmetry, i.e  $\mathcal{P} H \mathcal{P}^{-1} = -\tau_x H^* \tau_x = -H$  with  $\mathcal{P} = \tau_x \mathcal{K}$  and  $\mathcal{K}$  complex conjugation, as well as time reversal symmetry, i.e  $\mathcal{T} H \mathcal{T}^{-1} = H^* = H$  with  $\mathcal{T} = \mathcal{K}$  for this spinless case (for reference,  $\mathcal{T} = i\sigma_y \mathcal{K}$  for a 1/2-spin system). Once again, to understand why this is the case check.

### Topological invariant

#### 1. Majorana modes at a domain wall

Consider the case where we weld together two semi-infinite nanowires with one in it's trivial phase and the other in it's trivial phase. The spacial profile of the chemical potential  $\mu(x)$  would then approximately a Heaviside theta function from  $|\mu_{\text{left}}| > 2t$  to  $|\mu_{\text{right}}| < 2t$ , forming a doping domain wall at it's center. Hamiltonian wise, one just substitutes  $\mu \rightarrow \mu(x)$  directly into Eq.(47). What one obtains in this situation is a Majorana mode localized at the domain wall with its twin forming in the semi-infinite edge of the topological side.




Figure 10. needs caption

## 2. Kitaev ring

### B. SSH model

The most relevant references used for this section follow:

### C. Oreg-Lutchyn models

The Oreg-Lutchyn Majorana minimal model consists of a finite 1D semiconductor (SM) nanowire with strong spin-orbit coupling (SOC)  $\alpha$  and a tunable chemical potential  $\mu$ , in proximity of a superconductor (SC) of homogeneous pairing  $\Delta$ , having a magnetic field  $B_z$  applied along it's length, defined as the  $\hat{z}$  direction. The Rashba effect describes the coupling of an electric field  $E_x$  that breaks inversion symmetry breaking in the direction perpendicular to the wire, to the electron's spin, i.e  $\propto (i\vec{\nabla} \times \hat{x}) \cdot \vec{\sigma} = i\sigma_y \partial_z$  with  $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ . The Zeeman effect described the spin splitting due to the in-plane magnetic field  $B_z$ . The pairing term describes the Cooper pairs from BCS theory than could tunnel from the SM to the SC.

The tight-binding Hamiltonian describing such system can then be decomposed as

$$H = H_K + H_{SOC} + H_Z + H_{SC} \quad (59)$$

$$H_K = (2t - \mu) \sum_{i\sigma} c_{i\sigma}^\dagger c_{i\sigma} - t \sum_{\langle i,j \rangle \sigma} c_{i\sigma}^\dagger c_{j\sigma} \quad (60)$$

$$H_{SOC} = \frac{\alpha}{2a_0} \sum_{i\sigma} (c_{i+1\bar{\sigma}}^\dagger c_{i\sigma} + h.c.) \quad (61)$$

$$H_Z = V_Z \sum_i (c_{i\uparrow}^\dagger c_{i\uparrow} - c_{i\downarrow}^\dagger c_{i\downarrow}) \quad (62)$$

$$H_{SC} = \Delta (c_{i\downarrow}^\dagger c_{i\uparrow}^\dagger + h.c.) \quad (63)$$

with  $c_i^\dagger$  ( $c_i$ ) fermionic creation (annihilation) operators,  $\mu$  the chemical potential,  $t = \eta/a_0^2$  the hopping energy into  $\langle i,j \rangle$  nearest-neighbouring sites with  $a_0$  the lattice constant and  $\eta = \hbar^2/2m^*$  with  $m^*$  the effective mass of the electrons,  $V_Z = g_J \mu_B B_z / 2$  the Zeeman potential with  $g_J$  the Landé gyromagnetic moment and  $\mu_B$  Bohr's magneton,  $\alpha$  the Rashba SOC strength and  $\Delta$  proximity induced superconducting  $s$ -wave pairing.

A paragraph explaining the bands.




Figure 11.

A paragraph explaining the phase-diagram, pfaffian and band spectrum.




Figure 12.

*Bogoliubov-de Gennes Hamiltonian* Shown below are the broad strokes of a numerical implementation of the Hamiltonian in Julia using the Quantica.jl. However, prior to this implementation, we will be needing the Bogoliubov-de Gennes formalism. For this, need to double the degrees of freedom through the Nambu-spinor. In the so called unrotated-spin basis we define a Nambu spinor as

$$\check{c}_i^\dagger = \begin{pmatrix} c_i^\dagger & c_i \end{pmatrix} = \begin{pmatrix} c_{i\uparrow}^\dagger & c_{i\downarrow}^\dagger & c_{i\uparrow} & c_{i\downarrow} \end{pmatrix} \quad (64)$$

In this Nambu $\otimes$ spin orbital space the Hamiltonian in Eq.(59) reads

$$H = H_K + H_{SOC} + H_Z + H_{SC} \quad (65)$$

$$H_K = (2t - \mu) \sum_i \check{c}_i^\dagger [\tau_z \otimes \sigma_0] \check{c}_i - \frac{1}{2} t \sum_{\langle i,j \rangle} \check{c}_i^\dagger [\tau_z \otimes \sigma_0] \check{c}_j \quad (66)$$

$$H_{SOC} = \frac{\alpha}{2a_0} \sum_i \check{c}_i^\dagger [\tau_z \otimes i\sigma_y] \check{c}_{i+1} \quad (67)$$

$$H_Z = V_Z \sum_i \check{c}_i^\dagger [\tau_z \otimes \sigma_z] \check{c}_i \quad (68)$$

$$H_{SC} = \frac{1}{2} \Delta \sum_i \check{c}_i^\dagger [\tau_y \otimes \sigma_y] \check{c}_i \quad (69)$$

with  $\tau$  Pauli matrices in the particle-hole subspace and  $\sigma$  in the spin subspace.

To understand why this is the case check we show explicitly the derivation for the pairing term as an example. It reads:

$$\begin{aligned} \check{c}^\dagger [\tau_y \otimes \sigma_y] \check{c} &= \left( \begin{array}{cccc} c_\uparrow^\dagger & c_\downarrow^\dagger & c_\uparrow & c_\downarrow \end{array} \right) \left( \begin{array}{cc|cc} 0 & 0 & 0 & -1 \\ 0 & 0 & +1 & 0 \\ \hline 0 & +1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{array} \right) \left( \begin{array}{c} c_\uparrow \\ c_\downarrow \\ c_\uparrow \\ c_\downarrow \end{array} \right) \\ &= -c_\uparrow^\dagger c_\downarrow^\dagger + c_\downarrow^\dagger c_\uparrow^\dagger + c_\uparrow c_\downarrow - c_\downarrow c_\uparrow = 2 (c_\downarrow^\dagger c_\uparrow^\dagger + \text{h.c.}) \end{aligned} \quad (70)$$

where we the fermionic anti-commutation properties  $\{c_i, c_j^\dagger\} = \delta_{ij}$  and  $\{c_i, c_j\} = 0$ .

The remaining terms derivation is analogous but even simpler because there is will be no mixing of particle with particle-hole components; the holeonic terms will correspond to the negative of the electronic terms, meaning that one just needs to expand the space according to  $\tau_z \otimes$  the respective spin matrix. For the kinetic term there is no mixing of spin so it must trivially have the spin Pauli matrix  $\sigma_0$ . Similarly, for the Zeeman term there is only the same-spin mixing of the type  $\uparrow\uparrow - \downarrow\downarrow$  so it must have  $\sigma_z$ . As for the SOC term there is spin-mixing of opposing spins, so the options are either  $\sigma_x$  or  $i\sigma_y$  (with a  $i$  for it to be hermitian). One can check with the fermionic anti-commutation properties that it is indeed  $i\sigma_y$ .

*Alternative Nambu basis* It is common for people to define instead the Nambu spinor in a rotated basis as such

$$\bar{c}_i^\dagger = \left( \begin{array}{c} c_i^\dagger \\ [i\sigma_y c_i] \end{array} \right) = \left( \begin{array}{cc} c_{i\uparrow}^\dagger & c_{i\downarrow}^\dagger \\ \hline c_{i\downarrow} & -c_{i\uparrow} \end{array} \right) \quad (71)$$

As also explained in section II.C.1 of the previous part, these basis' operators relate to each other as

$$\bar{c}_i = \bar{\mathcal{U}} \check{c}_i \Leftrightarrow \check{c}_i = \bar{\mathcal{U}}^\dagger \bar{c}_i \quad (72)$$

$$\bar{c}_i^\dagger = \check{c}_i^\dagger \bar{\mathcal{U}}^\dagger \Leftrightarrow \check{c}_i^\dagger = \bar{c}_i^\dagger \bar{\mathcal{U}} \quad (73)$$

and, consequently, for a generic  $\check{M}$  matrix,

$$\bar{M} = \bar{\mathcal{U}} \check{M} \bar{\mathcal{U}}^\dagger \quad (74)$$

with  $\bar{\mathcal{U}}$  is a unitary matrix (i.e  $\bar{\mathcal{U}}^\dagger \bar{\mathcal{U}} = \bar{\mathcal{U}} \bar{\mathcal{U}}^\dagger = \mathbb{1}$ )

$$\bar{\mathcal{U}} = \begin{pmatrix} \sigma_0 & 0 \\ 0 & \imath\sigma_y \end{pmatrix} \quad (75)$$

Making use of Pauli matrices' property

$$\sigma_\alpha \sigma_\beta = \sigma = \sigma_0 \delta_{\alpha\beta} + i\varepsilon_{\alpha\beta\gamma} \sigma_\gamma \quad (76)$$

one can check that

$$H_K : \bar{\mathcal{U}} [\tau_z \otimes \sigma_0] \bar{\mathcal{U}}^\dagger = [\tau_z \otimes \sigma_0] \quad (77)$$

$$H_{SOC} : \bar{\mathcal{U}} [\tau_z \otimes \imath\sigma_y] \bar{\mathcal{U}}^\dagger = [\tau_z \otimes \imath\sigma_y] \quad (78)$$

$$H_Z : \bar{\mathcal{U}} [\tau_z \otimes \sigma_z] \bar{\mathcal{U}}^\dagger = [\tau_z \otimes \sigma_z] \quad (79)$$

$$H_{SC} : \bar{\mathcal{U}} [\tau_y \otimes \sigma_y] \bar{\mathcal{U}}^\dagger = [\tau_x \otimes \sigma_0] \quad (80)$$

meaning that, in this the rotated basis, only the pairing Hamiltonian has it's Pauli matrices changed. Concretely,

$$H_{SC} = \frac{1}{2}\Delta \sum_i \bar{\mathcal{C}}_i^\dagger [\tau_x \otimes \sigma_0] \bar{\mathcal{C}}_i \quad (81)$$

## VI. TOPOLOGICAL SUPERCONDUCTIVITY IN 2D MODELS

Need a intuitive and organized introduction relating all the nomenclatures "topological insulator", "Chern insulator" with the various effects. I still don't have a clear map of what's what and the subtle symmetry differences.

### A. Quantum hall effect

### B. Anomalous quantum Hall effect in the Haldane model

The quantum Hall effect without an external magnetic field is also referred to as the quantum anomalous Hall effect.

### C. Quantum spin Hall effect in the Kane-Mele model

#### D. Integer quantum Hall effect

#### E. Fractional quantum Hall effect

## Part III

# Topics on topological superconductor theory

## VII. DIRTY SUPERCONDUCTOR THEORY

### A. Green's function formulation

#### 1. Abrikosov-Gorkov Green's Functions

#### 2. Eilenberger-Larkin-Ovchinnikov equations

#### 3. Usadel diffusion equation

## VIII. SELF-CONSISTENT TOPOLOGICAL SUPERCONDUCTIVITY

### A. Self-consistent Hartree-Fock-Bogoliubov mean-field theory\*

Need to reorganize this chapter as to now have so many unnecessary derivations steps as I prefer to have them on the corresponding supplementary materials as to not bore the reader.

Take an interacting many-body system of electrons described by the Hamiltonian

$$\begin{aligned} H &= H_0 + H_{\text{int}} \\ &= \sum_{ij} c_i^\dagger H_0^{ij} c_j + \frac{1}{2} \sum_{ij} \sum_{i'j'} c_i^\dagger c_{i'} V_{j'i'}^{ij} c_j^\dagger c_{j'} \end{aligned} \quad (82)$$

with  $c^\dagger(c)$  fermionic creation (annihilation) operators and  $\mathbf{i} \equiv (i, s_i)$  composite degrees of freedom of purely spacial indices  $i$  and an additional orbital index at each site  $s_i$  such as spin. It follows directly from the composite notation that  $\sum_i = \sum_{s_i} \sum_i$  and  $\delta_{ij} = \delta_{ij} \delta_{s_i s_j}$ .

*Wannier spinless potential*

Let us now consider ultra-localized/Wannier orbitals such that the interaction tensor can approximately behave as  $V_{j'i'}^{ij} \approx v^{ij} \delta_{ii'} \delta_{jj'}$  mediated by a spinless electrostatic scalar potential  $v^{ij} \rightarrow v^{ij} = v(\vec{r}_i - \vec{r}_j)$  with  $v^{ij} = v^{ji}$ .

Moreover, a perturbation theory treatment of the interaction based on path integrals starts by casting the Hamiltonian into its normal-ordered form. Foremost, see that from the normal ordering of  $H_{\text{int}}$  a non-physical spurious self-interaction term  $1/2 c_i^\dagger c_i v^{ii}$  reveals itself explicitly, which should not take part since single electrons cannot interact with themselves. This term can be removed preemptively by incorporating it in Eq.(82) instead with a negative sign. Within the scope of this approximations the normal-ordered Hamiltonians takes the form

$$H = \sum_{ij} c_i^\dagger H_0^{ij} c_j + \frac{1}{2} \sum_{ij} c_i^\dagger c_j^\dagger v^{ij} c_j c_i \quad (83)$$

### Mean-field Hartree-Fock-Bogoliubov decoupling

We wish to derive from Eq.(83) the explicit form of the Hartree/electrostatic  $\Sigma_H$ , Fock/exchange  $\Sigma_F$  and Bogoliubov/pairing  $\Sigma_B$  self-energies between spatial sites  $i$  and  $j$ , being matrices over orbital space. For this, one must Hartree-Fock-Bogoliubov decouple the Hamiltonian in Eq.(83) such that it could be expressed as:

$$\begin{aligned} H &\approx H_0 + H_{\text{int}}^H + H_{\text{int}}^F + H_{\text{int}}^B \\ &= \sum_{ij} \left[ c_i^\dagger \left( H_0^{ij} + \Sigma_H^{ij} + \Sigma_F^{ij} \right) c_j + \frac{1}{2} \left( c_i^\dagger \Sigma_B^{ij} c_j^\dagger + h.c. \right) \right]. \end{aligned} \quad (84)$$

In particular, within a Nambu-spinor representation, which we will be presenting next, we should be able to make this Hamiltonian the form

$$\check{H} = \frac{1}{2} \sum_{ij} \check{c}_i \left( \check{H}_0^{ij} + \check{\Sigma}^{ij} \right) \check{c}_j, \quad (85)$$

having an inherent Bogoliubov-de-Gennes (BdG) symmetry.

#### Nambu representation

For this derivation we focus on the system's reduced density matrix (rDM) equation of motion (EoM). In normal systems, i.e non-superconducting systems, the rDM defined as  $\rho_{ee} = \langle c_j^\dagger c_i \rangle$  is sufficient, however, if one wishes to study superconducting systems, one must also account for the anomalous/pairing terms. For this, we introduce Nambu-spinors representation with doubling of degrees of freedom so that electron  $e$ , hole  $h$  become additional quantum numbers,

$$\check{c}_i^\dagger = \begin{pmatrix} c_i^\dagger & c_i \end{pmatrix} \text{ and } \check{c}_i = \begin{pmatrix} c_i \\ c_i^\dagger \end{pmatrix} \quad (86)$$

The corresponding Nambu rDM, can then be written as the direct/tensor product of this Nambu-spinors for each of the  $e, h$  combinations

$$\check{\rho}_{ij} = \langle \check{c}_j^\dagger \otimes \check{c}_i \rangle = \begin{pmatrix} \langle c_j^\dagger c_i \rangle & \langle c_j c_i \rangle \\ \langle c_j^\dagger c_i^\dagger \rangle & \langle c_j c_i^\dagger \rangle \end{pmatrix} \equiv \begin{pmatrix} \rho_{ee}^{ij} & \rho_{eh}^{ij} \\ \rho_{he}^{ij} & \rho_{hh}^{ij} \end{pmatrix} \quad (87)$$

Directly from the equal-time fermionic anti-commutator properties in Eqs.(??)-(??), this terms relate to themselves and to each other as

$$\rho_{ee}^{ij} = (\rho_{ee}^{ji})^\dagger \text{ and } \rho_{eh}^{ij} = -\rho_{eh}^{ji} \quad (88)$$

$$\rho_{hh}^{ij} = \delta_{ij} - \rho_{ee}^{ji} \text{ and } \rho_{he}^{ij} = (\rho_{eh}^{ji})^\dagger \quad (89)$$

Moreover, accounting for the additional spin orbital quantum number, the Nambu-spinor corresponds instead to the 4-spinor

$$\check{c}_i^\dagger = \begin{pmatrix} c_i^\dagger & c_i \end{pmatrix} = \begin{pmatrix} c_{i\uparrow}^\dagger & c_{i\downarrow}^\dagger & c_{i\uparrow} & c_{i\downarrow} \end{pmatrix} \quad (90)$$

such that each composite rDM is a matrix over orbital space, reading explicitly as

$$\check{\rho}_{ij} = \langle \check{c}_j^\dagger \otimes \check{c}_i \rangle = \begin{pmatrix} \langle c_j^\dagger \otimes c_i \rangle & \langle c_j \otimes c_i \rangle \\ \langle c_j^\dagger \otimes c_i^\dagger \rangle & \langle c_j \otimes c_i^\dagger \rangle \end{pmatrix} \equiv \begin{pmatrix} \rho_{ee}^{ij} & \rho_{eh}^{ij} \\ \rho_{he}^{ij} & \rho_{hh}^{ij} \end{pmatrix} \equiv \begin{pmatrix} \rho_{ee}^{i\uparrow j\uparrow} & \rho_{ee}^{i\uparrow j\downarrow} & \rho_{eh}^{i\uparrow j\uparrow} & \rho_{eh}^{i\uparrow j\downarrow} \\ \rho_{ee}^{i\downarrow j\uparrow} & \rho_{ee}^{i\downarrow j\downarrow} & \rho_{eh}^{i\downarrow j\uparrow} & \rho_{eh}^{i\downarrow j\downarrow} \\ \hline \rho_{he}^{i\uparrow j\uparrow} & \rho_{he}^{i\uparrow j\downarrow} & \rho_{hh}^{i\uparrow j\uparrow} & \rho_{hh}^{i\uparrow j\downarrow} \\ \rho_{he}^{i\downarrow j\uparrow} & \rho_{he}^{i\downarrow j\downarrow} & \rho_{hh}^{i\downarrow j\uparrow} & \rho_{hh}^{i\downarrow j\downarrow} \end{pmatrix} \quad (91)$$

and relating to themselves and to each other as

$$\boldsymbol{\rho}_{ee}^{ij} = (\boldsymbol{\rho}_{ee}^{ji})^\dagger \text{ and } \boldsymbol{\rho}_{eh}^{ij} = -\left(\boldsymbol{\rho}_{eh}^{ji}\right)^T \quad (92)$$

$$\boldsymbol{\rho}_{hh}^{ij} = \boldsymbol{\delta}_{ij} - (\boldsymbol{\rho}_{ee}^{ji})^T \text{ and } \boldsymbol{\rho}_{he}^{ij} = \left(\boldsymbol{\rho}_{eh}^{ji}\right)^\dagger \quad (93)$$

To clarify possible misinterpretations of the notation, see that the underlying electron-hole structure of an object is being concealed with the check notation while the underlying spin structure of an *object* is being concealed within the bold notation. The bold notation of a composite *index*  $\mathbf{i} \equiv (i, s_i)$  does not underlie the objects spin structure, so beware the differences between  $\rho_{ee}^{ij} = \rho_{ee}^{is_ijs_j}$  and  $\boldsymbol{\rho}_{ee}^{ij} = [\rho_{ee}^{i\uparrow j\uparrow} \ \rho_{ee}^{i\uparrow j\downarrow}; \ \rho_{ee}^{i\downarrow j\uparrow} \ \rho_{ee}^{i\downarrow j\downarrow}]$ . The notation  $\boldsymbol{\rho}_{ee}^{ij}$  is not applicable for now since we are considering only spin and no other orbital degree of freedom. See that, for example, in a spinless case we would write  $\check{\rho}^{ij} = [\rho_{ee}^{ij} \ \rho_{eh}^{ij}; \ \rho_{he}^{ij} \ \rho_{hh}^{ij}]$ .

### 1. Mean-field self-energy in Nambu space

We start by solving for the purely electronic rDM equation of motions in the Heisenberg picture of quantum mechanics, where the fermionic operators evolve accordingly to the Heisenberg equation. We have

$$\frac{d}{dt} \rho_{ee}^{ij} = \frac{i}{\hbar} \left\langle [H, c_j^\dagger] c_i \right\rangle + \frac{i}{\hbar} \left\langle c_j^\dagger [H, c_i] \right\rangle \quad (94)$$

with  $H$  the Hamiltonian in Eq.(83) and  $i$  the imaginary unit. The time dependency in the fermionic operators is being omitted for compactness.

Making use of the fermionic anti-commutator properties, these commutators read, respectively,

$$[H, c_j^\dagger] = \sum_{\alpha\beta} H_0^{\alpha\beta} [c_\alpha^\dagger c_\beta, c_j^\dagger] + \frac{1}{2} \sum_{\alpha\beta} v^{\alpha\beta} [c_\alpha^\dagger c_\beta^\dagger c_\beta c_\alpha, c_j^\dagger]$$

$$[H, c_i] = \sum_{\alpha\beta} H_0^{\alpha\beta} [c_\alpha^\dagger c_\beta, c_i] + \frac{1}{2} \sum_{\alpha\beta} v^{\alpha\beta} [c_\alpha^\dagger c_\beta^\dagger c_\beta c_\alpha, c_i]$$

The commutators reading

$$[c_\alpha^\dagger c_\beta, c_j^\dagger] = c_\alpha^\dagger \delta_{\beta j} \quad (95)$$

$$[c_\alpha^\dagger c_\beta, c_i] = -c_\beta \delta_{i\alpha} \quad (96)$$

$$[c_\alpha^\dagger c_\beta^\dagger c_\beta c_\alpha, c_j^\dagger] = c_\alpha^\dagger c_\beta^\dagger c_\beta \delta_{\alpha j} - c_\alpha^\dagger c_\beta^\dagger c_\alpha \delta_{\beta j} \quad (97)$$

$$[c_\alpha^\dagger c_\beta^\dagger c_\beta c_\alpha, c_i] = c_\alpha^\dagger c_\beta c_\alpha \delta_{\beta i} - c_\beta^\dagger c_\beta c_\alpha \delta_{i\alpha} \quad (98)$$

where we used that

$$[AB, Z] = A\{B, Z\} - \{Z, A\}B \quad (99)$$

$$\begin{aligned} [ABCD, Z] &= AB(C\{D, Z\} - \{Z, C\}D) \\ &\quad + (A\{B, Z\} - \{Z, A\}B)CD \end{aligned} \quad (100)$$

The complete Heisenberg commutators then yield

$$\begin{aligned} [H, c_j^\dagger] &= \sum_{\alpha} H_0^{\alpha j} c_\alpha^\dagger + \frac{1}{2} \sum_{\beta} v^{j\beta} c_j^\dagger c_\beta^\dagger c_\beta - \frac{1}{2} \sum_{\alpha} v^{\alpha j} c_\alpha^\dagger c_j^\dagger c_\alpha \\ &= \sum_{\alpha} H_0^{\alpha j} c_\alpha^\dagger + \frac{1}{2} \sum_{\alpha} v^{\alpha j} (c_j^\dagger c_\alpha^\dagger - c_\alpha^\dagger c_j^\dagger) c_\alpha \end{aligned} \quad (101)$$

$$\begin{aligned} [H, c_i] &= - \sum_{\beta} H_0^{i\beta} c_\beta + \frac{1}{2} \sum_{\alpha} v^{\alpha i} c_\alpha^\dagger c_i c_\alpha - \frac{1}{2} \sum_{\beta} v^{i\beta} c_\beta^\dagger c_\beta c_i \\ &= - \sum_{\alpha} H_0^{i\alpha} c_\alpha + \frac{1}{2} \sum_{\alpha} v^{\alpha i} c_\alpha^\dagger (c_i c_\alpha - c_\alpha c_i) \end{aligned} \quad (102)$$

Substituting Eqs.(101) and (102) back into Eq.(??) yields

$$\begin{aligned} -i\hbar \frac{d}{dt} \rho_{ee}^{ij} &= H_0^{\alpha j} \langle c_\alpha^\dagger c_i \rangle - H_0^{i\alpha} \langle c_j^\dagger c_\alpha \rangle \\ &\quad + \frac{1}{2} v^{\alpha j} (\langle c_j^\dagger c_\alpha^\dagger c_\alpha c_i \rangle - \langle c_\alpha^\dagger c_j^\dagger c_\alpha c_i \rangle) \\ &\quad + \frac{1}{2} v^{\alpha i} (\langle c_j^\dagger c_\alpha^\dagger c_i c_\alpha \rangle - \langle c_j^\dagger c_\alpha^\dagger c_\alpha c_i \rangle) \end{aligned}$$

Coming back to the rDM definitions and commutating the terms it reads

$$\begin{aligned} i\hbar \frac{d}{dt} \rho_{ee}^{ij} &= - H_0^{\alpha j} \rho_{ee}^{i\alpha} + H_0^{i\alpha} \rho_{ee}^{\alpha j} \\ &\quad - (v^{\alpha j} - v^{\alpha i}) \langle c_j^\dagger c_\alpha^\dagger c_\alpha c_i \rangle \end{aligned} \quad (103)$$

Notice, however, that the interaction term will give rise to expectation values of four-operators. For this, we introduce a mean-field approximation where we assume the two-particle expectation value to simply behave as a product of two one-particle expectation values. From this mean-field decoupling we can then make use of *Wick's theorem*, yielding

$$\langle c_j^\dagger c_\alpha^\dagger c_\alpha c_i \rangle \approx \rho_{he}^{\alpha j} \rho_{eh}^{i\alpha} - \rho_{ee}^{\alpha j} \rho_{ee}^{i\alpha} + \rho_{ee}^{ij} \rho_{ee}^{\alpha\alpha} \quad (104)$$

From this approximation, we defining the Hartree, Fock and Bogoliubov self-energies, themselves self-consistently dependent on the rDM, respectively as

$$\Sigma_H^{ij} = \delta_{ij} \sum_{\alpha} v^{i\alpha} \rho_{ee}^{\alpha\alpha} \quad (105)$$

$$\Sigma_F^{ij} = - v^{ij} \rho_{ee}^{ij} \quad (106)$$

$$\Sigma_B^{ij} = v^{ij} \rho_{eh}^{ij} \quad (107)$$

The interacting term of Eq.(??) yield

$$\begin{aligned}
& v^{\alpha j} \left[ \rho_{he}^{\alpha j} \rho_{eh}^{i\alpha} - \rho_{ee}^{\alpha j} \rho_{ee}^{i\alpha} + \rho_{ee}^{ij} \rho_{ee}^{\alpha\alpha} \right] \\
&= \left\{ v^{\alpha j} \rho_{he}^{\alpha j} \right\} \rho_{eh}^{i\alpha} - \left\{ v^{\alpha j} \rho_{ee}^{\alpha j} \right\} \rho_{ee}^{i\alpha} + \rho_{ee}^{ij} \left\{ v^{\alpha j} \rho_{ee}^{\alpha\alpha} \right\} \\
&= \rho_{eh}^{i\alpha} \left( -\Sigma_B^{\alpha j} \right)^{\dagger} + \rho_{ee}^{i\alpha} \Sigma_F^{\alpha j} + \rho_{ee}^{ij} \Sigma_H^{ii} \\
&= \left[ -\rho_{eh}^{ij} \left( -\Sigma_B^{ji} \right)^{\dagger} + \rho_{ee}^{ij} \Sigma_F^{ij} + \rho_{ee}^{ij} \Sigma_H^{ij} \delta_{ij} \right]^{s_i s_j} \tag{108}
\end{aligned}$$

$$\begin{aligned}
& -v^{\alpha i} s \left[ \rho_{he}^{\alpha j} \rho_{eh}^{i\alpha} - \rho_{ee}^{\alpha j} \rho_{ee}^{i\alpha} + \rho_{ee}^{ij} \rho_{ee}^{\alpha\alpha} \right] \\
&= - \left( \rho_{he}^{\alpha j} \left\{ v^{i\alpha} \rho_{eh}^{i\alpha} \right\} + \rho_{ee}^{\alpha j} \left\{ -v^{i\alpha} \rho_{ee}^{i\alpha} \right\} + \rho_{ee}^{ij} \left\{ v^{i\alpha} \rho_{ee}^{\alpha\alpha} \right\} \right) \\
&= - \left( \Sigma_B^{i\alpha} \rho_{he}^{\alpha j} + \Sigma_F^{i\alpha} \rho_{ee}^{\alpha j} + \Sigma_H^{ii} \rho_{ee}^{ij} \right) \\
&= - \left[ \Sigma_B^{ij} \rho_{he}^{ij} + \Sigma_F^{ij} \rho_{ee}^{ij} + \rho_{ee}^{ij} \Sigma_H^{ij} \delta_{ij} \right]^{s_i s_j} \tag{109}
\end{aligned}$$

where  $\lceil \cdot \rceil^{s_i s_j}$  is the element at position  $s_i s_j$ , e.g  $\lceil M \rceil^{\uparrow\uparrow} = M[1, 1]$ . Putting the three pieces together, and accounting for the complete spin structure, the purely electronic rDM EoM yields and shown in Eq.(112).

Note that, from the relations in Eqs.(88)-(89),

$$\Sigma_B^{ij} = \left( v^{ij} \rho_{eh}^{ij} \right) = v^{ij} \left( -\rho_{eh}^{ji} \right) = -\Sigma_B^{ji} \tag{110}$$

$$\left( \Sigma_B^{ji} \right)^{\dagger} = \left( v^{ji} \rho_{eh}^{ji} \right)^{\dagger} = \left( v^{ij} \rho_{he}^{ij} \right)^{\dagger} = -\left( \Sigma_B^{ij} \right)^{\dagger} \tag{111}$$

The purely electronic rDM EoM then yields

$$i\hbar \frac{d}{dt} \rho_{ee}^{ij} \approx \left[ \mathbf{H}_{HF}^{ij}, \rho_{ee}^{ij} \right] + \Sigma_B^{ij} \cdot \rho_{he}^{ij} - \rho_{eh}^{ij} \cdot \left( \Sigma_B^{ji} \right)^{\dagger} \tag{112}$$

where we have defined

$$\mathbf{H}_{HF}^{ij} = \mathbf{H}_0^{ij} + \Sigma_H^{ij} \delta_{ij} + \Sigma_F^{ij} \tag{113}$$

Analogously for the anomalous rDM EoM, we obtain

$$\frac{d}{dt} \rho_{eh}^{ij} = \frac{i}{\hbar} \langle [H, c_j] c_i \rangle + \frac{i}{\hbar} \langle c_j [H, c_i] \rangle \tag{114}$$

$$\langle [H, c_j] c_i \rangle = - \sum_{\alpha} H_0^{j\alpha} \rho_{eh}^{i\alpha} - \sum_{\alpha} v^{\alpha j} \langle c_{\alpha}^{\dagger} c_{\alpha} c_j c_i \rangle \tag{115}$$

$$\langle c_j [H, c_i] \rangle = - \sum_{\alpha} H_0^{i\alpha} \rho_{eh}^{\alpha j} - \sum_{\alpha} v^{\alpha i} \langle c_j c_{\alpha}^{\dagger} c_{\alpha} c_i \rangle \tag{116}$$

$$\langle c_{\alpha}^{\dagger} c_{\alpha} c_j c_i \rangle \approx \rho_{ee}^{\alpha\alpha} \rho_{eh}^{ij} - \rho_{ee}^{j\alpha} \rho_{eh}^{i\alpha} + \rho_{ee}^{i\alpha} \rho_{eh}^{j\alpha} \tag{117}$$

$$\langle c_j c_{\alpha}^{\dagger} c_{\alpha} c_i \rangle \approx \rho_{hh}^{\alpha j} \rho_{eh}^{i\alpha} - \rho_{eh}^{j\alpha} \rho_{ee}^{i\alpha} + \rho_{eh}^{ij} \rho_{ee}^{\alpha\alpha} \tag{118}$$

and finally

$$\begin{aligned}
i\hbar \frac{d}{dt} \rho_{eh}^{ij} &\approx \mathbf{H}_{HF}^{ij} \cdot \rho_{eh}^{ij} + \rho_{eh}^{ij} \cdot \left( \mathbf{H}_{HF}^{ij} \right)^T \\
&\quad + \Sigma_B^{ij} \cdot \rho_{hh}^{ij} - \rho_{ee}^{ij} \cdot \Sigma_B^{ij} \tag{119}
\end{aligned}$$

Finally, as introduced in Eqs.(59)-(??), we can represent the Nambu rDM EoM in terms of an effective Bogoliubov-de Gennes (BdG) Hamiltonian  $\tilde{H}$  as simply as

$$\imath\hbar\frac{d}{dt}\check{\rho}_{ij} = \left[ \left( \check{H}_0^{ij} + \check{\Sigma}^{ij} \right), \check{\rho}_{ij} \right] \quad (120)$$

$$\check{H}_0^{ij} = \begin{pmatrix} \mathbf{H}_0^{ij} & 0 \\ 0 & -(\mathbf{H}_0^{ji})^T \end{pmatrix} \quad (121)$$

$$\check{\Sigma}^{ij} = \begin{pmatrix} \Sigma_H^{ij} + \Sigma_F^{ij} & \Sigma_B^{ij} \\ (\Sigma_B^{ji})^\dagger & -(\Sigma_H^{ji} + \Sigma_F^{ji})^T \end{pmatrix} \quad (122)$$

It would be useful if we could express  $\check{\Sigma}$  more compactly in terms of  $\check{\rho}$ . For this, and accounting for their spin structure, we re-express the Hartree term as

$$\begin{aligned} \Sigma_H^{ij} &= \delta_{ij} \sum_{\alpha} v^{i\alpha} \rho_{ee}^{\alpha\alpha} \\ &= \delta_{ij} \sigma_0 \sum_{\alpha} v^{i\alpha} \sum_{s_{\alpha}} \rho_{ee}^{\alpha s_{\alpha} \alpha s_{\alpha}} \\ &= \delta_{ij} \sigma_0 \sum_{\alpha} v^{i\alpha} (\rho_{ee}^{\alpha\uparrow\alpha\uparrow} + \rho_{ee}^{\alpha\downarrow\alpha\downarrow}) \\ &= \delta_{ij} \sigma_0 \sum_{\alpha} v^{i\alpha} \text{Tr}(\rho_{ee}^{\alpha\alpha}) \end{aligned} \quad (123)$$

$$= \delta_{ij} \sigma_0 \sum_{\alpha} v^{i\alpha} \left[ \frac{1}{2} \text{Tr}([\tau_z \otimes \sigma_0] \check{\rho}^{\alpha\alpha}) + 1 \right] \quad (124)$$

$$\equiv \frac{1}{2} \delta_{ij} \sigma_0 \sum_{\alpha} v^{i\alpha} \text{Tr}([\tau_z \otimes \sigma_0] \check{\rho}^{\alpha\alpha}) \quad (125)$$

with  $\sigma$  and  $\tau$  as Pauli matrices but operating in their respectively spin and electron-hole subspaces the Hamiltonian and  $\check{\rho}$  (var rho instead of rho) a newly defined BdG symmetric object:

$$\begin{aligned} \check{\rho}^{ij} &= \begin{pmatrix} \rho_{ee}^{ij} & \rho_{eh}^{ij} \\ (\rho_{eh}^{ji})^\dagger & -(\rho_{ee}^{ji})^T \end{pmatrix} \\ &= \check{\rho}^{ij} - \left[ \frac{1}{2} (\tau_0 - \tau_z) \otimes \sigma_0 \right] \end{aligned} \quad (126)$$

Likewise, the Fock and Bogoliubov term, accounting for their spin structure, read as

$$\Sigma_F^{ij} = -v^{ij} \rho_{ee}^{ij} \quad (127)$$

$$\Sigma_B^{ij} = v^{ij} \rho_{eh}^{ij} \quad (128)$$

but can be re-expressed together in terms of the Nambu rDM as

$$\begin{aligned} \Sigma_F^{ij} + \Sigma_B^{ij} &= -v^{ij} [\tau_z \otimes \sigma_0] \check{\rho}_{ij} [\tau_z \otimes \sigma_0] \\ &= -v^{ij} [\tau_z \otimes \sigma_0] \check{\rho}_{ij} [\tau_z \otimes \sigma_0] \end{aligned} \quad (129)$$

We can then compactly rewrite the Nambu mean field Hamiltonian as

$$\begin{aligned} \check{\Sigma}^{ij} &= \frac{1}{2} \delta_{ij} [\tau_z \otimes \sigma_0] \sum_{\alpha} v^{i\alpha} \text{Tr}([\tau_z \otimes \sigma_0] \check{\rho}^{\alpha\alpha}) \\ &\quad - v^{ij} [\tau_z \otimes \sigma_0] \check{\rho}^{ij} [\tau_z \otimes \sigma_0] \end{aligned} \quad (130)$$

See that, apart from the Nambu/BdG symmetries of Eq.(122), there are additional constraint: the Hartree and Fock terms inherent the symmetry from Eq.(??), i.e  $\Sigma_{H/F}^{ji} = (\Sigma_{H/F}^{ij})^\dagger$  and the Bogoliubov term the symmetry from Eq.(??), i.e  $\Sigma_B^{ij} = -(\Sigma_B^{ji})^T$ . From these constrains one can then see that

$$\begin{aligned} & -[\tau_x \otimes \sigma_0] (\check{\Sigma}^{ji})^* [\tau_x \otimes \sigma_0] \\ &= \begin{pmatrix} -(\Sigma_{22}^{ji})^* & -(\Sigma_{21}^{ji})^* \\ -(\Sigma_{12}^{ji})^* & -(\Sigma_{11}^{ji})^* \end{pmatrix} = \check{\Sigma}^{ij} \end{aligned} \quad (131)$$

where  $\{\check{\Sigma}^{ij}\}^*$  terms reads explicitly as

$$\begin{aligned} & \begin{pmatrix} \left\{ [\Sigma_H^{ji}]^\dagger + [\Sigma_F^{ji}]^\dagger \right\}^* & \left\{ -[\Sigma_B^{ji}]^T \right\}^* \\ \left\{ (-[\Sigma_B^{ij}]^T)^\dagger \right\}^* & \left\{ -\left( [\Sigma_H^{ij}]^\dagger + [\Sigma_F^{ij}]^\dagger \right)^T \right\}^* \end{pmatrix} \\ &= \begin{pmatrix} (\Sigma_H^{ji} + \Sigma_F^{ji})^T & -(\Sigma_B^{ji})^\dagger \\ -\Sigma_B^{ij} & -(\Sigma_H^{ij} + \Sigma_F^{ij}) \end{pmatrix} \end{aligned} \quad (132)$$

All these symmetries relations can be summarized as

$$\check{\Sigma}^{ij} = (\check{\Sigma}^{ji})^\dagger \quad (133)$$

$$\check{\Sigma}^{ij} = -[\tau_x \otimes \sigma_0] (\check{\Sigma}^{ji})^* [\tau_x \otimes \sigma_0] \quad (134)$$

## 2. Orbital-dependent interactions

We now generalize the interaction model to include spin-spin interactions with orbital matrix elements  $Q_{s_i s'_i}$  with  $Q_{s_i s'_i} = Q_{s'_i s_i}$ . The charge-charge interaction is recovered with  $\check{Q} = [\tau_z \otimes \sigma_0]$ . In this context, see that the potential as it is written Eq.(??) is no longer true, but should read instead as

$$V_{j'i'}^{ij} \approx Q_{s_i s'_i} v^{ij} Q_{s_j s'_j} \delta_{ii'} \delta_{jj'} \quad (135)$$

As intended, the spurious interaction

$$\frac{1}{2} \sum_{is_i} \sum_{s'_j} \left( c_{is_i}^\dagger Q^2 |_{s_i s'_j} c_{is'_j} \right) v^{ii} \quad (136)$$

cancel out when writing the Hamiltonian it's normal-ordered form leaving us only with

$$H_{\text{int}} = \frac{1}{2} \sum_{is_i} \sum_{js_j} \sum_{s'_i s'_j} c_{is_i}^\dagger c_{js_j}^\dagger \left\{ Q_{s_i s'_i} v^{ij} Q_{s_j s'_j} \right\} c_{js'_j} c_{is'_i}$$

From Eq.(123), (127), (128), the Hartree-Fock-Bogoliubov self-energies considering orbital-dependent interactions read respectively as

$$\Sigma_H^{ij} = \delta_{ij} \mathbf{Q} \sum_{\alpha} v^{i\alpha} \text{Tr} (\rho_{ee}^{\alpha\alpha}) \quad (137)$$

$$\Sigma_F^{ij} = -v^{ij} \mathbf{Q} \rho_{ee}^{ij} \mathbf{Q}^T \quad (138)$$

$$\Sigma_B^{ij} = v^{ij} \mathbf{Q} \rho_{eh}^{ij} \mathbf{Q}^T \quad (139)$$

and thus, the Nambu self-energy becomes

$$\check{\Sigma}^{ij} = \frac{1}{2} \delta_{ij} \check{\mathbf{Q}} \sum_{\alpha} v^{i\alpha} \text{Tr} (\check{\mathbf{Q}} \check{\rho}^{\alpha\alpha}) - v^{ij} \check{\mathbf{Q}} \check{\rho}^{ij} \check{\mathbf{Q}} \quad (140)$$

### Alternate Nambu basis

One may define the Nambu spinor differently. For example, instead of  $\check{\mathbf{c}}_i^\dagger$  in Eq.(90), there is also common to encounter the so-called rotated basis where

$$\bar{\mathbf{c}}_i^\dagger = ( \mathbf{c}_i^\dagger \ [i\sigma_y \mathbf{c}_i] ) = ( c_{i\uparrow}^\dagger \ c_{i\downarrow}^\dagger \ c_{i\downarrow} \ -c_{i\uparrow} ) \quad (141)$$

These relate to the previous choice of basis as

$$\bar{\mathbf{c}}_i = \bar{\mathcal{U}} \check{\mathbf{c}}_i \Leftrightarrow \check{\mathbf{c}}_i = \bar{\mathcal{U}}^\dagger \bar{\mathbf{c}}_i \quad (142)$$

$$\bar{\mathbf{c}}_i^\dagger = \check{\mathbf{c}}_i^\dagger \bar{\mathcal{U}}^\dagger \Leftrightarrow \check{\mathbf{c}}_i^\dagger = \bar{\mathbf{c}}_i^\dagger \bar{\mathcal{U}} \quad (143)$$

with  $\bar{\mathcal{U}}$  is a unitary matrix (i.e  $\bar{\mathcal{U}}^\dagger \bar{\mathcal{U}} = \bar{\mathcal{U}} \bar{\mathcal{U}}^\dagger = \mathbb{1}$ ) reading

$$\bar{\mathcal{U}} = \begin{pmatrix} \sigma_0 & 0 \\ 0 & i\sigma_y \end{pmatrix} = \left( \begin{array}{cc|cc} +1 & 0 & 0 & 0 \\ 0 & +1 & 0 & 0 \\ \hline 0 & 0 & 0 & +1 \\ 0 & 0 & -1 & 0 \end{array} \right) \quad (144)$$

The rotated basis Nambu rDMs, explicitly reading

$$\begin{aligned} \bar{\rho}_{ij} &= \langle \bar{\mathbf{c}}_j^\dagger \otimes \bar{\mathbf{c}}_i \rangle \\ &= \begin{pmatrix} \langle \mathbf{c}_j^\dagger \otimes \mathbf{c}_i \rangle & \langle [i\sigma_y \mathbf{c}_j] \otimes \mathbf{c}_i \rangle \\ \langle \mathbf{c}_j^\dagger \otimes [i\sigma_y \mathbf{c}_i]^\dagger \rangle & \langle [i\sigma_y \mathbf{c}_j] \otimes [i\sigma_y \mathbf{c}_i]^\dagger \rangle \end{pmatrix} \\ &= \begin{pmatrix} \rho_{ee}^{i\uparrow j\uparrow} & \rho_{ee}^{i\uparrow j\downarrow} & \rho_{eh}^{i\uparrow j\downarrow} & -\rho_{eh}^{i\uparrow j\uparrow} \\ \rho_{ee}^{i\downarrow j\uparrow} & \rho_{ee}^{i\downarrow j\downarrow} & \rho_{eh}^{i\downarrow j\downarrow} & -\rho_{eh}^{i\downarrow j\uparrow} \\ \hline \rho_{he}^{i\downarrow j\uparrow} & \rho_{he}^{i\downarrow j\downarrow} & \rho_{hh}^{i\downarrow j\downarrow} & -\rho_{hh}^{i\downarrow j\uparrow} \\ -\rho_{he}^{i\uparrow j\uparrow} & -\rho_{he}^{i\uparrow j\downarrow} & -\rho_{hh}^{i\uparrow j\downarrow} & \rho_{hh}^{i\uparrow j\uparrow} \end{pmatrix} \end{aligned}$$

can then be expressed also in terms of  $\bar{\mathcal{U}}$  as

$$\bar{\rho}_{ij} = \langle \bar{\mathbf{c}}_j^\dagger \otimes \bar{\mathbf{c}}_i \rangle = \langle \check{\mathbf{c}}_i^\dagger \bar{\mathcal{U}}^\dagger \otimes \bar{\mathcal{U}} \check{\mathbf{c}}_i \rangle = \bar{\mathcal{U}} \check{\rho}_{ij} \bar{\mathcal{U}}^\dagger \quad (145)$$

Likewise, the matrix for the Nambu  $\mathbf{Q}$  becomes  $\bar{\mathbf{Q}} = \bar{\mathcal{U}} \check{\mathbf{Q}} \bar{\mathcal{U}}^\dagger$ . Since the last term of  $\check{\rho}$  in Eq.(126) transforms into itself, then it also follows that  $\bar{\rho} = \bar{\mathcal{U}} \check{\rho} \bar{\mathcal{U}}^\dagger$  and thus  $\bar{\Sigma}^{ij} = \bar{\mathcal{U}} \check{\Sigma}^{ij} \bar{\mathcal{U}}^\dagger$  as in Eq.(130):

$$\bar{\Sigma}^{ij} = \frac{1}{2} \delta_{ij} \bar{\mathbf{Q}} \sum_{\alpha} v^{i\alpha} \text{Tr} (\bar{\mathbf{Q}} \bar{\rho}^{\alpha\alpha}) - v^{ij} \bar{\mathbf{Q}} \bar{\rho}^{ij} \bar{\mathbf{Q}} \quad (146)$$

where we used that  $\bar{\mathcal{U}}$  is unitary and the cyclic property of the trace

$$\text{Tr}(\bar{\mathcal{U}}^\dagger \bar{Q} \bar{\varrho}^{\alpha\alpha} \bar{\mathcal{U}}) = \text{Tr}(\bar{\mathcal{U}} \bar{\mathcal{U}}^\dagger \bar{Q} \bar{\varrho}^{\alpha\alpha}) = \text{Tr}(\bar{Q} \bar{\varrho}^{\alpha\alpha}) \quad (147)$$

As a result of this transformation the Nambu symmetries expressed in Eq.(122) become

$$\bar{\Sigma}^{ij} = \begin{pmatrix} \Sigma_H^{ij} + \Sigma_F^{ij} & -i\Sigma_B^{ij}\sigma_y \\ (-i\Sigma_B^{ji}\sigma_y)^\dagger & -\sigma_y (\Sigma_H^{ji} + \Sigma_F^{ji})^T \sigma_y \end{pmatrix} \quad (148)$$

Using the fact that  $\bar{\mathcal{U}}$  is unitary, Eq.(134) then becomes

$$\begin{aligned} \bar{\Sigma}^{ij} &= \bar{\mathcal{U}} \check{\Sigma}^{ij} \bar{\mathcal{U}}^\dagger \\ &= - [\bar{\mathcal{U}} [\tau_x \otimes \sigma_0] \bar{\mathcal{U}}^\dagger] [\bar{\mathcal{U}} (\check{\Sigma}^{ij})^* \bar{\mathcal{U}}^\dagger] [\bar{\mathcal{U}} [\tau_x \otimes \sigma_0] \bar{\mathcal{U}}^\dagger] \\ &= - [\tau_y \otimes \sigma_y] (\bar{\Sigma}^{ij})^* [\tau_y \otimes \sigma_y] \end{aligned}$$

where

$$\begin{aligned} \bar{\mathcal{U}} [\tau_x \otimes \sigma_0] \bar{\mathcal{U}}^\dagger &= \begin{pmatrix} \sigma_0 & 0 \\ 0 & i\sigma_y \end{pmatrix} \begin{pmatrix} 0 & \sigma_0 \\ \sigma_0 & 0 \end{pmatrix} \begin{pmatrix} \sigma_0 & 0 \\ 0 & -i\sigma_y \end{pmatrix} \\ &= \begin{pmatrix} 0 & -i\sigma_y \\ i\sigma_y & 0 \end{pmatrix} = (-i\tau_y) \otimes (i\sigma_y) \end{aligned} \quad (149)$$

The Nambu symmetries relations for the rotated basis read as

$$\bar{\Sigma}^{ij} = (\bar{\Sigma}^{ji})^\dagger \quad (150)$$

$$\bar{\Sigma}^{ij} = - [\tau_y \otimes \sigma_y] (\bar{\Sigma}^{ij})^* [\tau_y \otimes \sigma_y] \quad (151)$$

### 3. Hubbard model

Consider, instead of the spinless potential from Eq.(??), a spinful Hubbard potential

$$v^{ij} = U\delta_{ij} \quad (152)$$

In this model, the original non-normal-ordered interaction Hamiltonian from Eq.(??) is instead

$$\begin{aligned} H_{\text{int}} &= \frac{1}{2} \sum_{ij} c_i^\dagger c_i v^{ij} c_j^\dagger c_j - \frac{1}{2} \sum_i c_i^\dagger c_i v^{ii} \\ &= \frac{1}{2} \sum_{is_i} \sum_{js_j} c_{is_i}^\dagger c_{is_i} \{U\delta_{ij}\} c_{js_j}^\dagger c_{js_j} - \frac{1}{2} \sum_{is_i} c_{is_i}^\dagger c_{is_i} \{U\delta_{ii}\} \\ &= \frac{1}{2} U \sum_{is_i} \left[ (c_{is_i}^\dagger c_{is_i} n_{i\uparrow} + c_{is_i}^\dagger c_{is_i} n_{i\downarrow}) - (n_{i\uparrow} + n_{i\downarrow}) \delta_{ii} \right] \\ &= \frac{1}{2} U \sum_i (n_{i\uparrow} n_{i\downarrow} + n_{i\downarrow} n_{i\uparrow}) \end{aligned} \quad (153)$$

where  $n_{is_i} = c_{is_i}^\dagger c_{is_i}$  is the number operator and

$$(n_{i\uparrow} n_{i\uparrow} - n_{i\uparrow} \delta_{ii}) + (n_{i\downarrow} n_{i\downarrow} - n_{i\downarrow} \delta_{ii}) = 0 \quad (154)$$

because  $n_{is_i} n_{is_i} = (n_{is_i})^2 = n_{is_i}$  (there is or there is not an electron at site  $i$  and spin  $s_i$ , i.e  $n_{is_i} = 0$  or 1). Using that

$$\begin{aligned} n_{is_i} n_{i\bar{s}_i} &= c_{is_i}^\dagger \left\{ c_{is_i} c_{i\bar{s}_i}^\dagger \right\} c_{i\bar{s}_i} = c_{is_i}^\dagger \left( \delta_{ii} \delta_{s_i \bar{s}_i} - c_{i\bar{s}_i}^\dagger c_{is_i} \right) c_{i\bar{s}_i} \\ &= - \left\{ c_{is_i}^\dagger c_{i\bar{s}_i}^\dagger \right\} \left\{ c_{is_i} c_{i\bar{s}_i} \right\} = -c_{i\bar{s}_i}^\dagger \left\{ c_{is_i}^\dagger c_{i\bar{s}_i} \right\} c_{is_i} \\ &= n_{i\bar{s}_i} n_{is_i}, \end{aligned} \quad (155)$$

one can simply write

$$H_{\text{int}} \rightarrow H_U = U \sum_i n_{i\uparrow} n_{i\downarrow} \quad (156)$$

Also, see that by definition of the model, spurious interaction do not take part so there is no need to subtracting them preemptively. The Nambu self-energy term then reads

$$\begin{aligned} \check{\Sigma}_U^{ij} &= U \delta_{ij} \left( \frac{1}{2} [\tau_z \otimes \sigma_0] \text{Tr} ([\tau_z \otimes \sigma_0] \check{\varrho}^{ii}) \right. \\ &\quad \left. - [\tau_z \otimes \sigma_0] \check{\varrho}^{ij} [\tau_z \otimes \sigma_0] \right) \end{aligned} \quad (157)$$

#### 4. The gap equation accounting for Zeeman effect

#### B. Self-consistent Majorana nanowire

#### C. Self-consistent Bernal graphene

## Part IV Appendices

#### D. Peierls substitution

The Peierls substitution method is a widely common approximation for describing tight-binding models in the presence of a slowly varying magnetic vector potential  $\mathbf{A}(\mathbf{r})$ . In a tight-binding scheme it straightforwardly introduces a phase  $\theta$  to hopping strength, dubbed the Peierls phase, such that the tight-binding hopping Hamiltonian reads instead

$$H_{\text{hop}} = t_{ij} e^{i\theta} c_j^\dagger c_i + h.c. \quad (158)$$

which directly follows from the gauge invariance of the time-dependent Schrodinger equation.

**Proof:** Consider the time-dependent Schrodinger equation describing such a system at the continuum limit, dubbed the Hofstadter Hamiltonian, reads

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}, t) = \left[ \frac{i\hbar \nabla_{\mathbf{r}} - \frac{e}{c} \mathbf{A}(\mathbf{r})}{2m} + eU(\mathbf{r}, t) \right] \psi(\mathbf{r}, t). \quad (159)$$

with  $U(\mathbf{r})$  a generic scalar potential, for example the crystal lattice potential landscape. Furthermore, consider that one adds a local phase shift to the wavefunction as

$$\psi(\mathbf{r}, t) \rightarrow e^{\frac{ie}{\hbar c} \Lambda(\mathbf{r}, t)} \psi(\mathbf{r}, t) \quad (160)$$

Substituting this ansatz directly into the time-dependent Schrödinger equation one obtains

$$e^{\frac{ie}{\hbar c} \Lambda} \left( i\hbar \frac{\partial}{\partial t} - \frac{e}{c} \frac{\partial \Lambda}{\partial t} \right) \psi = e^{\frac{ie}{\hbar c}} \frac{1}{2m} \left( -i\hbar \nabla - \frac{e}{c} \mathbf{A} + \frac{e}{c} \nabla \Lambda + 2meU \right)^2 \psi$$

where we have omitted the spacial and temporal dependency for simplicity. See that if one now defines the potentials as

$$\begin{aligned} \mathbf{A} &\rightarrow \mathbf{A} + \nabla \Lambda \\ U &\rightarrow U + \frac{1}{c} \frac{\partial \Lambda}{\partial t} \end{aligned}$$

one recovers the original equation. This means that applying the gauge transformation (meaning that there exists other physical descriptions of the system that leaves the free energy unchanged) to  $\mathbf{A}$  and  $U$  is equivalent to multiplying the state by a phase factor, albeit one that changes in space and time.

## E. Overview of simpler systems

### 1. Linear lattice

### 2. Square lattice

## F. Overview of graphene systems

### 1. Monolayer graphene

Hexagonal boron nitride (hBN) is a 2D material composed of a simple layer of alternating boron and nitrogen atoms disposed in a planar honeycomb lattice, as shown in Fig.(14)(a). The Bravais lattice

$$\mathbf{r}_i = n_{i1}\mathbf{a}_1 + n_{i2}\mathbf{a}_2, \quad n_{i1}, n_{i2} \in \mathbb{Z} \quad (161)$$

is generated by the real vectors basis

$$\mathbf{a}_1 = a_0 \begin{bmatrix} +\sin(30^\circ) \\ +\cos(30^\circ) \end{bmatrix} \text{ and } \mathbf{a}_2 = a_0 \begin{bmatrix} +\sin(30^\circ) \\ -\cos(30^\circ) \end{bmatrix}. \quad (162)$$

where  $\sin(30^\circ) = 1/2$  and  $\cos(30^\circ) = \sqrt{3}/2$ . In each diamond shaped Wigner-Seitz primitive cell (depicted in yellow), we have one boron atom and one nitride atom, which we designate as sub-lattices  $A$  (depicted in red) and  $B$  (depicted in blue) respectively. The atoms within the central primitive cell are located at

$$\mathbf{s}_A = \frac{a_0}{\sqrt{3}} \begin{bmatrix} 0 \\ -1/2 \end{bmatrix} \text{ and } \mathbf{s}_B = \frac{a_0}{\sqrt{3}} \begin{bmatrix} 0 \\ +1/2 \end{bmatrix}. \quad (163)$$

where the origin is defined at the midpoint between the atoms. For each site  $A$ , the position of the nearest-neighbors (NN) in the sites  $B$  are given by

$$\boldsymbol{\delta}_1 = \frac{a_0}{\sqrt{3}} \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad \boldsymbol{\delta}_2 = \frac{a_0}{\sqrt{3}} \begin{bmatrix} +\sin(60^\circ) \\ -\cos(60^\circ) \end{bmatrix} \text{ and } \boldsymbol{\delta}_3 = \frac{a_0}{\sqrt{3}} \begin{bmatrix} -\sin(60^\circ) \\ -\cos(60^\circ) \end{bmatrix}. \quad (164)$$

where  $\sin(60^\circ) = \sqrt{3}/2$  and  $\cos(60^\circ) = 1/2$ . All these vectors are shown in Fig.(14)(a) within the real space lattice. Furthermore, from the real lattice basis vectors, in order to fulfill  $\mathbf{a}_i \cdot \mathbf{b}_j = 2\pi\delta_{ij}$ , the reciprocal lattice basis vectors follow as

$$\mathbf{b}_1 = \frac{2\pi}{a_0} \begin{bmatrix} +\cos(30^\circ) \\ -\sin(30^\circ) \end{bmatrix} \text{ and } \mathbf{b}_2 = \frac{2\pi}{a_0} \begin{bmatrix} +\cos(30^\circ) \\ +\sin(30^\circ) \end{bmatrix}. \quad (165)$$

These are also shown in Fig.(14)(b) together with the first zone of Brillouin, formed by the area enclosed by the intersection of their bisectrices. The high-simmetry points are  $\Gamma$ , the origin, the Dirac points  $K_\pm$  and  $M$  read as

$$\mathbf{\Gamma} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad K_\pm = \pm \frac{4\pi}{3a_0} \begin{bmatrix} 1 \\ 0 \end{bmatrix} \text{ and } M = \frac{2\pi}{a_0} \begin{bmatrix} +\cos(30^\circ)/2 \\ +\sin(30^\circ)/2 \end{bmatrix} \quad (166)$$

where the  $K$  point is found such that  $(\mathbf{M} + K_{k_x} \hat{\mathbf{M}}_\perp)_{k_y} = 0$ , with  $\hat{\mathbf{M}}_\perp$  the unit vector in the perpendicular direction to  $\mathbf{M}$ . In far right side of Fig.(13), we make a note that the discretized grid it's in the Bloch momentums basis  $\{\phi_1, \phi_2\}$ , i.e in the direction of the reciprocal lattice vectors, and not simply in the reciprocal space  $\{k_x, k_y\}$ . In the Bloch momentums basis the Dirac points would reads as  $K_\pm = 2\pi/3a_0 [\pm 1, \mp 1]$ .




Figure 13.

Let us consider the nearest-neighbors (NN) tight-binding model, written in real space as

$$H_{\text{TB}}(\mathbf{R}) = \sum_i \epsilon_A a_{\mathbf{r}_i}^\dagger a_{\mathbf{r}_i} + \sum_i \epsilon_B b_{\mathbf{r}_i}^\dagger b_{\mathbf{r}_i} - t \sum_{\langle i,j \rangle} (a_{\mathbf{r}_i}^\dagger b_{\mathbf{r}_i+\delta_j} + b_{\mathbf{r}_j}^\dagger a_{\mathbf{r}_i-\delta_j}), \quad (167)$$

where the operators  $a_{\mathbf{r}_i}^\dagger (a_{\mathbf{r}_i})$  create (annihilate) an electron in the sub-lattice  $A$  in a given Bravais lattice site  $\mathbf{r}_i$ , the operators  $b_{\mathbf{r}_i}^\dagger (b_{\mathbf{r}_i})$  the same but for sub-lattice  $B$ ,  $\epsilon_A$  and  $\epsilon_B$  are the onsite energies of site  $A$  and  $B$  respectively, and  $t$  is the hopping strength between nearest-neighbouring sites  $A$  and  $B$  and back, denoted with  $\langle i,j \rangle$ .

Expressing the creation/annihilation operators as their Fourier counterparts,

$$a_{\mathbf{R}_i} = \frac{1}{\sqrt{V}} \sum_{\mathbf{k}} e^{i\mathbf{k}\cdot(\mathbf{R}_i+\mathbf{s}_A)} a_{\mathbf{k}} \quad \text{and} \quad b_{\mathbf{R}_i} = \frac{1}{\sqrt{V}} \sum_{\mathbf{k}} e^{i\mathbf{k}\cdot(\mathbf{R}_i+\mathbf{s}_B)} b_{\mathbf{k}}, \quad (168)$$

and using the identity  $\delta(\mathbf{k} - \mathbf{k}') = 1/N \sum_i e^{-i\mathbf{R}_i \cdot (\mathbf{k} - \mathbf{k}')}$ , we obtain the Hamiltonian in reciprocal space,

$$H_{\text{TB}}(\mathbf{R}) = \sum_{\mathbf{k}} \epsilon_A a_{\mathbf{k}}^\dagger a_{\mathbf{k}} + \sum_{\mathbf{k}} \epsilon_B b_{\mathbf{k}}^\dagger b_{\mathbf{k}} - t \sum_{\mathbf{k}} \left( \gamma_{\mathbf{k}} a_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \gamma_{\mathbf{k}}^\dagger b_{\mathbf{k}}^\dagger a_{\mathbf{k}} \right), \quad (169)$$

where  $\gamma_{\mathbf{k}} = \sum_{\langle j \rangle} \exp(+i\mathbf{k} \cdot \boldsymbol{\delta}_j)$  is complex number. If we now define a row vector  $c_{\mathbf{k}}^\dagger = [a_{\mathbf{k}}^\dagger \ b_{\mathbf{k}}^\dagger]$  we can rewrite the system's Hamiltonian as  $H_{\mathbf{R}}^{\text{TB}} = \sum_{\mathbf{k}} c_{\mathbf{k}}^\dagger H_{\mathbf{k}}^{\text{TB}} c_{\mathbf{k}}$  with

$$H_{\text{TB}}(\mathbf{k}) = \begin{bmatrix} \epsilon_A & -t\gamma_{\mathbf{k}} \\ -t\gamma_{\mathbf{k}}^\dagger & \epsilon_B \end{bmatrix}. \quad (170)$$

Within this simplified tight-binding model, the expression for the electronic two-band structure can easily be obtained analytically by diagonalizing the matrix in Eq.(170), yielding

$$E_{\text{TB}}^{\pm}(\mathbf{k}) = \pm \sqrt{\epsilon^2 + t^2 \left[ 3 + 2 \cos(a_0 k_x) + 4 \cos\left(\frac{a_0 \sqrt{3}}{2} k_y\right) \cos\left(\frac{a_0}{2} k_x\right) \right]}, \quad (171)$$

having defined the zero point energy at  $(\epsilon_A + \epsilon_B)/2$  and defined  $\epsilon \equiv (\epsilon_A - \epsilon_B)/2$  at the middle of the gap such that  $\epsilon_A = \epsilon$  and  $\epsilon_B = -\epsilon$ . The valence band corresponds to the  $E_{\text{TB}}^-(\mathbf{k})$  dispersion while the  $E_{\text{TB}}^+(\mathbf{k})$  corresponds to the conduction band, as shown in Fig.(14)(c). The band structure is accompanied by the density of states  $\text{DoS}(E) = \sum_{\mathbf{k}} \delta(E - E(\mathbf{k}))$ .

Notice that, if  $\epsilon_A = \epsilon_B$ , as is the case for graphene, we obtain  $\epsilon = 0$  and the band dispersion closes in a linear fashion at the so called Dirac points. In hBN, the electronic band dispersion is also at its minimum near these points but has instead a parabolic shape. In either case, this points represent a fundamental symmetry of the system, called valley parity. To see why the dispersion is parabolic at these valley points, we Taylor series expand the exponential of  $\gamma_{\mathbf{k}}$  in Eq.(??) near  $\mathbf{k} \rightarrow \mathbf{K} + \mathbf{p}$  with  $\mathbf{p} \rightarrow 0$ . We obtain  $\exp(+i\mathbf{p} \cdot \boldsymbol{\delta}_j) \approx 1 + i\mathbf{p} \cdot \boldsymbol{\delta}_j$ . Now, since  $\sum_{\langle j \rangle} \exp(+i\mathbf{K} \cdot \boldsymbol{\delta}_j) = 0$  we are left with  $\gamma_{\mathbf{K} + \mathbf{p}} \simeq i\mathbf{p} \cdot \sum_{\langle j \rangle} \exp(+i\mathbf{K} \cdot \boldsymbol{\delta}_j) \boldsymbol{\delta}_j = -\sqrt{3}a_0/2(p_x - ip_y)$ . Invoking the Pauli matrices definitions, from Eq.(170) we can write the TB Hamiltonian  $H_{\text{TB}}^{\mathbf{k}}$  in this low-energy regime as

$$H_{\text{TB}}(\mathbf{K} + \mathbf{p}) = \epsilon \sigma_z + t \frac{\sqrt{3}a_0}{2} (\mathbf{p} \cdot \boldsymbol{\sigma}), \quad (172)$$

which clearly resembles the 2D Dirac Hamiltonian,  $H_{\text{Dirac}} = \sigma_z mc^2 + c(\mathbf{p} \cdot \boldsymbol{\sigma})$  with  $\epsilon$  taking the role of the rest mass energy  $mc^2$  and instead with a velocity  $v_F = t\sqrt{3}a_0/2$ , termed the *Fermi velocity*, as a replacement to the velocity of light  $c$ . Notice that, for the case of graphene, since  $\epsilon = 0$ , the electrons would behave as if they are massless. In this limit, the hBN low-energy dispersion can be written as the typical relativistic dispersion relation

$$E_{\text{TB}}(\mathbf{K} + \mathbf{p}) = \pm \sqrt{p^2 v_F^2 + m_{\text{eff}}^2 v_F^4}. \quad (173)$$

where  $m_{\text{eff}}$  is the effective mass of the electron at a given point near the valleys.

*Refazer esta figura em Quantica para aprender a fazer densidade de estados. Falar das singularidades de van Hove.*




Figure 14. hBN electronic band structure from a nearest-neighbor tight-binding model accompanied by the density of the states. The dispersion goes along the symmetry path  $\mathbf{k} : \Gamma \rightarrow K \rightarrow M \rightarrow \Gamma$  and was calculated using  $\epsilon_g = 7.8\text{eV}$  for the energy gap,  $t = 3.1\text{eV}$  for the hopping parameter and  $a_0 = 1.42\sqrt{3}\text{\AA}$  for the honeycomb lattice length.

## 2. Bilayer Bernal graphene

Consider a bilayer graphene model depicted in Fig.(15).




Figure 15. (a) Top view of the bilayer graphene (b) Side view of the dotted region in (a)

The tight-binding Hamiltonian of such a model reads

$$\begin{aligned}
H_{\text{BLG}} &= H_{\text{intralayer}} + H_{\text{interlayer}} = (H_{\text{top}} + H_{\text{bot}}) + (H_{\gamma_1} + H_{\gamma_3} + H_{\gamma_4}) \\
H_{\text{top}} &= \sum_i (\epsilon_A - \mu) c_i^\dagger a_i + \sum_i (\epsilon_B - \mu) b_i^\dagger b_i - \gamma_0 \sum_{\langle i,j \rangle} (a_i^\dagger b_j + h.c.) \\
H_{\text{bot}} &= \sum_i (\epsilon_C - \mu) c_i^\dagger c_i + \sum_i (\epsilon_D - \mu) d_i^\dagger d_i - \gamma_0 \sum_{\langle i,j \rangle} (c_i^\dagger d_j + h.c.) \\
H_{\gamma_1} &= +\gamma_1 \sum_{\langle i,j \rangle} (b_i^\dagger c_j + h.c.) \\
H_{\gamma_3} &= -\gamma_3 \sum_{\langle i,j \rangle} (a_i^\dagger d_j + h.c.) \\
H_{\gamma_4} &= +\gamma_4 \sum_{\langle i,j \rangle} (b_i^\dagger d_j + h.c.) + t_4 \sum_{\langle i,j \rangle} (a_i^\dagger C_j + h.c.)
\end{aligned}$$

Here, a site located at  $\mathbf{r}_i$  is indexed by the side index  $i$  and its next nearest neighbors located at  $\mathbf{r}_j$  are indexed with the site index  $j$ . Of course,  $\mathbf{r}_j$  depends on the kind of hopping in questions: for  $\gamma_0$  it's  $\mathbf{r}_j = \mathbf{r}_i + \boldsymbol{\delta}_j$  with  $j = 1, 2, 3$ , for  $\gamma_1$  it's  $\mathbf{r}_j = \mathbf{r}_i \pm a_z \hat{\mathbf{z}}$ , and for  $\gamma_3$  and  $\gamma_4$  it's  $\mathbf{r}_j = \mathbf{r}_i + \boldsymbol{\delta}_j \pm a_z \hat{\mathbf{z}}$ . Moreover, let us consider an electric field  $\mathbf{E}$  uniform in the plane  $xOy$  and growing along the  $\hat{\mathbf{z}}$ , described by the tight-binding Hamiltonian

$$H_E = \sum_i E_i (f_{i\uparrow}^\dagger f_{i\uparrow} - f_{i\downarrow}^\dagger f_{i\downarrow})$$

where  $E_i = E \times z_i$  is the amplitude of the electric field at position  $\mathbf{r}_i$ , only really dependent on  $z_i$ , and  $f_i^\dagger = [f_{i\uparrow}^\dagger \ f_{i\downarrow}^\dagger]$  is a generic fermionic operator. Since in our bilayer model the bottom layer is situated at  $z = 0$  we redefine  $E(a_z) = E$ , such that

$$H_{\text{BLG}+} = E \sum_i \left\{ (a_{i\uparrow}^\dagger a_{i\uparrow} - a_{i\downarrow}^\dagger a_{i\downarrow}) + (b_{i\uparrow}^\dagger b_{i\uparrow} - b_{i\downarrow}^\dagger b_{i\downarrow}) \right\}$$




Figure 16. (a, b) Bandstructure along symmetry path  $\Gamma \rightarrow K_+ \rightarrow M$  and (c) trigonal warping of BLG around the Dirac point  $K_+$ .

#### *Armchair and Zigzag configurations*

##### 3. Twisted bilayer graphene

##### 4. Kekulé modulation




Figure 17. Caption

## Part V

# Bibliography

As you been noticing, there is no references whatsoever within the text. Because these are personal journal notes, I preferred to avoid the hassle of managing minor references altogether, since, to be honest, one just skims through those (at best). I will, however, cite the actually relevant chapters/sections of papers/notes/courses/book that I closely followed for the making of this notes.

For Part I "Introduction to superconductivity theory" I mostly followed **Tinkham's "Introduction to superconductivity"** at [Dover publisher link] and **Rashid's "Lecture notes: Superconductivity"** at [personal website link] for both chapter structure and content. For Part II "Introduction to topological superconductivity" I closely followed **Akhmerov's "Online course on topology in condensed matter"** at <https://topocondmat.org/> and **Asboth's "A short course on topological insulators"** at arXiv:1509.02295.. for both chapter structure and content. Moreover for "Concepts of symmetry and topology" and the "SSH chain model" I additionally refer to **Tiago Antão's Master thesis on "Disorder and Topology in Spin Systems"** at [uni repositiorium link] For the "Kitaev model" and "Oreg-Lutchyn model" I refer, respectively, to the original papers of **Kitaev's "Unpaired Majorana fermions in quantum wires"** at Phys.-Usp. 44 131, and **Lutchyn's "Majorana Fermions and a Topological Phase Transition in Semiconductor-Superconductor Heterostructures"** at Phys. Rev. Lett. 105, 077001 and **Oreg's "Helical liquids and Majorana bound states in quantum wires"** at Phys. Rev. Lett. 105, 177002.

- BCS theory from Fernandes's Lecture Notes: BCS theory of superconductivity