Hybrid versus intrinsic Majorana nanowires, a self-consistent Hartree-Fock-Bogoliubov analysis

Francisco Lobo[1], Elsa Prada[1], Pablo San-Jose[1]
[1] *Instituto de Ciencia de Materiales de Madrid (ICMM), CSIC, Madrid, Spain*
(Dated: July 2, 2024)

Not yet worthy nor ready for the abstract.

I. INTRODUCTION

Not yet worthy nor ready for the introduction.

II. SELF-CONSISTENT HARTREE-FOCK-BOGOLIUBOV THEORY

Consider a generic tight-binding quartic Hamiltonian,

$$\hat{H} = \sum_{\alpha\beta} h_{\alpha\beta} c^\dagger_\alpha c_\beta + \frac{1}{2}\sum_{\alpha\beta\gamma\delta} V^{\alpha\beta}_{\gamma\delta} c^\dagger_\alpha c^\dagger_\beta c_\gamma c_\delta, \tag{1}$$

with $c^\dagger_\alpha(c_\alpha)$ fermionic creation (annihilation) operators of site and spin degree of freedom $\alpha = (i, \sigma_i)$, and $V^{\alpha\beta}_{\gamma\delta}$ a spinless electrostatic potential tensor of symmetries $V^{\alpha\beta}_{\gamma\delta} = V^{\beta\alpha}_{\delta\gamma}$ and $(V^{\alpha\beta}_{\gamma\delta})^* = V^{\gamma\delta}_{\alpha\beta}$ following directly from $V(\mathbf{r}-\mathbf{r}') = V(\mathbf{r}'-\mathbf{r})$. We fabricate a doubling of the degrees of freedom by introducing the Nambu-spinor $\check{c}_a = (c_a \; c^\dagger_a)$, constraining a particle-hole symmetry. In this Nambu representation the system's reduced density matrix (rDM) corresponds to the tensor product $\rho_{ab} = \langle \check{c}^\dagger_b \otimes \check{c}_a \rangle$, reading explicitly as

$$\rho_{ab} = \begin{pmatrix} \langle c^\dagger_b c_a\rangle & \langle c_b c_a\rangle \\ \langle c^\dagger_b c^\dagger_a\rangle & \langle c_b c^\dagger_a\rangle \end{pmatrix} = \begin{pmatrix} \rho^{\rm ee}_{ab} & \rho^{\rm eh}_{ab} \\ \rho^{\rm he}_{ab} & \rho^{\rm hh}_{ab} \end{pmatrix}, \tag{2}$$

with it's terms relating to each other directly from the fermionic anti-commutator properties.

Performing a mean-field Hartree-Fock decoupling as follows $\langle c^\dagger_\alpha c^\dagger_\beta c_\gamma c_\delta\rangle \approx \rho^{\rm he}_{\beta\alpha}\rho^{\rm eh}_{\delta\gamma} - \rho^{\rm ee}_{\gamma\alpha}\rho^{\rm ee}_{\delta\beta} + \rho^{\rm ee}_{\delta\alpha}\rho^{\rm ee}_{\gamma\beta}$, the Nambu rDM equation of motion can be shown to read simply as $i\hbar\dot{\rho}_{ab} = \left[H^{\rm BdG}_{ab}, \rho_{ab}\right]$ with

$$H^{\rm BdG}_{ab} = \begin{pmatrix} h^{\rm HF}_{ab} & \Sigma^\Delta_{ab} \\ (\Sigma^\Delta_{ba})^* & -h^{\rm HF}_{ba} \end{pmatrix} \tag{3}$$

the effective Bogoliubov-de Gennes (BdG) Hamiltonian, $h_{\rm HF}(\rho) = h + \mathbf{\Sigma}_H(\rho) + \mathbf{\Sigma}_F(\rho)$ the Hartree-Fock Hamiltonian, having defined the Hartree, Fock and pairing self-energies as:

$$\begin{matrix} \Sigma^H_{ab} = \sum_{\gamma\delta} V^{\delta a}_{b\gamma}\rho^{\rm ee}_{\gamma\delta} \\ \Sigma^F_{ab} = -\sum_{\gamma\delta} V^{a\delta}_{b\gamma}\rho^{\rm ee}_{\gamma\delta} \end{matrix} \quad \text{and} \quad \Delta_{ab} = \sum_{\gamma\delta} V^{ab}_{\delta\gamma}\rho^{\rm eh}_{\gamma\delta} \tag{4}$$

Note the pairing symmetry $\Sigma^\Delta_{ab} = -\Sigma^\Delta_{ba}$.

We build upon these results by further realizing the mean-field self-energy as a spin-1/2 Hubbard term in the Wannier limit, i.e considering the approximation of ultra-localized orbitals $V^{ij}_{kl} \approx V^{ij}\delta_{il}\delta_{jk}$ with a spinless potential $V^{i\sigma_i j\sigma_j} = V^{ij}$, followed by the exclusion of all offsite term contribution through $V^{ij} = U\delta_{ij}$ with $U < 0$ an attractive Hubbard potential. The Nambu-spinor follows as $\check{\boldsymbol{c}}_i = (c_{i\uparrow} \; c_{i\downarrow} \; c^\dagger_{i\uparrow} \; c^\dagger_{i\downarrow})$. Defining a new BdG symmetric rDM object as

$$\tilde{\boldsymbol{\rho}}_{ij} = \begin{pmatrix} \boldsymbol{\rho}^{\rm ee}_{ij} & -\boldsymbol{\rho}^{\rm eh}_{ij} \\ -\left(\boldsymbol{\rho}^{\rm eh}_{ji}\right)^* & -\boldsymbol{\rho}^{\rm ee}_{ji} \end{pmatrix}, \tag{5}$$

allows one to write the total onsite self-energy of the system in the compact form

$$\mathbf{\Sigma}_{ij} = \frac{1}{2}U\left[{\rm Tr}\left(\left[\tau_z\otimes\sigma_0\right]\tilde{\boldsymbol{\rho}}_{ii}\right)\left[\tau_z\otimes\sigma_0\right] - 2\tilde{\boldsymbol{\rho}}_{ij}\right]\delta_{ij}. \tag{6}$$

with both σ and τ being Pauli matrices operating in their respectively spin and electron-hole subspaces and the Kronecker deltas δ_{ij} denoting on-site terms and $\delta^\pm_{ij\pm1}$ the 1st neighbors hoppings (front and back). Note that the underlying spin structure is being concealed within the bold notation.

III. NANOWIRE MODELS

First and foremost, we showcase Oreg-Lutchyn minimal Majorana model [?] as our model's groundwork. Oreg-Lutchyn minimal model consists of a finite one-dimensional nanowire in the presence of an external Zeeman field $V_{\rm Z}$ and spin-orbit coupling (SOC) effect, being described by the Hamiltonian

$$\hat{\boldsymbol{H}}_0 = \sum_{\sigma_i\sigma_j} c^\dagger_{\sigma_i}\left(-\eta\partial^2_x - \mu + \alpha_{\rm R} i\sigma_y\partial_x + V_{\rm Z}\sigma_z\right)c_{\sigma_j}, \tag{7}$$

with $\eta = \hbar^2/2m^*$, m^* the electron's effective mass, and μ the chemical potential. The SOC effect due an externally applied transverse magnetic field of Rashba strength $\alpha_{\rm R}$ and the Zeeman splitting $V_{\rm Z} = g_J\mu_B B_x/2$ due to a in-plane magnetic field B_x, with g_J the Landé $g$yromagnetic factor of the semiconductor and μ_B Bohr's magneton. Furthermore, the minimal model then considers a s-type superconductive (SC) proximity effect of uniform pairing Δ_0 reading as $(\Delta_0 c^\dagger_\uparrow c^\dagger_\downarrow + h.c)/2$.

The major step up to the minimal model is that we do not consider an homogeneous superconductive proximity effect term but instead self-consistently calculate for the induced pairing (but also magnetization and doping).

As derived in the previous section, this is realized in a Hartree-Fock-Bogoliubov mean-field theory. Within the Nambu tight-binding scheme, this correction amounts to the substitution $\Delta_0 \left[\tau_y \otimes \sigma_y\right] \delta_{ij} \to \boldsymbol{\Sigma}_{ij}$, with $\boldsymbol{\Sigma}_{ij}$ made explicit in Eq.(6). The full Hamiltonian matrix reads explicitly as

$$\boldsymbol{H} = \left(f\delta_{ij} - t\delta_{ij\pm1}\right)\left[\tau_z \otimes \sigma_0\right] + V_{\mathrm{Z}}\left[\tau_z \otimes \sigma_z\right]\delta_{ij} + \frac{\alpha_{\mathrm{R}}}{2a_0}\delta^{\pm}_{ij\pm1}\left[\tau_z \otimes i\sigma_y\right] + \boldsymbol{\Sigma}_{ij} \tag{8}$$

with $f = 2t - \mu$ the band filling ($\mu = 0$ and $\mu = 2t$ corresponding to half and zero filling respectively), $t = \eta/a_0^2$ the hopping parameter and a_0 the real-space lattice spacing.

Schematic models of the hybrid and intrinsic self-consistent Majorana nanowires are shown in Fig.(1).



Figure 1. **Self-consistent finite nanowire models. (a) Hybrid model.** A depleted semiconducting (SM) nanowire in a doped s-wave superconducting (SC) nanowire substrate. Both nanowires are subject to an externally applied magnetic field B but only the SM exhibits spin-orbit coupling. Crucially, only the SC nanowire contains the self-consistent Hubbard term U_{SC} (therewith $U_{\mathrm{SM}} = 0$), thus serving as the sole source of pairing within the system. The proximity effect is modeled by a hopping with intensity $\tau < t$. Moreover, note that considering $\mu_{\mathrm{SC}} \gg \mu_{\mathrm{SM}}$ is crucial for an effective proximity effect because we do not account for disorder and the parallel momentum must be conserved across the interface. **(b) Intrinsic model.** Just a doped semiconductor wire (no SC subtract) with strong SOC subjected to an externally applied magnetic field B. The superconductivity arrives intrinsically due to the self-consistent Hubbard term U.

IV. RESULTS

Somewhat worthy, but better if you write it. Here are some not so obvious small talking points such that you don't forget them:

1. Mention zero-temperature somewhere
2. Bruno mentioned to me after the presentation that the $\tilde{\rho}$ form could be more elegant, so maybe I will look into it.
3. Shallow mode μ shift.
4. Self-consistent routine.
   - Full spectrum diagonalization for finite case
   - Schur's factorization for infinite case
   - Anderson acceleration method



Figure 2. **Self-consistent nanowire models results.** The white boundary line denotes the abrupt change in the Majorana number $\mathcal{M}$ marking the trivial-topological phase transition. Common to all models, we have set the lattice spacing to $a = 10$nm, the finite nanowire length to $L = 2000$nm, the effective mass to $m^* = 0.023m_e$ and the SOC Rashba strength to $\alpha = 40$meV. For the hybrid case the SC Hubbard strength is $U_{\mathrm{SC}} = -30$meV with the SC chemical potential $\mu_{\mathrm{SC}} = 5$meV and SC-SM hopping coupling $\tau = 0.8t$. For the intrinsic case the Hubbard strength is $U = -8$meV.

5. Analytical BCS gap equation:

   Furthermore, we compare the μ_0 linecuts to the analytical BCS gap equation, generalized to account for Zeeman but not SOC, derived in appendix:

$$\Delta_0 = 2\epsilon_0 \exp\left(-\sqrt{\frac{\pi^2}{2}\frac{\hbar^2}{m_e e}}\sqrt{\frac{\mu}{m_0}}\frac{1}{|U|a_0}\right), \tag{9}$$

   It is important to note that Eq.(9) assumes $\mu \gg \Delta$. See that the actual spectral gap Ω (show in dashed black) does not always correspond to the pairing potential Δ_0. Generally it reads $\Omega = \max(0, |\Delta_0| - |V_{\mathrm{Z}}|)$ since for $|V_{\mathrm{Z}}| > \Delta_0$ the gap has to close although the pairing remains finite. Moreover, see that one always needs $\mu > 0$ or else the pairing becomes non-real. In the particular case $V_{\mathrm{Z}} = 0$ with $\mu < 0$ the gap must correspond instead to $\Omega = -\mu$.

6. Direct analysis from the figure.

- Topological invariant, OL hyperboloid
- Self-induced dopping shifts μ_0^{hyb} and $\mu_0^{\mathbf{int}}$
- Mirror-symmetric holeonic phase-transition hyperboloid
- Insulator region in the phase diagram
- No Majorana oscillation because large L

7. Refer to the multi-mode intrinsic results.

V. CONCLUSIONS

Not yet worthy nor ready for the conclusions.

ACKNOWLEDGMENTS

We are deeply grateful to our colleague Rui E. Silva and Bruno Amorim who provided useful insight and expertise that greatly assisted the research of this paper.

[0.0] Lutchyn, Roman M., Jay D. Sau, and S. Das Sarma. "Majorana fermions and a topological phase transition in semiconductor-superconductor heterostructures." Physical review letters 105.7 (2010): 077001.

[0.1] Oreg, Yuval, Gil Refael, and Felix Von Oppen. "Helical liquids and Majorana bound states in quantum wires." Physical review letters 105.17 (2010): 177002.

[0.2] Chamon, C., et al. "Quantizing Majorana fermions in a superconductor." Physical Review B 81.22 (2010): 224515

[0.3] Prada, Elsa, et al. "From Andreev to Majorana bound states in hybrid superconductor–semiconductor nanowires." Nature Reviews Physics 2.10 (2020): 575-594..

[0.4] Avila, J., et al. "Superconducting islands with semiconductor-nanowire-based topological Josephson junctions." arXiv preprint arXiv:2003.02852 (2020).

[0.5] Sau, Jay D., et al. "Generic new platform for topological quantum computation using semiconductor heterostructures." Physical review letters 104.4 (2010): 040502.

[0.6] Fu, Liang, and Charles L. Kane. "Superconducting proximity effect and Majorana fermions at the surface of a topological insulator." Physical review letters 100.9 (2008): 096407.

[1.1] Nambu, Yoichiro. "Quasi-particles and gauge invariance in the theory of superconductivity." Physical Review 117.3 (1960): 648.

APPENDIX

Appendix A: Analytical gap equation

The main objective of this section is to derive the typical textbook BCS gap equation for an infinite system without Zeeman or SOC [??] from our Hartree-Fock-Bogoliubov approach. We generalize the result for a Zeeman field, accounting not only for the self-consistent pairing but also the self-consistently induced magnetization and doping. We will not, however, account for SOC effects.

Foremost, let us reparametrize the self-energy of Eq.(6) in terms of the self-consistently induced magnetization $\delta V_{\rm Z}$, doping $\delta\mu$ and pairing Δ (assumed to be real) as $\mathbf{\Sigma}_{ii} = (\delta V_{\rm Z}\left[\tau_z\otimes\sigma_z\right] - \delta\mu\left[\tau_z\otimes\sigma_0\right]) - \Delta\left[\tau_y\otimes\sigma_y\right]$. Comparing this rewritten form with the original one, we can identifying the off-diagonal block $\Sigma^{\rm he}_{ii}$ as being $-i\Delta\sigma_y = U\rho^{\rm eh}_{i\uparrow i\downarrow}$, noting that $\rho^{\rm eh}_{i\uparrow i\downarrow} = -\rho^{\rm eh}_{i\downarrow i\uparrow}$.

In another context, consider the Hamiltonian (no SOC) $H_0(k) = \sum_s c^\dagger_{ks}\left[(\epsilon_k-\mu)\delta_{ss'} + V_Z\sigma_z\right]c_{ks'}$ with s spin and k momentum. For the Hamiltonian at hand we have $\rho^{\rm eh}_{k\uparrow\downarrow} = \frac{1}{N}\sum_k\left[u^*_{k+}v_{k+}f(\tilde{E}_{k+}) + u^*_{k-}v_{k-}f(\tilde{E}_{k-})\right]$ where $\tilde{E}_{k\pm} = \tilde{V}_{\rm Z}\pm\tilde{E}_k$ with $\tilde{E}_k = \sqrt{\Delta^2+\tilde{\epsilon}^2_k}$ (no $\pm$ subindex) are its eigenvalues and $(u_{k\pm},\, v_{k\pm})$ the (normalized) eigenvectors of the $\uparrow\downarrow$ block; $\tilde{\epsilon}_k = \epsilon_k - \tilde{\mu}$, $\tilde{\mu} = \mu + \delta\mu$ and $\tilde{V}_{\rm Z} = V_{\rm Z} + \delta V_{\rm Z}$ already account for the U-induced shift and f is the Fermi-Dirac distribution. It is critical to realized that is in this generalization we have additional, implicit, self-consistent condition for both $\delta V_{\rm Z}$ and $\delta\mu$. These are obtained in a similar fashion to the pairing term but considering instead the diagonal elements of $\mathbf{\Sigma}_{ii}$. The self-consistent system of three equation for pairing Δ, induced magnetization $\delta V_{\rm Z}$ and doping $\delta\mu$ read as

$$\Delta = -U\sum_k\frac{\Delta}{4\tilde{E}_k}\left[\tanh\left(\frac{\tilde{V}_{\rm Z}+\tilde{E}_k}{2k_BT}\right) - \tanh\left(\frac{\tilde{V}_{\rm Z}-\tilde{E}_k}{2k_BT}\right)\right] \tag{10}$$

$$\delta V_{\rm Z} = -\frac{U}{2N}\sum_k\frac{\sinh\left(\frac{\tilde{V}_{\rm Z}}{k_BT}\right)}{\cosh\left(\frac{\tilde{V}_{\rm Z}}{k_BT}\right)+\cosh\left(\frac{\tilde{E}_k-\tilde{V}_{\rm Z}}{k_BT}\right)} \tag{11}$$

$$\delta\mu = \frac{U}{4} + \frac{U}{4N}\sum_k\frac{\tilde{\epsilon}_k}{2\tilde{E}_k}\left[\tanh\left(\frac{\tilde{V}_{\rm Z}+\tilde{E}_k}{2k_BT}\right) - \tanh\left(\frac{\tilde{V}_{\rm Z}-\tilde{E}_k}{2k_BT}\right)\right] \tag{12}$$

See that, if one ignores in Eq.(10) Zeeman all together, $V_{\rm Z}$ and $\delta V_{\rm Z}$, but also the self-consistently doping $\delta\mu$ it reduces to the traditional gap equation [??] as intended. Note that this has a nonzero solution only if the interaction is attractive i.e $U<0$. Although these analytical expressions can be obtained, the generalized analytical solution of the self-consistent system of equation is neither trivial nor worth the trouble for our purposes. We can, however, solve the conventional gap equation in the zero-temperature, small-gap limit, which offers a sufficiently accurate analytical comparison to our numerical results. Thereby, we approximate the dispersion to $\epsilon_k = \pm v_{\rm F}(k\mp k_{\rm F})$ with $v_{\rm F}$ and $k_{\rm F}$ the Fermi velocity and wavevector respectively, and mutate the $\frac{1}{N}\sum_k$ sum into an integral over ϵ_k, introducing also a fitted phenomenological cutoff ϵ_0. In the limit where $\Delta\ll\epsilon_0$ and for $U<0$ the gap equation simplifies to

$$\Delta = 2\epsilon_0\exp\left(-\sqrt{\frac{\pi^2}{2}\frac{\hbar^2}{m_e e}}\sqrt{\frac{\mu}{m_0}}\frac{1}{|U|a_0}\right). \tag{13}$$

Even though we neglected Zeeman this solution is still valid for finite $|V_{\rm Z}|\le\Delta$ since the tanh factors remain equal to 1 throughout the integration interval. Nevertheless, the generalization of Eq.(13) for $|V_{\rm Z}|>\Delta$ can be found from Eq.(10) within the same regime of approximation and expressed in terms of the previous $\Delta_0\equiv\Delta(V_{\rm Z}=0)$ solution as

$$\Delta = \frac{1}{2\epsilon_0}\sqrt{\Delta_0\left(2\left|V_{\rm Z}\right|-\Delta_0\right)}. \tag{14}$$

It is important to note that the actual spectral gap Ω does not always correspond to the pairing potential Δ_0. Generally it reads $\Omega = \max(0, |\Delta_0| - |V_{\rm Z}|)$ since for $|V_{\rm Z}|>\Delta_0$ the gap has to close although the pairing remains finite. Moreover, see that one always needs $\mu>0$ or else the pairing becomes non-real. In the particular case $V_{\rm Z}=0$ with $\mu<0$ the gap must correspond instead to $\Omega=-\mu$.