

Methods and Applications of CRISPR-Mediated Base Editing in Eukaryotic Genomes

Gaelen T. Hess,^{1,2} Josh Tycko,^{1,2} David Yao,^{1,2} and Michael C. Bassik^{1,*}

¹Department of Genetics and Stanford University Chemistry, Engineering, and Medicine for Human Health (ChEM-H), Stanford, CA, USA

²These authors contributed equally

*Correspondence: bassik@stanford.edu

<https://doi.org/10.1016/j.molcel.2017.09.029>

The past several years have seen an explosion in development of applications for the CRISPR-Cas9 system, from efficient genome editing, to high-throughput screening, to recruitment of a range of DNA and chromatin-modifying enzymes. While homology-directed repair (HDR) coupled with Cas9 nuclease cleavage has been used with great success to repair and re-write genomes, recently developed base-editing systems present a useful orthogonal strategy to engineer nucleotide substitutions. Base editing relies on recruitment of cytidine deaminases to introduce changes (rather than double-stranded breaks and donor templates) and offers potential improvements in efficiency while limiting damage and simplifying the delivery of editing machinery. At the same time, these systems enable novel mutagenesis strategies to introduce sequence diversity for engineering and discovery. Here, we review the different base-editing platforms, including their deaminase recruitment strategies and editing outcomes, and compare them to other CRISPR genome-editing technologies. Additionally, we discuss how these systems have been applied in therapeutic, engineering, and research settings. Lastly, we explore future directions of this emerging technology.

Introduction

The ability to precisely edit genomic DNA using the CRISPR-Cas9 system has revolutionized the field of genome engineering (reviewed in Doudna and Charpentier, 2014; Hsu et al., 2014; Komor et al., 2017a; Sander and Joung, 2014; Shalem et al., 2015; Sternberg and Doudna, 2015; Wang et al., 2016). This targeted nuclease system was discovered as a bacterial adaptive immune system where foreign DNA is incorporated into an array of clustered regularly interspaced short palindromic repeats (CRISPRs) (Barrangou, 2013; Ishino et al., 1987; Jansen et al., 2002; Mojica et al., 2000). The short segments of foreign DNA captured between repeats are then used to generate guide RNAs, composed of a targeting CRISPR RNA (crRNA) and trans-activating CRISPR RNA (tracrRNA). These complex with Cas9 endonuclease, which is targeted to the foreign DNA via the crRNA sequence and a required protospacer-adjacent motif (PAM) to digest the foreign DNA (Brouns et al., 2008; Garneau et al., 2010; Marraffini and Sontheimer, 2008). Significant work has been done to reveal the underlying mechanisms of this process (Deltcheva et al., 2011; Gasiunas et al., 2012; Jiang and Doudna, 2017; Jinek et al., 2012; Nishimasu et al., 2014) and enable its transfer to eukaryotic systems (Cong et al., 2013; Jinek et al., 2013; Mali et al., 2013). In most applications, single guide RNAs (sgRNAs) (Jinek et al., 2012), which combine the crRNA and tracrRNA, are used to target the Cas9 nuclease to a specific site in the genome, where it introduces a double-strand break (DSB).

Once a DSB is formed by Cas9 and an sgRNA, it can be repaired through non-homologous end-joining (NHEJ) or homology-directed repair (HDR) (Jasin and Haber, 2016). NHEJ introduces insertions or deletions at the damage site (Cong et al., 2013; Jinek et al., 2013; Mali et al., 2013) and is useful for generating frameshifts in coding regions causing effective gene

knockout (Shalem et al., 2014, 2015; Wang et al., 2014) or disrupting transcription factor binding motifs (Canver et al., 2015; Ernst et al., 2016; Rajagopal et al., 2016; Sanjana et al., 2016) and splicing elements (Charton et al., 2016; Kapahnke et al., 2016; Mou et al., 2017). Alternatively, templated repair of the DSB can be used to introduce specific changes through HDR in the presence of either single- or double-stranded DNA donor templates that encode the desired sequence. Cas9-directed HDR, which utilizes homologous recombination and other repair pathways, has enabled dramatic improvements in endogenous tagging of genes (Bak and Porteus, 2017; Lackner et al., 2015; Leonetti et al., 2016), generation of in-frame protein variants (Findlay et al., 2014; Garst et al., 2017; Ryan et al., 2014), and clinical correction of deleterious mutations (Dever et al., 2016; DeWitt et al., 2016; Yang et al., 2016b). Significant optimization of Cas9-mediated genome editing has allowed for single-nucleotide editing efficiencies reaching 60% or more in cell culture systems (Richardson et al., 2016). However, this strategy generally depends on inducing a DSB; therefore, small indels are an unintended by-product, which can have major negative functional consequences if the targeted gene is knocked out.

In order to limit DNA damage during editing and eliminate the need to deliver an HDR donor template, a number of groups have recently developed CRISPR base-editing systems. These make use of Cas9 variants, cytidine deaminases, and manipulation of DNA repair pathways to achieve specific editing outcomes. Importantly, base-editing systems enable both highly targeted single-base changes (see Precision Base Editing) and local sequence diversification (see Localized Sequence Diversification). Here we review the different base editor systems, including their components, and how they affect the repair of the deaminated base with respect to efficiency and possible base




Figure 1. Molecular Components and Mechanisms for Base Editing

(A) The molecular components of the base editor (BE), Target-AID, Targeted AID-mediated Mutagenesis (TAM), and CRISPR-X technologies are shown. The BE3, YEE-BE3, and Target-AID systems employ the Cas9 D10A (purple) nickase along with a cytidine deaminase (red) and a uracil DNA glycosylase inhibitor (yellow). The BE4 system also uses the Gam protein (pink). The TAM and CRISPR-X systems used dCas9 (green) to recruit variants of the deaminase AID (AIDx or MS2-AID*Δ). Below each technology, the targeting window is indicated. The increased spectrum of base substitutions that can be achieved with TAM and CRISPR-X is represented by the multi-color editing windows.

(B) For base editors, the mechanisms of DNA repair are critical for the eventual editing outcome. After the initial deamination (C to U) conversion, uracil DNA glycosylase can remove the base and perform error-free (correctly replacing U with C) or error-prone repair, generating different substitutions (left side).

(legend continued on next page)

substitutions. Given the therapeutic potential for base editing, current approaches for identifying off-targets of the system are discussed in detail. Lastly, we review the diverse applications of the base editors and highlight future directions for the field.

Section 1: Repurposing Deaminases for Base Editing and Diversification

Natural Roles of Deaminases

Deaminases play important roles in diverse processes including musculoskeletal myogenesis (Hsieh et al., 2014), central nervous system development and function (Li and Church, 2013), and innate and adaptive immune pathways (Hamilton et al., 2010; Keegan et al., 2001; Samuel, 2012; Vieira and Soares, 2013). The apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like (APOBEC) family of cytidine deaminases (Chiu and Greene, 2006; Navaratnam and Sarwar, 2006) restrict viral infection by editing viral genomes. For example, APOBEC3G was identified as a cellular factor capable of restricting HIV infection (Sheehy et al., 2002; Yu, 2006); it deaminates HIV viral cDNA (Harris et al., 2003; Lecossier et al., 2003), rendering it incapable of replication. APOBEC3 proteins have also been shown to restrict the activity of endogenous retroelements (Esnault et al., 2005; Koito and Ikeda, 2013; Muckenfuss et al., 2006). Deaminases also enable functional diversification: APOBEC1 was the first RNA-editing enzyme identified (Powell et al., 1987) and functions in lipid homeostasis by editing apolipoprotein B (apoB) mRNA, converting a glutamine to a stop codon. This creates a truncated APOB48 protein specifically in the small intestine that transports dietary lipids, while the full-length APOB100 form retains the ability to bind the LDL receptor and performs a more general role in lipid transport around the body (Chan, 1992).

Activation-induced cytidine deaminase (AID) is responsible for antibody diversification through somatic hypermutation and class switch recombination. During somatic hypermutation, AID is specifically targeted to the Ig locus, where it creates diverse mutations that are then subjected to selection through antigen binding; this allows for maturation of both affinity and specificity. Loss of AID function prevents class switch recombination and somatic hypermutation, which causes a form of hyper-immunoglobulin M syndrome where low-affinity antibodies are generated (Revy et al., 2000; Xu et al., 2012). Promiscuous targeting of AID outside the Ig locus has been associated with B cell lymphomas (Küppers et al., 1999; Xu et al., 2012) and has been shown to be required for development of germinal-center-derived B cell lymphomas in several mouse models (Pasqualucci et al., 2008; Ramiro et al., 2006). Intriguingly, the process of somatic hypermutation mediated by AID biases repair pathways to error-prone outcomes through mechanisms that are incompletely understood (Larson et al., 2005; Odegard and Schatz, 2006; Reynaud et al., 2003).

These deaminases share a common mechanism by which they modify cytidine to uridine. RNA targets of cytidine deamination do not undergo further processing, but for DNA, the conversion generates a DNA lesion that can be repaired through several pathways: (1) uridine can be replaced by a thymidine during DNA replication, (2) base excision repair (BER) can excise the uridine allowing the insertion of an alternative nucleotide, or (3) mismatch repair (MMR) by way of translesion synthesis, where an error-prone polymerase goes through the lesion, increasing the possibility of mutations at nearby nucleotides. Incorporation of uracil is common, and its repair is often performed in an error-free manner; however, the potential for error-prone repair facilitates antiviral defense and functional diversification.

Precision Base Editing with BE3/4 and Target-AID

The ability to manipulate the damage and repair pathways utilized by cytidine deaminases, combined with the precise targeting of Cas9, is a powerful strategy to introduce specific point mutations. Cas9-targeted deamination was first demonstrated in the pioneering work of the Liu group to develop the Base Editor (BE) system (Komor et al., 2016). Since a major goal of this work was to induce base changes without double-strand DNA breaks, the initial BE1 system included only rat deaminase APOBEC1 (rAPOBEC1) fused to deactivated Cas9 (dCas9), in which the two nuclease domains of Cas9 are inactivated by mutation of D10A and either H840A or N863A (Jinek et al., 2012; Qi et al., 2013). Transfection of BE1 and a targeted sgRNA successfully converted cytidines to thymidines within a window of –16 to +12 bases from the PAM of the sgRNA (Figure 1A).

An optimized BE2 system resulted from the addition of the uracil DNA glycosylase inhibitor (UGI) (Figure 1A) from the *Bacillus subtilis* bacteriophage PBS1 (Mol et al., 1995). UGI directly binds and inhibits uracil DNA glycosylase, thus blocking uridine excision and the ensuing BER pathway. This improvement increased base-editing efficiency of the C>T substitution 3-fold by effectively disfavoring error-free repair (Figure 1B). Further improvement was realized by changing the dCas9 to a “nickase” Cas9 D10A, which nicks the strand opposite the deaminated cytidine. In principle this should initiate long-patch BER, where the deaminated strand is preferentially used to template the repair to produce a U:A base pair. This intermediate is then converted to T:A during DNA replication (Heller and Mariani, 2006; Robertson et al., 2009; Zharkov, 2008) (Figure 1B). The optimized BE3 editor consists of rAPOBEC1 fused to the N terminus of nickase Cas9 D10A and a UGI fused to the C terminus. This iteration, BE3, achieved a 6-fold increase in efficiency over BE2, with up to 37% of targeted alleles edited. It is worth noting that the optimized BE3 system also exhibited a slightly increased indel frequency of 1.1%, as compared to < 0.1% for the BE2 system, since nicks can lead to NHEJ at a low rate (Certo et al., 2011).

A similar system, Target-AID (Nishida et al., 2016), uses nickase Cas9 D10A to recruit the cytidine deaminase PmCDA1 (from

Alternatively, a uracil DNA glycosylase inhibitor (UGI) can be used to block base excision by uracil N-glycosidase (UNG). Nicking via a Cas9 nickase of the non-deaminated strand promotes long-patch BER where the deaminated strand is preferred for templated repair, generating a U:A base pair that is then resolved as a T:A base pair. Upon formation of the abasic site, an AP lyase can remove the site, forming a ssDNA break. This excision in conjunction with the Cas9 D10A can create a DSB. Addition of the bacteriophage protein Gam reduces the frequency of indel formation by directly binding the DSB, which may select against cells with DSBs. Endogenous repair machinery is labeled in black and represented with black triangles. Exogenous components of the base-editing tools are illustrated and labeled in color.

Box 1. Quantifying Mutation Frequency

Quantifying base-editing mutation rates is a challenging problem, especially at the low frequencies induced over the large sequence space generated by diversifying base editors (analogous to the problems encountered when sequencing the B cell receptor repertoire). In addition to sequencing assays, fluorescent and drug resistance assays have been used to estimate the rate of mutations.

MEASUREMENT BY PHENOTYPE

In a number of studies, change in a gene's phenotype has been used as a proxy for mutation rate. For example, both the TAM and CRISPR-X systems were initially quantitated by following GFP fluorescence (by repair of a stop codon or loss of fluorescence, respectively). Similarly, on-target editing by the Target-AID system has been measured in *S. cerevisiae* by following loss of function in the CAN1 gene (causing resistance to canavanine) or ADE1 gene (red pigment upon loss of function). In this study, loss of LYP1, whose knockout provides protection against S-aminoethyl-L-cysteine, was used as a separate marker for non-specific mutations in the genome. In mammalian cells, the HPRT gene was targeted, where its loss of function conferred resistance to 6TG (Nishida et al., 2016). Although these various phenotypic measurements provide a rapid method for assessing mutation frequency, it is important to note that this may be inaccurate in some cases because not all mutations lead to a measurable phenotype; conversely, the phenotype may require all alleles to be mutated simultaneously. As an example of this discrepancy, the rescue of GFP fluorescence using the TAM system was observed in 4%–5% of cells, but sequencing of the GFP loci revealed more than 20% of the loci contained at least one mutation (Ma et al., 2016).

MEASUREMENT BY SEQUENCING

High-throughput sequencing of the locus can address these issues and has been used to measure the frequency of mutations induced by base editors. Sequencing data are typically filtered for read quality and overall alignment to the regions targeted for mutagenesis before quantifying the frequency of mutation at each base position in the targeting window, when compared to the unmutated gene or parent population.

One key limitation to high-throughput sequencing approaches for detecting mutations is the inherent error rate. Errors can be introduced in a number of ways: (1) SNPs or sequence heterogeneity may already exist in some cultured cell lines, (2) PCR error can occur during the preparation of sequencing libraries, or (3) bases can be incorrectly read during sequencing. Error rates for high-throughput sequencing systems are typically well below 1% (Reuter et al., 2015), but still limit the detection frequency of mutations. These errors can be estimated and accounted for by preparing and sequencing unmutated samples in parallel. For example, in the CRISPR-X system, mutation frequencies for individual bases were calculated by normalizing to the average frequency of mutations in non-mutated samples. The standard deviation of the mutation frequency for non-mutated samples was used to estimate the noise in sequencing. The calculated standard deviation of the mutation frequency at a single base was 0.05%, allowing for the detection of mutations as low as 0.05%. Using this background, mutation frequency was quantified at each base position in the 100 bp targeting window for the 30 sgRNAs tested, and ranged as high as 20% (Hess et al., 2016). While it may be possible to detect even more rare mutations, these may require sequencing samples with high coverage over the targeted area in order to confidently distinguish them from errors in sequencing.

A second limitation is the ability to detect co-occurring mutations when multiple sgRNAs are introduced over a large region (>600 bp). While long-read technologies exist, they are limited in throughput, and for highly diverse populations, the lack of sequencing depth can misrepresent the population. Alternatively, by measuring the relative frequencies of mutations in the population after each round of selection, it may be possible to determine which mutations co-occur (Sakoparnig et al., 2015; Wang et al., 2012; Yeang et al., 2008; Zhang et al., 2014).

A final consideration, especially for experiments with diversifying base edits, is the number of cells that should be maintained for the experiment. This number plays a critical role in experimental design to avoid bottlenecking the population and to determine the sequencing depth needed to assay the population. Estimating diversity through sequencing is a complex problem due to sequencing errors, as described above. Significant work has been done on this problem when estimating the diversity of the antibody repertoire (reviewed in Benichou et al., 2012; Robins, 2013). This undertaking has proved challenging due to the complexity of the library and the need to separate the mutation and selection processes. However, through a number of statistical models (Elhanati et al., 2015; Greiff et al., 2015; Mora et al., 2010) and other strategies (reviewed in Hoehn et al., 2016), the potential diversity of the antibody repertoire has been estimated to have a lower bound of 10^{18} (Elhanati et al., 2015), though in fact there may only be $3\text{--}9 \times 10^6$ different antibodies among $\sim 10^9$ B cells in the human body (Arnaout et al., 2011). While quantifying mutation frequency over a single sgRNA-targeted region is somewhat simpler, significant variability has been observed in terms of the range of mutations that can be induced with diversifying base editors. For example, for the TAM and CRISPR-X systems, mutation frequencies at single bases for a single sgRNA targeting an integrated GFP locus were found to range between $\sim 1\%$ and 25% (Hess et al., 2016; Ma et al., 2016). Given the challenge of quantifying diversity in a starting population of mutated cells for diversifying

(Continued on next page)

Box 1. Continued

base editors, it is helpful to estimate the total diversity created by a single sgRNA. For example, between 20% and 80% of all possible single-base mutations were detectable above background sequencing noise within the +20 to +40 bp mutation hotspot, measured in a population (0.5–1.5 million cells extracted) with an average of 4.5 million sequencing reads using the CRISPR-X system (Hess et al., 2016). This calculation provides an estimate of the number of possible variants made by a single sgRNA. In this way, mutation frequency, size of the mutagenized region, population size, and sequencing read depth can be taken into account when designing selection experiments.

sea lamprey) to a gene of interest. Using this system to target the ADE1 gene in yeast, ~15%–50% of cells could be induced to exhibit the red colony phenotype for the mutated ADE1 gene (Box 1). When this system was transported into mammalian cells using a dCas9-PmCDA1 fusion to target the HPRT gene, a smaller fraction (~2%) of cells exhibited resistance to 6-Thioguanine (6TG). Introducing the Cas9 D10A nickase increased the frequency of resistant cells to 35%, but subsequent targeted sequencing revealed that many of these mutations were deletions. To address this, the authors added UGI to block the removal of the uracil base and observed a 2- to 3-fold increase of mutation frequency and reduction in deletions. Further characterization through deep sequencing at other targeted loci indicated mutation frequencies of 15%–55% localized to –19 to –15 bases with respect to the PAM, similar to those observed in yeast. Thus, while BE3 and Target-AID use different cytidine deaminases and protein fusion arrangement (Figure 1A), both demonstrate the utility of an opposite-strand nick and UGI to increase C>T base-editing efficiency.

Within a year of its publication, BE3 has been widely applied to edit diverse cell, plant, and animal genomes (see [Applications of Precision Editing](#)), but further development of base editor tools has continued in parallel. The Liu group has now introduced BE4, with optimized longer linkers and two fused copies of UGI to further reduce uracil N-glycosylase (UNG)-mediated BER (Komor et al., 2017b) (Figure 1A). This design decision is well supported by experiments in UNG knockout cells that clearly demonstrate UNG is necessary for generating most undesirable substitutions and indels. The additional fusion of the DNA end-binding Gam protein (from Mu bacteriophage) (d'Adda di Fagnola et al., 2003) to BE4 helped moderately reduce undesirable indels. In theory, this works by Gam binding the double-stranded breaks that occasionally result from the combination of an endogenous DNA-apurinic or apyrimidinic site lyase (AP lyase) nick on the target strand and a Cas9 D10A nick on the non-target strand (Figure 1B). The authors speculate that Gam competes with error-prone DSB repair processes and induces cell death, thus removing indels from the surviving population of cells. In a head-to-head experiment with six sites in HEK293T cells, BE4 generally outperformed BE3 and Target-AID, although Target-AID is optimal at more PAM-distal Cs because it has a different targeting window. When compared with BE4 containing SpCas9 D10A, BE4 with SaCas9 D10A appeared to maintain consistently high editing efficiency at cytosines within –19 to –9 bases with respect to the SaCas9 PAM. Therefore, the choice of optimal precision base editor for a given target site will likely depend on PAM availability, position of target C relative to PAM, acceptability of indels in population, and delivery modality constraints.

Localized Sequence Diversification with CRISPR-X and TAM

While BE4 and Target-AID perform a specific C>T conversion, an alternative approach is to use deamination as a means of creating a diverse library of point mutations localized to a targeted region of the genome. This is similar to the process of somatic hypermutation during antibody affinity maturation, normally performed by AID. Two technologies, Targeted AID-mediated Mutagenesis (TAM) (Ma et al., 2016) and CRISPR-X (Hess et al., 2016), use this strategy to generate localized sequence diversity through base editing.

In the TAM system (Ma et al., 2016), human AID is fused to the C terminus of dCas9 (Figure 1A). When this system was targeted to a copy of GFP containing a premature stop codon with a pool of sgRNAs, 2% of cells exhibited GFP fluorescence, indicating successful editing. This efficiency could be doubled by removing the C terminus of AID (AIDx), which includes a nuclear export signal (NES) that normally limits AID activity. Sequencing of the targeted GFP locus revealed mutation frequencies > 20%, with transitions and transversions from cytidine and guanine to the other three bases with only a slight bias for C>T and G>A transitions. Coexpression of UGI resulted in an increase in frequency of mutation when using single sgRNAs, but limited mutations to C>T transitions within a window of –16 to –12 bases with respect to the PAM of the sgRNA.

In the CRISPR-X system (Hess et al., 2016), dCas9 is used to recruit a hyperactive variant of the AID enzyme. Three mutations (K10E, T82I, and E156G) were previously identified in a bacterial screen that increased the deamination activity of AID (Wang et al., 2009); this variant was combined with a deletion of the NES to generate AID*Δ. Instead of fusing the enzyme to dCas9, AID*Δ was fused to MS2 protein and recruited to the locus via two MS2 RNA hairpins inserted into the sgRNA backbone (Konermann et al., 2015) (Figure 1A). Compared to native AID without an NES (AIDΔ), which induced mutations within a targeting window of +12 to +32 bp from the PAM site, CRISPR-X using AID*Δ induced a ~2- to 6-fold increased frequency of mutations (> 20%) over a larger window of –50 to +50 bp. Similar to the TAM system, the conversion of C to A, G, or T and G to A, C, or T showed only a slight preference for C>T. Mutations at A and T bases were also observed, but at a much lower frequency. Interestingly, using CRISPR-X, the position of targeted mutations appears to be sensitive to the direction of transcription, with the region of highest mutation downstream of the PAM site with respect to the direction of transcription. This is consistent with previous observations that mutagenesis by AID depends on interactions with the transcription machinery (Chaudhuri et al., 2003). Nonetheless, when CRISPR-X was

Table 1. Efficiency of Precision Base Editors in Diverse Model Systems

Gene (Variant)	Cell Type	Delivery Method	Base Editor	Substitution Frequency	Indel Frequency	Comparison to			Reference
						Active Cas9 (With or Without Donor)	Substitution Frequency	Indel Frequency	
APOE4 (C158R)	Mouse astrocyte	Plasmid nucleofection	BE3	58%–75%	4%–6%	SpCas9 and donor 200-nt ssDNA	0.1%–0.3%	26%–40%	Komor et al., 2016
TP53 (Y163C)	HCC1954 human breast cancer cell line	Plasmid nucleofection	BE3	3%–8%	<0.7%	SpCas9 and donor 200-nt ssDNA	0%	6%–8%	Komor et al., 2016
6 loci (incl. <i>EMX1</i> , <i>FANCF</i> , <i>RNF2</i>)	HEK293T cells	Plasmid lipofection	BE3	5%–37%	1%–6%	SpCas9 and donor 200-nt ssDNA	<2%	2%–7%	Komor et al., 2016
6 loci (incl. <i>EMX1</i> , <i>FANCF</i> , <i>RNF2</i>)	HEK293T cells	Plasmid lipofection	BE4	20%–50%	0.5%–3%	N/A	N/A	N/A	Komor et al., 2017b
6 loci (incl. <i>EMX1</i> , <i>FANCF</i> , <i>RNF2</i>)	HEK293T cells	Plasmid lipofection	Target-AID	5%–50%	0.5%–3%	N/A	N/A	N/A	Komor et al., 2017b
4 loci (incl. <i>FANCF</i>)	HEK293T cells	Plasmid lipofection	SaBE4-Gam	40%–65%	0.5%–4%	N/A	N/A	N/A	Komor et al., 2017b
4 loci (incl. <i>EMX1</i> , <i>FANCF</i> , <i>VEGFA</i>)	HEK293T cells	RNP lipofection	BE3	5%–65%	5%	N/A	N/A	N/A	Rees et al., 2017
<i>TYR1</i> , <i>TYR2</i> , <i>TYR3</i>	Zebrafish embryo	RNP lipid nanoparticle injection	BE3	0%–5%	<2%	N/A	N/A	N/A	Rees et al., 2017
<i>VEGFA</i> site 2	Murine NIH/3T3 cells	Plasmid lipofection	BE3	10%–85%	N/R	N/A	N/A	N/A	Rees et al., 2017
<i>VEGFA</i> site 2	Murine NIH/3T3 cells	RNP lipofection	BE3	5%–45%	N/R	N/A	N/A	N/A	Rees et al., 2017
<i>VEGFA</i> site 2	Mouse inner ear	RNP lipid nanoparticle injection	BE3	<1.5%	<0.1%	N/A	N/A	N/A	Rees et al., 2017
<i>Dmd</i> , <i>Tyr</i>	Mouse embryo	RNP electroporation	BE3	56%–69%	N/R	N/A	N/A	N/A	Kim et al., 2017b
<i>Dmd</i> , <i>Tyr</i>	Mouse embryo	mRNA microinjection	BE3	44%–100%	N/R	N/A	N/A	N/A	Kim et al., 2017b
<i>Tyr</i>	Mouse embryo	mRNA microinjection	HF-BE2	11.6%–50%	4.5%	N/A	N/A	N/A	Liang et al., 2017
<i>HPRT</i> , <i>EMX1</i> , <i>Efemp1</i> , <i>MGAT1</i>	CHO cells	Plasmid lipofection and G418 selection	Target-AID	17%–55%	4%–12%	SpCas9 without donor	0.1%–0.6%	71%–93%	Nishida et al., 2016
<i>Can1</i>	<i>Saccharomyces cerevisiae</i> BY4741	Transformation and selection for mutations	Target-AID	96%	0.2%	SpCas9 without donor	<5%	5%–40%	Nishida et al., 2016
<i>PCSK9</i>	Adult mouse liver	Adenoviral vector	BE3	19%–34%	1%–2%	SpCas9 without donor	N/A	50%	Chadwick et al., 2017; Ding et al., 2014
<i>FTIP1e</i> and <i>ALS</i>	Rice calli	<i>Agrobacterium</i> transfection and selection for mutations	Target-AID	34%–89%	11%–66%	N/A	N/A	N/A	Shimatani et al., 2017
<i>NRT1.1B</i> and <i>SLR1</i>	Rice calli	<i>Agrobacterium</i> transformation and selection for delivery	BE3 without UGI	2.7%–13.3%	8.9%–10.8%	N/A	N/A	N/A	Lu and Zhu, 2017

(Continued on next page)

Table 1. Continued

Gene (Variant)	Cell Type	Delivery Method	Base Editor	Substitution Frequency	Indel Frequency	Comparison to Active Cas9 (With or Without Donor)	Substitution Frequency	Indel Frequency	Reference
<i>DELLA</i> and <i>ETR1</i>	Tomato	<i>Agrobacterium</i> transformation and selection for delivery	Target-AID	33%–85%	14%–69%	SpCas9 without donor	0.5%–2%	25%–44%	Shimatani et al., 2017
<i>OsCDC48</i>	Rice calli	<i>Agrobacterium</i> transformation and selection for delivery	BE3	43.5%	N/R	N/A	N/A	N/A	Zong et al., 2017
<i>TaLOX2</i>	Wheat immature embryo cells	Biolistic gold-DNA particle bombardment without selection	BE3	1.3%	N/R	N/A	N/A	N/A	Zong et al., 2017
<i>ZmCENH3</i>	Maize immature embryo cells	<i>Agrobacterium</i> transformation and selection for delivery	BE3	10.1%	N/R	N/A	N/A	N/A	Zong et al., 2017

A summary of base-editing experiments over the past 2 years. Efficiency data are presented as the approximate minimum and maximum across loci and replicates in a given model system. Where data are available, base editing is compared with nuclease-active Cas9. Quantification of substitution and indel frequencies in studies using cell populations were based on high-throughput sequencing; studies using whole organisms were based on genotyping. RNP, ribonucleoprotein complex of base editor and sgRNA; N/R, not reported; N/A, not applicable.

targeted to regions upstream of annotated transcription start sites, mutations could be detected at similar frequencies to coding regions.

Although the designs of each of these base-editing systems are different, a common thread is the ability to generate point mutations at cytosine and guanine nucleotides with a low rate of indels, in contrast with HDR-based methods, which often generate more indels than substitutions. This is critical when modifying protein-coding genes where indels could generate frameshifts that make truncated or nonsense protein products. The diverse editing outcomes generated by these systems enable both precision editing of single C>T bases as well as sequence diversification; each technology facilitates a range of possible applications.

Section 2: Applications of Base Editing

Though base-editing technologies are still new, there have already been many successful applications *in vitro* and *in vivo*, ranging from precise therapeutic and agronomic editing to broad screens for directed evolution and mapping of protein-drug interactions (Figure 2). These pioneering experiments have been conducted in a wide panel of cell lines and organisms, summarized in Tables 1 and 2. In this section, we review how current base-editing systems have been applied.

Applications of Precision Editing

As a Therapeutic Platform. The precise conversion of cytidine to thymidine by the BE system lends itself as a potential therapeutic approach for the targeted insertion of beneficial mutations. Komor et al. first demonstrated the potential use of this system by correcting Alzheimer's Disease-associated mutations in APOE4 in mouse astrocytes and TP53 mutations in a mouse breast cancer line (Komor et al., 2016). Since this initial demonstration, there have been many successful examples of base editing to substitute a desired mutation (Table 1). Base-editing systems, most notably BE3, have been used in a wide range of mammalian cell culture models including HEK293Ts (Chadwick et al., 2017; Kim et al., 2017a, 2017c; Komor et al., 2016), human iPSCs (Chadwick et al., 2017), CHO cells (Nishida et al., 2016), U2OS cells (Kim et al., 2017c; Komor et al., 2016), HCC1954 cells (Komor et al., 2016), mouse Neuro-2a cells (Chadwick et al., 2017), and mouse astrocytes (Komor et al., 2016). There have also been successful demonstrations of base editing *in vivo* via delivery by adenoviral vector into mouse liver (Chadwick et al., 2017) and by RNP delivery into the mouse inner ear (Rees et al., 2017). Zebrafish (Rees et al., 2017; Zhang et al., 2017) and mouse embryos (Kim et al., 2017b; Liang et al., 2017) have also been successfully base edited, with the latter being reintroduced into surrogate mothers and giving rise to properly edited offspring.

Komor et al. estimated that 3,000 genetic variants listed in ClinVar would be correctable with a C>T substitution. However, given PAM sequence requirements, narrow base editing windows relative to the guide-targeted PAM sequence, and the potential need to avoid editing multiple cytidines in the editing window, they further estimated that only 300–900 would be targetable (Komor et al., 2016). To address these limitations, the BE system has been fused to an array of Cas9 variants with varied PAM-sequence specificities derived from

Box 2. Specificity and Off-Target Base Editing

Unintended off-target mutations pose a challenge to the application of any genome-editing technology. Researchers measuring and improving base-editing specificity can employ some of the same tools developed for preceding technologies, including computational prediction of off-targets and targeted deep sequencing to detect off-target mutations (reviewed in [Hsu et al., 2014](#); [Tsai and Joung, 2016](#); [Tycko et al., 2016](#)). However, distinct properties of base editors also create a need for distinct specificity methods and tools.

PROXIMAL OFF-TARGETS

Base editing at additional cytosines proximal to the targeted base may be unintended and considered as off-target mutations. One solution is to more tightly restrict the targetable window. Optimizing the rAPOBEC1 with a mutant deaminase domain effectively reduced the window from 4–6 bases to as little as 1–2 bases ([Kim et al., 2017c](#)). A triple mutant rAPOBEC1 (YEE-BE3) achieves this extreme precision at the cost of 2.9-fold lower average maximal editing than BE3, while double mutant YE1-BE3 retained maximal editing with a targeting window of 2–3 bases. Notably, the restricted targeting window decreases the likelihood of finding an appropriate NGG PAM site for a given target. However, base editors can be built with a variety of engineered *S. pyogenes* Cas9 PAM variants ([Kleinstiver et al., 2015b, 2016](#)) or *S. aureus* Cas9 and its engineered PAM variants ([Kleinstiver et al., 2015a](#)) to widen the number of available targeting sites ([Kim et al., 2017c](#); [Komor et al., 2017b](#)). Careful experimental design can sometimes obviate the need for these engineered variants, as proximal off-target mutations are relatively inconsequential in coding regions if the off-target mutation creates a synonymous codon ([Komor et al., 2016](#)).

GENOME-WIDE OFF-TARGETS

Base editors can also induce distal off-target mutagenesis at other binding sites across the genome. In fact, as Cas9 binding is more promiscuous than Cas9 cleavage ([Wu et al., 2014](#)), one might expect less specificity from base editors than active Cas9 nuclease. However, head-to-head comparisons across a range of mismatched target sites show generally similar specificity rules, with notable exceptions in which their off-targets do not coincide ([Kim et al., 2017a](#)). For example, BE3 appears more tolerant of DNA-sgRNA mismatch bulges than Cas9. Unsurprisingly, Cas9 is more prone to off-target effects at sites without editable cytosines in the BE3-targetable window as the base editor induces minimal mutagenesis of non-cytosine bases.

These exceptions, along with the fact that most off-target detection methods rely on DSBs and would not be applicable to base editors, motivate the development of new methods for the unbiased detection of base editing genome-wide. The developers of Digenome-seq, an *in vitro* genome-wide detection method ([Kim et al., 2015](#)), recently modified the protocol so it could be applied to base editors ([Kim et al., 2017a](#)). Their modified protocol took advantage of the deaminated uracil intermediate, which rarely occurs in the genome. They applied the USER enzymes (a uracil DNA glycosylase and DNA glycosylase-lyase endonuclease VIII) to cleave off the base-edited uracils on one strand while the Cas9 D10A nicks the opposite strand. This generates an enrichment of free DNA ends at the targeted site, which can be computationally identified in aligned whole-genome sequencing data. Given that this process requires uracil DNA glycosylase activity to cleave the DNA, BE3 was modified so that it no longer included the uracil glycosylase inhibitor. Across seven different guide RNAs, they found C-to-U conversions at only 18 ± 9 sites in the human genome. These off-targets were often different from those of active Cas9.

Standard whole-genome sequencing (WGS) can also be used to measure editing genome-wide in cells. WGS was applied in the smaller yeast genome and detected a slight increase in single-nucleotide variants, which could be attributed to the deaminase but not the Cas9 ([Nishida et al., 2016](#)). When WGS was applied in a F0 base-edited mouse, no off-target mutations were discovered, demonstrating the high specificity of base editing in some applications ([Kim et al., 2017b](#)). While WGS is appropriate for clonal or largely homogeneous genome mixtures, the modified Digenome-seq *in vitro* method is more sensitive to rare off-targets, detecting sites with substitution frequencies as low as 0.1% of alleles. Nonetheless, there remains a need for improved methods for unbiased off-target detection in cells and tissues with increased sensitivity to detect variants in heterogeneous populations.

MINIMIZING OFF-TARGET EDITING

Optimizing sgRNA design can also lower the frequency of off-targets at mismatched sites. While there are a number of active nuclease Cas9 sgRNA design tools that can predict off-targets, most do not explicitly consider the targetable windows of base editors as a design constraint. At least one tool (<https://www.benchling.com>) allows users to rank base-editing sgRNAs by their predicted off target effects, but there remains a need for data-driven models of base-editing specificity. It is important to consider not only mismatched sites, but also sites with small indels or non-canonical PAMs ([Leenay et al., 2016](#)). Ideally, sgRNAs should also be designed to be specific in the actual genome under experimentation and not merely in the reference genome ([Yang et al., 2014](#)). Beyond careful sgRNA selection, specificity may be improved by modifying the sgRNA construct. For example,

(Continued on next page)

Box 2. Continued

extending the sgRNA with two additional guanines (ggX20) maintained on-target efficiency while improving specificity in one study (Kim et al., 2017a).

Cas9 fusion proteins can also be optimized for improved specificity. The engineered rAPOBEC1 variants with restricted targeting windows decrease off-target editing at mismatched sites (Kim et al., 2017a). Alternatively, installing the BE3 fusion onto an engineered “high fidelity” HF-Cas9 (Kleinstiver et al., 2016) reduces off-target effects, though this results in somewhat reduced on-target efficiency (Rees et al., 2017). However, delivering BE3 as a purified ribonucleoprotein (RNP) resulted in even greater specificity than HF-BE3 plasmid delivery while maintaining on-target efficiency (Rees et al., 2017), which is consistent with previous demonstrations of Cas9 RNP specificity (Kim et al., 2014; Zuris et al., 2015).

Overall, off-targets can be minimized by iterating a process of: (1) selecting specific sgRNAs (Doench et al., 2016; Perez et al., 2017; Scott and Zhang, 2017), (2) employing optimized Cas9 variants and delivery methods, (3) detecting mutations with unbiased genome-wide methods, and (4) validating off-target edits with sensitive targeted sequencing.

S. pyogenes (“NGG”) or *S. aureus* (“NGRRRT”) or evolved artificially (“NGA,” VQR-Cas9; “NGAG,” EQR-Cas9; “NGCG,” VRER-Cas9; “NNNRRT,” SaKKH-Cas9), thus allowing the BE system to target 2,200 of these C:T or G:A point mutations (Kim et al., 2017c; Komor et al., 2017b). The space of targetable mutations has been further expanded by the fusion of cytidine deaminases to other programmable DNA binding proteins such as zinc fingers and TALENs, which are not restricted by PAM sites (Yang et al., 2016a).

Base-editing systems may offer several advantages over traditional HDR-based CRISPR-Cas9 genome-editing methods as a therapeutic tool. Importantly, these systems exhibit significantly lower indel rates, which decreases the chance of disrupting the function of the target gene through NHEJ (Komor et al., 2016). More recently, base-editing constructs (e.g., YEE-BE3) have been engineered to decrease the number of undesirable substitutions generated proximal to the target site by narrowing the editing window to 1 or 2 bases (Kim et al., 2017c) (Box 2). Comparisons between the frequency of off-target mutations generated using base editing versus cleavage with active Cas9 nuclease suggest that base editing may have greater specificity (Kim et al., 2017a, 2017c). To further reduce non-specific mutations, BE3 has been fused to a high-fidelity SpCas9 (HF-Cas9) (Kleinstiver et al., 2016), which has reduced affinity to off-target mismatch sites, resulting in the creation of a HF-BE3 with lower non-specific editing rates (Rees et al., 2017). However, this benefit came at the cost of lower on-target editing rates.

Additional engineering has been done to control the dynamics of base editing. The Liu group embedded self-cleaving catalytic RNA aptazymes in the sgRNA, which block the sgRNA spacer sequence until exposure to a ligand (Tang et al., 2017). This makes base editing responsive to a ligand, although the system exhibited some activity without ligand and was somewhat less efficient than standard sgRNAs. Additional means of inducing Cas9 activity, such as light (Nihongaki et al., 2015; Polstein and Gersbach, 2015), cell surface interactions (Dingal et al., 2017), and other small molecules (Gao et al., 2016; Senturk et al., 2017; Zetsche et al., 2015b), should be transferrable to base editors and enable fine control of editing dynamics.

Notably, the unbiased, sensitive, genome-wide off-target detection methods in cells (i.e., GUIDE-seq [Tsai et al., 2015]) and tissues (i.e., BLESS [Ran et al., 2015]) rely on double-stranded breaks and are not compatible with base editors in their

current form. While base editor off-targets can be characterized with *in vitro* assays (i.e., Digenome-seq [Kim et al., 2017a]), whole-genome sequencing, and targeted sequencing, unbiased methods have not been developed for detecting off-target base edits in cells and tissues. Additional improvements to specificity and off-target detection would accelerate translation of safe and effective base-editing therapeutics.

Engineering Agricultural Traits. The recent demonstrations of efficient base editing in crops (Lu and Zhu, 2017; Shimatani et al., 2017; Zong et al., 2017) (Figure 2A) provide a new method to select for and fine-tune desirable crop traits that are attributable to point mutations, such as 14 key agronomic traits in rice (Huang et al., 2010) and the flavor of tomatoes (Tieman et al., 2017). This can be further extended to other agricultural demands, including improving nutrition and production of crops (reviewed in Baltes et al., 2017). In rice, multiplex base editing of *ALS* and *FTIP1e* generated double mutants resistant to two herbicides (Shimatani et al., 2017). Interestingly, despite using similar Target-AID constructs, the rate of indels has been significantly higher in crops (Lu and Zhu, 2017; Shimatani et al., 2017) (Table 1) than in yeast or mammalian cells (Nishida et al., 2016). Nonetheless, indel rates are still lower than using traditional HDR-based CRISPR-Cas9 gene editing (Li et al., 2013; Mao et al., 2013). Base-editing tools have also been successfully used in wheat, maize (Zong et al., 2017), and *Arabidopsis* (Chen et al., 2017). Delivery of base-editing systems to plants has been performed using either transformed agrobacterium (Hooymaas and Schilperoort, 1992; Voytas, 2013) or biolistic particle bombardment (Sanford, 1988). Although delivery of DNA via agrobacterium is more controllable for single copy insertions, the particle bombardment strategy can be used to deliver the system to a wider range of organisms, including those that are not compatible with agrobacterium colonization (Nam et al., 1997). Both methods have been effective (Table 1) in a variety of plants. Given these initial successes, it is likely that base editing may be broadly useful in other varieties of crops, plants (Baltes et al., 2017), and animals (Petersen, 2017) as well.

Applications of Diversifying Editing

Evolving Protein Function and Structure. Base-editing systems have been used for protein evolution and saturation mutagenesis (Table 2 and Figure 2B) and present a number of useful features when compared to traditional methods such as error-prone PCR, recombination, and DNA synthesis (reviewed in Packer




Figure 2. Applications of Base-Editing Technologies

Base editing has been applied in two main areas: precision editing and screening applications.

(A) For precision editing, corrective or beneficial conversion of cytidines to thymidines has been demonstrated in many systems. Multiple delivery targets and methods that have been used are shown (see also Table 1). Precision editing has also been used for agricultural engineering using agrobacterium or biolistic DNA particle bombardment to deliver base editors to tomato, maize, wheat, and rice plants.

(B) Diversifying base editors have been used to evolve and study protein function and structure. In these experiments, sgRNAs tiling a region are used to generate targeted mutations, and the subsequent population of variants is screened for specific phenotypes (i.e., drug resistance/fluorescence), followed by targeted sequencing of the locus. To validate identified mutations, enriched variants are individually installed, and drug resistance or the activity of the individual clones is validated.

(C) The precise base-editing system can also be used for screening applications. Targeting the BE3 system to Arg, Gln, and Trp codons can edit each into stop codons, generating truncated gene products (CRISPR-STOP, iSTOP). This platform provides a potential alternative to Cas9 gene screens without inducing double-strand breaks.

and Liu, 2015). More recently, active Cas9 has been used together with a synthetic library of oligonucleotide HDR templates (Findlay et al., 2014; Garst et al., 2017; Ryan et al., 2014) to maximize variant sequence space by allowing complete control of all base substitutions at a specified locus. However, these techniques require laborious synthesis of donor templates for each region and carry the risk of DNA damage and indel generation by active Cas9. The ability to efficiently create diverse

populations of mutations at precise regions within endogenous genes using diversifying base editors eliminates the need to separately synthesize and deliver a donor library—which may be constrained by donor length, sequence complexity, and library size. Modifying the endogenous gene obviates the need to use artificial promoters to express a variant library, which may change the native regulation of protein function and localization. Proteins evolved in bacteria or yeast may also not

Table 2. Comparison of Diversifying Base Edits in Screening Applications.

Gene	Selection Strategy	Cell Type	Delivery	Base Editor	Number of sgRNA	Number of Significant Hits/Variants	Result	Reference
PSMB5	Bortezomib resistance	K562 cells	Lentiviral vector	CRISPR-X	143 targeting exons; 705 safe-targeting controls	11 variants	Discovery of 2 known and 9 novel cancer drug resistance mutations	Hess et al., 2016
wtGFP	Fluorescence shift; cell sorting	K562 cells	Plasmid nucleofection or lentiviral vector	CRISPR-X	4 targeting wtGFP; 4 safe-targeting controls	1 variant	Recovery of known S65T substitution that changes fluorescence spectra to EGFP	Hess et al., 2016
BCR-ABL Kinase Domain	Imatinib resistance	K562 cells	Plasmid nucleofection	TAM	16 targeting exon 6; 3 targeting AAVS1	6 variants	Discovery of 1 known and 5 novel cancer drug resistance mutations	Ma et al., 2016
HPRT, Puromycin, mCherry	6TG selection, Puromycin selection, and mCherry FACS	HEK293T cells	Lentiviral vector	BE3	3 targeting HPRT; 4 targeting Puromycin; 4 targeting mCherry; 16 controls	7 sgRNA	Demonstration of CRISPR-STOP approach in a functional screen	Kuscu et al., 2017

A summary of base-editing screens. Selection strategy, cell type, delivery method, and variant of base editor used are shown. The composition of the libraries and control libraries are described along with the number of identified variants or sgRNAs with phenotypes. A short summary of results for each screen is included.

function appropriately when transferred back into mammalian systems (Way et al., 2014; Yuen et al., 2006). At the same time, the targeted nature of the base-editing approach decreases the number of cells and reduces sequencing cost that would otherwise be needed to analyze a mutagenized genome generated with radiation or chemical mutagenesis.

AID has evolved as an efficient generator of genetic diversity. B cell lymphoma lines with misregulated AID-dependent somatic hypermutation have been used to evolve antibodies or fluorescent proteins (Arakawa et al., 2008; Lim et al., 2016; Wang et al., 2004), but the promiscuous mutagenesis in this system is relatively inefficient. In contrast, by directly recruiting AID, diversifying base-editing systems (CRISPR-X and TAM) can facilitate efficient, targeted mutagenesis of key structural or functional domains of endogenous targets through carefully designed sgRNA pools. For example, the CRISPR-X platform has been used to evolve wtGFP to EGFP using a targeted pool of sgRNAs. In this proof-of-principle study, dCas9-expressing K562 cells were electroporated with MS2-AID*Δ and sgRNAs targeting wtGFP near known spectrum-shifting mutation sites; cells with shifted emission spectra were sorted by FACS, and the GFP locus was sequenced. The identified residue change (S65T) matched the previously known mutation to confer the wtGFP to EGFP spectral shift (Heim and Tsien, 1996). The broad editing window of CRISPR-X and its ability to generate diverse base transitions and transversions are well-suited for the directed evolution of proteins (Hess et al., 2016).

In addition to the evolution of protein function, base-editing systems can also be used to map key structural features required for protein-protein and protein-drug interactions. For example, base editing across full protein coding regions performed using the TAM and CRISPR-X platforms has enabled the identification of known and novel drug-resistant gene variants, such as mutations in *PSMB5* that confer bortezomib resistance (Hess et al., 2016) and mutations in the *BCR-ABL* kinase domain that confer imatinib resistance (Ma et al., 2016). Identifying variant-drug interactions can improve molecular diagnosis to inform clinical treatment, lend insight toward a mechanistic understanding of drug resistance, and facilitate drug design. Other groups have successfully used active Cas9 to mutagenize protein-coding regions, wherein error-prone NHEJ mechanisms create gene variants that confer novel phenotypes such as drug resistance (Donovan et al., 2017; Ipsaro et al., 2017). While using active Cas9 for mutagenesis is straightforward, a large portion of NHEJ-generated mutations are frameshifts and multiple base indels, which may be either highly useful or undesirable depending on the application (Shalem et al., 2014; van Overbeek et al., 2016; Wang et al., 2014). Therefore, this approach may require a larger starting cell population as well as a strong selectable phenotype to enrich for single point mutants. Thus, diversifying base editors present a useful alternative for broad mutagenesis screens by minimizing indels and creating more varied point mutations.

Screening Gene Knockouts with CRISPR-STOP and iSTOP. Precision base editing can also be used to screen the effects of gene knockouts. The CRISPR-STOP and iSTOP systems use BE3 to efficiently generate early stop codons at arginine, glutamine, and tryptophan residues (Figure 2C), effectively

causing gene inactivation (Billon et al., 2017; Kuscu et al., 2017). In addition to transcriptional repression using CRISPRi (Gilbert et al., 2013, 2014), CRISPR-STOP and iSTOP may be an attractive alternative approach for disrupting the function of genes present at a high copy number, where active Cas9 nuclease can sometimes create heterogeneous repair outcomes and trigger excessive DNA damage and cell death (Aguirre et al., 2016; Morgens et al., 2017; Munoz et al., 2016; Wang et al., 2015). In contrast with CRISPRi screens, CRISPR-STOP and iSTOP generate gene knockouts via early truncation at targeted loci rather than transcriptional knockdown, with potentially different phenotypic consequences. While genome-wide screens with CRISPR-STOP and iSTOP have not yet been conducted, these tools may have advantages where a defined protein truncation product is desired and it is necessary to avoid DSBs.

Section 3: Future Directions and Areas for Improvement

While base editing has already been successful in initial studies, several technological improvements would enable additional applications: increased editing specificity, decreased PAM site dependencies, and an expansion of control over the types and frequencies of base transitions and transversions. Nevertheless, current tools enable significant extension of current applications and exploration of new ground. In this final section, we briefly discuss how base-editing systems might be improved, current limitations for their delivery *in vivo*, and future possibilities for base editing in therapy and engineering.

Future Improvements to Base-Editing Tools

Additional engineering to the enzymes recruited by dCas9 may increase the versatility of base-editing technology. Specifically, engineering the deaminase or DNA repair machinery may make it possible to direct different specific conversion events (e.g., convert C>A instead of C>T). It should also be noted that the deaminases comprise a large family of diverse enzymes. In addition to the APOBEC family and AID, these include adenosine (Alseth et al., 2014; Bass, 2002) and guanine deaminases (Roy and Roy, 1967; Yuan et al., 1999). For example, ADA is an adenosine deaminase involved in the purine salvage process that converts adenosine to inosine; loss of this enzyme causes autosomal recessive severe combined immunodeficiency (SCID) (Hershfield, 1998; Hirschhorn, 1990). Recruitment of adenosine deaminases would expand the range of targetable bases to allow conversion of adenine (and thymidine as its reverse complement) to alternative bases. Furthermore, ADAR enzymes (Li and Church, 2013; Nishikura, 2010), adenosine deaminases able to act on RNA, open up the potential for post-transcriptional editing of mRNA (Hanswillemenke et al., 2015; McMahon et al., 2016; Montiel-Gonzalez et al., 2013; Schneider et al., 2014) using a modified Cas9 system capable of targeting RNA (Nelles et al., 2016) or the C2c2 CRISPR effector (Abudayyeh et al., 2016; East-Seletsky et al., 2016), which can be targeted to RNA.

New base-editing Cas9 variants (alternate PAM sites [Kim et al., 2017c], Cpf1 [Zetsche et al., 2015a], high-fidelity variants [Rees et al., 2017]) may each have different susceptibilities to off-target effects and will need to be carefully characterized. Although significant work has been done to quantify the on-target and off-target effects for active Cas9 (reviewed in Hsu et al., 2014; Tsai and Joung, 2016; Tycko et al., 2016), similar

methods for base editing are only emerging (Box 2). These new methods will be critical to benchmark improvements in base-editing platforms.

Therapeutic Delivery of CRISPR Base Editors

Therapeutic base editing will depend on efficient delivery. RNP lipofection (Rees et al., 2017), mRNA microinjection (Kim et al., 2017b), and administration of adenoviral vectors (Chadwick et al., 2017) have been used for making therapeutic precision edits (Table 1). A comparison of the plasmid and RNP delivery methods found that RNPs had higher specificity (Rees et al., 2017). Another study that compared mRNA injections and RNPs found that mutant ratios among blastocysts and transferred embryos that gave rise to offspring were both efficient (Kim et al., 2017b). Given that the choice of delivery method may influence editing efficiency and specificity, it will be important to systematically and directly compare delivery methods for each application. Notably, adenovirus has been superseded by adeno-associated virus (AAV) in *in vivo* gene therapy pipelines because of AAV's favorable efficiency and immunogenicity profile (Valdmanis and Kay, 2017). However, base editors are large fusion proteins that exceed the packaging capacity of AAV. Further efforts to reduce the size (Friedland et al., 2015; Zetsche et al., 2015b) and limit immunogenicity of base-editing components or vehicles (Abudayyeh et al., 2016; Bessis et al., 2004; Chew et al., 2016; Hirsch et al., 2016; Wang et al., 2017; Yin et al., 2016; Zuris et al., 2015) may also improve their clinical utility.

Identifying and Characterizing Pathogenic Variants

Base editing can efficiently install mutations and generate variant libraries to screen for specific phenotypes; combining these two current applications may facilitate high-throughput characterization of known and putatively pathogenic SNPs/SNVs. While the majority of known disease-associated gene variants are point mutations (Landrum et al., 2016), there is often insufficient statistical power from human subject studies to establish a causal relationship between specific variants and their putative phenotypes (Strande et al., 2017). In cases where base editors can target and correct the mutations, such edits can rescue the wild-type phenotype *in vitro* or in animal models, validating the mutation. Recently, active Cas9 has also been used to validate possible causal variants affecting *MYB* expression through saturation mutagenesis (Canver et al., 2017). In addition to characterizing variants, base editors with broad, diversifying properties can be used to identify known and novel variants that affect measurable phenotypes (e.g., alternative splicing behaviors or gene expression levels). The information generated from these approaches could be used to support a causal variant-phenotype relationship or improve our understanding of disease mechanisms and identify key functional residues.

Investigating the Non-coding Genome

Base-editing systems may also be well-suited to investigate sequence-function relationships in enhancers and other non-coding regions. Previously, active Cas9 and CRISPRi have been used to tile enhancer regions to identify important regulatory elements that control gene expression (Canver et al., 2015, 2017; Diao et al., 2017; Fulco et al., 2016; Gasperini et al., 2017; Korkmaz et al., 2016; Sanjana et al., 2016; Xie et al., 2017). Precise base-editing systems, and those with

broader editing windows like CRISPR-X, may be similarly applied to interrogate regulatory elements through saturation mutagenesis. While base editors used in such a way may identify the sequence determinants of a regulatory element's function at single-nucleotide resolution, the narrower functional and mutagenic windows may practically constrain the amount of sequence space that can be screened. Mutations generated by current base editors may not necessarily disrupt regulatory function, whereas active Cas9 and CRISPRi may be more readily able to inhibit activity—although repressing enhancers and other regulatory elements with CRISPRi may also vary in efficiency depending on the chosen repressor and surrounding epigenetic signature.

Mapping Protein-Protein and Protein-Drug Interactions

The CRISPR-X and TAM base-editing systems have been used to identify mutations that confer drug resistance. Both systems may be extended to map and characterize protein-protein and protein-ligand interactions to determine biological function. As with the identification of bortezomib- and imatinib-resistant mutants, an efficient interaction screen would likely require a selectable phenotype amenable to high-throughput screening, such as cell growth and survival, reporter expression, or two-component reporter systems such as the kinase substrate sensor (KISS) (Lievens et al., 2014).

One of the benefits of Cas9-based technologies is the straightforward path to simultaneously target multiple sites in the genome. Indeed, both Target-AID and CRISPR-X systems have been used to mutagenize two loci within a cell (Hess et al., 2016; Nishida et al., 2016). This capacity opens up a number of possibilities for future applications. Because the repair mechanism for base editing does not involve DSBs—a requirement for HDR-mediated editing—multiple edits can be made without risk of translocations or paired deletions between the cut sites. Simultaneous mutagenesis of two sites should make it possible to study protein-protein interactions through co-evolution.

Base Editing for Directed Evolution and Synthetic Biology

While base editing can be used to investigate molecular interactions, it can also be used for the directed evolution of improved biomolecules. As an initial proof-of-principle, wild-type GFP was evolved to EGFP using CRISPR-X (Hess et al., 2016). This strategy can be used in both industrial engineering and synthetic biology applications (Cobb et al., 2013). For example, metabolic engineering (Jakociunas et al., 2017) of microbes or yeast can be accomplished by generating focused libraries of mutants *in situ* to refine the function of a desired enzyme or pathway. The base-editing diversification strategies provide a straightforward means of generating new libraries without requiring the library to be removed from the organism, randomized through gene shuffling or error-prone PCR, and replaced. The endeavor to engineer promoters, sensors, or other regulatory elements and functional circuits may also benefit from base-editing tools, which can mutate these components within mammalian cells in their native genomic context (Lienert et al., 2014; Mathur et al., 2017).

Conclusion

In summary, base editing presents a useful alternative to HDR-mediated genome editing with active Cas9 nuclease, avoiding

DSBs and the resulting indels. Capable of generating either precise C>T edits or targeted diversification of a genomic region, base-editing systems have been applied to therapeutic correction of deleterious mutations, engineering of agricultural crops, and the study and evolution of protein structure and function. As with any emerging technology, base-editing systems will require additional fine tuning and characterization, especially for therapeutic applications of precise base editing. Nonetheless, base editing is an exciting new addition to the CRISPR toolbox, and its potential in genome engineering and biology is just beginning to be realized.

ACKNOWLEDGMENTS

We thank Laure Frésard, Stephen Montgomery, Karlene Cimprich, Billy Li, Andy Fire, members of the Boyd lab, and members of the Bassik lab for helpful discussions. This work was funded by the NIH Director's New Innovator Award Program (project no. 1DP2HD084069-01), NIH T32HG000044 (G.T.H., J.T., and D.Y.), and a National Science Foundation (NSF) Graduate Research Fellowship (grant DGE-1656518) to J.T. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the NSF. Stanford University has filed a patent application based on the CRISPR-X system (U.S. provisional patent application no. 62/376,681).

REFERENCES

- Abudayyeh, O.O., Gootenberg, J.S., Konermann, S., Joung, J., Slaymaker, I.M., Cox, D.B., Shmakov, S., Makarova, K.S., Semenova, E., Minakhin, L., et al. (2016). C2c2 is a single-component programmable RNA-guided RNA-targeting CRISPR effector. *Science* 353, aaf5573.
- Aguirre, A.J., Meyers, R.M., Weir, B.A., Vazquez, F., Zhang, C.Z., Ben-David, U., Cook, A., Ha, G., Harrington, W.F., Doshi, M.B., et al. (2016). Genomic Copy Number Dictates a Gene-Independent Cell Response to CRISPR/Cas9 Targeting. *Cancer Discov.* 6, 914–929.
- Alseth, I., Dalhus, B., and Bjørås, M. (2014). Inosine in DNA and RNA. *Curr. Opin. Genet. Dev.* 26, 116–123.
- Arakawa, H., Kudo, H., Batrak, V., Caldwell, R.B., Rieger, M.A., Ellwart, J.W., and Buerstedde, J.M. (2008). Protein evolution by hypermutation and selection in the B cell line DT40. *Nucleic Acids Res.* 36, e1.
- Arnaout, R., Lee, W., Cahill, P., Honan, T., Sparrow, T., Weiland, M., Nusbaum, C., Rajewsky, K., and Koralov, S.B. (2011). High-resolution description of antibody heavy-chain repertoires in humans. *PLoS ONE* 6, e22365.
- Bak, R.O., and Porteus, M.H. (2017). CRISPR-Mediated Integration of Large Gene Cassettes Using AAV Donor Vectors. *Cell Rep.* 20, 750–756.
- Baltes, N.J., Gil-Humanes, J., and Voytas, D.F. (2017). Genome Engineering and Agriculture: Opportunities and Challenges. *Prog. Mol. Biol. Transl. Sci.* 149, 1–26.
- Barrangou, R. (2013). CRISPR-Cas systems and RNA-guided interference. *Wiley Interdiscip. Rev. RNA* 4, 267–278.
- Bass, B.L. (2002). RNA editing by adenosine deaminases that act on RNA. *Annu. Rev. Biochem.* 71, 817–846.
- Benichou, J., Ben-Hamo, R., Louzoun, Y., and Efroni, S. (2012). Rep-Seq: uncovering the immunological repertoire through next-generation sequencing. *Immunology* 135, 183–191.
- Bessis, N., GarciaCozar, F.J., and Boissier, M.C. (2004). Immune responses to gene therapy vectors: influence on vector function and effector mechanisms. *Gene Ther.* 11 (Suppl 1), S10–S17.
- Billon, P., Bryant, E.E., Joseph, S.A., Nambiar, T.S., Hayward, S.B., Rothstein, R., and Ciccia, A. (2017). CRISPR-Mediated Base Editing Enables Efficient Disruption of Eukaryotic Genes through Induction of STOP Codons. *Mol. Cell*, 1068–1079.

- Brouns, S.J., Jore, M.M., Lundgren, M., Westra, E.R., Slijkhuis, R.J., Snijders, A.P., Dickman, M.J., Makarova, K.S., Koonin, E.V., and van der Oost, J. (2008). Small CRISPR RNAs guide antiviral defense in prokaryotes. *Science* 321, 960–964.
- Canver, M.C., Smith, E.C., Sher, F., Pinello, L., Sanjana, N.E., Shalem, O., Chen, D.D., Schupp, P.G., Vinjamur, D.S., Garcia, S.P., et al. (2015). BCL11A enhancer dissection by Cas9-mediated in situ saturating mutagenesis. *Nature* 527, 192–197.
- Canver, M.C., Lessard, S., Pinello, L., Wu, Y., Ilboudo, Y., Stern, E.N., Needleman, A.J., Galactéros, F., Brugnara, C., Kutlar, A., et al. (2017). Variant-aware saturating mutagenesis using multiple Cas9 nucleases identifies regulatory elements at trait-associated loci. *Nat. Genet.* 49, 625–634.
- Certo, M.T., Ryu, B.Y., Annis, J.E., Garibov, M., Jarjour, J., Rawlings, D.J., and Scharenberg, A.M. (2011). Tracking genome engineering outcome at individual DNA breakpoints. *Nat. Methods* 8, 671–676.
- Chadwick, A.C., Wang, X., and Musunuru, K. (2017). In Vivo Base Editing of PCSK9 (Proprotein Convertase Subtilisin/Kexin Type 9) as a Therapeutic Alternative to Genome Editing. *Arterioscler. Thromb. Vasc. Biol.* 37, 1741–1747.
- Chan, L. (1992). Apolipoprotein B, the major protein component of triglyceride-rich and low density lipoproteins. *J. Biol. Chem.* 267, 25621–25624.
- Charton, K., Suel, L., Henriques, S.F., Moussu, J.P., Bovolenta, M., Taille-pierre, M., Becker, C., Lipson, K., and Richard, I. (2016). Exploiting the CRISPR/Cas9 system to study alternative splicing in vivo: application to titin. *Hum. Mol. Genet.* 25, 4518–4532.
- Chaudhuri, J., Tian, M., Khuong, C., Chua, K., Pinaud, E., and Alt, F.W. (2003). Transcription-targeted DNA deamination by the AID antibody diversification enzyme. *Nature* 422, 726–730.
- Chen, Y., Wang, Z., Ni, H., Xu, Y., Chen, Q., and Jiang, L. (2017). CRISPR/Cas9-mediated base-editing system efficiently generates gain-of-function mutations in Arabidopsis. *Sci. China Life Sci.* 60, 520–523.
- Chew, W.L., Tabebordbar, M., Cheng, J.K., Mali, P., Wu, E.Y., Ng, A.H., Zhu, K., Wagers, A.J., and Church, G.M. (2016). A multifunctional AAV-CRISPR-Cas9 and its host response. *Nat. Methods* 13, 868–874.
- Chiu, Y.L., and Greene, W.C. (2006). Multifaceted antiviral actions of APOBEC3 cytidine deaminases. *Trends Immunol.* 27, 291–297.
- Cobb, R.E., Sun, N., and Zhao, H. (2013). Directed evolution as a powerful synthetic biology tool. *Methods* 60, 81–90.
- Cong, L., Ran, F.A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P.D., Wu, X., Jiang, W., Marraffini, L.A., and Zhang, F. (2013). Multiplex genome engineering using CRISPR/Cas systems. *Science* 339, 819–823.
- d'Adda di Fagnagna, F., Weller, G.R., Doherty, A.J., and Jackson, S.P. (2003). The Gam protein of bacteriophage Mu is an orthologue of eukaryotic Ku. *EMBO Rep.* 4, 47–52.
- Deltcheva, E., Chylinski, K., Sharma, C.M., Gonzales, K., Chao, Y., Pirzada, Z.A., Eckert, M.R., Vogel, J., and Charpentier, E. (2011). CRISPR RNA maturation by trans-encoded small RNA and host factor RNase III. *Nature* 471, 602–607.
- Dever, D.P., Bak, R.O., Reinisch, A., Camarena, J., Washington, G., Nicolas, C.E., Pavel-Dinu, M., Saxena, N., Wilkens, A.B., Mantri, S., et al. (2016). CRISPR/Cas9 β -globin gene targeting in human haematopoietic stem cells. *Nature* 539, 384–389.
- DeWitt, M.A., Magis, W., Bray, N.L., Wang, T., Berman, J.R., Urbinati, F., Heo, S.J., Mitros, T., Muñoz, D.P., Boffelli, D., et al. (2016). Selection-free genome editing of the sickle mutation in human adult hematopoietic stem/progenitor cells. *Sci. Transl. Med.* 8, 360ra134.
- Diao, Y., Fang, R., Li, B., Meng, Z., Yu, J., Qiu, Y., Lin, K.C., Huang, H., Liu, T., Marina, R.J., et al. (2017). A tiling-deletion-based genetic screen for cis-regulatory element identification in mammalian cells. *Nat. Methods* 14, 629–635.
- Ding, Q., Strong, A., Patel, K.M., Ng, S.L., Gosis, B.S., Regan, S.N., Cowan, C.A., Rader, D.J., and Musunuru, K. (2014). Permanent alteration of PCSK9 with in vivo CRISPR-Cas9 genome editing. *Circ. Res.* 115, 488–492.
- Dingal, P.D.P., Kipniss, N.H., Labanieh, L., Gao, Y., and Qi, L.S. (2017). Engineering Cell Sensing and Responses Using a GPCR-Coupled CRISPR-Cas System. *bioRxiv*. <http://dx.doi.org/10.1101/152496>.
- Doench, J.G., Fusi, N., Sullender, M., Hegde, M., Vaimberg, E.W., Donovan, K.F., Smith, I., Tothova, Z., Wilen, C., Orchard, R., et al. (2016). Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. *Nat. Biotechnol.* 34, 184–191.
- Donovan, K.F., Hegde, M., Sullender, M., Vaimberg, E.W., Johannessen, C.M., Root, D.E., and Doench, J.G. (2017). Creation of Novel Protein Variants with CRISPR/Cas9-Mediated Mutagenesis: Turning a Screening By-Product into a Discovery Tool. *PLoS ONE* 12, e0170445.
- Doudna, J.A., and Charpentier, E. (2014). Genome editing. The new frontier of genome engineering with CRISPR-Cas9. *Science* 346, 1258096.
- East-Seletsky, A., O'Connell, M.R., Knight, S.C., Burstein, D., Cate, J.H., Tjian, R., and Doudna, J.A. (2016). Two distinct RNase activities of CRISPR-C2c2 enable guide-RNA processing and RNA detection. *Nature* 538, 270–273.
- Elhanati, Y., Sethna, Z., Marcou, Q., Callan, C.G., Jr., Mora, T., and Walczak, A.M. (2015). Inferring processes underlying B-cell repertoire diversity. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 370, 370.
- Ernst, J., Melnikov, A., Zhang, X., Wang, L., Rogov, P., Mikkelsen, T.S., and Kellis, M. (2016). Genome-scale high-resolution mapping of activating and repressive nucleotides in regulatory regions. *Nat. Biotechnol.* 34, 1180–1190.
- Esnault, C., Heidmann, O., Delebecque, F., Dewannieux, M., Ribet, D., Hance, A.J., Heidmann, T., and Schwartz, O. (2005). APOBEC3G cytidine deaminase inhibits retrotransposition of endogenous retroviruses. *Nature* 433, 430–433.
- Findlay, G.M., Boyle, E.A., Hause, R.J., Klein, J.C., and Shendure, J. (2014). Saturation editing of genomic regions by multiplex homology-directed repair. *Nature* 513, 120–123.
- Friedland, A.E., Baral, R., Singhal, P., Loveluck, K., Shen, S., Sanchez, M., Marco, E., Gotta, G.M., Maeder, M.L., Kennedy, E.M., et al. (2015). Characterization of Staphylococcus aureus Cas9: a smaller Cas9 for all-in-one adeno-associated virus delivery and paired nickase applications. *Genome Biol.* 16, 257.
- Fulco, C.P., Munschauer, M., Anyoha, R., Munson, G., Grossman, S.R., Perez, E.M., Kane, M., Cleary, B., Lander, E.S., and Engreitz, J.M. (2016). Systematic mapping of functional enhancer-promoter connections with CRISPR interference. *Science* 354, 769–773.
- Gao, Y., Xiong, X., Wong, S., Charles, E.J., Lim, W.A., and Qi, L.S. (2016). Complex transcriptional modulation with orthogonal and inducible dCas9 regulators. *Nat. Methods* 13, 1043–1049.
- Garneau, J.E., Dupuis, M.-E., Villion, M., Romero, D.A., Barrangou, R., Boyaval, P., Fremaux, C., Horvath, P., Magadán, A.H., and Moineau, S. (2010). The CRISPR/Cas bacterial immune system cleaves bacteriophage and plasmid DNA. *Nature* 468, 67–71.
- Garst, A.D., Bassalo, M.C., Pines, G., Lynch, S.A., Halweg-Edwards, A.L., Liu, R., Liang, L., Wang, Z., Zeitoun, R., Alexander, W.G., and Gill, R.T. (2017). Genome-wide mapping of mutations at single-nucleotide resolution for protein, metabolic and genome engineering. *Nat. Biotechnol.* 35, 48–55.
- Gasiunas, G., Barrangou, R., Horvath, P., and Siksnys, V. (2012). Cas9-crRNA ribonucleoprotein complex mediates specific DNA cleavage for adaptive immunity in bacteria. *Proc. Natl. Acad. Sci. USA* 109, E2579–E2586.
- Gasperini, M., Findlay, G.M., McKenna, A., Milbank, J.H., Lee, C., Zhang, M.D., Cusanovich, D.A., and Shendure, J. (2017). CRISPR/Cas9-Mediated Scanning for Regulatory Elements Required for HPRT1 Expression via Thousands of Large, Programmed Genomic Deletions. *Am. J. Hum. Genet.* 101, 192–205.
- Gilbert, L.A., Larson, M.H., Morsut, L., Liu, Z., Brar, G.A., Torres, S.E., Stern-Ginossar, N., Brandman, O., Whitehead, E.H., Doudna, J.A., et al. (2013). CRISPR-mediated modular RNA-guided regulation of transcription in eukaryotes. *Cell* 154, 442–451.
- Gilbert, L.A., Horlbeck, M.A., Adamson, B., Villalta, J.E., Chen, Y., Whitehead, E.H., Guimaraes, C., Panning, B., Ploegh, H.L., Bassik, M.C., et al. (2014). Genome-Scale CRISPR-Mediated Control of Gene Repression and Activation. *Cell* 159, 647–661.

- Greiff, V., Bhat, P., Cook, S.C., Menzel, U., Kang, W., and Reddy, S.T. (2015). A bioinformatic framework for immune repertoire diversity profiling enables detection of immunological status. *Genome Med.* 7, 49.
- Hamilton, C.E., Papavasiliou, F.N., and Rosenberg, B.R. (2010). Diverse functions for DNA and RNA editing in the immune system. *RNA Biol.* 7, 220–228.
- Hanswillemenke, A., Kuzdere, T., Vogel, P., Jékely, G., and Stafforst, T. (2015). Site-Directed RNA Editing in Vivo Can Be Triggered by the Light-Driven Assembly of an Artificial Riboprotein. *J. Am. Chem. Soc.* 137, 15875–15881.
- Harris, R.S., Bishop, K.N., Sheehy, A.M., Craig, H.M., Petersen-Mahrt, S.K., Watt, I.N., Neuberger, M.S., and Malim, M.H. (2003). DNA deamination mediates innate immunity to retroviral infection. *Cell* 113, 803–809.
- Heim, R., and Tsien, R.Y. (1996). Engineering green fluorescent protein for improved brightness, longer wavelengths and fluorescence resonance energy transfer. *Curr. Biol.* 6, 178–182.
- Heller, R.C., and Marians, K.J. (2006). Replisome assembly and the direct restart of stalled replication forks. *Nat. Rev. Mol. Cell Biol.* 7, 932–943.
- Hershfield, M.S. (1998). Adenosine deaminase deficiency: clinical expression, molecular basis, and therapy. *Semin. Hematol.* 35, 291–298.
- Hess, G.T., Frésard, L., Han, K., Lee, C.H., Li, A., Cimprich, K.A., Montgomery, S.B., and Bassik, M.C. (2016). Directed evolution using dCas9-targeted somatic hypermutation in mammalian cells. *Nat. Methods* 13, 1036–1042.
- Hirsch, M.L., Wolf, S.J., and Samulski, R.J. (2016). Delivering Transgenic DNA Exceeding the Carrying Capacity of AAV Vectors. *Methods Mol. Biol.* 1382, 21–39.
- Hirschhorn, R. (1990). Adenosine deaminase deficiency. *Immunodef. Rev.* 2, 175–198.
- Hoehn, K.B., Fowler, A., Lunter, G., and Pybus, O.G. (2016). The Diversity and Molecular Evolution of B-Cell Receptors during Infection. *Mol. Biol. Evol.* 33, 1147–1157.
- Hooykaas, P.J., and Schilperoort, R.A. (1992). Agrobacterium and plant genetic engineering. *Plant Mol. Biol.* 19, 15–38.
- Hsieh, C.L., Liu, H., Huang, Y., Kang, L., Chen, H.W., Chen, Y.T., Wee, Y.R., Chen, S.J., and Tan, B.C. (2014). ADAR1 deaminase contributes to scheduled skeletal myogenesis progression via stage-specific functions. *Cell Death Differ.* 21, 707–719.
- Hsu, P.D., Lander, E.S., and Zhang, F. (2014). Development and applications of CRISPR-Cas9 for genome engineering. *Cell* 157, 1262–1278.
- Huang, X., Wei, X., Sang, T., Zhao, Q., Feng, Q., Zhao, Y., Li, C., Zhu, C., Lu, T., Zhang, Z., et al. (2010). Genome-wide association studies of 14 agronomic traits in rice landraces. *Nat. Genet.* 42, 961–967.
- Ipsaro, J.J., Shen, C., Arai, E., Xu, Y., Kinney, J.B., Joshua-Tor, L., Vakoc, C.R., and Shi, J. (2017). Rapid generation of drug-resistance alleles at endogenous loci using CRISPR-Cas9 indel mutagenesis. *PLoS ONE* 12, e0172177.
- Ishino, M., Sawada, Y., Yaegashi, T., Demura, M., and Fujinaga, K. (1987). Nucleotide sequence of the adenovirus type 40 inverted terminal repeat: close relation to that of adenovirus type 5. *Virology* 156, 414–416.
- Jakobciunas, T., Jensen, M.K., and Keasling, J.D. (2017). System-level perturbations of cell metabolism using CRISPR/Cas9. *Curr. Opin. Biotechnol.* 46, 134–140.
- Jansen, R., Embden, J.D., Gaastera, W., and Schouls, L.M. (2002). Identification of genes that are associated with DNA repeats in prokaryotes. *Mol. Microbiol.* 43, 1565–1575.
- jasin, M., and Haber, J.E. (2016). The democratization of gene editing: Insights from site-specific cleavage and double-strand break repair. *DNA Repair (Amst.)* 44, 6–16.
- Jiang, F., and Doudna, J.A. (2017). CRISPR-Cas9 Structures and Mechanisms. *Annu. Rev. Biophys.* 46, 505–529.
- Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J.A., and Charpentier, E. (2012). A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* 337, 816–821.
- Jinek, M., East, A., Cheng, A., Lin, S., Ma, E., and Doudna, J. (2013). RNA-programmed genome editing in human cells. *eLife* 2, e00471.
- Kapahnke, M., Banning, A., and Tikkanen, R. (2016). Random Splicing of Several Exons Caused by a Single Base Change in the Target Exon of CRISPR/Cas9 Mediated Gene Knockout. *Cells* 5, E45.
- Keegan, L.P., Gallo, A., and O'Connell, M.A. (2001). The many roles of an RNA. *Nat. Rev. Genet.* 2, 869–878.
- Kim, S., Kim, D., Cho, S.W., Kim, J., and Kim, J.S. (2014). Highly efficient RNA-guided genome editing in human cells via delivery of purified Cas9 ribonucleoproteins. *Genome Res.* 24, 1012–1019.
- Kim, D., Bae, S., Park, J., Kim, E., Kim, S., Yu, H.R., Hwang, J., Kim, J.I., and Kim, J.S. (2015). Digenome-seq: genome-wide profiling of CRISPR-Cas9 off-target effects in human cells. *Nat. Methods* 12, 237–243.
- Kim, D., Lim, K., Kim, S.T., Yoon, S.H., Kim, K., Ryu, S.M., and Kim, J.S. (2017a). Genome-wide target specificities of CRISPR RNA-guided programmable deaminases. *Nat. Biotechnol.* 35, 475–480.
- Kim, K., Ryu, S.M., Kim, S.T., Baek, G., Kim, D., Lim, K., Chung, E., Kim, S., and Kim, J.S. (2017b). Highly efficient RNA-guided base editing in mouse embryos. *Nat. Biotechnol.* 35, 435–437.
- Kim, Y.B., Komor, A.C., Levy, J.M., Packer, M.S., Zhao, K.T., and Liu, D.R. (2017c). Increasing the genome-targeting scope and precision of base editing with engineered Cas9-cytidine deaminase fusions. *Nat. Biotechnol.* 35, 371–376.
- Kleistiver, B.P., Prew, M.S., Tsai, S.Q., Nguyen, N.T., Topkar, V.V., Zheng, Z., and Joung, J.K. (2015a). Broadening the targeting range of Staphylococcus aureus CRISPR-Cas9 by modifying PAM recognition. *Nat. Biotechnol.* 33, 1293–1298.
- Kleistiver, B.P., Prew, M.S., Tsai, S.Q., Topkar, V.V., Nguyen, N.T., Zheng, Z., Gonzales, A.P., Li, Z., Peterson, R.T., Yeh, J.R., et al. (2015b). Engineered CRISPR-Cas9 nucleases with altered PAM specificities. *Nature* 523, 481–485.
- Kleistiver, B.P., Pattanayak, V., Prew, M.S., Tsai, S.Q., Nguyen, N.T., Zheng, Z., and Joung, J.K. (2016). High-fidelity CRISPR-Cas9 nucleases with no detectable genome-wide off-target effects. *Nature* 529, 490–495.
- Koito, A., and Ikeda, T. (2013). Intrinsic immunity against retrotransposons by APOBEC cytidine deaminases. *Front. Microbiol.* 4, 28.
- Komor, A.C., Kim, Y.B., Packer, M.S., Zuris, J.A., and Liu, D.R. (2016). Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage. *Nature* 533, 420–424.
- Komor, A.C., Badran, A.H., and Liu, D.R. (2017a). CRISPR-Based Technologies for the Manipulation of Eukaryotic Genomes. *Cell* 169, 559.
- Komor, A.C., Zhao, K.T., Packer, M.S., Gaudelli, N.M., Waterbury, A.L., Koblan, L.W., Kim, Y.B., Badran, A.H., and Liu, D.R. (2017b). Improved base excision repair inhibition and bacteriophage Mu Gam protein yields C:G-to-T:A base editors with higher efficiency and product purity. *Sci. Adv.* 3, eaao4774.
- Konermann, S., Brigham, M.D., Trevino, A.E., Joung, J., Abudayyeh, O.O., Barcena, C., Hsu, P.D., Habib, N., Gootenberg, J.S., Nishimasu, H., et al. (2015). Genome-scale transcriptional activation by an engineered CRISPR-Cas9 complex. *Nature* 517, 583–588.
- Korkmaz, G., Lopes, R., Ugalde, A.P., Nevedomskaya, E., Han, R., Myacheva, K., Zwart, W., Elkon, R., and Agami, R. (2016). Functional genetic screens for enhancer elements in the human genome using CRISPR-Cas9. *Nat. Biotechnol.* 34, 192–198.
- Küppers, R., Klein, U., Hansmann, M.L., and Rajewsky, K. (1999). Cellular origin of human B-cell lymphomas. *N. Engl. J. Med.* 341, 1520–1529.
- Kuscu, C., Parlak, M., Tufan, T., Yang, J., Szlachta, K., Wei, X., Mammadov, R., and Adli, M. (2017). CRISPR-STOP: gene silencing through base-editing-induced nonsense mutations. *Nat. Methods* 14, 710–712.
- Lackner, D.H., Carré, A., Guzzardo, P.M., Banning, C., Mangena, R., Henley, T., Oberndorfer, S., Gapp, B.V., Nijman, S.M., Brummelkamp, T.R., and Burckstummer, T. (2015). A generic strategy for CRISPR-Cas9-mediated gene tagging. *Nat. Commun.* 6, 10237.

- Landrum, M.J., Lee, J.M., Benson, M., Brown, G., Chao, C., Chitipiralla, S., Gu, B., Hart, J., Hoffman, D., Hoover, J., et al. (2016). ClinVar: public archive of interpretations of clinically relevant variants. *Nucleic Acids Res.* **44** (D1), D862–D868.
- Larson, E.D., Cummings, W.J., Bednarski, D.W., and Maizels, N. (2005). MRE11/RAD50 cleaves DNA in the AID/UNG-dependent pathway of immunoglobulin gene diversification. *Mol. Cell* **20**, 367–375.
- Lecossier, D., Bouchonnet, F., Clavel, F., and Hance, A.J. (2003). Hypermutation of HIV-1 DNA in the absence of the Vif protein. *Science* **300**, 1112.
- Leenay, R.T., Maksimchuk, K.R., Slotkowski, R.A., Agrawal, R.N., Goma, A.A., Briner, A.E., Barrangou, R., and Beisel, C.L. (2016). Identifying and visualizing functional PAM diversity across CRISPR-Cas systems. *Mol. Cell* **62**, 137–147.
- Leonetti, M.D., Sekine, S., Kamiyama, D., Weissman, J.S., and Huang, B. (2016). A scalable strategy for high-throughput GFP tagging of endogenous human proteins. *Proc. Natl. Acad. Sci. USA* **113**, E3501–E3508.
- Li, J.B., and Church, G.M. (2013). Deciphering the functions and regulation of brain-enriched A-to-I RNA editing. *Nat. Neurosci.* **16**, 1518–1522.
- Li, J.F., Norville, J.E., Aach, J., McCormack, M., Zhang, D., Bush, J., Church, G.M., and Sheen, J. (2013). Multiplex and homologous recombination-mediated genome editing in Arabidopsis and Nicotiana benthamiana using guide RNA and Cas9. *Nat. Biotechnol.* **31**, 688–691.
- Liang, P., Sun, H., Sun, Y., Zhang, X., Xie, X., Zhang, J., Zhang, Z., Chen, Y., Ding, C., Xiong, Y., et al. (2017). Effective gene editing by high-fidelity base editor 2 in mouse zygotes. *Protein Cell* **8**, 601–611.
- Lienert, F., Lohmueller, J.J., Garg, A., and Silver, P.A. (2014). Synthetic biology in mammalian cells: next generation research tools and therapeutics. *Nat. Rev. Mol. Cell Biol.* **15**, 95–107.
- Lievens, S., Gerlo, S., Lemmens, I., De Clercq, D.J., Risseuw, M.D., Vanderroost, N., De Smet, A.S., Ruyssinck, E., Chevet, E., Van Calenberg, S., and Tavernier, J. (2014). Kinase Substrate Sensor (KISS), a mammalian in situ protein interaction sensor. *Mol. Cell. Proteomics* **13**, 3332–3342.
- Lim, A.W., Williams, G.T., Rada, C., and Sale, J.E. (2016). Directed evolution of human scFvs in DT40 cells. *Protein Eng. Des. Sel.* **29**, 39–48.
- Lu, Y., and Zhu, J.K. (2017). Precise Editing of a Target Base in the Rice Genome Using a Modified CRISPR/Cas9 System. *Mol. Plant* **10**, 523–525.
- Ma, Y., Zhang, J., Yin, W., Zhang, Z., Song, Y., and Chang, X. (2016). Targeted AID-mediated mutagenesis (TAM) enables efficient genomic diversification in mammalian cells. *Nat. Methods* **13**, 1029–1035.
- Mali, P., Yang, L., Esvelt, K.M., Aach, J., Guell, M., DiCarlo, J.E., Norville, J.E., and Church, G.M. (2013). RNA-guided human genome engineering via Cas9. *Science* **339**, 823–826.
- Mao, Y., Zhang, H., Xu, N., Zhang, B., Gou, F., and Zhu, J.K. (2013). Application of the CRISPR-Cas system for efficient genome engineering in plants. *Mol. Plant* **6**, 2008–2011.
- Marraffini, L.A., and Sontheimer, E.J. (2008). CRISPR interference limits horizontal gene transfer in staphylococci by targeting DNA. *Science* **322**, 1843–1845.
- Mathur, M., Xiang, J.S., and Smolke, C.D. (2017). Mammalian synthetic biology for studying the cell. *J. Cell Biol.* **216**, 73–82.
- McMahon, A.C., Rahman, R., Jin, H., Shen, J.L., Fieldsend, A., Luo, W., and Rosbash, M. (2016). TRIBE: Hijacking an RNA-Editing Enzyme to Identify Cell-Specific Targets of RNA-Binding Proteins. *Cell* **165**, 742–753.
- Mojica, F.J., Díez-Villaseñor, C., Soria, E., and Juez, G. (2000). Biological significance of a family of regularly spaced repeats in the genomes of Archaea, Bacteria and mitochondria. *Mol. Microbiol.* **36**, 244–246.
- Mol, C.D., Arvai, A.S., Sanderson, R.J., Slupphaug, G., Kavli, B., Krokan, H.E., Mosbaugh, D.W., and Tainer, J.A. (1995). Crystal structure of human uracil-DNA glycosylase in complex with a protein inhibitor: protein mimicry of DNA. *Cell* **82**, 701–708.
- Montiel-Gonzalez, M.F., Vallecillo-Viejo, I., Yudowski, G.A., and Rosenthal, J.J. (2013). Correction of mutations within the cystic fibrosis transmembrane conductance regulator by site-directed RNA editing. *Proc. Natl. Acad. Sci. USA* **110**, 18285–18290.
- Mora, T., Walczak, A.M., Bialek, W., and Callan, C.G., Jr. (2010). Maximum entropy models for antibody diversity. *Proc. Natl. Acad. Sci. USA* **107**, 5405–5410.
- Morgens, D.W., Wainberg, M., Boyle, E.A., Ursu, O., Araya, C.L., Tsui, C.K., Haney, M.S., Hess, G.T., Han, K., Jeng, E.E., et al. (2017). Genome-scale measurement of off-target activity using Cas9 toxicity in high-throughput screens. *Nat. Commun.* **8**, 15178.
- Mou, H., Smith, J.L., Peng, L., Yin, H., Moore, J., Zhang, X.O., Song, C.Q., Sheel, A., Wu, Q., Ozata, D.M., et al. (2017). CRISPR/Cas9-mediated genome editing induces exon skipping by alternative splicing or exon deletion. *Genome Biol.* **18**, 108.
- Muckenfuss, H., Hamdorf, M., Held, U., Perkovic, M., Löwer, J., Cichutek, K., Flory, E., Schumann, G.G., and Munk, C. (2006). APOBEC3 proteins inhibit human LINE-1 retrotransposition. *J. Biol. Chem.* **281**, 22161–22172.
- Munoz, D.M., Cassiani, P.J., Li, L., Billy, E., Korn, J.M., Jones, M.D., Golji, J., Ruddy, D.A., Yu, K., McAllister, G., et al. (2016). CRISPR Screens Provide a Comprehensive Assessment of Cancer Vulnerabilities but Generate False-Positive Hits for Highly Amplified Genomic Regions. *Cancer Discov.* **6**, 900–913.
- Nam, J., Matthysse, A.G., and Gelvin, S.B. (1997). Differences in susceptibility of Arabidopsis ecotypes to crown gall disease may result from a deficiency in T-DNA integration. *Plant Cell* **9**, 317–333.
- Navaratnam, N., and Sarwar, R. (2006). An overview of cytidine deaminases. *Int. J. Hematol.* **83**, 195–200.
- Nelles, D.A., Fang, M.Y., O'Connell, M.R., Xu, J.L., Markmiller, S.J., Doudna, J.A., and Yeo, G.W. (2016). Programmable RNA Tracking in Live Cells with CRISPR/Cas9. *Cell* **165**, 488–496.
- Nihongaki, Y., Yamamoto, S., Kawano, F., Suzuki, H., and Sato, M. (2015). CRISPR-Cas9-based photoactivatable transcription system. *Chem. Biol.* **22**, 169–174.
- Nishida, K., Arazoe, T., Yachie, N., Banno, S., Kakimoto, M., Tabata, M., Mochizuki, M., Miyabe, A., Araki, M., Hara, K.Y., et al. (2016). Targeted nucleotide editing using hybrid prokaryotic and vertebrate adaptive immune systems. *Science* **353**, aaf8729.
- Nishikura, K. (2010). Functions and regulation of RNA editing by ADAR deaminases. *Annu. Rev. Biochem.* **79**, 321–349.
- Nishimasu, H., Ran, F.A., Hsu, P.D., Konermann, S., Shehata, S.I., Dohmae, N., Ishitani, R., Zhang, F., and Nureki, O. (2014). Crystal structure of Cas9 in complex with guide RNA and target DNA. *Cell* **156**, 935–949.
- Odegard, V.H., and Schatz, D.G. (2006). Targeting of somatic hypermutation. *Nat. Rev. Immunol.* **6**, 573–583.
- Packer, M.S., and Liu, D.R. (2015). Methods for the directed evolution of proteins. *Nat. Rev. Genet.* **16**, 379–394.
- Pasqualucci, L., Bhagat, G., Jankovic, M., Compagno, M., Smith, P., Muramatsu, M., Honjo, T., Morse, H.C., 3rd, Nussenzweig, M.C., and Dalla-Favera, R. (2008). AID is required for germinal center-derived lymphomagenesis. *Nat. Genet.* **40**, 108–112.
- Perez, A.R., Pritykin, Y., Vidigal, J.A., Chhangawala, S., Zamparo, L., Leslie, C.S., and Ventura, A. (2017). GuideScan software for improved single and paired CRISPR guide RNA design. *Nat. Biotechnol.* **35**, 347–349.
- Petersen, B. (2017). Basics of genome editing technology and its application in livestock species. *Reprod. Domest. Anim.* **52** (Suppl 3), 4–13.
- Polstein, L.R., and Gersbach, C.A. (2015). A light-inducible CRISPR-Cas9 system for control of endogenous gene activation. *Nat. Chem. Biol.* **11**, 198–200.
- Powell, L.M., Wallis, S.C., Pease, R.J., Edwards, Y.H., Knott, T.J., and Scott, J. (1987). A novel form of tissue-specific RNA processing produces apolipoprotein-B48 in intestine. *Cell* **50**, 831–840.

- Qi, L.S., Larson, M.H., Gilbert, L.A., Doudna, J.A., Weissman, J.S., Arkin, A.P., and Lim, W.A. (2013). Repurposing CRISPR as an RNA-guided platform for sequence-specific control of gene expression. *Cell* 152, 1173–1183.
- Rajagopal, N., Srinivasan, S., Kooshesh, K., Guo, Y., Edwards, M.D., Banerjee, B., Syed, T., Emons, B.J., Gifford, D.K., and Sherwood, R.I. (2016). High-throughput mapping of regulatory DNA. *Nat. Biotechnol.* 34, 167–174.
- Ramiro, A.R., Jankovic, M., Callen, E., Difilippantonio, S., Chen, H.T., McBride, K.M., Eisenreich, T.R., Chen, J., Dickins, R.A., Lowe, S.W., et al. (2006). Role of genomic instability and p53 in AID-induced c-myc-Igh translocations. *Nature* 440, 105–109.
- Ran, F.A., Cong, L., Yan, W.X., Scott, D.A., Gootenberg, J.S., Kriz, A.J., Zetsche, B., Shalem, O., Wu, X., Makarova, K.S., et al. (2015). In vivo genome editing using *Staphylococcus aureus* Cas9. *Nature* 520, 186–191.
- Rees, H.A., Komor, A.C., Yeh, W.H., Caetano-Lopes, J., Warman, M., Edge, A.S.B., and Liu, D.R. (2017). Improving the DNA specificity and applicability of base editing through protein engineering and protein delivery. *Nat. Commun.* 8, 15790.
- Reuter, J.A., Spacek, D.V., and Snyder, M.P. (2015). High-throughput sequencing technologies. *Mol. Cell* 58, 586–597.
- Revy, P., Muto, T., Levy, Y., Geissmann, F., Plebani, A., Sanal, O., Catalan, N., Forveille, M., Dufourcq-Labeolouse, R., Gennery, A., et al. (2000). Activation-induced cytidine deaminase (AID) deficiency causes the autosomal recessive form of the Hyper-IgM syndrome (HIGM2). *Cell* 102, 565–575.
- Reynaud, C.A., Aoufouchi, S., Faili, A., and Weill, J.C. (2003). What role for AID: mutator, or assembler of the immunoglobulin mutasome? *Nat. Immunol.* 4, 631–638.
- Richardson, C.D., Ray, G.J., Bray, N.L., and Corn, J.E. (2016). Non-homologous DNA increases gene disruption efficiency by altering DNA repair outcomes. *Nat. Commun.* 7, 12463.
- Robertson, A.B., Klungland, A., Rognes, T., and Leiros, I. (2009). DNA repair in mammalian cells: Base excision repair: the long and short of it. *Cell. Mol. Life Sci.* 66, 981–993.
- Robins, H. (2013). Immunosequencing: applications of immune repertoire deep sequencing. *Curr. Opin. Immunol.* 25, 646–652.
- Roy, J.E., and Roy, K.L. (1967). The mechanism and specificity of guanine deaminase. *Can. J. Biochem.* 45, 1263–1269.
- Ryan, O.W., Skerker, J.M., Maurer, M.J., Li, X., Tsai, J.C., Poddar, S., Lee, M.E., DeLoache, W., Dueber, J.E., Arkin, A.P., and Cate, J.H. (2014). Selection of chromosomal DNA libraries using a multiplex CRISPR system. *eLife* 3.
- Sakoparnig, T., Fried, P., and Beerenwinkel, N. (2015). Identification of constrained cancer driver genes based on mutation timing. *PLoS Comput. Biol.* 11, e1004027.
- Samuel, C.E. (2012). ADARs: viruses and innate immunity. *Curr. Top. Microbiol. Immunol.* 353, 163–195.
- Sander, J.D., and Joung, J.K. (2014). CRISPR-Cas systems for editing, regulating and targeting genomes. *Nat. Biotechnol.* 32, 347–355.
- Sanford, J.C. (1988). The biolistic process. *Trends Biotechnol.* 6, 299–302.
- Sanjana, N.E., Wright, J., Zheng, K., Shalem, O., Fontanillas, P., Joung, J., Cheng, C., Regev, A., and Zhang, F. (2016). High-resolution interrogation of functional elements in the noncoding genome. *Science* 353, 1545–1549.
- Schneider, M.F., Wettengel, J., Hoffmann, P.C., and Stafforst, T. (2014). Optimal guideRNAs for re-directing deaminase activity of hADAR1 and hADAR2 in trans. *Nucleic Acids Res.* 42, e87.
- Scott, D.A., and Zhang, F. (2017). Implications of human genetic variation in CRISPR-based therapeutic genome editing. *Nat. Med.* 23, 1095–1101.
- Senturk, S., Shirole, N.H., Nowak, D.G., Corbo, V., Pal, D., Vaughan, A., Tuveson, D.A., Trotman, L.C., Kinney, J.B., and Sordella, R. (2017). Rapid and tunable method to temporally control gene editing based on conditional Cas9 stabilization. *Nat. Commun.* 8, 14370.
- Shalem, O., Sanjana, N.E., Hartenian, E., Shi, X., Scott, D.A., Mikkelsen, T., Heckl, D., Ebert, B.L., Root, D.E., Doench, J.G., and Zhang, F. (2014). Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science* 343, 84–87.
- Shalem, O., Sanjana, N.E., and Zhang, F. (2015). High-throughput functional genomics using CRISPR-Cas9. *Nat. Rev. Genet.* 16, 299–311.
- Sheehy, A.M., Gaddis, N.C., Choi, J.D., and Malim, M.H. (2002). Isolation of a human gene that inhibits HIV-1 infection and is suppressed by the viral Vif protein. *Nature* 418, 646–650.
- Shimatani, Z., Kashojiya, S., Takayama, M., Terada, R., Arazoe, T., Ishii, H., Teramura, H., Yamamoto, T., Komatsu, H., Miura, K., et al. (2017). Targeted base editing in rice and tomato using a CRISPR-Cas9 cytidine deaminase fusion. *Nat. Biotechnol.* 35, 441–443.
- Sternberg, S.H., and Doudna, J.A. (2015). Expanding the Biologist's Toolkit with CRISPR-Cas9. *Mol. Cell* 58, 568–574.
- Strande, N.T., Riggs, E.R., Buchanan, A.H., Ceyhan-Birsoy, O., DiStefano, M., Dwight, S.S., Goldstein, J., Ghosh, R., Seifert, B.A., Sneddon, T.P., et al. (2017). Evaluating the Clinical Validity of Gene-Disease Associations: An Evidence-Based Framework Developed by the Clinical Genome Resource. *Am. J. Hum. Genet.* 100, 895–906.
- Tang, W., Hu, J.H., and Liu, D.R. (2017). Aptazyme-embedded guide RNAs enable ligand-responsive genome editing and transcriptional activation. *Nat. Commun.* 8, 15939.
- Tieman, D., Zhu, G., Resende, M.F., Jr., Lin, T., Nguyen, C., Bies, D., Rambla, J.L., Beltran, K.S., Taylor, M., Zhang, B., et al. (2017). A chemical genetic roadmap to improved tomato flavor. *Science* 355, 391–394.
- Tsai, S.Q., and Joung, J.K. (2016). Defining and improving the genome-wide specificities of CRISPR-Cas9 nucleases. *Nat. Rev. Genet.* 17, 300–312.
- Tsai, S.Q., Zheng, Z., Nguyen, N.T., Liebers, M., Topkar, V.V., Thapar, V., Wyvekens, N., Khayter, C., Iafrate, A.J., Le, L.P., et al. (2015). GUIDE-seq enables genome-wide profiling of off-target cleavage by CRISPR-Cas nucleases. *Nat. Biotechnol.* 33, 187–197.
- Tycko, J., Myer, V.E., and Hsu, P.D. (2016). Methods for Optimizing CRISPR-Cas9 Genome Editing Specificity. *Mol. Cell* 63, 355–370.
- Valdmanis, P.N., and Kay, M.A. (2017). Future of rAAV Gene Therapy: Platform for RNAi, Gene Editing, and Beyond. *Hum. Gene Ther.* 28, 361–372.
- van Overbeek, M., Capurso, D., Carter, M.M., Thompson, M.S., Frias, E., Russ, C., Reece-Hoyes, J.S., Nye, C., Gradia, S., Vidal, B., et al. (2016). DNA Repair Profiling Reveals Nonrandom Outcomes at Cas9-Mediated Breaks. *Mol. Cell* 63, 633–646.
- Vieira, V.C., and Soares, M.A. (2013). The role of cytidine deaminases on innate immune responses against human viral infections. *BioMed Res. Int.* 2013, 683095.
- Voytas, D.F. (2013). Plant genome engineering with sequence-specific nucleases. *Annu. Rev. Plant Biol.* 64, 327–350.
- Wang, L., Jackson, W.C., Steinbach, P.A., and Tsien, R.Y. (2004). Evolution of new nonantibody proteins via iterative somatic hypermutation. *Proc. Natl. Acad. Sci. USA* 101, 16745–16749.
- Wang, M., Yang, Z., Rada, C., and Neuberger, M.S. (2009). AID upmutants isolated using a high-throughput screen highlight the immunity/cancer balance limiting DNA deaminase activity. *Nat. Struct. Mol. Biol.* 16, 769–776.
- Wang, C., Huang, R., He, B., and Du, Q. (2012). Improving the thermostability of alpha-amylase by combinatorial coevolving-site saturation mutagenesis. *BMC Bioinformatics* 13, 263.
- Wang, T., Wei, J.J., Sabatini, D.M., and Lander, E.S. (2014). Genetic screens in human cells using the CRISPR-Cas9 system. *Science* 343, 80–84.
- Wang, T., Birsoy, K., Hughes, N.W., Krupczak, K.M., Post, Y., Wei, J.J., Lander, E.S., and Sabatini, D.M. (2015). Identification and characterization of essential genes in the human genome. *Science* 350, 1096–1101.
- Wang, H., La Russa, M., and Qi, L.S. (2016). CRISPR/Cas9 in Genome Editing and Beyond. *Annu. Rev. Biochem.* 85, 227–264.
- Wang, H.X., Li, M., Lee, C.M., Chakraborty, S., Kim, H.W., Bao, G., and Leong, K.W. (2017). CRISPR/Cas9-Based Genome Editing for Disease Modeling and

Therapy: Challenges and Opportunities for Nonviral Delivery. *Chem. Rev.* 117, 9874–9906.

Way, J.C., Collins, J.J., Keasling, J.D., and Silver, P.A. (2014). Integrating biological redesign: where synthetic biology came from and where it needs to go. *Cell* 157, 151–161.

Wu, X., Scott, D.A., Kriz, A.J., Chiu, A.C., Hsu, P.D., Dadon, D.B., Cheng, A.W., Trevino, A.E., Konermann, S., Chen, S., et al. (2014). Genome-wide binding of the CRISPR endonuclease Cas9 in mammalian cells. *Nat. Biotechnol.* 32, 670–676.

Xie, S., Duan, J., Li, B., Zhou, P., and Hon, G.C. (2017). Multiplexed Engineering and Analysis of Combinatorial Enhancer Activity in Single Cells. *Mol. Cell* 66, 285–299.

Xu, Z., Zan, H., Pone, E.J., Mai, T., and Casali, P. (2012). Immunoglobulin class-switch DNA recombination: induction, targeting and beyond. *Nat. Rev. Immunol.* 12, 517–531.

Yang, L., Grishin, D., Wang, G., Aach, J., Zhang, C.-Z., Chari, R., Homsy, J., Cai, X., Zhao, Y., Fan, J.-B., et al. (2014). Targeted and genome-wide sequencing reveal single nucleotide variations impacting specificity of Cas9 in human stem cells. *Nat. Commun.* 5, 5507.

Yang, L., Briggs, A.W., Chew, W.L., Mali, P., Guell, M., Aach, J., Goodman, D.B., Cox, D., Kan, Y., Lesha, E., et al. (2016a). Engineering and optimising deaminase fusions for genome editing. *Nat. Commun.* 7, 13330.

Yang, Y., Wang, L., Bell, P., McMenamin, D., He, Z., White, J., Yu, H., Xu, C., Morizono, H., Musunuru, K., et al. (2016b). A dual AAV system enables the Cas9-mediated correction of a metabolic liver disease in newborn mice. *Nat. Biotechnol.* 34, 334–338.

Yeang, C.H., McCormick, F., and Levine, A. (2008). Combinatorial patterns of somatic gene mutations in cancer. *FASEB J.* 22, 2605–2622.

Yin, H., Song, C.Q., Dorkin, J.R., Zhu, L.J., Li, Y., Wu, Q., Park, A., Yang, J., Suresh, S., Bizhanova, A., et al. (2016). Therapeutic genome editing by combined viral and non-viral delivery of CRISPR system components in vivo. *Nat. Biotechnol.* 34, 328–333.

Yu, X.F. (2006). Innate cellular defenses of APOBEC3 cytidine deaminases and viral counter-defenses. *Curr. Opin. HIV AIDS* 1, 187–193.

Yuan, G., Bin, J.C., McKay, D.J., and Snyder, F.F. (1999). Cloning and characterization of human guanine deaminase. Purification and partial amino acid sequence of the mouse protein. *J. Biol. Chem.* 274, 8175–8180.

Yuen, C.M., Rodda, S.J., Vokes, S.A., McMahon, A.P., and Liu, D.R. (2006). Control of transcription factor activity and osteoblast differentiation in mammalian cells using an evolved small-molecule-dependent intein. *J. Am. Chem. Soc.* 128, 8939–8946.

Zetsche, B., Gootenberg, J.S., Abudayyeh, O.O., Slaymaker, I.M., Makarova, K.S., Essletzbichler, P., Volz, S.E., Joung, J., van der Oost, J., Regev, A., et al. (2015a). Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system. *Cell* 163, 759–771.

Zetsche, B., Volz, S.E., and Zhang, F. (2015b). A split-Cas9 architecture for inducible genome editing and transcription modulation. *Nat. Biotechnol.* 33, 139–142.

Zhang, J., Wu, L.Y., Zhang, X.S., and Zhang, S. (2014). Discovery of co-occurring driver pathways in cancer. *BMC Bioinformatics* 15, 271.

Zhang, Y., Qin, W., Lu, X., Xu, J., Huang, H., Bai, H., Li, S., and Lin, S. (2017). Programmable base editing of zebrafish genome using a modified CRISPR-Cas9 system. *Nat. Commun.* 8, 118.

Zharkov, D.O. (2008). Base excision DNA repair. *Cell. Mol. Life Sci.* 65, 1544–1565.

Zong, Y., Wang, Y., Li, C., Zhang, R., Chen, K., Ran, Y., Qiu, J.L., Wang, D., and Gao, C. (2017). Precise base editing in rice, wheat and maize with a Cas9-cytidine deaminase fusion. *Nat. Biotechnol.* 35, 438–440.

Zuris, J.A., Thompson, D.B., Shu, Y., Guiling, J.P., Bessen, J.L., Hu, J.H., Maeder, M.L., Joung, J.K., Chen, Z.Y., and Liu, D.R. (2015). Cationic lipid-mediated delivery of proteins enables efficient protein-based genome editing in vitro and in vivo. *Nat. Biotechnol.* 33, 73–80.