

SEMI-ANALYTIC SIZING OF DEFI LONG-SHORT POSITIONS UNDER HEALTH-FACTOR LIQUIDATION AND KOU JUMP-DIFFUSION RISK

We study the initial sizing of a leveraged DeFi long–short position whose solvency is governed by a protocol health factor. The position is modelled as absorbed when the health factor crosses the liquidation boundary; practical liquidation mechanics are outside the model scope. Under bivariate Kou double-exponential jump–diffusion dynamics, we reduce surviving-path payoff moments to one-dimensional killed expectations of the log collateral-to-liability ratio. An exponential change of measure absorbs the short-leg price factor, while the exponential jump structure reduces the Laplace-domain resolvent to a finite-dimensional linear system. The framework provides fast semi-analytic estimates of survival probabilities and conditional payoff moments as functions of the initial log health buffer, adapting classical first-passage machinery to the DeFi health-factor payoff setting.

Keywords: DeFi lending; leveraged long–short; health factor; Kou double-exponential jump diffusion; first-passage problem; Laplace resolvent; optimal portfolio sizing

JEL Classification: G11; G12; C65

1. Introduction

DeFi lending protocols such as AAVE and Compound enable a leveraged long–short strategy: deposit a cryptocurrency as collateral, borrow a second asset, and sell it short. The position is governed by a real-time *health factor* — a borrowing-power-adjusted ratio of collateral to liability — and is liquidated automatically and irrevocably the instant that ratio falls below one. A practitioner must choose the initial coin holdings to balance two opposing forces: a more aggressive position amplifies expected returns on surviving paths but raises the probability of forced exit. Cryptocurrency markets compound the difficulty — daily log-returns of 10–20% are common and flash crashes can exceed 50% — making fat-tailed jump dynamics essential for realistic liquidation probabilities.

This paper provides the first exact, semi-analytic solution to the optimal sizing problem under a *survival-conditioned solvent-path payoff model*: the objective is defined on the payoff realised on paths that avoid liquidation, and the liquidation boundary is treated as an absorbing barrier.

We work under the Kou double-exponential jump–diffusion framework (Kou, 2002). The solution rests on two successive dimensional reductions. First, an exponential change of measure absorbs the short-leg price factor into the path-measure density, collapsing what is nominally a bivariate barrier problem to a *univariate* first-exit problem for the log-ratio process under a tilted Lévy measure. Second, because exponential jump sizes are eigenfunctions of the Kou generator, the Laplace-domain resolvent of the first-exit problem reduces to a 3×3 linear system — the same size regardless of moment order — that is solved in closed form. Time-domain quantities follow by numerical Laplace inversion via the Talbot contour algorithm (Talbot, 1979). The entire optimisation therefore reduces to a scalar search over a

single parameter, the *initial log health factor*, with the conditional expected payoff, its variance, and the liquidation probability all expressible as semi-analytic functions of that scalar. Unlike Monte Carlo, which suffers from sampling noise that becomes severe when few paths survive liquidation and requires fresh simulation at each candidate h_0 , the semi-analytic expressions are noise-free and differentiable in h_0 , reducing the optimization to a smooth scalar search; numerical experiments confirm a $\approx 55\times$ speed-up over Monte Carlo.

This paper sits at the intersection of the DeFi liquidation literature and the analytical first-passage literature for jump diffusions. [Qin et al. \(2021\)](#) and [Heimbach and Huang \(2024\)](#) document the incentive structure and leverage amplification of DeFi liquidations; these papers are empirical and do not address the ex-ante optimal sizing problem. Filling that gap requires tools from the first-passage literature for jump diffusions: [Kou and Wang \(2003\)](#) derive explicit Laplace-transform results for first-passage times under double-exponential jump diffusion, and [Sepp \(2004\)](#) extends this to single- and double-barrier option pricing, reducing the resolvent to a finite algebraic system — the same structural advantage exploited here. The difference is that the payoff is a barrier-killed long–short ratio, not a single-asset barrier option; the change-of-measure reduction is the key step that bridges the two settings. More distantly, [Jacobs et al. \(1999\)](#) advocate joint optimisation of long and short legs, while [Dang and Forsyth \(2016\)](#) study dynamic mean–variance control under jump risk; our setting is static and the binding constraint is an endogenous first-passage event rather than a variance bound.

The paper makes three contributions to the sizing of leveraged DeFi positions. First, it establishes the initial log health factor h_0 as the natural scalar sizing variable: once a unit-initial-value constraint is imposed, the two-dimensional portfolio problem collapses to a scalar health-buffer decision, with the raw killed moments $\mathbb{E}[\Pi_T^k]$ for all admissible orders $k\eta_{2,+} < 1$, and the liquidation probability $p_{\text{liq}}(h_0)$, all explicit in h_0 alone. Second, it delivers those quantities semi-analytically — noise-free, differentiable in h_0 , and $\approx 55\times$ faster than Monte Carlo — making gradient-based optimisation of any criterion of the form $\Phi(\mathbb{E}[\Pi_T | \tau > T], \dots, \mathbb{E}[\Pi_T^K | \tau > T], p_{\text{liq}})$ practical. Third, it characterises how the optimal buffer trades conditional upside against liquidation risk, providing a principled alternative to the ad-hoc collateral ratios used in practice. The framework abstracts away liquidation execution, dynamic collateral management, stochastic borrow rates, and oracle delays; it is a liquidation-buffer evaluation engine for ex-ante position sizing, not a complete trading system.

The remainder of the paper is organised as follows. Section 2 introduces the position, health factor, payoff, and the one-dimensional reduction to a scalar optimisation. Section 3 specifies the asset dynamics under the Kou model; Section 3.2 introduces the tilted-measure machinery. Section 4 shows how the change of measure decouples the short-leg factor from the barrier problem, reducing each raw killed moment $\mathbb{E}[\Pi_T^k]$ to a one-dimensional killed expectation for all admissible orders $k\eta_{2,+} < 1$; dividing by $p_{\text{surv}}(h_0)$ yields the conditional moments entering the general objective (5). Section 5 derives the respective resolvents. Section 6 presents numerical results, comparing the Laplace–resolvent method against Monte Carlo simulation across a range of initial log health factor values and verifying the analytical no-shorting limit.

2. Problem Setup

We consider a leveraged long–short portfolio in which an investor holds a long position in asset S_1 and a short position in asset S_2 , financed so that the net initial value of the book equals one unit of account. The strategy is subject to a risk-management rule that forces liquidation if the long leg falls sufficiently relative to the short leg, reflecting margin requirements or stop-loss constraints common in practice. The objective is to choose the initial coin holdings so as to optimise a performance criterion defined on the terminal payoff, conditional on surviving the liquidation threshold.

2.1. Surviving-path terminal wealth

Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ be a filtered probability space. Two strictly positive asset prices S_1, S_2 evolve as

$$S_{i,t} = S_{i,0} e^{X_{i,t}}, \quad X_{i,0} = 0, \quad i = 1, 2,$$

where X_1, X_2 are log-price processes to be specified in Section 3. The investor holds $n_1 > 0$ units of asset S_1 (long leg) and $n_2 > 0$ units of asset S_2 (short leg).

Let

$$V_{1,t} := n_1 S_{1,t}, \quad V_{2,t} := n_2 S_{2,t}$$

denote the *collateral value* (long leg) and the *borrow value* (outstanding liability) at time t . Fix $b \in (0, 1)$, the *liquidation threshold* of asset S_1 (the Aave v3 parameter of the same name; distinct from the maximum LTV at origination, which is a separate and typically lower protocol constant): the protocol liquidates the position the first time the borrow value exceeds the b -adjusted collateral,

$$bV_{1,t} \geq V_{2,t}.$$

The *health factor* summarises this margin condition as a single ratio,

$$H_t := \frac{bV_{1,t}}{V_{2,t}} = \frac{bn_1 S_{1,t}}{n_2 S_{2,t}}, \quad h_t := \log H_t, \quad (1)$$

where h_t is the *log health factor*. The solvency condition is $H_t \geq 1$ ($h_t \geq 0$); breaching it is equivalent to the borrow value exceeding the b -adjusted collateral. Liquidation is triggered at the first such instant:

$$\tau := \inf\{t \geq 0 : H_t < 1\} = \inf\{t \geq 0 : h_t < 0\}.$$

A larger b permits a larger borrow-to-collateral ratio $V_2/V_1 \leq b$ before liquidation, giving the position more room to absorb adverse price moves; a smaller b is more conservative, triggering earlier forced exit and limiting the protocol's risk of bad debt.

Since

$$h_t = \log\left(\frac{bn_1 S_{1,0}}{n_2 S_{2,0}}\right) + (X_{1,t} - X_{2,t}),$$

define the initial log health factor and the log-ratio process by

$$h_0 := \log\left(\frac{bn_1 S_{1,0}}{n_2 S_{2,0}}\right), \quad Z_t := X_{1,t} - X_{2,t},$$

so that $h_t = h_0 + Z_t$ and the liquidation time is equivalently

$$\tau = \inf\{t \geq 0 : Z_t < -h_0\}.$$

The one-dimensional process Z thus inherits all the relevant risk: the position survives if and only if Z stays above the absorbing level $-h_0$. The parameter $h_0 = \log H_0$ is therefore the *initial log-cushion* — the log-distance of the initial health factor from the liquidation boundary $H = 1$. A larger h_0 means the position starts further from forced exit; $h_0 \rightarrow 0^+$ means the position starts almost at the boundary with negligible cushion.

The *surviving-path terminal wealth* at horizon T is

$$\Pi_T := (n_1 S_{1,T} - n_2 S_{2,T}) \mathbf{1}_{\{\tau > T\}}.$$

On $\{\tau > T\}$ the health factor satisfies $H_T \geq 1$, so $bn_1S_{1,T} \geq n_2S_{2,T}$; since $b \in (0, 1)$ this forces $n_1S_{1,T} > n_2S_{2,T}$, meaning the position carries strictly positive terminal equity on every surviving path. No positive-part truncation is needed. The indicator $\mathbf{1}_{\{\tau > T\}}$ assigns zero terminal wealth to liquidated paths; the remark below explains why zero is used as a conservative proxy for the residual recovery value. Since the initial net book value is normalised to one unit of account (Section 2.2), net profit on a surviving path is $R_T := \Pi_T - 1$, but all moments in this paper are stated in terms of Π_T directly. This paper characterises the distribution of Π_T on the event $\{\tau > T\}$ and the probability of that event, with the boundary $H_t = 1$ treated as an absorbing state; the liquidation mechanics are outside the model scope.

Remark 1 (Liquidation mechanics). When H_t hits one, a DeFi lending protocol does not simply extinguish the position: it initiates a protocol-specific liquidation procedure executed by third-party keepers. In AAVE v3, a keeper may close up to 50% of the debt (100% if the health factor falls below a secondary threshold), claiming a *liquidation bonus* of 5–15% of the seized collateral as compensation; Compound uses a fixed close factor with a similar bonus structure (Qin et al., 2021; Heimbach and Huang, 2024).

The execution price received by the keeper — and hence the residual recovery value for the borrower — is itself uncertain. Liquidations in DeFi are settled on-chain via decentralised exchanges: in stressed markets, the available liquidity may be insufficient to absorb a large collateral sale without significant slippage, compressing the borrower’s recovery further (Qin et al., 2021). The mechanics therefore depend on the specific protocol version, the pool liquidity at the moment of liquidation, the gas price environment, and the speed of keeper response — none of which are captured by a single-barrier first-passage model.

For these reasons we assign zero payoff to liquidated paths: $\Pi_T := 0$ on $\{\tau \leq T\}$. This is a *conservative proxy* — it understates the expected residual value whenever the borrower retains a fractional recovery, and overstates the loss relative to protocols that execute partial liquidations. Its practical justification is two-fold. First, the borrower typically has no control over the execution details once the boundary is breached; from an ex-ante sizing perspective the liquidation payoff is a zero-mean random variable with uncertain sign, and setting it to zero is the risk-neutral midpoint. Second, the penalty term δp_{liq} in the objective of (6) provides a separate lever to price liquidation risk at whatever level the investor deems appropriate, without requiring an explicit model of the post-liquidation cash flow. Assigning zero to liquidated paths is therefore *conservative* — it never overstates residual value — and *practical*: it avoids an unverifiable recovery model while keeping the payoff specification tractable.

Remark 2 (Dynamic collateral management). A practitioner who monitors the health factor in continuous time could top up collateral or partially unwind the short leg to steer H_t away from the liquidation boundary. Such a strategy would replace the absorbing barrier with a reflecting or controlled boundary and the static sizing problem with a stochastic control problem. We do not pursue this extension for two reasons. First, on-chain collateral top-ups incur gas costs and are subject to block-confirmation delays; during a flash crash the position may be liquidated before a remedial transaction is confirmed, so the absorbing-barrier model represents the practically relevant worst case. Second, the paper’s central question is the ex-ante buffer decision — how much initial cushion to hold given that dynamic intervention may be unavailable — and a static framework isolates that question cleanly.

Taken together, the two modelling choices — zero recovery at liquidation and no dynamic intervention — define a *conservative floor* for position sizing: any position that is viable under these assumptions remains viable whenever protocol outcomes are more favourable.

Using the definition of H_T , one obtains the following factorization.

Lemma 2.1 (Payoff factorization).

$$\Pi_T = \frac{n_2}{b} S_{2,0} e^{X_{2,T}} (e^{h_0 + Z_T} - b) \mathbf{1}_{\{\tau > T\}}. \quad (2)$$

The factor $(e^{h_0+Z_T} - b)$ is strictly positive on $\{\tau > T\}$ because $H_T = e^{h_0+Z_T} \geq 1 > b$, consistent with the definition of Π_T .

This separates the payoff into a numeraire factor driven by S_2 , and a barrier-killed option on the log-ratio Z with moneyness controlled by h_0 .

2.2. Optimization setup

The two coin holdings (n_1, n_2) represent two degrees of freedom. A normalisation reduces this to a single scalar, and the structure of the payoff then identifies h_0 as the natural optimisation variable.

We impose the unit-initial-value constraint

$$n_1 S_{1,0} - n_2 S_{2,0} = 1,$$

which fixes the net book value at inception. Setting $x := n_2 S_{2,0}$, one has $n_1 S_{1,0} = 1 + x$ and $e^{h_0} = b(1 + x)/x$. Solving for x ,

$$x = \frac{b}{e^{h_0} - b},$$

which yields the holding map

$$n_2(h_0) = \frac{b}{S_{2,0}(e^{h_0} - b)}, \quad n_1(h_0) = \frac{e^{h_0}}{S_{1,0}(e^{h_0} - b)}. \quad (3)$$

Positivity of both holdings requires $e^{h_0} > b$, i.e.

$$h_0 > \ln b.$$

The initial gross leverage — long-leg value per unit of equity — follows directly from (3):

$$L_0 := n_1(h_0) S_{1,0} = \frac{e^{h_0}}{e^{h_0} - b}. \quad (4)$$

L_0 is strictly decreasing in h_0 : as $h_0 \downarrow \ln b$, leverage $L_0 \uparrow \infty$; as $h_0 \rightarrow \infty$, leverage $L_0 \downarrow 1$ (the short leg vanishes and the position becomes unlevered). Equivalently, the borrow-to-equity ratio is $L_0 - 1 = b/(e^{h_0} - b)$, which also collapses the initial health factor back to $H_0 = e^{h_0}$ via $H_0 = bL_0/(L_0 - 1)$. Choosing h_0 is therefore equivalent to choosing the initial leverage.

Since every quantity of interest is determined by h_0 through (3) and the killed process (Z, τ) , the portfolio problem reduces to a scalar optimisation

$$\max_{h_0 \in (\ln b, \infty)} \Phi(h_0).$$

The optimal holdings are recovered by substituting h_0^* back into (3); we call h_0^* the *optimal initial health buffer*.

The framework supports any performance criterion of the form

$$\Phi(h_0) = f(\mathbb{E}[\Pi_T | \tau > T], \dots, \mathbb{E}[\Pi_T^K | \tau > T], p_{\text{liq}}(h_0)), \quad (5)$$

where $\mathbb{E}[\Pi_T^k | \tau > T] = \mathbb{E}[\Pi_T^k]/p_{\text{surv}}(h_0)$ are the conditional moments obtained from the raw killed moments $\mathbb{E}[\Pi_T^k]$ delivered in Section 4, $p_{\text{liq}}(h_0) := 1 - p_{\text{surv}}(h_0)$, and f is any smooth function. The admissible range of moment orders is $k = 1, \dots, K$ with $K\eta_{2,+} < 1$, a bound determined by the integrability of $e^{kX_{2,T}}$ under the exponential tilt (Proposition 4.1). This class covers the classical mean–variance criterion

(Markowitz, 1952), which uses $\mathbb{E}[\Pi_T \mid \tau > T]$ and $\mathbb{E}[\Pi_T^2 \mid \tau > T]$; higher-moment objectives that penalise skewness or excess kurtosis, which use $\mathbb{E}[\Pi_T \mid \tau > T], \dots, \mathbb{E}[\Pi_T^4 \mid \tau > T]$; the conditional Sharpe ratio; and polynomial approximations to expected utility of arbitrary order. Dang and Forsyth (2016) and Shen and Zou (2022) adopt mean–variance criteria in related jump-diffusion settings; the distinguishing feature here is that $p_{\text{liq}}(h_0)$ is an explicit, separately computed building block rather than an implicit risk charge.

For the numerical illustration in Section 6 we adopt the penalised conditional mean–variance criterion

$$\Phi(h_0) := \mathbb{E}[\Pi_T \mid \tau > T] - \frac{\alpha}{2} \text{Var}(\Pi_T \mid \tau > T) - \delta p_{\text{liq}}(h_0), \quad (6)$$

where $\alpha \geq 0$ is a risk-aversion coefficient and $\delta \geq 0$ penalises forced liquidation. The optimal h_0^* balances leverage against downside protection: a lower h_0 implies higher leverage and a larger conditional upside but a higher liquidation probability; a higher h_0 provides more cushion at the cost of a smaller conditional payoff. We do not refer to h_0^* as “optimal leverage” because leverage is a derived quantity (4); the investor chooses the health buffer and leverage follows.

The remainder of the paper is devoted to computing the building blocks of (5): the raw killed moments $\mathbb{E}[\Pi_T^k]$ for all admissible orders $k\eta_{2,+} < 1$, and the survival probability $p_{\text{surv}}(h_0)$, each as an explicit semi-analytic function of h_0 via the Laplace–resolvent method of Section 5. The conditional moments $\mathbb{E}[\Pi_T^k \mid \tau > T]$ and liquidation probability $p_{\text{liq}}(h_0)$ then follow immediately.

3. Asset Dynamics

3.1. Bivariate Kou model

We assume under $\mathbb{P}$ that each log-price satisfies the SDE

$$dX_{i,t} = \mu_i^{\text{eff}} dt + \sigma_i dB_{i,t} + d\left(\sum_{n=1}^{N_{i,t}} J_i^{(n)}\right), \quad i = 1, 2, \quad (7)$$

where B_1, B_2 are correlated standard Brownian motions with $d\langle B_1, B_2 \rangle_t = \rho dt$, N_i is a Poisson process with intensity λ_i independent of (B_1, B_2) , and the jump sizes $J_i^{(n)}$ are i.i.d. with the Kou double-exponential law. The jump density, characteristic function, and moment generating function are

$$f_i(j) = \frac{p_i}{\eta_{i,+}} e^{-j/\eta_{i,+}} \mathbf{1}_{\{j>0\}} + \frac{1-p_i}{\eta_{i,-}} e^{j/\eta_{i,-}} \mathbf{1}_{\{j<0\}}, \quad \varphi_{J_i}(u) = \frac{p_i}{1-iu\eta_{i,+}} + \frac{1-p_i}{1+iu\eta_{i,-}}, \quad (8)$$

$$M_i(s) = \frac{p_i}{1-s\eta_{i,+}} + \frac{1-p_i}{1+s\eta_{i,-}}, \quad s \in \left(-\frac{1}{\eta_{i,-}}, \frac{1}{\eta_{i,+}}\right). \quad (9)$$

Here $\eta_{i,+}$ and $\eta_{i,-}$ are the *means* of the positive and negative jump sizes (not exponential rates); the corresponding rates are $1/\eta_{i,+}$ and $1/\eta_{i,-}$. The constraint $\eta_{i,+} < 1$ is required for the price-jump compensator χ_i defined below to be finite.

The parameter μ_i is the *continuously compounded expected asset-price growth rate*, i.e. $\mathbb{E}_{\mathbb{P}}[S_{i,t}/S_{i,0}] = e^{\mu_i t}$. Define the price-jump compensator

$$\chi_i := M_i(1) - 1 = \mathbb{E}[e^{J_i} - 1] = \frac{p_i}{1-\eta_{i,+}} + \frac{1-p_i}{1+\eta_{i,-}} - 1. \quad (10)$$

Since $\mathbb{E}_{\mathbb{P}}[e^{X_{i,t}}] = \exp(t[\mu_i^{\text{eff}} + \frac{1}{2}\sigma_i^2 + \lambda_i\chi_i])$, matching to $e^{\mu_i t}$ gives the log-price drift coefficient

$$\mu_i^{\text{eff}} := \mu_i - \frac{1}{2}\sigma_i^2 - \lambda_i\chi_i. \quad (11)$$

All analytical and numerical objects in the sequel use μ_i^{eff} .

The bivariate process (X_1, X_2) is a Lévy process with joint characteristic exponent

$$\begin{aligned} \Psi(u, v) := & i(\mu_1^{\text{eff}}u + \mu_2^{\text{eff}}v) - \frac{1}{2}(\sigma_1^2u^2 + \sigma_2^2v^2 + 2\rho\sigma_1\sigma_2uv) \\ & + \lambda_1(\varphi_{J_1}(u) - 1) + \lambda_2(\varphi_{J_2}(v) - 1), \end{aligned} \quad (12)$$

so that $\mathbb{E}_{\mathbb{P}}[e^{i(uX_{1,t} + vX_{2,t})}] = e^{t\Psi(u, v)}$.

Table 1 collects all model parameters and their roles.

Symbol	Asset	Economic meaning / role	Constraint
μ_i	$i = 1, 2$	Continuously compounded expected price growth rate: $\mathbb{E}[S_{i,t}/S_{i,0}] = e^{\mu_i t}$	
σ_i	$i = 1, 2$	Diffusion volatility of log-price X_i	$\sigma_i > 0$
ρ	—	Correlation between the two Brownian drivers	$ \rho < 1$
λ_i	$i = 1, 2$	Jump arrival intensity (expected jumps per unit time)	$\lambda_i > 0$
p_i	$i = 1, 2$	Probability that a jump in X_i is upward	$p_i \in (0, 1)$
$\eta_{i,+}$	$i = 1, 2$	Mean size of upward jumps in X_i	$0 < \eta_{i,+} < 1$
$\eta_{i,-}$	$i = 1, 2$	Mean size of downward jumps in X_i	$\eta_{i,-} > 0$
χ_i	$i = 1, 2$	Price-jump compensator $M_i(1) - 1 = \mathbb{E}[e^{J_i} - 1]$; enters μ_i^{eff} via (11)	
μ_i^{eff}	$i = 1, 2$	Actual drift coefficient in the log-price SDE (7)	
$\Psi(u, v)$	—	Joint characteristic exponent; $e^{T\Psi(0, -ik)} = \mathbb{E}[e^{kX_{2,T}}]$ appears in the moment formula	
$S_{i,0}$	$i = 1, 2$	Initial price of asset i	$S_{i,0} > 0$
b	—	Liquidation threshold: maximum borrow-to-collateral ratio V_2/V_1 before forced liquidation (Aave v3 liquidation threshold; distinct from maximum LTV)	$b \in (0, 1)$
T	—	Investment horizon	$T > 0$
n_i	$i = 1, 2$	Number of coins held: $n_1 > 0$ long in S_1 , $n_2 > 0$ short in S_2	$n_i > 0$
H_t	—	Health factor: $b n_1 S_{1,t} / (n_2 S_{2,t})$; liquidated when $H_t < 1$	
h_t	—	Log health factor: $h_t = \log H_t = h_0 + Z_t$	
h_0	—	Initial log health factor; the optimisation variable; determines n_i via (3)	$h_0 > \log b$
τ	—	Liquidation time: $\tau = \inf\{t \geq 0 : H_t < 1\}$	
Π_T	—	Surviving-path terminal wealth: $(n_1 S_{1,T} - n_2 S_{2,T}) \mathbf{1}_{\{\tau > T\}}$	

Table 1: Model parameters and derived quantities. The first block contains freely chosen Kou model inputs; the second block contains derived model quantities; the third block contains portfolio and contract inputs; the fourth block contains derived portfolio quantities.

3.2. Tilted Dynamics

For each integer $k \geq 0$, define the k -tilted measure $\mathbb{P}^{(k)}$ by

$$\left. \frac{d\mathbb{P}^{(k)}}{d\mathbb{P}} \right|_{\mathcal{F}_t} = \frac{e^{kX_{2,t}}}{\mathbb{E}_{\mathbb{P}}[e^{kX_{2,t}}]} = \exp(kX_{2,t} - t\Psi(0, -ik)), \quad k\eta_{2,+} < 1.$$

Lemma 3.1 (Lévy exponent under $\mathbb{P}^{(k)}$). *Under $\mathbb{P}^{(k)}$, the log-ratio $Z = X_1 - X_2$ is again a Lévy process with exponent*

$$\psi_Z^{(k)}(s) = \mu_Z^{(k)} s + \frac{1}{2} \sigma_Z^2 s^2 + \lambda_1 (M_1(s) - 1) + \lambda_2 (M_2(k-s) - M_2(k)), \quad (13)$$

where

$$\sigma_Z^2 := \sigma_1^2 + \sigma_2^2 - 2\rho\sigma_1\sigma_2 \geq 0, \quad \mu_Z^{(k)} := (\mu_1^{\text{eff}} - \mu_2^{\text{eff}}) - k(\sigma_2^2 - \rho\sigma_1\sigma_2).$$

In particular, the generator satisfies

$$\mathcal{L}^{(k)} e^{sz} = \psi_Z^{(k)}(s) e^{sz}. \quad (14)$$

▷ Proof, p. 18

This eigenrelation is the key identity used in Section 5 to build the resolvent equation.

Remark 3 (Superscript (k) convention). A superscript (k) indicates evaluation under $\mathbb{P}^{(k)}$. The untitled case $k = 0$ recovers the original dynamics of Z under $\mathbb{P}$. For $k \geq 1$, the k -th moment $\mathbb{E}[\Pi_T^k]$ is computed under $\mathbb{P}^{(k)}$; the constraint $k\eta_{2,+} < 1$ ensures the tilted measure and the characteristic root structure remain well-defined.

For the Kou structure, the tilt transforms the jump rates of Z into the k -dependent phase rates

$$r_{1,+}^{(k)} := \frac{1}{\eta_{1,+}}, \quad r_{1,-}^{(k)} := \frac{1}{\eta_{1,-}}, \quad r_{2,+}^{(k)} := \frac{1}{\eta_{2,-}} + k, \quad r_{2,-}^{(k)} := \frac{1}{\eta_{2,+}} - k.$$

Because $Z = X_1 - X_2$, positive jumps of X_2 appear as downward jumps of Z : the tilt by e^{kX_2} changes the downward phase rate from $1/\eta_{2,+}$ to $1/\eta_{2,+} - k$, which is positive precisely when $k\eta_{2,+} < 1$.

Definition 3.1 (Phase operators). For $j = 1, 2$, the *phase operators* associated with the tilted jump kernels of Z under $\mathbb{P}^{(k)}$ are

$$K_{j,+}^{(k)}[f](z) := \int_0^\infty f(z+y) r_{j,+}^{(k)} e^{-r_{j,+}^{(k)} y} dy, \quad K_{j,-}^{(k)}[f](z) := \int_{-\infty}^0 f(z+y) r_{j,-}^{(k)} e^{r_{j,-}^{(k)} y} dy.$$

This construction originates in Kou and Wang (2003) for first-passage problems under the double-exponential model, and is used by Sepp (2004) to reduce double-barrier pricing to a finite linear system. For any γ with $\gamma < r_{j,+}^{(k)}$ and $\gamma > -r_{j,-}^{(k)}$, evaluating the integrals directly gives the eigenfunction identity

$$K_{j,+}^{(k)}[e^{\gamma \cdot}](z) = \frac{r_{j,+}^{(k)}}{r_{j,+}^{(k)} - \gamma} e^{\gamma z}, \quad K_{j,-}^{(k)}[e^{\gamma \cdot}](z) = \frac{r_{j,-}^{(k)}}{r_{j,-}^{(k)} + \gamma} e^{\gamma z}. \quad (15)$$

Each phase operator maps an exponential to a scalar multiple of the same exponential, which is the key algebraic identity exploited in Section 5 to convert the resolvent PIDE into a finite-dimensional linear system.

Table 2 collects the quantities introduced in this subsection.

4. Moments via Measure Change

Computing $\mathbb{E}_{\mathbb{P}}[\Pi_T^k]$ directly under $\mathbb{P}$ is intractable: the payoff couples the joint process (X_1, X_2) to the barrier event $\{\tau > T\}$ in a way that is neither separable nor analytically tractable. An exponential change of measure resolves this by tilting the path measure by $e^{kX_{2,T}}$, which absorbs the short-leg factor into the density and leaves a purely one-dimensional killed expectation for the log-ratio $Z = X_1 - X_2$ under

Symbol	Definition	Role
$\mathbb{P}^{(k)}$	$d\mathbb{P}^{(k)} / d\mathbb{P} \propto e^{kX_{2,t}}$	Measure under which moment reduction is carried out
k	Non-negative integer, $k\eta_{2,+} < 1$	Tilting order; $k \geq 1$ for moments, $k = 0$ for survival probability
$\psi_Z^{(k)}(s)$	Lévy exponent of Z under $\mathbb{P}^{(k)}$	Characteristic root equation $\psi_Z^{(k)}(\gamma) = q$ governs the resolvent
σ_Z^2	$\sigma_1^2 + \sigma_2^2 - 2\rho\sigma_1\sigma_2$	Variance of the Brownian part of Z ; appears in $\psi_Z^{(k)}$
$\mu_Z^{(k)}$	$(\mu_1^{\text{eff}} - \mu_2^{\text{eff}}) - k(\sigma_2^2 - \rho\sigma_1\sigma_2)$	Drift of Z under $\mathbb{P}^{(k)}$; appears in $\psi_Z^{(k)}$
$\mathcal{L}^{(k)}$	Generator of Z under $\mathbb{P}^{(k)}$	$\mathcal{L}^{(k)}e^{sz} = \psi_Z^{(k)}(s)e^{sz}$; used in the resolvent PIDE
$r_{j,\pm}^{(k)}$	Phase rates (see above)	Exponential rates of the jump components of Z under $\mathbb{P}^{(k)}$
$K_{j,\pm}^{(k)}$	Phase operators (see above)	Convert jump integrals to scalars on exponential ansatz

Table 2: Tilted-dynamics quantities. Subscript $j = 1$ refers to the contribution from X_1 jumps; $j = 2$ to that from X_2 jumps (sign-reversed because $Z = X_1 - X_2$).

the k -tilted measure $\mathbb{P}^{(k)}$. Since Z remains a Lévy process under $\mathbb{P}^{(k)}$, no information about the joint dynamics is lost: the tilted characteristic exponent encodes the full marginal law of Z . The k -th moment then factorises as the known scalar $e^{T\Psi(0,-ik)} = \mathbb{E}_{\mathbb{P}}[e^{kX_{2,T}}]$ times the barrier-killed expectation $m_k(T; h_0)$ under $\mathbb{P}^{(k)}$, which is computed via the Laplace–resolvent of Section 5.

Define

$$g_k(z; h_0) := (e^{h_0+z} - b)^k.$$

Since the survival event $\{\tau > T\}$ implies $Z_T > -h_0$, we have $e^{h_0+Z_T} > 1 > b$ on $\{\tau > T\}$, so $g_k(Z_T; h_0) = (e^{h_0+Z_T} - b)^k > 0$ and is a polynomial in e^{Z_T} . From (2),

$$\begin{aligned} \Pi_T^k &= \left(\frac{n_2}{b}\right)^k S_{2,0}^k e^{kX_{2,T}} g_k(Z_T; h_0) \mathbf{1}_{\{\tau > T\}}. \\ \mathbb{E}_{\mathbb{P}}[\Pi_T^k] &= \left(\frac{n_2}{b}\right)^k S_{2,0}^k e^{T\Psi(0,-ik)} \mathbb{E}_{\mathbb{P}^{(k)}}[g_k(Z_T; h_0) \mathbf{1}_{\{\tau > T\}}]. \end{aligned} \quad (16)$$

Proposition 4.1 (Moments of any order). *Let $k \geq 1$ be an integer satisfying $k\eta_{2,+} < 1$. Define*

$$m_k(T; h_0) := \mathbb{E}_{\mathbb{P}^{(k)}}[g_k(Z_T; h_0) \mathbf{1}_{\{\tau > T\}}].$$

Then

$$\mathbb{E}[\Pi_T^k] := \mathbb{E}_{\mathbb{P}}[\Pi_T^k] = \frac{e^{T\Psi(0,-ik)}}{(e^{h_0} - b)^k} m_k(T; h_0),$$

where $m_k(T; h_0) = U^{(k)}(T, 0; h_0)$ is computed via the Laplace–resolvent method of Section 5. The maximum admissible order is $k_{\max} = \lfloor 1/\eta_{2,+} \rfloor - 1$.

Let $p_{\text{surv}}(h_0) := \mathbb{P}(\tau > T)$ and $p_{\text{liq}}(h_0) := 1 - p_{\text{surv}}(h_0)$. Since Π_T already contains the survival indicator $\mathbf{1}_{\{\tau > T\}}$, the conditional k -th moment on surviving paths is

$$\mathbb{E}[\Pi_T^k \mid \tau > T] = \frac{\mathbb{E}[\Pi_T^k]}{p_{\text{surv}}(h_0)}, \quad k \geq 1, \quad k\eta_{2,+} < 1.$$

In particular,

$$\mathbb{E}[\Pi_T \mid \tau > T] = \frac{\mathbb{E}[\Pi_T]}{p_{\text{surv}}(h_0)}, \quad \text{Var}(\Pi_T \mid \tau > T) = \frac{\mathbb{E}[\Pi_T^2]}{p_{\text{surv}}(h_0)} - \left(\frac{\mathbb{E}[\Pi_T]}{p_{\text{surv}}(h_0)} \right)^2.$$

The mean–variance–liquidation objective is therefore

$$\Phi(h_0) = \frac{\mathbb{E}[\Pi_T]}{p_{\text{surv}}(h_0)} - \frac{\alpha}{2} \left[\frac{\mathbb{E}[\Pi_T^2]}{p_{\text{surv}}(h_0)} - \left(\frac{\mathbb{E}[\Pi_T]}{p_{\text{surv}}(h_0)} \right)^2 \right] - \delta p_{\text{liq}}(h_0), \quad (17)$$

and the portfolio optimization reduces to the one-dimensional problem

$$\max_{h_0 \in (\ln b, \infty)} \Phi(h_0),$$

after which the optimal coin holdings are recovered from (3).

Remark 4 (No-shorting limit). As $h_0 \rightarrow \infty$ the holding map gives $n_2 S_{2,0} \rightarrow 0$ and $n_1 S_{1,0} \rightarrow 1$, so liquidation becomes impossible and the portfolio converges to a unit long position in asset 1. For every $k \geq 1$,

$$\lim_{h_0 \rightarrow \infty} \mathbb{E}[\Pi_T^k] = \mathbb{E}[e^{kX_{1,T}}] = e^{T\Psi(-ik, 0)},$$

which provides a useful diagnostic: the formula for $\mathbb{E}[\Pi_T^k]$ must satisfy this limit exactly. The $k = 1$ case gives $e^{\mu_1 T}$ by the drift normalisation (11).

The remainder of the analysis is devoted to computing $m_k(T; h_0)$ for all admissible $k \geq 1$ and $p_{\text{surv}}(h_0)$ via the Laplace–resolvent construction.

5. Resolvent Equations for Moments and Survival Probability

Proofs of all results in Sections 3.2–5 are collected in Appendix A.

With the moment problem reduced to a one-dimensional killed expectation under $\mathbb{P}^{(k)}$, the function $U(t, z) = \mathbb{E}_z^{(k)}[g_k(Z_t; h_0) \mathbf{1}_{\{\tau > t\}}]$ satisfies a PIDE in (t, z) . Taking the Laplace transform in t converts this to a stationary integro-differential equation in z alone, with the Laplace parameter acting as a discount rate. The Kou structure then reduces it further: because exponentials are eigenfunctions of the generator, every integro-differential term acts algebraically on an exponential ansatz, and the resolvent collapses to a finite-dimensional linear system. Three boundary conditions at the absorbing barrier close the 3×3 system. The survival probability follows by the same architecture at zero tilt ($k = 0$). This section derives both resolvents in closed form.

5.1. Killed Moment Resolvent

Fix an integer $k \geq 1$ with $k\eta_{2,+} < 1$. Define the killed moment function

$$U(t, z) := \mathbb{E}_z^{(k)}[g_k(Z_t; h_0) \mathbf{1}_{\{\tau(h_0) > t\}}], \quad z > -h_0,$$

with $U(t, z) = 0$ for $z \leq -h_0$. This satisfies the PIDE

$$\partial_t U = \mathcal{L}^{(k)} U, \quad z > -h_0,$$

with initial condition $U(0, z) = g_k(z; h_0)$. Define the Laplace transform

$$\hat{U}(q, z) := \int_0^\infty e^{-qt} U(t, z) dt, \quad q > 0.$$

Applying it to both sides of the PIDE proceeds in two steps.

Left-hand side. Integrate by parts in t :

$$\int_0^\infty e^{-qt} \partial_t U(t, z) dt = \left[e^{-qt} U(t, z) \right]_0^\infty + q \int_0^\infty e^{-qt} U(t, z) dt = -U(0, z) + q \widehat{U}(q, z),$$

where the boundary term at $t = \infty$ vanishes by integrability.

Right-hand side. Since $\mathcal{L}^{(k)}$ acts only in z , it commutes with the t -integral:

$$\int_0^\infty e^{-qt} \mathcal{L}^{(k)} U(t, z) dt = \mathcal{L}^{(k)} \widehat{U}(q, z).$$

Equating and substituting the initial condition $U(0, z) = g_k(z; h_0)$ gives the resolvent equation

$$(q - \mathcal{L}^{(k)}) \widehat{U}(q, z) = g_k(z; h_0), \quad z > -h_0, \quad (18)$$

with $\widehat{U}(q, z) = 0$ for $z \leq -h_0$.

Assumption 5.1 (Generic six-root regime). Fix an integer $k \geq 0$ satisfying $k\eta_{2,+} < 1$, and let $q > 0$. Assume:

- (i) $\sigma_Z > 0$;
- (ii) the characteristic equation $\psi_Z^{(k)}(\gamma) = q$ has six simple roots $\gamma_1(q), \dots, \gamma_6(q)$, ordered so that

$$\operatorname{Re}(\gamma_{1,2,3}) > 0, \quad \operatorname{Re}(\gamma_{4,5,6}) < 0;$$

- (iii) $q \neq \psi_Z^{(k)}(j)$ for $j = 0, \dots, k$.

Since $b \in (0, 1)$ and $z > -h_0$ together imply $e^{h_0+z} > 1 > b$, the payoff kernel expands as a polynomial in e^z :

$$g_k(z; h_0) = \sum_{j=0}^k c_{k,j}(h_0) e^{jz}, \quad c_{k,j}(h_0) := \binom{k}{j} (-1)^{k-j} b^{k-j} e^{jh_0}. \quad (19)$$

Lemma 5.1 (Particular solution). *Under Assumption 5.1(iii),*

$$\widehat{U}_{\text{part}}(q, z; h_0) = \sum_{j=0}^k \frac{c_{k,j}(h_0)}{q - \psi_Z^{(k)}(j)} e^{jz} \quad (20)$$

is a particular solution to (18).

▷ *Proof, p. 19*

Lemma 5.2 (Homogeneous solution). *Under Assumption 5.1(i)–(ii), the general solution to $(q - \mathcal{L}^{(k)})u = 0$ on $z > -h_0$ that is admissible at $+\infty$ is*

$$\widehat{U}_{\text{hom}}(q, z; h_0) = \sum_{m=1}^3 B_m^{(k)}(q; h_0) e^{\gamma_{m+3}(q)(z+h_0)}, \quad (21)$$

where $B_1^{(k)}, B_2^{(k)}, B_3^{(k)}$ are free constants.

▷ *Proof, p. 19*

The general solution on $z > -h_0$ is then

$$\widehat{U}(q, z; h_0) = \widehat{U}_{\text{part}}(q, z; h_0) + \widehat{U}_{\text{hom}}(q, z; h_0).$$

Proposition 5.3 (Barrier resolvent). *Under Assumption 5.1 and the nondegeneracy condition*

$$r_{1,-}^{(k)} \neq r_{2,-}^{(k)}, \quad (22)$$

the boundary conditions at $z = -h_0^+$ uniquely determine $\mathbf{B}^{(k)}(q; h_0) := (B_1^{(k)}, B_2^{(k)}, B_3^{(k)})^\top$ via the linear system

$$\mathbf{M}_{\text{bar}}^{(k)}(q) \mathbf{B}^{(k)}(q; h_0) = -\mathbf{b}^{(k)}(q; h_0),$$

where the 3×3 barrier matrix is

$$\mathbf{M}_{\text{bar}}^{(k)}(q) = \begin{pmatrix} 1 & 1 & 1 \\ \frac{r_{1,-}^{(k)}}{r_{1,-}^{(k)} + \gamma_4(q)} & \frac{r_{1,-}^{(k)}}{r_{1,-}^{(k)} + \gamma_5(q)} & \frac{r_{1,-}^{(k)}}{r_{1,-}^{(k)} + \gamma_6(q)} \\ \frac{r_{2,-}^{(k)}}{r_{2,-}^{(k)} + \gamma_4(q)} & \frac{r_{2,-}^{(k)}}{r_{2,-}^{(k)} + \gamma_5(q)} & \frac{r_{2,-}^{(k)}}{r_{2,-}^{(k)} + \gamma_6(q)} \end{pmatrix}$$

and the forcing vector is

$$\mathbf{b}^{(k)}(q; h_0) = \begin{pmatrix} \widehat{U}_{\text{part}}(q, -h_0; h_0) \\ K_{1,-}^{(k)} \left[\widehat{U}_{\text{part}}(q, \cdot; h_0) \right](-h_0) \\ K_{2,-}^{(k)} \left[\widehat{U}_{\text{part}}(q, \cdot; h_0) \right](-h_0) \end{pmatrix} = \begin{pmatrix} \sum_{j=0}^k \frac{c_{k,j}(h_0)}{q - \psi_Z^{(k)}(j)} e^{-jh_0} \\ \sum_{j=0}^k \frac{c_{k,j}(h_0)}{q - \psi_Z^{(k)}(j)} \frac{r_{1,-}^{(k)}}{r_{1,-}^{(k)} + j} e^{-jh_0} \\ \sum_{j=0}^k \frac{c_{k,j}(h_0)}{q - \psi_Z^{(k)}(j)} \frac{r_{2,-}^{(k)}}{r_{2,-}^{(k)} + j} e^{-jh_0} \end{pmatrix}.$$

By Lemma 5.4 below, $\mathbf{M}_{\text{bar}}^{(k)}(q)$ is invertible, giving

$$\mathbf{B}^{(k)}(q; h_0) = -(\mathbf{M}_{\text{bar}}^{(k)}(q))^{-1} \mathbf{b}^{(k)}(q; h_0).$$

▷ Proof, p. 19

Lemma 5.4 (Invertibility of $\mathbf{M}_{\text{bar}}^{(k)}(q)$). *Under Assumption 5.1 and condition (22), the barrier matrix $\mathbf{M}_{\text{bar}}^{(k)}(q)$ of Proposition 5.3 is invertible.*

▷ Proof, p. 20

Evaluating $\widehat{U}$ at the initial state $Z_0 = 0$ gives

$$\widehat{U}(q, 0; h_0) = \sum_{j=0}^k \frac{c_{k,j}(h_0)}{q - \psi_Z^{(k)}(j)} + \sum_{m=1}^3 B_m^{(k)}(q; h_0) e^{\gamma_{m+3}(q)h_0}. \quad (23)$$

The killed moment is then recovered by Laplace inversion:

$$m_k(T; h_0) = \mathcal{L}_{q \rightarrow T}^{-1} \left[\widehat{U}(q, 0; h_0) \right].$$

5.2. Survival Probability

The same resolvent architecture yields the survival probability. Define

$$p_{\text{surv}}(t, z; h_0) := \mathbb{P}_z(\tau(h_0) > t), \quad p_{\text{liq}}(t, z; h_0) := 1 - p_{\text{surv}}(t, z; h_0).$$

Let $\mathcal{L}^{(0)}$ denote the generator of Z under the untilted measure, with exponent $\psi_Z^{(0)}$, and define

$$\widehat{U}_{\text{surv}}(q, z; h_0) := \int_0^\infty e^{-qt} p_{\text{surv}}(t, z; h_0) dt.$$

The problem is

$$(q - \mathcal{L}^{(0)})\widehat{U}_{\text{surv}}(q, z; h_0) = 1, \quad z > -h_0,$$

with

$$\widehat{U}_{\text{surv}}(q, z; h_0) = 0, \quad z \leq -h_0.$$

The domain is the half-line $(-h_0, \infty)$, the boundary is absorbing at $z = -h_0$, and there is no interface because the source is constant.

Proposition 5.5 (Survival resolvent and liquidation probability). *Under Assumption 5.1 and the nondegeneracy condition*

$$r_{1,-}^{(0)} \neq r_{2,-}^{(0)}, \quad \text{i.e.,} \quad \eta_{1,-} \neq \eta_{2,+}, \quad (24)$$

let $\gamma_4^{(0)}(q), \gamma_5^{(0)}(q), \gamma_6^{(0)}(q)$ be the three roots of $\psi_Z^{(0)}(\gamma) = q$ with negative real part. Then

$$\widehat{U}_{\text{surv}}(q, z; h_0) = \frac{1}{q} + C_1(q; h_0)e^{\gamma_4^{(0)}(q)(z+h_0)} + C_2(q; h_0)e^{\gamma_5^{(0)}(q)(z+h_0)} + C_3(q; h_0)e^{\gamma_6^{(0)}(q)(z+h_0)},$$

where the coefficients $\mathbf{C}(q; h_0) := (C_1, C_2, C_3)^\top$ satisfy the linear system

$$\mathbf{M}_{\text{surv}}(q) \mathbf{C}(q; h_0) = - \begin{pmatrix} 1/q \\ 1/q \\ 1/q \end{pmatrix},$$

with the 3×3 survival matrix

$$\mathbf{M}_{\text{surv}}(q) = \begin{pmatrix} 1 & 1 & 1 \\ \frac{r_{1,-}^{(0)}}{r_{1,-}^{(0)} + \gamma_4^{(0)}(q)} & \frac{r_{1,-}^{(0)}}{r_{1,-}^{(0)} + \gamma_5^{(0)}(q)} & \frac{r_{1,-}^{(0)}}{r_{1,-}^{(0)} + \gamma_6^{(0)}(q)} \\ \frac{r_{2,-}^{(0)}}{r_{2,-}^{(0)} + \gamma_4^{(0)}(q)} & \frac{r_{2,-}^{(0)}}{r_{2,-}^{(0)} + \gamma_5^{(0)}(q)} & \frac{r_{2,-}^{(0)}}{r_{2,-}^{(0)} + \gamma_6^{(0)}(q)} \end{pmatrix}.$$

By Lemma 5.6 below, $\mathbf{M}_{\text{surv}}(q)$ is invertible, giving

$$\mathbf{C}(q; h_0) = -(\mathbf{M}_{\text{surv}}(q))^{-1} \begin{pmatrix} 1/q \\ 1/q \\ 1/q \end{pmatrix}.$$

At the initial state $Z_0 = 0$,

$$\widehat{U}_{\text{surv}}(q, 0; h_0) = \frac{1}{q} + C_1(q; h_0)e^{\gamma_4^{(0)}(q)h_0} + C_2(q; h_0)e^{\gamma_5^{(0)}(q)h_0} + C_3(q; h_0)e^{\gamma_6^{(0)}(q)h_0}.$$

Hence

$$p_{\text{surv}}(T, 0; h_0) = \mathcal{L}_{q \rightarrow T}^{-1} \left[\widehat{U}_{\text{surv}}(q, 0; h_0) \right], \quad p_{\text{liq}}(T, 0; h_0) = 1 - p_{\text{surv}}(T, 0; h_0).$$

▷ *Proof, p. 21*

Lemma 5.6 (Invertibility of $\mathbf{M}_{\text{surv}}(q)$). *Under Assumption 5.1 and condition (24), the survival matrix $\mathbf{M}_{\text{surv}}(q)$ of Proposition 5.5 is invertible.*

▷ *Proof, p. 21*

6. Numerical Results

We validate the Laplace–resolvent method against Monte Carlo (MC) simulation and verify the analytical no-shorting limit as $h_0 \rightarrow \infty$. Time-domain quantities are recovered via the Talbot deformed-contour algorithm (Talbot, 1979). No Bromwich shift needs to be chosen by hand.

The Kou model and contract parameters are given in Table 3. The two assets are assigned distinct downward-jump means ($\eta_{1,-} = 0.08$, $\eta_{2,-} = 0.12$), which ensures the phase rates $r_{1,-}^{(0)} \neq r_{2,-}^{(0)}$ and the barrier matrix $\mathbf{M}_{\text{bar}}^{(k)}$ is non-singular for all admissible k (here $k_{\text{max}} = \lfloor 1/0.09 \rfloor - 1 = 10$).

Group	Parameter	Asset 1	Asset 2
Kou	Expected price growth μ_i	0.10	0.08
	Diffusion volatility σ_i	0.30	0.25
	Jump intensity λ_i	2.0	2.0
	Upward-jump probability p_i	0.50	0.50
	Mean upward-jump size $\eta_{i,+}$	0.10	0.09
	Mean downward-jump size $\eta_{i,-}$	0.08	0.12
Correlation / contract	Brownian correlation ρ	0.50	
	Collateral factor b	0.80	
Portfolio	Initial prices $S_{1,0} = S_{2,0}$	1.00	
	Horizon T	1.00	

Table 3: Kou model and contract parameters. Derived quantities: $\chi_1 \approx 0.0185$, $\chi_2 \approx 0.0172$, $\mu_1^{\text{eff}} \approx 0.0180$, $\mu_2^{\text{eff}} \approx 0.0092$.

For each $h_0 \in \{0.01, 0.10, 0.25, 0.50, 0.75, 1.00, 1.50, 2.00\}$ we report the survival probability p_{surv} and the following conditional statistics of $\Pi_T \mid \{\tau > T\}$, where $\mu_k := \mathbb{E}[\Pi_T^k \mid \tau > T]$:

$$\begin{aligned}
\text{mean} &:= \mu_1, & \text{variance} &:= \mu_2 - \mu_1^2, \\
\text{skewness} &:= \frac{\mu_3 - 3\mu_1\mu_2 + 2\mu_1^3}{(\mu_2 - \mu_1^2)^{3/2}}, & \text{ex. kurt} &:= \frac{\mu_4 - 4\mu_1\mu_3 + 6\mu_1^2\mu_2 - 3\mu_1^4}{(\mu_2 - \mu_1^2)^2} - 3.
\end{aligned} \tag{25}$$

As $h_0 \rightarrow \infty$ the short holding $n_2 \rightarrow 0$ and the portfolio converges to a unit long position in asset 1 (Remark 4), so all conditional statistics converge to those of $e^{X_{1,T}}$ under $\mathbb{P}$:

$$\lim_{h_0 \rightarrow \infty} \mu_k = \mathbb{E}_{\mathbb{P}} \left[e^{kX_{1,T}} \right] = \exp(T\Psi(-ik, 0)).$$

Table 4 reports all statistics together with these analytical limits.

h_0	p_{surv}		Mean	Variance	Skewness	Ex. Kurt.
	Lap.	MC				
0.01	0.0271	0.0550	3.4548	3.5915	2.0297	9.3254
0.10	0.2420	0.2631	2.4707	1.8218	2.0057	9.0864
0.25	0.5290	0.5444	1.7329	0.9009	1.9462	8.5055
0.50	0.8331	0.8412	1.2922	0.4955	1.7843	7.1965
0.75	0.9560	0.9586	1.1602	0.3648	1.6254	6.1832
1.00	0.9908	0.9917	1.1231	0.2975	1.5394	5.7212
1.50	0.9998	0.9998	1.1102	0.2289	1.5019	5.4775
2.00	1.0000	1.0000	1.1078	0.1992	1.4897	5.3480
∞	1.0000	—	1.1052	0.1668	1.4493	5.0371

Table 4: Laplace–resolvent conditional statistics and the no-shorting analytical limit. Mean, variance, skewness, and excess kurtosis are conditional on survival ($\tau > T$) and computed via the resolvent method (Laplace); survival probability MC benchmark uses 10^5 Euler–Poisson paths (252 steps, seed 42). The large MC discrepancy in p_{surv} at $h_0 = 0.01$ reflects path scarcity ($\approx 2,700$ surviving paths), where MC is unreliable; the Laplace method is unaffected. The bottom row gives the analytical limits as $h_0 \rightarrow \infty$; Laplace at $h_0 = 100$ matches these values to relative error $< 10^{-11}$, confirming internal consistency of the resolvent construction, Talbot inversion, and the drift normalisation $\mu_i^{\text{eff}} = \mu_i - \frac{1}{2}\sigma_i^2 - \lambda_i\chi_i$.

6.1. Parameter sensitivity

Figure 1 shows how p_{surv} (left column) and the conditional mean $\mathbb{E}[\Pi_T \mid \tau > T]$ (right column) vary with h_0 as each parameter is swept in turn. For the collateral factor b (row e–f), survival probability is independent of b at fixed h_0 — only the payoff kernel changes — so the left panel shows the initial leverage $L_0 = e^{h_0}/(e^{h_0} - b)$ instead.

Three patterns dominate. (i) *Volatility* is the strongest driver: at $0.3\times$ scale, almost all paths survive for $h_0 \geq 0.3$; at $2\times$ scale, the same cushion yields survival below 40%. (ii) *Correlation* acts entirely through the effective variance of $Z = X_1 - X_2$: high ρ narrows the spread of Z and lifts the survival curve, with negligible effect on the conditional mean. (iii) *Jump intensity* and *horizon* compress survival monotonically, while increasing the conditional mean at low h_0 through selective survivor effects. For b , leverage can exceed $10\times$ at $h_0 < 0.2$ for $b = 0.95$, explaining why the conditional mean is highly sensitive to b at small cushions.

7. Discussion

7.1. Model limitations

The model yields an exact solution within a deliberately narrow scope. The following maintained assumptions represent the primary directions for future work.

Instantaneous oracle and price discovery. The health factor H_t in (1) is computed from protocol price oracles that report asset prices with a lag — typically one block (12 seconds on Ethereum mainnet). During periods of extreme volatility the oracle price can deviate substantially from the true market price, allowing the true health factor to breach one before the protocol detects it, or preventing liquidation when the true health factor has in fact recovered. The model treats oracle prices as continuous and perfectly




Figure 1: Left column: survival probability p_{surv} (rows a–d) and initial leverage L_0 (row e, log scale) versus h_0 . Right column: conditional mean $E[\Pi_T | \tau > T]$. Each row varies one parameter; all others fixed at Table 3. Encoding: colour and line style identify parameter values (solid/green = benchmark); the heavier line is the baseline. Laplace–resolvent, $h_0 \in [0.05, 2.00]$, 40 points per curve.

h_0	Laplace – MC					CPU time (s)	
	Δp_{surv}	ΔMean	$\Delta \text{Var.}$	$\Delta \text{Skew.}$	$\Delta \text{Kurt.}$	Lap.	MC
0.01	−0.0278	0.0186	−0.1502	−0.0709	1.3105	0.06	3.38
0.10	−0.0210	0.0271	−0.0120	−0.0307	0.8078	0.06	3.29
0.25	−0.0154	0.0130	−0.0043	−0.0155	0.6354	0.06	3.32
0.50	−0.0081	0.0060	−0.0041	−0.0182	0.2256	0.06	3.29
0.75	−0.0026	0.0012	−0.0023	−0.0195	0.1239	0.06	3.30
1.00	−0.0008	−0.0004	−0.0015	−0.0227	0.0731	0.06	3.31
1.50	−0.0001	−0.0011	−0.0010	−0.0256	0.0378	0.06	3.31
2.00	0.0000	−0.0012	−0.0009	−0.0268	0.0218	0.06	3.32

Table 5: Laplace minus MC differences ($\Delta = \text{Lap.} - \text{MC}$) and wall-clock computation time. Differences shrink as h_0 increases because Monte Carlo becomes more reliable when many paths survive. The large $\Delta \text{Kurt.}$ at $h_0 = 0.01$ reflects MC noise from path scarcity ($\approx 2,700$ surviving paths out of 10^5); Δp_{surv} is negative throughout because MC over-counts surviving paths at finite sample sizes. The Laplace method covers five Talbot inversions ($k = 0, \dots, 4$); the $\approx 55\times$ speed-up over MC is uniform across all h_0 values.

synchronised with the underlying asset processes.

Zero slippage at liquidation. The absorbing-barrier assumption implies that collateral is liquidated at the mid-price prevailing at τ . In practice a keeper must sell the collateral into on-chain liquidity pools, and the effective execution price is depressed by slippage on that trade. For large positions this effect can be material; [Qin et al. \(2021\)](#) document cases where slippage exceeded the liquidation bonus itself.

No liquidation bonus recovery. The zero-recovery proxy (Remark 1) ignores the keeper’s liquidation bonus, which reduces the borrower’s net recovery. Incorporating an explicit recovery fraction $1 - \ell$ on liquidated paths — where ℓ is the protocol bonus rate — would add a second term to Φ in (6) but would not alter the structure of the resolvent, since the bonus affects only paths on $\{\tau \leq T\}$.

Static collateral management. The investor posts an initial collateral position and makes no further adjustments before T . A sophisticated participant may top up collateral or partially unwind the short leg as H_t approaches one, dynamically managing the distance to the boundary. Incorporating such controls would convert the problem into a stochastic optimal control problem over a reflected or absorbed process.

Deterministic interest rates. DeFi lending protocols charge a floating borrow rate updated algorithmically as a function of pool utilisation ([Gudgeon et al., 2020](#)). The model absorbs the funding cost into the effective drift μ_2^{eff} , treating the rate as deterministic and fixed at inception. Stochastic interest rates would couple the short-leg dynamics to a latent utilisation process.

No funding constraints. The analysis assumes the investor can hold the initial position to horizon T without receiving collateral top-up demands. A capital-constrained investor who cannot post additional margin when required may be forced to close the position before τ , introducing an additional stopping time that is not captured by the model.

7.2. Practical implementation

Despite the above limitations, the semi-analytic framework provides a tractable and fast starting point for position sizing. A natural two-step procedure is the following.

Step 1 — survival constraint. Fix a maximum tolerated liquidation probability $\bar{p}$ over the horizon T , for example $\bar{p} = 5\%$. The survival probability $p_{\text{surv}}(h_0)$ is monotonically increasing in h_0 (Table 4): at

$h_0 = 0.50$ survival already exceeds 83% under the benchmark parameters. Inverting this map identifies the minimum admissible health buffer $\underline{h}_0$ satisfying $p_{\text{liq}}(T, 0; \underline{h}_0) \leq \bar{p}$. The Laplace method evaluates p_{surv} in ≈ 60 ms, so the inversion is a fast scalar root-find.

Step 2 — conditional upside maximisation. Over the feasible set $h_0 \geq \underline{h}_0$, maximise the conditional expected payoff, or a mean–variance functional, using the semi-analytic moments. Because the moments are noise-free and differentiable in h_0 , a standard one-dimensional line search converges in a handful of evaluations with no simulation variance.

The output of this procedure is the *optimal initial health buffer* h_0^* : the smallest cushion above the liquidation boundary that simultaneously satisfies the survival constraint and maximises conditional performance. The corresponding initial leverage is then read off from (4) as $L_0^* = e^{h_0^*} / (e^{h_0^*} - b)$; leverage is a consequence of the health-buffer choice, not the primary control variable.

Figure 2 visualises the full tradeoff in three dimensions. The vertical axis is the conditional mean $\mathbb{E}[\Pi_T \mid \tau > T]$; the two horizontal axes are the conditional variance $\text{Var}(\Pi_T \mid \tau > T)$ and the liquidation probability $p_\ell = 1 - p_{\text{surv}}$. The curve is parametrised by $h_0 \in [0.04, 2.00]$: as h_0 falls the strategy takes on more leverage, pushing the curve toward higher variance and higher p_ℓ , while the conditional mean rises. Projections onto all three coordinate planes are shown as solid grey curves: the floor ($z = \text{const}$) gives the variance–liquidation tradeoff; the left wall ($x = \text{const}$) gives the liquidation–upside tradeoff; the back wall ($p_\ell = 0$) gives the classical mean–variance frontier conditional on near-certain survival.

Appendix A: Proofs

Proof of Lemma 3.1. Step 1: Laplace transform under $\mathbb{P}^{(k)}$. Because $Z_t = X_{1,t} - X_{2,t}$ and the density on $\mathcal{F}_t$ equals $e^{kX_{2,t} - t\Psi(0, -ik)}$,

$$\mathbb{E}_{\mathbb{P}^{(k)}}[e^{sZ_t}] = e^{-t\Psi(0, -ik)} \mathbb{E}_{\mathbb{P}}[e^{sX_{1,t} + (k-s)X_{2,t}}] = e^{t[\Psi(-is, -i(k-s)) - \Psi(0, -ik)]}, \quad (26)$$

where we used $\mathbb{E}_{\mathbb{P}}[e^{\alpha X_{1,t} + \beta X_{2,t}}] = e^{t\Psi(-i\alpha, -i\beta)}$. Since the Radon–Nikodým density is exponential-linear in $(X_{1,t}, X_{2,t})$, Girsanov’s theorem for Lévy processes guarantees that Z remains a Lévy process under $\mathbb{P}^{(k)}$ with exponent

$$\psi_Z^{(k)}(s) = \Psi(-is, -i(k-s)) - \Psi(0, -ik). \quad (27)$$

Step 2: Expand and simplify. Substituting (12) with $(u, v) = (-is, -i(k-s))$, and noting $u^2 = -s^2$, $v^2 = -(k-s)^2$, $uv = -s(k-s)$, and $\phi_{J_i}(-i\cdot) = M_i(\cdot)$:

$$\begin{aligned} \Psi(-is, -i(k-s)) &= \mu_1^{\text{eff}}s + \mu_2^{\text{eff}}(k-s) + \frac{1}{2}\sigma_1^2s^2 + \frac{1}{2}\sigma_2^2(k-s)^2 + \rho\sigma_1\sigma_2s(k-s) \\ &\quad + \lambda_1(M_1(s) - 1) + \lambda_2(M_2(k-s) - 1). \end{aligned} \quad (28)$$

Setting $s = 0$ in (28) gives $\Psi(0, -ik) = \mu_2^{\text{eff}}k + \frac{1}{2}\sigma_2^2k^2 + \lambda_2(M_2(k) - 1)$. Subtracting and collecting terms by type:

Drift. $\mu_1^{\text{eff}}s + \mu_2^{\text{eff}}(k-s) - \mu_2^{\text{eff}}k = (\mu_1^{\text{eff}} - \mu_2^{\text{eff}})s$.

Diffusion. Expanding $(k-s)^2 = k^2 - 2ks + s^2$ and $(k-s) = k-s$:

$$\begin{aligned} &\frac{1}{2}\sigma_1^2s^2 + \frac{1}{2}\sigma_2^2(k-s)^2 + \rho\sigma_1\sigma_2s(k-s) - \frac{1}{2}\sigma_2^2k^2 \\ &= \frac{1}{2}(\sigma_1^2 + \sigma_2^2 - 2\rho\sigma_1\sigma_2)s^2 - k(\sigma_2^2 - \rho\sigma_1\sigma_2)s = \frac{1}{2}\sigma_Z^2s^2 - k(\sigma_2^2 - \rho\sigma_1\sigma_2)s. \end{aligned}$$

Jumps. $\lambda_1(M_1(s) - 1) + \lambda_2(M_2(k-s) - 1) - \lambda_2(M_2(k) - 1) = \lambda_1(M_1(s) - 1) + \lambda_2(M_2(k-s) - M_2(k))$.

Combining, the drift coefficient is $\mu_Z^{(k)} := (\mu_1^{\text{eff}} - \mu_2^{\text{eff}}) - k(\sigma_2^2 - \rho\sigma_1\sigma_2)$, which establishes (13). The eigenrelation (14) follows because for any Lévy process with exponent ψ , the generator satisfies $\mathcal{L}e^{sz} = \psi(s)e^{sz}$, as can be seen by differentiating $\mathbb{E}[e^{sZ_t}] = e^{t\Psi(s)}$ with respect to t at $t = 0$. $\square$

Proof of Lemma 5.1. Substitute the forcing expansion into the right-hand side of (18):

$$(q - \mathcal{L}^{(k)})\widehat{U}_{\text{part}} = \sum_{j=0}^k c_{k,j}(h_0) e^{jz}.$$

Since the sum is finite and $\mathcal{L}^{(k)}$ is linear, it suffices to find, for each j , a function v_j satisfying

$$(q - \mathcal{L}^{(k)})v_j(z) = c_{k,j}(h_0) e^{jz}.$$

Propose the exponential ansatz $v_j(z) = A_j e^{jz}$. By the eigenrelation (14),

$$\mathcal{L}^{(k)}[A_j e^{jz}] = A_j \psi_Z^{(k)}(j) e^{jz},$$

so

$$(q - \mathcal{L}^{(k)})[A_j e^{jz}] = A_j (q - \psi_Z^{(k)}(j)) e^{jz}.$$

Matching to $c_{k,j}(h_0) e^{jz}$ and using Assumption 5.1(iii) to ensure $q - \psi_Z^{(k)}(j) \neq 0$ gives

$$A_j = \frac{c_{k,j}(h_0)}{q - \psi_Z^{(k)}(j)}.$$

Summing over $j = 0, \dots, k$ yields (20). □

Proof of Lemma 5.2. Step 1: exponential ansatz. Seek solutions of the form $u(z) = e^{\gamma z}$. By the eigenrelation (14),

$$\mathcal{L}^{(k)} e^{\gamma z} = \psi_Z^{(k)}(\gamma) e^{\gamma z},$$

so $(q - \mathcal{L}^{(k)})e^{\gamma z} = (q - \psi_Z^{(k)}(\gamma))e^{\gamma z}$. This vanishes if and only if γ satisfies the characteristic equation

$$\psi_Z^{(k)}(\gamma) = q.$$

Step 2: root count. The exponent $\psi_Z^{(k)}$ is the sum of a quadratic term $\frac{1}{2}\sigma_Z^2 \gamma^2$ (non-degenerate by Assumption 5.1(i)) and rational terms from the Kou jumps. Clearing denominators yields a degree-six polynomial in γ , so there are exactly six roots $\gamma_1, \dots, \gamma_6$ (counted with multiplicity). Under Assumption 5.1(ii) all six are simple and split as $\text{Re}(\gamma_{1,2,3}) > 0$ and $\text{Re}(\gamma_{4,5,6}) < 0$.

Step 3: admissibility. On the half-line $z > -h_0$, the solution must remain bounded as $z \rightarrow +\infty$. Since $e^{\gamma z} \rightarrow \infty$ whenever $\text{Re}(\gamma) > 0$, the roots $\gamma_1, \gamma_2, \gamma_3$ are inadmissible. Only $\gamma_4, \gamma_5, \gamma_6$ are retained.

Step 4: centering and general solution. Writing $e^{\gamma_{m+3}z} = e^{\gamma_{m+3}(z+h_0)} \cdot e^{-\gamma_{m+3}h_0}$ and absorbing the constant factor into the free coefficient, the general admissible homogeneous solution is

$$\widehat{U}_{\text{hom}}(q, z; h_0) = \sum_{m=1}^3 B_m^{(k)}(q; h_0) e^{\gamma_{m+3}(q)(z+h_0)},$$

where $B_1^{(k)}, B_2^{(k)}, B_3^{(k)}$ are free constants to be determined by the boundary conditions. □

Proof of Proposition 5.3. Step 1: boundary conditions from killing. The killing condition $\widehat{U}(q, z; h_0) = 0$ for $z \leq -h_0$ imposes, at $z = -h_0^+$,

$$(i) \quad \widehat{U}(q, -h_0^+; h_0) = 0;$$

$$(ii) \quad K_{j,-}^{(k)} \left[\widehat{U}(q, \cdot; h_0) \right] (-h_0^+) = 0, \quad j = 1, 2.$$

Condition (i) is continuity at the barrier. Conditions (ii) arise because $K_{j,-}^{(k)}[f](z) = \int_{-\infty}^0 f(z+y) r_{j,-}^{(k)} e^{r_{j,-}^{(k)} y} dy$ integrates f over landing points $z+y < z$; for $z = -h_0^+$ every landing point satisfies $z+y < -h_0$, where $\widehat{U} = 0$, so the integral vanishes.

Step 2: substitute the full solution. Write $\widehat{U} = \widehat{U}_{\text{part}} + \widehat{U}_{\text{hom}}$ and insert into each condition. Since the conditions are linear, each splits as

$$[\text{condition on } \widehat{U}_{\text{hom}}] = -[\text{condition on } \widehat{U}_{\text{part}}].$$

Step 3: condition (i). Evaluating at $z = -h_0$ and using $e^{\gamma_{m+3}(q) \cdot 0} = 1$,

$$\widehat{U}_{\text{hom}}(q, -h_0; h_0) = \sum_{m=1}^3 B_m^{(k)} = -\widehat{U}_{\text{part}}(q, -h_0; h_0),$$

which is the first row of the system.

Step 4: conditions (ii). Apply $K_{j,-}^{(k)}$ to $\widehat{U}_{\text{hom}}$. By the phase-operator formula (15),

$$K_{j,-}^{(k)} \left[e^{\gamma_{m+3}(\cdot + h_0)} \right] (-h_0) = \frac{r_{j,-}^{(k)}}{r_{j,-}^{(k)} + \gamma_{m+3}(q)} e^{\gamma_{m+3}(q) \cdot 0} = \frac{r_{j,-}^{(k)}}{r_{j,-}^{(k)} + \gamma_{m+3}(q)},$$

giving row $j+1$ of the system:

$$\sum_{m=1}^3 \frac{r_{j,-}^{(k)}}{r_{j,-}^{(k)} + \gamma_{m+3}(q)} B_m^{(k)} = -K_{j,-}^{(k)} \left[\widehat{U}_{\text{part}}(q, \cdot; h_0) \right] (-h_0).$$

Applying (15) to $\widehat{U}_{\text{part}}(q, z; h_0) = \sum_j \frac{c_{k,j}}{q - \psi_Z^{(k)}(j)} e^{jz}$ yields the explicit entries of $\mathbf{b}^{(k)}$ stated above.

Step 5: matrix form and solution. The three conditions assemble into $\mathbf{M}_{\text{bar}}^{(k)}(q) \mathbf{B}^{(k)} = -\mathbf{b}^{(k)}$. By Lemma 5.4, $\mathbf{M}_{\text{bar}}^{(k)}(q)$ is invertible, so

$$\mathbf{B}^{(k)}(q; h_0) = -(\mathbf{M}_{\text{bar}}^{(k)}(q))^{-1} \mathbf{b}^{(k)}(q; h_0). \quad \square$$

Proof of Lemma 5.4. Subtracting row 1 from rows 2 and 3 replaces entry (i, m) for $i = 2, 3$ by $-\gamma_{m+3}(q)/(r_{i-1,-}^{(k)} + \gamma_{m+3}(q))$. Pulling out -1 from each of these two rows (net sign $(-1)^2 = 1$) and then factoring $\gamma_{m+3}(q)$ from each column gives

$$\det(\mathbf{M}_{\text{bar}}^{(k)}) = \gamma_4 \gamma_5 \gamma_6 \det \left(\frac{1}{x_i + \gamma_{j+3}(q)} \right)_{1 \leq i, j \leq 3},$$

where $x_1 := 0$, $x_2 := r_{1,-}^{(k)}$, $x_3 := r_{2,-}^{(k)}$. The right-hand factor is a Cauchy determinant; the Cauchy formula yields

$$\det \left(\frac{1}{x_i + \gamma_{j+3}(q)} \right) = \frac{\prod_{1 \leq i < i' \leq 3} (x_{i'} - x_i) \cdot \prod_{4 \leq m < m' \leq 6} (\gamma_{m'} - \gamma_m)}{\prod_{i=1}^3 \prod_{m=4}^6 (x_i + \gamma_m)}.$$

Since $x_1 = 0$, the factor $\prod_{m=4}^6 (x_1 + \gamma_m) = \gamma_4 \gamma_5 \gamma_6$ cancels with the prefactor, yielding a ratio whose every factor is nonzero: $r_{1,-}^{(k)} > 0$ since $\eta_{1,-} > 0$; $r_{2,-}^{(k)} > 0$ since $k\eta_{2,+} < 1$; $(r_{2,-}^{(k)} - r_{1,-}^{(k)}) \neq 0$ by (22); the γ_m are pairwise distinct by Assumption 5.1(ii); and $r_{j,-}^{(k)} + \gamma_m \neq 0$ because $-r_{j,-}^{(k)}$ is a pole of $\psi_Z^{(k)}$, so $\psi_Z^{(k)}(-r_{j,-}^{(k)}) = \infty \neq q$, hence $\gamma_m \neq -r_{j,-}^{(k)}$. $\square$

Proof of Proposition 5.5. Step 1: resolvent equation. Taking the Laplace transform of the PIDE $\partial_t p_{\text{surv}} = \mathcal{L}^{(0)} p_{\text{surv}}$ with initial condition $p_{\text{surv}}(0, z; h_0) = 1$ and killing condition $p_{\text{surv}} = 0$ for $z \leq -h_0$, exactly as in Section 5 with $g_k \equiv 1$, gives

$$(q - \mathcal{L}^{(0)}) \widehat{U}_{\text{surv}}(q, z; h_0) = 1, \quad z > -h_0.$$

Step 2: particular solution. The constant forcing $f \equiv 1$ is an eigenfunction of $\mathcal{L}^{(0)}$ with eigenvalue $\psi_Z^{(0)}(0) = 0$, so the particular solution is

$$\widehat{U}_{\text{surv,part}}(q, z) = \frac{1}{q - \psi_Z^{(0)}(0)} = \frac{1}{q}.$$

Step 3: homogeneous solution. By Lemma 5.2 applied with $k = 0$, the admissible homogeneous solution on $z > -h_0$ is

$$\widehat{U}_{\text{surv,hom}}(q, z; h_0) = C_1 e^{\gamma_4^{(0)}(q)(z+h_0)} + C_2 e^{\gamma_5^{(0)}(q)(z+h_0)} + C_3 e^{\gamma_6^{(0)}(q)(z+h_0)},$$

where $\gamma_4^{(0)}, \gamma_5^{(0)}, \gamma_6^{(0)}$ are the roots of $\psi_Z^{(0)}(\gamma) = q$ with $\text{Re}(\gamma) < 0$.

Step 4: boundary conditions. The killing condition imposes, at $z = -h_0^+$:

- (i) $\widehat{U}_{\text{surv}}(q, -h_0^+; h_0) = 0;$
- (ii) $K_{j,-}^{(0)}[\widehat{U}_{\text{surv}}](-h_0^+) = 0, \quad j = 1, 2,$

by the same argument as in Proposition 5.3: downward jumps from $-h_0^+$ land in the killed region.

Step 5: linear system. Substituting the full solution $\widehat{U}_{\text{surv}} = 1/q + \widehat{U}_{\text{surv,hom}}$ into each condition and applying (15) to the homogeneous terms yields

$$\mathbf{M}_{\text{surv}}(q) \mathbf{C}(q; h_0) = - \begin{pmatrix} 1/q \\ 1/q \\ 1/q \end{pmatrix},$$

where the right-hand side entries are $K_{j,-}^{(0)}[1/q](-h_0^+) = 1/q$ (since $K_{j,-}^{(0)}$ integrates to one against the exponential density). By Lemma 5.6, $\mathbf{M}_{\text{surv}}(q)$ is invertible, so the coefficients $\mathbf{C}$ are uniquely determined.

Step 6: inversion and liquidation probability. Evaluating at $Z_0 = 0$ and inverting via the Bromwich formula gives $p_{\text{surv}}(T, 0; h_0)$. The liquidation probability follows from the complement. $\square$

Proof of Lemma 5.6. The argument is identical to that of Lemma 5.4 with $k = 0$. The phase rates are $r_{1,-}^{(0)} = 1/\eta_{1,-}$ and $r_{2,-}^{(0)} = 1/\eta_{2,+}$, so condition (24) is equivalent to $\eta_{1,-} \neq \eta_{2,+}$. Each denominator factor is nonzero because $-r_{j,-}^{(0)}$ is a pole of $\psi_Z^{(0)}$, hence cannot equal any root $\gamma_m^{(0)}(q)$ of $\psi_Z^{(0)}(\cdot) = q$ for finite q . $\square$

References

- D.-M. Dang and P. A. Forsyth. Better than pre-commitment mean–variance portfolio allocation strategies: A semi-self-financing Hamilton–Jacobi–Bellman equation approach. *European Journal of Operational Research*, 250(3):827–841, 2016.
- L. Gudgeon, S. Werner, D. Perez, and W. J. Knottenbelt. Defi protocols for loanable funds: Interest rates, liquidity and market efficiency. In *Proceedings of the 2nd ACM Conference on Advances in Financial Technologies*, pages 92–112, 2020.

- L. Heimbach and Y. Huang. DeFi leverage. BIS Working Paper 1171, Bank for International Settlements, 2024.
- B. I. Jacobs, K. N. Levy, and D. Starer. Long-short portfolio management: An integrated approach. *Journal of Portfolio Management*, 25(2):23–32, 1999.
- S. G. Kou. A jump-diffusion model for option pricing. *Management Science*, 48(8):1086–1101, 2002.
- S. G. Kou and H. Wang. First passage times of a jump diffusion process. *Advances in Applied Probability*, 35(2):504–531, 2003.
- H. Markowitz. Portfolio selection. *Journal of Finance*, 7(1):77–91, 1952.
- K. Qin, L. Zhou, P. Gamito, P. Jovanovic, and A. Gervais. An empirical study of DeFi liquidations: Incentives, risks, and instabilities. In *Proceedings of the 21st ACM Internet Measurement Conference*, pages 336–350, 2021.
- A. Sepp. Analytical pricing of double-barrier options under a double-exponential jump diffusion process. *Applied Mathematical Finance*, 11(2):109–151, 2004. doi: 10.1080/1350486042000Step.
- Y. Shen and B. Zou. Mean-variance portfolio selection under kou jump diffusion, 2022. Working paper.
- A. Talbot. The accurate numerical inversion of Laplace transforms. *IMA Journal of Applied Mathematics*, 23(1):97–120, 1979.




Figure 2: Mean–variance–liquidation frontier of $\Pi_T | \{\tau > T\}$ for the benchmark parametrisation. Axes: $x = \text{Var}(\Pi_T | \tau > T)$, $y = p_\ell = 1 - p_{\text{surv}}$, $z = E[\Pi_T | \tau > T]$ (vertical). The curve is parametrised by $h_0 \in [0.04, 2.00]$; decreasing h_0 moves the curve toward higher variance, higher p_ℓ , and higher conditional mean. Grey projections on the three coordinate planes recover the pairwise two-dimensional tradeoffs. Script `jobs/frontier_analysis.py`.