



# Hamiltonian fluid dynamics and hydrodynamic stability

James Hawley

University of Waterloo

August 14, 2015

# Contents

Hamiltonian Fluid Dynamics

Stability Theory

Quasigeostrophic Model

Stability of Zonal Flows

References

# Classical Mechanics

# Classical Mechanics

## Newtonian Dynamics

# Classical Mechanics

## Newtonian Dynamics

- $m \frac{d^2 \mathbf{q}}{dt} = \mathbf{F}$
- $\mathbf{F} = -\nabla \Pi$
- $\mathbf{p} = \frac{d}{dt}(m\mathbf{q})$

# Classical Mechanics

## Newtonian Dynamics

- $m \frac{d^2 \mathbf{q}}{dt} = \mathbf{F}$
- $\mathbf{F} = -\nabla \Pi$
- $\mathbf{p} = \frac{d}{dt}(m\mathbf{q})$

## Hamiltonian Dynamics

# Classical Mechanics

## Newtonian Dynamics

- $m \frac{d^2 \mathbf{q}}{dt} = \mathbf{F}$
- $\mathbf{F} = -\nabla \Pi$
- $\mathbf{p} = \frac{d}{dt}(m\mathbf{q})$

## Hamiltonian Dynamics

- $\frac{d\mathbf{p}}{dt} = -\frac{\partial H}{\partial \mathbf{q}}$
- $\frac{d\mathbf{q}}{dt} = \frac{\partial H}{\partial \mathbf{p}}$
- $H = \mathbf{p}^2/2m + \Pi(\mathbf{q})$

# Fluids



# Fluids

## Navier-Stokes Equations

# Fluids

## Navier-Stokes Equations

- $\rho \frac{d\mathbf{u}}{dt} = \nabla \cdot \boldsymbol{\sigma} + \mathbf{F}$
- $\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{u} = 0$
- System of coupled, nonlinear, partial differential equations

# Fluids

## Navier-Stokes Equations

- $\rho \frac{d\mathbf{u}}{dt} = \nabla \cdot \boldsymbol{\sigma} + \mathbf{F}$
- $\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{u} = 0$
- System of coupled, nonlinear, partial differential equations

## Hamiltonian

# Fluids

## Navier-Stokes Equations

- $\rho \frac{d\mathbf{u}}{dt} = \nabla \cdot \boldsymbol{\sigma} + \mathbf{F}$
- $\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{u} = 0$
- System of coupled, nonlinear, partial differential equations

## Hamiltonian

- $\frac{d\mathbf{p}}{dt} = -\frac{\delta \mathcal{H}}{\delta \mathbf{q}}$
- $\frac{d\mathbf{q}}{dt} = \frac{\delta \mathcal{H}}{\delta \mathbf{p}}$
- $\mathcal{H} = \iint \frac{1}{2} \nabla \psi \cdot \nabla \psi \, dA - \sum_n \lambda_n \oint \mathbf{n} \cdot \nabla \psi \, ds$

# Hamiltonian Fluid Dynamics

# Hamiltonian Fluid Dynamics

System of PDEs

$$\mathbf{0} = \mathbf{F} \left( \mathbf{q}, \frac{\partial}{\partial x_i}, \frac{\partial}{\partial t} \right)$$

# Hamiltonian Fluid Dynamics

System of PDEs

$$\mathbf{0} = \mathbf{F} \left( \mathbf{q}, \frac{\partial}{\partial x_i}, \frac{\partial}{\partial t} \right)$$

is *Hamiltonian* if  $\exists \mathcal{H}, J$  such that

# Hamiltonian Fluid Dynamics

System of PDEs

$$\mathbf{0} = \mathbf{F} \left( \mathbf{q}, \frac{\partial}{\partial x_i}, \frac{\partial}{\partial t} \right)$$

is *Hamiltonian* if  $\exists \mathcal{H}, J$  such that

$$\mathcal{H} = \int_{\Omega} H(\mathbf{q}) d\mathbf{x}$$

is conserved, and solutions  $\mathbf{q}(x_i, t)$  satisfy



# Hamiltonian Fluid Dynamics

System of PDEs

$$\mathbf{0} = \mathbf{F} \left( \mathbf{q}, \frac{\partial}{\partial x_i}, \frac{\partial}{\partial t} \right)$$

is *Hamiltonian* if  $\exists \mathcal{H}, J$  such that

$$\mathcal{H} = \int_{\Omega} H(\mathbf{q}) d\mathbf{x}$$

is conserved, and solutions  $\mathbf{q}(x_i, t)$  satisfy

$$\mathbf{q}_t = \mathbf{J} \frac{\delta \mathcal{H}}{\delta \mathbf{q}}$$

# Variational Derivatives

# Variational Derivatives

$$\mathcal{F}(\mathbf{q}) = \int_{\Omega} F(\mathbf{q}) d\mathbf{x}$$

# Variational Derivatives

$$\mathcal{F}(\mathbf{q}) = \int_{\Omega} F(\mathbf{q}) d\mathbf{x}$$


$$\lim_{\epsilon \rightarrow 0} \frac{\partial}{\partial \epsilon} \mathcal{F}(\mathbf{q} + \epsilon \delta \mathbf{q}) \equiv \left\langle \frac{\delta \mathcal{F}}{\delta \mathbf{q}}, \delta \mathbf{q} \right\rangle \equiv \int_{\Omega} \frac{\delta \mathcal{F}}{\delta \mathbf{q}} \delta \mathbf{q} d\mathbf{x}$$

# Poisson Bracket

$$\begin{aligned}\frac{\partial \mathcal{F}}{\partial t} &= \int_{\Omega} \frac{\partial F}{\partial t} d\mathbf{x} \\ &= \int_{\Omega} \frac{\delta \mathcal{F}}{\delta \mathbf{q}}^{\top} \mathbf{q}_t d\mathbf{x} \\ &= \int_{\Omega} \frac{\delta \mathcal{F}}{\delta \mathbf{q}}^{\top} \mathbf{J} \frac{\delta \mathcal{H}}{\delta \mathbf{q}} d\mathbf{x} \\ &= \left\langle \frac{\delta \mathcal{F}}{\delta \mathbf{q}}, \mathbf{J} \frac{\delta \mathcal{H}}{\delta \mathbf{q}} \right\rangle \\ &\equiv \{\mathcal{F}, \mathcal{H}\}\end{aligned}$$




# Quasigeostrophic Model






# Quasigeostrophic Model



$$\psi_n = -U_n y$$

## Quasigeostrophic Model



$$\psi_n = -U_n y$$

$$q_1 = \frac{1}{\alpha_1} \nabla^2 \psi_1 + (\psi_2 - \psi_1) = (U_1 - U_2) y$$

$$q_2 = \frac{1}{\alpha_2} \nabla^2 \psi_2 + (\psi_1 - \psi_2) = (U_2 - U_1) y$$





## Quasigeostrophic Model

$$0 = \frac{\partial q_1}{\partial t} + D(\psi_1, q_1)$$

$$0 = \frac{\partial q_2}{\partial t} + D(\psi_2, q_2)$$

$$D(a, b) = \frac{\partial a}{\partial x} \frac{\partial b}{\partial y} - \frac{\partial a}{\partial y} \frac{\partial b}{\partial x}$$



## Quasigeostrophic Model

$$0 = \frac{\partial q_1}{\partial t} + D(\psi_1, q_1)$$

$$0 = \frac{\partial q_2}{\partial t} + D(\psi_2, q_2)$$

$$D(a, b) = \frac{\partial a}{\partial x} \frac{\partial b}{\partial y} - \frac{\partial a}{\partial y} \frac{\partial b}{\partial x}$$

$$\mathcal{H} = \frac{1}{2} \iint \sum_{n=1}^2 \left[ \frac{1}{\alpha_n} \vec{\nabla} \psi_n \cdot \vec{\nabla} \psi_n + 2C_n(q_n) \right] + (\psi_1 - \psi_2)^2 dA$$

# Second Variation

## Second Variation

$$\delta^2 \mathcal{H} = \iint \sum_n \left[ \frac{1}{\alpha_n} \vec{\nabla} \delta \psi_n \cdot \vec{\nabla} \delta \psi_n + \psi'_n \cdot (\delta q_n)^2 \right] + (\delta \psi_1 - \delta \psi_2)^2 dA$$

## Second Variation

$$\delta^2 \mathcal{H} = \iint \sum_n \left[ \frac{1}{\alpha_n} \vec{\nabla} \delta \psi_n \cdot \vec{\nabla} \delta \psi_n + \psi'_n \cdot (\delta q_n)^2 \right] + (\delta \psi_1 - \delta \psi_2)^2 dA$$

$$U_1(U_1 - U_2) = 0$$

$$U_2(U_2 - U_1) = 0$$

## References

1. G. Swaters. "Introduction to Hamiltonian Fluid Dynamics and Stability Theory". Monographs and Surveys in Pure and Applied Mathematics (1999).
2. M. E. McIntyre and T. G. Shepherd. "An exact local conservation theorem for finite-amplitude disturbances to non-parallel shear flows, with remarks on Hamiltonian structure and on Arnolds stability theorems". Journal of Fluid Mechanics (1987), vol. 181, pp. 527-565.
3. T. G. Shepherd. "Symmetries, conservation laws, and Hamiltonian structure in geophysical fluid dynamics". Advances in Geophysics (1990), vol. 32, pp. 287-338.
4. J. Pedlosky. "Geophysical Fluid Dynamics". Springer Study Edition (1992).