

Hamiltonian fluid dynamics and stability

James Hawley

University of Waterloo

August 14, 2015

Contents

Hamiltonian Dynamics

Stability

Quasigeostrophic Model

Stability of Zonal Flows

References



Classical Mechanics

Classical Mechanics

Newtonian Dynamics

Classical Mechanics

Newtonian Dynamics

- $m \frac{d^2 \mathbf{q}}{dt} = \mathbf{F}$
- $\mathbf{F} = -\nabla \Pi$
- Second order ODE

Classical Mechanics

Newtonian Dynamics

- $m \frac{d^2 \mathbf{q}}{dt} = \mathbf{F}$
- $\mathbf{F} = -\nabla \Pi$
- Second order ODE

Hamiltonian Dynamics

Classical Mechanics

Newtonian Dynamics

- $m \frac{d^2 \mathbf{q}}{dt^2} = \mathbf{F}$
- $\mathbf{F} = -\nabla \Pi$
- Second order ODE

Hamiltonian Dynamics

- $\frac{d\mathbf{p}}{dt} = -\frac{\partial H}{\partial \mathbf{q}}$
- $\frac{d\mathbf{q}}{dt} = \frac{\partial \mathcal{H}}{\partial \mathbf{p}}$
- System of first order ODEs

Fluids

Fluids

Navier-Stokes Equations

Fluids

Navier-Stokes Equations

- $\rho \frac{d\mathbf{u}}{dt} = \nabla \cdot \boldsymbol{\sigma} + \mathbf{F}$
- $\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{u} = 0$
- System of coupled, nonlinear, PDEs

Fluids

Navier-Stokes Equations

- $\rho \frac{d\mathbf{u}}{dt} = \nabla \cdot \boldsymbol{\sigma} + \mathbf{F}$
- $\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{u} = 0$
- System of coupled, nonlinear, PDEs

Hamiltonian

Fluids

Navier-Stokes Equations

- $\rho \frac{d\mathbf{u}}{dt} = \nabla \cdot \boldsymbol{\sigma} + \mathbf{F}$
- $\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{u} = 0$
- System of coupled, nonlinear, PDEs

Hamiltonian

- $\frac{d\mathbf{p}}{dt} = -\frac{\delta \mathcal{H}}{\delta \mathbf{q}}$
- $\frac{d\mathbf{q}}{dt} = \frac{\delta \mathcal{H}}{\delta \mathbf{p}}$
- More analytical tools to work with

Hamiltonian Fluid Dynamics

Hamiltonian Fluid Dynamics

System of PDEs

$$\mathbf{0} = \mathbf{F} \left(\mathbf{q}, \frac{\partial}{\partial x_i}, \frac{\partial}{\partial t} \right)$$

Hamiltonian Fluid Dynamics

System of PDEs

$$\mathbf{0} = \mathbf{F} \left(\mathbf{q}, \frac{\partial}{\partial x_i}, \frac{\partial}{\partial t} \right)$$

is *Hamiltonian* if $\exists \mathcal{H}, J$ such that

Hamiltonian Fluid Dynamics

System of PDEs

$$\mathbf{0} = \mathbf{F} \left(\mathbf{q}, \frac{\partial}{\partial x_i}, \frac{\partial}{\partial t} \right)$$

is *Hamiltonian* if $\exists \mathcal{H}, J$ such that

$$\mathcal{H} = \int_{\Omega} H(\mathbf{q}) d\mathbf{x}$$

is conserved, and solutions $\mathbf{q}(x_i, t)$ satisfy

Hamiltonian Fluid Dynamics

System of PDEs

$$\mathbf{0} = \mathbf{F} \left(\mathbf{q}, \frac{\partial}{\partial x_i}, \frac{\partial}{\partial t} \right)$$

is *Hamiltonian* if $\exists \mathcal{H}, J$ such that

$$\mathcal{H} = \int_{\Omega} H(\mathbf{q}) d\mathbf{x}$$

is conserved, and solutions $\mathbf{q}(x_i, t)$ satisfy

$$\mathbf{q}_t = \mathbf{J} \frac{\delta \mathcal{H}}{\delta \mathbf{q}}$$

Derivatives

Derivatives

$$\mathcal{F}(\mathbf{q}) = \int_{\Omega} F(\mathbf{q}) d\mathbf{x}$$

Derivatives

$$\mathcal{F}(\mathbf{q}) = \int_{\Omega} F(\mathbf{q}) d\mathbf{x}$$


$$\lim_{\epsilon \rightarrow 0} \frac{\partial}{\partial \epsilon} \mathcal{F}(\mathbf{q} + \epsilon \delta \mathbf{q}) \equiv \left\langle \frac{\delta \mathcal{F}}{\delta \mathbf{q}}, \delta \mathbf{q} \right\rangle \equiv \int_{\Omega} \frac{\delta \mathcal{F}}{\delta \mathbf{q}} \delta \mathbf{q} d\mathbf{x}$$

Poisson Bracket

$$\begin{aligned}\frac{\partial \mathcal{F}}{\partial t} &= \int_{\Omega} \frac{\partial F}{\partial t} d\mathbf{x} \\ &= \int_{\Omega} \frac{\delta \mathcal{F}}{\delta \mathbf{q}}^{\top} \mathbf{q}_t d\mathbf{x} \\ &= \int_{\Omega} \frac{\delta \mathcal{F}}{\delta \mathbf{q}}^{\top} \mathbf{J} \frac{\delta \mathcal{H}}{\delta \mathbf{q}} d\mathbf{x} \\ &= \left\langle \frac{\delta \mathcal{F}}{\delta \mathbf{q}}, \mathbf{J} \frac{\delta \mathcal{H}}{\delta \mathbf{q}} \right\rangle \\ &\equiv \{\mathcal{F}, \mathcal{H}\}\end{aligned}$$




Quasigeostrophic Model






Quasigeostrophic Model



$$\psi_n = -U_n y$$



Quasigeostrophic Model



$$\psi_n = -U_n y$$

$$q_1 = \frac{1}{\alpha_1} \nabla^2 \psi_1 + (\psi_2 - \psi_1) = (U_1 - U_2) y$$

$$q_2 = \frac{1}{\alpha_2} \nabla^2 \psi_2 + (\psi_1 - \psi_2) = (U_2 - U_1) y$$



Quasigeostrophic Model

$$0 = \frac{\partial q_1}{\partial t} + D(\psi_1, q_1)$$

$$0 = \frac{\partial q_2}{\partial t} + D(\psi_2, q_2)$$

$$D(a, b) = \frac{\partial a}{\partial x} \frac{\partial b}{\partial y} - \frac{\partial a}{\partial y} \frac{\partial b}{\partial x}$$



Quasigeostrophic Model

$$0 = \frac{\partial q_1}{\partial t} + D(\psi_1, q_1)$$

$$0 = \frac{\partial q_2}{\partial t} + D(\psi_2, q_2)$$

$$D(a, b) = \frac{\partial a}{\partial x} \frac{\partial b}{\partial y} - \frac{\partial a}{\partial y} \frac{\partial b}{\partial x}$$

$$\mathcal{H} = \frac{1}{2} \iint \sum_{n=1}^2 \left[\frac{1}{\alpha_n} \vec{\nabla} \psi_n \cdot \vec{\nabla} \psi_n + 2C_n(q_n) \right] + (\psi_1 - \psi_2)^2 dA$$

Second Variation

Second Variation

$$\delta^2 \mathcal{H} = \iint \sum_n \left[\frac{1}{\alpha_n} \vec{\nabla} \delta \psi_n \cdot \vec{\nabla} \delta \psi_n + \psi'_n \cdot (\delta q_n)^2 \right] + (\delta \psi_1 - \delta \psi_2)^2 dA$$

Second Variation

$$\delta^2 \mathcal{H} = \iint \sum_n \left[\frac{1}{\alpha_n} \vec{\nabla} \delta \psi_n \cdot \vec{\nabla} \delta \psi_n + \psi'_n \cdot (\delta q_n)^2 \right] + (\delta \psi_1 - \delta \psi_2)^2 dA$$

$$U_n \frac{dq_n}{dy} > 0, \quad n = 1, 2$$

Second Variation

$$\delta^2 \mathcal{H} = \iint \sum_n \left[\frac{1}{\alpha_n} \vec{\nabla} \delta \psi_n \cdot \vec{\nabla} \delta \psi_n + \psi'_n \cdot (\delta q_n)^2 \right] + (\delta \psi_1 - \delta \psi_2)^2 dA$$

$$U_n \frac{dq_n}{dy} > 0, \quad n = 1, 2$$

$$U_1(U_1 - U_2) > 0$$

$$U_2(U_2 - U_1) > 0$$

References

1. G. Swaters. "Introduction to Hamiltonian Fluid Dynamics and Stability Theory". Monographs and Surveys in Pure and Applied Mathematics (1999).
2. M. E. McIntyre and T. G. Shepherd. "An exact local conservation theorem for finite-amplitude disturbances to non-parallel shear flows, with remarks on Hamiltonian structure and on Arnolds stability theorems". Journal of Fluid Mechanics (1987), vol. 181, pp. 527-565.
3. T. G. Shepherd. "Symmetries, conservation laws, and Hamiltonian structure in geophysical fluid dynamics". Advances in Geophysics (1990), vol. 32, pp. 287-338.
4. Jacques Vanneste. "Explosive resonant interaction of baroclinic Rossby waves and stability of multilayer quasi-geostrophic flow". Journal of Fluid Mechanics (1995), vol. 291, pp. 83-107.