

Insulin sensitivity in Chinese ovo-lactovegetarians compared with omnivores.

AIM: To compare the insulin sensitivity indices between Chinese vegetarians and omnivores. **METHODS:** The study included 36 healthy volunteers (vegetarian, n=19; omnivore, n=17) who had normal fasting plasma glucose levels. Each participant completed an insulin suppression test. We compared steady-state plasma glucose (SSPG), fasting insulin, the homeostasis model assessment for insulin sensitivity (HOMA-IR and HOMA %S) and beta-cell function (HOMA %beta) between the groups. We also tested the correlation of SSPG with years on a vegetarian diet. **RESULTS:** The omnivore subjects were younger than the vegetarians (55.7 $\pm$ 3.7 vs 58.6 $\pm$ 3.6 year of age, P=0.022). There was no difference between the two groups in sex, blood pressure, renal function tests and lipid profiles. The omnivores had higher serum uric acid levels than vegetarians (5.25 $\pm$ 0.84 vs 4.54 $\pm$ 0.75 mg/dl, P=0.011). The results of the indices were different between omnivores and vegetarians (SSPG (mean $\pm$ s.d.) 105.4 $\pm$ 10.2 vs 80.3 $\pm$ 11.3 mg/dl, P<0.001; fasting insulin, 4.06 $\pm$ 0.77 vs 3.02 $\pm$ 1.19 microU/ml, P=0.004; HOMA-IR, 6.75 $\pm$ 1.31 vs 4.78 $\pm$ 2.07, P=0.002; HOMA %S, 159.2 $\pm$ 31.7 vs 264.3 $\pm$ 171.7%, P=0.018) except insulin secretion index, HOMA %beta (65.6 $\pm$ 18.0 vs 58.6 $\pm$ 14.8%, P=0.208). We found a clear linear relation between years on a vegetarian diet and SSPG (r=-0.541, P=0.017). **CONCLUSIONS:** The vegetarians were more insulin sensitive than the omnivore counterparts. The degree of insulin sensitivity appeared to be correlated with years on a vegetarian diet.