

A multicomponent intervention reduces body weight and cardiovascular risk at a GEICO corporate site.

.

PURPOSE: To determine whether a multicomponent nutrition intervention program at a corporate site reduces body weight and improves other cardiovascular risk factors in overweight individuals. DESIGN: Prospective clinical intervention study. SUBJECTS/SETTING: Employees of the Government Employees Insurance Company (GEICO) (N = 113), aged 21 to 65 years, with a body mass index ≥ 25 kg/m² and/or previous diagnosis of type 2 diabetes. INTERVENTION: A 22-week intervention including a low-fat, vegan diet. MEASURES: Changes in body weight, anthropometric measures, blood pressure, lipid profile, and dietary intake. ANALYSIS: Multivariate analyses of variance were calculated for clinical and nutrient measures, followed by univariate analyses of variance, to determine the significance of differences between groups in changes over time. RESULTS: Intervention-group participants experienced greater weight changes compared with control-group participants (mean, -5.1 [SE, .6] kg vs. +.1 [SE, .6] kg, $p < .0001$), as well as greater changes in waist circumference (mean, -4.7 [SE, .6] cm vs. +.8 [SE, .6] cm, $p < .0001$) and waist-to-hip ratio (mean, -.006 [SE, .003] vs. +.014 [SE, .005], $p = .0007$). Weight loss of 5% of body weight was more frequently observed in the intervention group (48.5%) compared with the control group (11.1%) ($\chi^2[1, N = 113] = 16.99$, $p < .0001$). CONCLUSIONS: Among individuals volunteering for a 22-week worksite research study, an intervention using a low-fat, vegan diet effectively reduced body weight and waist circumference.