

Egg consumption and risk of type 2 diabetes in older adults

Background: Type 2 diabetes (T2D) remains an important public health issue in the United States. There are limited and inconsistent data on the association between egg consumption and fasting glucose or incident diabetes. **Objectives:** We assessed the association between egg intake and incident diabetes in older adults. **Design:** In this prospective study of 3898 men and women from the Cardiovascular Health Study (1989–2007), we assessed egg consumption by using a picture-sorted food questionnaire and ascertained incident T2D annually by using information on hypoglycemic agents and plasma glucose. We used Cox proportional hazards models to estimate adjusted relative risks. **Results:** During a mean follow-up of 11.3 y, 313 new cases of T2D occurred. Crude incidence rates of T2D were 7.39, 6.83, 7.00, 6.72, and 12.20 per 1000 person-years in people who reported egg consumption of never, <1 egg/mo, 1–3 eggs/mo, 1–4 eggs/wk, and almost daily, respectively. In multivariable-adjusted models, there was no association between egg consumption and increased risk of T2D in either sex and overall. In a secondary analysis, dietary cholesterol was not associated with incident diabetes (P for trend = 0.47). In addition, egg consumption was not associated with clinically meaningful differences in fasting glucose, fasting insulin, or measures of insulin resistance despite small absolute analytic differences that were significant. **Conclusion:** In this cohort of older adults with limited egg intake, there was no association between egg consumption or dietary cholesterol and increased risk of incident T2D.