

An Analysis of Sigmoid function

Sigmoid function

Sigmoid function -- Part 1

where S is an increasing sigmoid function, $f(x)$ is a transformation of the independent variable, and A and B are constants controlling scaling and translation.

Sigmoid function -- Part 2

$f(x) = \frac{1}{2} \left(1 + e^{-2mx} \right)$, $|x| < 1$ $\operatorname{sgn}(x) |x| \geq 1$ $= \tanh\left(\frac{m}{2} (1 - x^2)\right)$, $|x| < 1$ $\operatorname{sgn}(x) |x| \geq 1$ using the hyperbolic tangent mentioned above. Here, m is a free parameter encoding the slope at $x = 0$, which must be greater than or equal to $\sqrt{3}$ because any smaller value will result in a function with multiple inflection points, which is therefore not a true sigmoid. This function is unusual because it actually attains the limiting values of -1 and 1 within a finite range, meaning that its value is constant at -1 for all $x \leq -1$ and at 1 for all $x \geq 1$. Nonetheless, it is smooth (infinitely differentiable, C^∞) everywhere, including at $x = \pm 1$.

Sigmoid function -- Part 3

1st kind A unitary sigmoid function of the first kind is a bounded increasing function that approaches its lower and upper asymptotes monotonically, without oscillation. This class includes many of the standard sigmoid functions used in statistics, biomathematics, and engineering, such as the logistic function and related generalizations.

Sigmoid function -- Part 4

M0: Construction method for sigmoid functions not evident or intuitive M1: Inverse of singularity functions M2: Sigmoid functions of embedded positive functions M3: Rising a sigmoid function to the power M4: Exponentiating a sigmoid function M5: Symmetric sigmoid functions derived from asymmetric ones M6: Sigmoid functions of the reciprocal independent variable M7: Embedding a sigmoid function into other function M8: Sum of sigmoid functions M9: Multiplication of sigmoid functions M10: Integral of the product of an increasing and a decreasing function M11: Derivation from lambda (bell-shaped) functions

M12: Integration of lambda (bell-shaped) function M13: Integration of the sum of lambda (bell-shaped) functions M14: Integration of the product of two lambda (bell-shaped) functions M15: Integration of the difference of two shifted sigmoid functions M16: Integration of the product of two shifted sigmoid functions M17: Convolution of sigmoid functions M18: Integration of the product of lambda and sigmoid function M19: Solutions of ordinary differential equations M20: Solutions of partial differential equation (PDE) M21: Solutions of functional differential equation (FDE) M22: Sum of a sigmoid function and some derivatives M23: Combination of sigmoid functions, its derivative and integral M24: Filtering sigmoid functions M25: Special cases of Gauss hypergeometric functions M26: Feedback closed-loop systems M27: Recursive functions M28: Recursive time-delayed feed-forward loops M29: Solutions of stochastic differential equation M30: Chaotic sigmoid functions Consult reference for more details.

Sigmoid function -- Part 5

Gudermannian function $f(x) = \operatorname{gd}(x) = \int_0^x \operatorname{sech} t \, dt = 2 \arctan(\tanh(x/2))$ $f(x) = \operatorname{gd}(x) = \int_0^x \operatorname{sech} t \, dt = 2 \arctan(\tanh(x/2))$

Sigmoid function -- Part 6

Up to shifts and scaling, many sigmoids are special cases of $f(x) = \phi(\phi(x, \beta), \alpha)$, where $\phi(x, \lambda) = \frac{1}{1 + \lambda x}$ $\lambda \neq 0$ $e^{-x/\lambda} = 0$ is the inverse of the negative Box-Cox transformation, and $\alpha < 1$ and $\beta < 1$ are shape parameters. Smooth transition function normalized to $(-1, 1)$:

Sigmoid function -- Part 7

3rd kind A unitary sigmoid function of the third kind is a bounded increasing function that oscillates near both the lower and upper asymptotes. These functions retain the global shape of a sigmoid curve but exhibit oscillatory behavior in the vicinity of both limiting states.

Sigmoid function -- Part 8

A sigmoid function is any mathematical function whose graph has a characteristic S-shaped or sigmoid curve. A common example of a sigmoid function is the logistic function. Other sigmoid functions are given in the Examples section. In some

An Analysis of Sigmoid function

Sigmoid function

fields, most notably in the context of artificial neural networks, the term "sigmoid function" is used as a synonym for "logistic function". Special cases of sigmoid functions include the Gompertz curve (used in modeling systems that saturate at large values of x) and the ogive curve (used in the spillway of some dams). Sigmoid functions have domain of all real numbers, with return (response) value commonly monotonically increasing but could be decreasing. Sigmoid functions most often show a return value (y axis) in the range 0 to 1. Another commonly used range is from -1 to 1. There is also the Heaviside step function, which instantaneously transitions between 0 and 1. A wide variety of sigmoid functions including the logistic and hyperbolic tangent functions have been used as the activation function of artificial neurons. Sigmoid curves are also common in statistics as cumulative distribution functions (which go from 0 to 1), such as the integrals of the logistic density, the normal density, and Student's t probability density functions. The logistic sigmoid function is invertible, and its inverse is the logit function.

Sigmoid function -- Part 9

Theory In mathematics, a unitary sigmoid function is a bounded sigmoid-type function normalized to the unit range, typically with lower and upper asymptotes at 0 and 1. The theory proposed by Grebenc distinguishes three kinds of unitary sigmoid functions according to their asymptotic behavior and the presence or absence of oscillation near the asymptotes. A general form of a unitary sigmoid function is

Sigmoid function -- Part 10

Hyperbolic tangent (shifted and scaled version of the logistic function, above) $f(x) = \tanh x = \frac{e^x - e^{-x}}{e^x + e^{-x}}$

$$f(x) = \tanh x = \frac{e^x - e^{-x}}{e^x + e^{-x}}$$