

A review of cases of human cysticercosis in Canada.

OBJECTIVE: Review of human cysticercosis in Canada, to estimate the magnitude of the disease and to describe the pattern of disease expression in this country.

METHODS: MEDLINE and manual search of case reports and case series of patients with cysticercosis diagnosed in Canada. ed data included year of diagnosis, citizenship status, clinical manifestations, and form of cysticercosis.

FINDINGS: A total of 21 articles reporting 60 patients were found. Forty (67%) of these patients were diagnosed in the past two decades. Most cases came from Ontario (n=43) and Quebec (n=14). Immigrants accounted for 96% of the 28 cases in whom citizenship information was available. Neurocysticercosis was observed in 55 patients, and isolated compromise of striated muscles in the remaining five. Seizures was the primary or sole manifestation of the disease in 72% of patients, and most of them had parenchymal brain cysticerci (either viable cysts or calcifications). Two of seven patients were positive for *Taenia* eggs. In no case were household contacts of the patients investigated for taeniasis.

CONCLUSIONS: An increasing number of patients with cysticercosis have been reported from Canada in the past two decades, suggesting that the prevalence of this parasitic disease may be on the rise. While most cases occur in immigrants, it is possible that at least some of these patients had acquired the disease in Canada.