

Cherry Consumption and the Risk of Recurrent Gout Attacks

Objective To study the relation between cherry intake and the risk of recurrent gout attacks among individuals with gout. **Methods** We conducted a case-crossover study to examine associations of a set of putative risk factors with recurrent gout attacks. Individuals with gout were prospectively recruited and followed online for one year. Participants were asked about the following information when experiencing a gout attack: the onset date of the gout attack, symptoms and signs, medications (including anti-gout medications), and potential risk factors (including daily intake of cherries and cherry extract) during the 2-day period prior to the gout attack. We assessed the same exposure information over 2-day control periods. We estimated the risk of recurrent gout attacks related to cherry intake using conditional logistic regression. **Results** Our study included 633 individuals with gout. Cherry intake over a 2-day period was associated with a 35% lower risk of gout attacks compared with no intake (multivariate odds ratio [OR] = 0.65, 95% CI: 0.50-0.85). Cherry extract intake showed a similar inverse association (multivariate OR=0.55, 95% CI: 0.30-0.98). The effect of cherry intake persisted across subgroups by sex, obesity status, purine intake, alcohol use, diuretic use, and use of anti-gout medications. When cherry intake was combined with allopurinol use, the risk of gout attacks was 75% lower than periods without either exposure (OR=0.25, 95% CI: 0.15-0.42). **Conclusions** These findings suggest that cherry intake is associated with a lower risk of gout attacks.