

Longitudinal Study of the Contamination of Air and of Soil Surfaces in the Vicinity of Pig Barns by Livestock-Associated Methicillin-Resistant *Staphylococcus aureus*

.

During 1 year, samples were taken on 4 days, one sample in each season, from pigs, the floor, and the air inside pig barns and from the ambient air and soil at different distances outside six commercial livestock-associated methicillin-resistant *Staphylococcus aureus* (LA-MRSA)-positive pig barns in the north and east of Germany. LA-MRSA was isolated from animals, floor, and air samples in the barn, showing a range of airborne LA-MRSA between 6 and 3,619 CFU/m³ (median, 151 CFU/m³). Downwind of the barns, LA-MRSA was detected in low concentrations (11 to 14 CFU/m³) at distances of 50 and 150 m; all upwind air samples were negative. In contrast, LA-MRSA was found on soil surfaces at distances of 50, 150, and 300 m downwind from all barns, but no statistical differences could be observed between the proportions of positive soil surface samples at the three different distances. Upwind of the barns, positive soil surface samples were found only sporadically. Significantly more positive LA-MRSA samples were found in summer than in the other seasons both in air and soil samples upwind and downwind of the pig barns. spa typing was used to confirm the identity of LA-MRSA types found inside and outside the barns. The results show that there is regular airborne LA-MRSA transmission and deposition, which are strongly influenced by wind direction and season, of up to at least 300 m around positive pig barns. The described boot sampling method seems suitable to characterize the contamination of the vicinity of LA-MRSA-positive pig barns by the airborne route.