

Introduction to Machine Learning for Social Scientists

Class 5: Intro to GLM

Edgar Franco Vivanco

Stanford University
Department of Political Science

edgarf1@stanford.edu

Summer 2018

Homework 3 Due Friday July 20th at midnight

Available tomorrow Start early!

In-class midterm Monday July 23rd

Final project

We'll talk more about this after midterm but you should start forming teams: 6 teams of 5 in total

All teams should be organized by **Wed July 25th**

Issues with the Room!

Next Monday old room, this one during the rest of the quarter

Today's Goals

1. Key concepts

- ▶ Training and test sets
- ▶ Linear Probability Model
- ▶ Logit function and logit inverse function, logistic regression


2. Key techniques and R functions

- ▶ glm
- ▶ Natural logarithm, log
- ▶ table

Divide your data

- ▶ **Training set:** A set of examples used to fit the parameters and learn. These are already classified by human coders, or produced in a semi-automated way, to create a 'gold standard'.
Conventionally 80% of available data.
- ▶ **Test set:** A set that follows the same probability distribution and is used to test the model.
Conventionally 20% of available data.

Divide your data



R!



Regression vs Classification

1. *Regression*

- Quantitative responses
- Example: Age, height, salary, price, vote share, etc.

Regression vs Classification

1. *Regression*

- Quantitative responses
- Example: Age, height, salary, price, vote share, etc.

2. *Classification*

- Qualitative responses
- Example: Election result (win, lose), fake news (yes, no, maybe).

Regression vs Classification

1. *Regression*

- Quantitative responses
- Example: Age, height, salary, price, vote share, etc.

2. *Classification*

- Qualitative responses
- Example: Election result (win, lose), fake news (yes, no, maybe).

NOTE: The distinction is not always that clear-cut: Logistic regression (a type of non-linear regression) is often used for classification.

Classification

For simplicity, in this class we are going to classify binary outcomes.

- Vote: Yay / Nay?

Classification

For simplicity, in this class we are going to classify binary outcomes.

- Vote: Yay / Nay?
- Email: Spam / Not Spam?

Classification

For simplicity, in this class we are going to classify binary outcomes.

- Vote: Yay / Nay?
- Email: Spam / Not Spam?
- Online transaction: Faudulent (Yes / No)?


Classification

For simplicity, in this class we are going to classify binary outcomes.

- Vote: Yay / Nay?
- Email: Spam / Not Spam?
- Online transaction: Faudulent (Yes / No)?


$$y \in \{0, 1\} \quad \left[\begin{array}{l} 0 : \text{"Negative class"} \\ 1 : \text{"Positive class"} \end{array} \right]$$

Predicting qualitative responses




- $y \in \{0, 1\}$, X continuous

Predicting qualitative responses




- ▶ $y \in \{0, 1\}$, X continuous
- ▶ We can always fit a linear model as we did before

Predicting qualitative responses




- ▶ $y \in \{0, 1\}$, X continuous
- ▶ We can always fit a linear model as we did before
- ▶ We call this the linear probability model because we can associate its predictions to probabilities.

Predicting qualitative responses




- ▶ $y \in \{0, 1\}$, X continuous
- ▶ We can always fit a linear model as we did before
- ▶ We call this the linear probability model because we can associate its predictions to probabilities.
- ▶ **PROBLEMS?**

Predicting qualitative responses




- ▶ $y \in \{0, 1\}$, X continuous
- ▶ We can always fit a linear model as we did before
- ▶ We call this the linear probability model because we can associate its predictions to probabilities.
- ▶ **PROBLEMS?**
- ▶ Predictions smaller than zero and larger than 1!

Predicting qualitative responses




- ▶ Logistic model
- ▶ Estimates the probability of yes:
 $Pr(\hat{Vote}_i = 1|x_i)$
using a logarithmic transformation

Predicting qualitative responses




- ▶ Logistic model
- ▶ Estimates the probability of yes:
 $Pr(\hat{Vote}_i = 1|x_i)$
- ▶ The linear model predictions intersect when the probability equals 0.5.

Predicting qualitative responses




- ▶ Logistic model
- ▶ Estimates the probability of yes:
 $Pr(\text{Vote}_i = 1|x_i)$
- ▶ The linear model predictions intersect when the probability equals 0.5.
- ▶ We can associate each point to a probability

Predicting qualitative responses



- Logistic model usually more stable to outliers

Predicting qualitative responses



- Logistic model usually more stable to outliers

A Brief Reminder About (Natural) Logarithms

Logarithm \log is a **class** of functions.

A Brief Reminder About (Natural) Logarithms

Logarithm \log is a **class** of functions.

- $\log_e z = x$ if $e^x = z$.

A Brief Reminder About (Natural) Logarithms

Logarithm \log is a **class** of functions.

- $\log_e z = x$ if $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log = \log_e$

A Brief Reminder About (Natural) Logarithms

Logarithm \log is a **class** of functions.

- $\log_e z = x$ if $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log = \log_e$
- $\log e = 1$ (because $e^1 = e$)

A Brief Reminder About (Natural) Logarithms

Logarithm \log is a **class** of functions.

- $\log_e z = x$ if $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log = \log_e$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

A Brief Reminder About (Natural) Logarithms

Logarithm \log is a **class** of functions.

- $\log_e z = x$ if $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log = \log_e$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Some rules of logarithms

A Brief Reminder About (Natural) Logarithms

Logarithm \log is a **class** of functions.

- $\log_e z = x$ if $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log = \log_e$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Some rules of logarithms

- $\exp(\log(a)) = e^{\log(a)} = a$

A Brief Reminder About (Natural) Logarithms

Logarithm \log is a **class** of functions.

- $\log_e z = x$ if $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log = \log_e$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Some rules of logarithms

- $\exp(\log(a)) = e^{\log(a)} = a$
- $\log(a \times b) = \log(a) + \log(b)$ (!!!!!)

A Brief Reminder About (Natural) Logarithms

Logarithm \log is a **class** of functions.

- $\log_e z = x$ if $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log = \log_e$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Some rules of logarithms

- $\exp(\log(a)) = e^{\log(a)} = a$
- $\log(a \times b) = \log(a) + \log(b)$ (!!!!!)
- $\log\left(\frac{a}{b}\right) = \log(a) - \log(b)$

A Brief Reminder About (Natural) Logarithms

Logarithm \log is a **class** of functions.

- $\log_e z = x$ if $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log = \log_e$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Some rules of logarithms

- $\exp(\log(a)) = e^{\log(a)} = a$
- $\log(a \times b) = \log(a) + \log(b)$ (!!!!!)
- $\log\left(\frac{a}{b}\right) = \log(a) - \log(b)$
- $\log(a^b) = b \log(a)$

A Brief Reminder About (Natural) Logarithms

Logarithm \log is a **class** of functions.

- $\log_e z = x$ if $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log = \log_e$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Some rules of logarithms

- $\exp(\log(a)) = e^{\log(a)} = a$
- $\log(a \times b) = \log(a) + \log(b)$ (!!!!!)
- $\log\left(\frac{a}{b}\right) = \log(a) - \log(b)$
- $\log(a^b) = b \log(a)$
- $\log(1) = 0$

A Brief Reminder About (Natural) Logarithms

Logarithm \log is a **class** of functions.

- $\log_e z = x$ if $e^x = z$.
- We'll call \log_e **natural logarithm**. And we'll assume $\log = \log_e$
- $\log e = 1$ (because $e^1 = e$)
- $\log_{10} 1000 = 3$ (because $10^3 = 1000$)

Some rules of logarithms

- $\exp(\log(a)) = e^{\log(a)} = a$
- $\log(a \times b) = \log(a) + \log(b)$ (!!!!!)
- $\log\left(\frac{a}{b}\right) = \log(a) - \log(b)$
- $\log(a^b) = b \log(a)$
- $\log(1) = 0$

Call $p_i = \Pr(\text{Vote}_i = 1 | \mathbf{x}_i)$

Call $p_i = \Pr(\text{Vote}_i = 1 | \mathbf{x}_i)$

$$\text{Vote}_i \sim \text{Bernoulli}(p_i)$$

Call $p_i = \Pr(\text{Vote}_i = 1 | \mathbf{x}_i)$

$$\text{Vote}_i \sim \text{Bernoulli}(p_i)$$

$$p_i = f(\beta \cdot \mathbf{x}_i)$$

Call $p_i = \Pr(\text{Vote}_i = 1 | \mathbf{x}_i)$

$$\text{Vote}_i \sim \text{Bernoulli}(p_i)$$

$$p_i = f(\beta \cdot \mathbf{x}_i)$$

$$\log \left(\frac{p_i}{1 - p_i} \right) = \beta \cdot \mathbf{x}_i$$

Call $p_i = \Pr(\text{Vote}_i = 1 | \mathbf{x}_i)$

$$\text{Vote}_i \sim \text{Bernoulli}(p_i)$$

$$p_i = f(\beta \cdot \mathbf{x}_i)$$

$$\log\left(\frac{p_i}{1 - p_i}\right) = \beta \cdot \mathbf{x}_i$$

$$p_i = \frac{\exp(\beta \cdot \mathbf{x}_i)}{1 + \exp(\beta \cdot \mathbf{x}_i)}$$

Call $p_i = \Pr(\text{Vote}_i = 1 | \mathbf{x}_i)$

$$\text{Vote}_i \sim \text{Bernoulli}(p_i)$$

$$p_i = f(\beta \cdot \mathbf{x}_i)$$

$$\log \left(\frac{p_i}{1 - p_i} \right) = \beta \cdot \mathbf{x}_i$$

$$\begin{aligned} p_i &= \frac{\exp(\beta \cdot \mathbf{x}_i)}{1 + \exp(\beta \cdot \mathbf{x}_i)} \\ &= \frac{1}{1 + \exp(-\beta \cdot \mathbf{x}_i)} \end{aligned}$$

Call $p_i = \Pr(\text{Vote}_i = 1 | \mathbf{x}_i)$

$$\text{Vote}_i \sim \text{Bernoulli}(p_i)$$

$$p_i = f(\beta \cdot \mathbf{x}_i)$$

$$\log\left(\frac{p_i}{1 - p_i}\right) = \beta \cdot \mathbf{x}_i$$

$$\begin{aligned} p_i &= \frac{\exp(\beta \cdot \mathbf{x}_i)}{1 + \exp(\beta \cdot \mathbf{x}_i)} \\ &= \frac{1}{1 + \exp(-\beta \cdot \mathbf{x}_i)} \end{aligned}$$

Important functions:

Call $p_i = \Pr(\text{Vote}_i = 1 | \mathbf{x}_i)$

$$\text{Vote}_i \sim \text{Bernoulli}(p_i)$$

$$p_i = f(\beta \cdot \mathbf{x}_i)$$

$$\log\left(\frac{p_i}{1 - p_i}\right) = \beta \cdot \mathbf{x}_i$$

$$\begin{aligned} p_i &= \frac{\exp(\beta \cdot \mathbf{x}_i)}{1 + \exp(\beta \cdot \mathbf{x}_i)} \\ &= \frac{1}{1 + \exp(-\beta \cdot \mathbf{x}_i)} \end{aligned}$$

Important functions:

$$\text{odds}(p) = \frac{p}{1 - p}$$

Call $p_i = \Pr(\text{Vote}_i = 1 | \mathbf{x}_i)$

$$\text{Vote}_i \sim \text{Bernoulli}(p_i)$$

$$p_i = f(\beta \cdot \mathbf{x}_i)$$

$$\log \left(\frac{p_i}{1 - p_i} \right) = \beta \cdot \mathbf{x}_i$$

$$\begin{aligned} p_i &= \frac{\exp(\beta \cdot \mathbf{x}_i)}{1 + \exp(\beta \cdot \mathbf{x}_i)} \\ &= \frac{1}{1 + \exp(-\beta \cdot \mathbf{x}_i)} \end{aligned}$$

Important functions:

$$\text{odds}(p) = \frac{p}{1 - p}$$

$$\log \text{ odds or logit}(p) = \log \left(\frac{p}{1 - p} \right)$$

Call $p_i = \Pr(\text{Vote}_i = 1 | \mathbf{x}_i)$

$$\text{Vote}_i \sim \text{Bernoulli}(p_i)$$

$$p_i = f(\beta \cdot \mathbf{x}_i)$$

$$\log \left(\frac{p_i}{1 - p_i} \right) = \beta \cdot \mathbf{x}_i$$

$$\begin{aligned} p_i &= \frac{\exp(\beta \cdot \mathbf{x}_i)}{1 + \exp(\beta \cdot \mathbf{x}_i)} \\ &= \frac{1}{1 + \exp(-\beta \cdot \mathbf{x}_i)} \end{aligned}$$

Important functions:

$$\text{odds}(p) = \frac{p}{1 - p}$$

$$\log \text{ odds or logit}(p) = \log \left(\frac{p}{1 - p} \right)$$

$$\text{logistic function or logit}^{-1}(a) = \frac{1}{1 + \exp(-a)}$$

More on tutorial

R!



Some Context: Iraq Vote



- ▶ In 2002 President George Bush announced the Joint Resolution to Authorize the Use of United States Armed Forces Against Iraq.
- ▶ Congressional opposition.

NEXT

- ▶ Assessing model predictions for classification
 - ▶ Precision
 - ▶ Accuracy
 - ▶ Recall