

Global Type built according to the proof of Theorem 5.8 in [1]

Franco Barbanera and Mariangiola Dezani-Ciancaglini

We recall the definition of $\mathcal{G}(\mathcal{D}, \mathbf{G}_1, \mathbf{G}_2, \mathbf{G}_3)$ as given in Theorem 5.8 of [1] by cases: first on the last applied rule in the compliance derivation \mathcal{D} and then on the shapes of the global types \mathbf{G}_1 , \mathbf{G}_2 and \mathbf{G}_3 .

Note that the h , v and w in the definition below are the names of the participants acting as interfaces in the composition dealt with in the proof of Theorem 5.8. Such names have to be explicitly provided as extra arguments in any actual implementation of the function \mathcal{G} .

$$\text{Case } \mathcal{D} = \begin{array}{c} \dots \\ [\text{COMP-0}] \\ \dots \end{array} \quad \mathcal{G}(\mathcal{D}, \mathbf{G}_1, \mathbf{G}_2, \mathbf{G}_3) = \mathbf{G}_1$$

$$\text{Case } \mathcal{D} = \begin{array}{c} \mathcal{D}_i \quad \forall i \in I \\ [\text{COMP-O/I-L}] \\ \dots \end{array}$$

$$\mathcal{G}(\mathcal{D}, \mathbf{G}_1, \mathbf{G}_2, \mathbf{G}_3) = \begin{cases} \mathbf{q} \rightarrow \mathbf{v} : \{\lambda_i. \widehat{\mathbf{G}}_i\}_{i \in I'} & \begin{array}{l} \text{if } \mathbf{G}_1 = \mathbf{h} \rightarrow \mathbf{p} : \{\lambda_j. \mathbf{G}'_j\}_{j \in J} \quad I \subseteq J \\ \mathbf{G}_2 = \mathbf{q} \rightarrow \mathbf{v} : \{\lambda_i. \mathbf{G}''_i\}_{i \in I'} \quad I' \subseteq I \\ \text{where } \widehat{\mathbf{G}}_i = \mathbf{v} \rightarrow \mathbf{h} : \lambda_i. \mathbf{h} \rightarrow \mathbf{p} : \lambda_i. \mathcal{G}(\mathcal{D}_i, \mathbf{G}'_i, \mathbf{G}''_i, \mathbf{G}_3) \end{array} \\ \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l. \mathcal{G}(\mathcal{D}, \mathbf{G}'_l, \mathbf{G}_2, \mathbf{G}_3)\}_{l \in L} & \text{if } \mathbf{G}_1 = \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l. \mathbf{G}'_l\}_{l \in L} \quad \mathbf{h} \notin \{\mathbf{t}, \mathbf{u}\} \\ \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l. \mathcal{G}(\mathcal{D}, \mathbf{G}_1, \mathbf{G}''_l, \mathbf{G}_3)\}_{l \in L} & \begin{array}{l} \text{if } \mathbf{G}_1 = \mathbf{h} \rightarrow \mathbf{p} : \{\lambda_j. \mathbf{G}'_j\}_{j \in J} \\ \mathbf{G}_2 = \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l. \mathbf{G}''_l\}_{l \in L} \quad \mathbf{v} \notin \{\mathbf{t}, \mathbf{u}\} \end{array} \end{cases}$$

$$\text{Case } \mathcal{D} = \begin{array}{c} \mathcal{D}_i \quad \forall i \in I \\ [\text{COMP-I/O-L}] \\ \dots \end{array}$$

$$\mathcal{G}(\mathcal{D}, \mathbf{G}_1, \mathbf{G}_2, \mathbf{G}_3) = \begin{cases} \mathbf{p} \rightarrow \mathbf{h} : \{\lambda_i. \widehat{\mathbf{G}}_i\}_{i \in I'} & \begin{array}{l} \text{if } \mathbf{G}_1 = \mathbf{p} \rightarrow \mathbf{h} : \{\lambda_j. \mathbf{G}'_j\}_{j \in J} \quad I \subseteq J \\ \mathbf{G}_2 = \mathbf{v} \rightarrow \mathbf{q} : \{\lambda_i. \mathbf{G}''_i\}_{i \in I'} \quad I' \subseteq I \\ \text{where } \widehat{\mathbf{G}}_i = \mathbf{h} \rightarrow \mathbf{v} : \lambda_i. \mathbf{v} \rightarrow \mathbf{q} : \lambda_i. \mathcal{G}(\mathcal{D}_i, \mathbf{G}'_i, \mathbf{G}''_i, \mathbf{G}_3) \end{array} \\ \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l. \mathcal{G}(\mathcal{D}, \mathbf{G}'_l, \mathbf{G}_2, \mathbf{G}_3)\}_{l \in L} & \text{if } \mathbf{G}_1 = \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l. \mathbf{G}'_l\}_{l \in L} \quad \mathbf{h} \notin \{\mathbf{t}, \mathbf{u}\} \\ \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l. \mathcal{G}(\mathcal{D}, \mathbf{G}_1, \mathbf{G}''_l, \mathbf{G}_3)\}_{l \in L} & \begin{array}{l} \text{if } \mathbf{G}_1 = \mathbf{h} \rightarrow \mathbf{p} : \{\lambda_j. \mathbf{G}'_j\}_{j \in J} \\ \mathbf{G}_2 = \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l. \mathbf{G}''_l\}_{l \in L} \quad \mathbf{v} \notin \{\mathbf{t}, \mathbf{u}\} \end{array} \end{cases}$$

$$\text{Case } \mathcal{D} = \frac{[\text{COMP-O/I-R}] \frac{\mathcal{D}_i \quad \forall i \in I}{\dots}}{\dots}$$

$$\mathcal{G}(\mathcal{D}, G_1, G_2, G_3) = \begin{cases} \mathbf{q} \rightarrow \mathbf{w} : \{\lambda_i.\widehat{G}_i\}_{i \in I'} & \text{if } G_1 = \mathbf{h} \rightarrow \mathbf{p} : \{\lambda_j.G'_j\}_{j \in J} \quad I \subseteq J \\ & G_3 = \mathbf{q} \rightarrow \mathbf{v} : \{\lambda_i.G'''_i\}_{i \in I'} \quad I' \subseteq I \\ & \text{where } \widehat{G}_i = \mathbf{w} \rightarrow \mathbf{h} : \lambda_i.\mathbf{h} \rightarrow \mathbf{p} : \lambda_i.\mathcal{G}(\mathcal{D}_i, G'_i, G''_i, G_3) \\ \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l.\mathcal{G}(\mathcal{D}, G'_l, G_2, G_3)\}_{l \in L} & \text{if } G_1 = \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l.G'_l\}_{l \in L} \quad \mathbf{h} \notin \{\mathbf{t}, \mathbf{u}\} \\ \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l.\mathcal{G}(\mathcal{D}, G_1, G''_l, G_3)\}_{l \in L} & \text{if } G_1 = \mathbf{h} \rightarrow \mathbf{p} : \{\lambda_j.G'_j\}_{j \in J} \\ & G_2 = \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l.G''_l\}_{l \in L} \quad \mathbf{v} \notin \{\mathbf{t}, \mathbf{u}\} \end{cases}$$

$$\text{Case } \mathcal{D} = \frac{[\text{COMP-I/O-R}] \frac{\mathcal{D}_i \quad \forall i \in I}{\dots}}{\dots}$$

$$\mathcal{G}(\mathcal{D}, G_1, G_2, G_3) = \begin{cases} \mathbf{p} \rightarrow \mathbf{h} : \{\lambda_i.\widehat{G}_i\}_{i \in I'} & \text{if } G_1 = \mathbf{p} \rightarrow \mathbf{h} : \{\lambda_j.G'_j\}_{j \in J} \quad I \subseteq J \\ & G_3 = \mathbf{v} \rightarrow \mathbf{q} : \{\lambda_i.G'''_i\}_{i \in I'} \quad I' \subseteq I \\ & \text{where } \widehat{G}_i = \mathbf{h} \rightarrow \mathbf{v} : \lambda_i.\mathbf{v} \rightarrow \mathbf{q} : \lambda_i.\mathcal{G}(\mathcal{D}_i, G'_i, G_2, G''') \\ \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l.\mathcal{G}(\mathcal{D}, G'_l, G_2, G_3)\}_{l \in L} & \text{if } G_1 = \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l.G'_l\}_{l \in L} \quad \mathbf{h} \notin \{\mathbf{t}, \mathbf{u}\} \\ \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l.\mathcal{G}(\mathcal{D}, G_1, G_2, G''')\}_{l \in L} & \text{if } G_1 = \mathbf{h} \rightarrow \mathbf{p} : \{\lambda_j.G'_j\}_{j \in J} \\ & G_3 = \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_l.G'''_l\}_{l \in L} \quad \mathbf{w} \notin \{\mathbf{t}, \mathbf{u}\} \end{cases}$$

$$\text{Case } \mathcal{D} = \frac{[\text{COMP-O-A}] \frac{\mathcal{D}_i \quad \forall i \in I}{\dots}}{\dots}$$

$$\mathcal{G}(\mathcal{D}, G_1, G_2, G_3) = \begin{cases} \mathbf{h} \rightarrow \mathbf{p} : \{\lambda_i.\mathcal{G}(\mathcal{D}_i, G'_i, G_2, G_3)\}_{i \in I} & \text{if } G_1 = \mathbf{h} \rightarrow \mathbf{p} : \{\lambda_i.G'_i\}_{i \in I} \\ \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_j.\mathcal{G}(\mathcal{D}, G'_j, G_2, G_3)\}_{j \in J} & \text{if } G_1 = \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_j.G'_j\}_{j \in J} \quad \mathbf{h} \notin \{\mathbf{t}, \mathbf{u}\} \end{cases}$$

$$\text{Case } \mathcal{D} = \frac{[\text{COMP-O-A}] \frac{\mathcal{D}_i \quad \forall i \in I}{\dots}}{\dots}$$

$$\mathcal{G}(\mathcal{D}, G_1, G_2, G_3) = \begin{cases} \mathbf{p} \rightarrow \mathbf{h} : \{\lambda_i.\mathcal{G}(\mathcal{D}_i, G'_i, G_2, G_3)\}_{i \in I} & \text{if } G_1 = \mathbf{p} \rightarrow \mathbf{h} : \{\lambda_i.G'_i\}_{i \in I} \\ \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_j.\mathcal{G}(\mathcal{D}, G'_j, G_2, G_3)\}_{j \in J} & \text{if } G_1 = \mathbf{t} \rightarrow \mathbf{u} : \{\lambda_j.G'_j\}_{j \in J} \quad \mathbf{h} \notin \{\mathbf{t}, \mathbf{u}\} \end{cases}$$

References

- [1] Franco Barbanera and Mariangiola Dezani-Ciancaglini. Partial typing for open compliance in multiparty sessions. submitted, 2024.