

# Solved selected problems of Classical Mechanics - Gregory

Franco Zacco

## Chapter 2 - Velocity, acceleration and scalar angular velocity

*Solution.* **2.3** Given that acceleration is constant then the instant acceleration  $a$  is equal to the average acceleration  $a_{avg}$  so assuming that we start from  $t = 0$  at  $x = 0$  and the final velocity is  $v$  then we have that

$$\begin{aligned}a &= \frac{\Delta v}{\Delta t} \\a &= \frac{v - u}{t - 0} \\at &= v - u \\v &= u + at\end{aligned}\tag{1}$$

Now integrating both sides of the equation with respect to  $t$ :

$$\begin{aligned}\int_0^t v dt &= \int_0^t u dt + \int_0^t at dt \\ \int_0^t dx &= \int_0^t u dt + \int_0^t at dt \\ x(t) - x(0) &= ut + \frac{1}{2}at^2 \\ x &= ut + \frac{1}{2}at^2\end{aligned}\tag{2}$$

From (1) we get that  $t = \frac{v-u}{a}$  and replacing in (2) we have

$$\begin{aligned}x &= u \frac{(v - u)}{a} + \frac{1}{2}a \frac{(v - u)^2}{a^2} \\ ax &= uv - u^2 + \frac{1}{2}(v^2 - uv + u^2) \\ ax &= \frac{v^2}{2} - \frac{u^2}{2}\end{aligned}$$

$$2ax = v^2 - u^2$$

$$v^2 = u^2 + 2ax$$

Finally if we replace  $t = 11.4$  s and  $v = 116$  mph  $= 168.96$  ft/s in (1) we get that  $a = 14.82$  ft/s<sup>2</sup> but then if we replace the same values in (2) assuming  $x = 0.25$  mi  $= 1320$  ft we get that  $a = 20.31$  ft/s<sup>2</sup> which means that the acceleration is not constant because otherwise it must satisfy both equations.  $\square$

*Solution.* **2.6** First we calculate the following

$$r = be^{\Omega t} \quad \theta = \Omega t$$

$$\dot{r} = b\Omega e^{\Omega t} \quad \dot{\theta} = \Omega$$

$$\ddot{r} = b\Omega^2 e^{\Omega t} \quad \ddot{\theta} = 0$$

Now replacing in  $\mathbf{v} = \dot{r}\hat{\mathbf{r}} + (r\dot{\theta})\hat{\boldsymbol{\theta}}$  and in  $\mathbf{a} = (\ddot{r} - r\dot{\theta}^2)\hat{\mathbf{r}} + (r\ddot{\theta} + 2\dot{r}\dot{\theta})\hat{\boldsymbol{\theta}}$  we get that

$$\mathbf{v} = b\Omega e^{\Omega t}\hat{\mathbf{r}} + (be^{\Omega t}\Omega)\hat{\boldsymbol{\theta}}$$

And

$$\begin{aligned} \mathbf{a} &= (b\Omega^2 e^{\Omega t} - b\Omega^2 e^{\Omega t})\hat{\mathbf{r}} + (be^{\Omega t}0 + 2b\Omega^2 e^{\Omega t})\hat{\boldsymbol{\theta}} \\ &= 2b\Omega^2 e^{\Omega t}\hat{\boldsymbol{\theta}} \end{aligned}$$

Then,

$$|v| = \sqrt{2}b\Omega^2 e^{\Omega t}$$

$$|a| = 2b\Omega^2 e^{\Omega t}$$

So by using the formula for the dot product we compute  $\cos(\alpha) = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|}$  as

$$\begin{aligned} \cos(\alpha) &= \frac{2b^2\Omega^3 e^{2\Omega t}}{(\sqrt{2}b\Omega^2 e^{\Omega t})(2b\Omega^2 e^{\Omega t})} \\ &= \frac{1}{\sqrt{2}} \end{aligned}$$

Finally, this means that  $\alpha = \pi/4$ .  $\square$

*Solution. 2.9* From  $\mathbf{v} = \dot{r}\hat{\mathbf{r}} + (r\dot{\theta})\hat{\boldsymbol{\theta}}$  we calculate the velocity vector where

$$\begin{aligned}\dot{r} &= \frac{2b}{\tau} - \frac{2b}{\tau^2}t \\ \dot{\theta} &= \frac{1}{\tau}\end{aligned}$$

Then the velocity vector is

$$\mathbf{v} = \frac{2b}{\tau^2}(\tau - t)\hat{\mathbf{r}} + \frac{bt}{\tau^3}(2\tau - t)\hat{\boldsymbol{\theta}}$$

Now let's compute  $|\mathbf{v}|^2$

$$\begin{aligned}|\mathbf{v}|^2 &= \frac{4b^2}{\tau^4}\tau^2 - \frac{8b^2\tau}{\tau^4}t + \frac{4b^2}{\tau^4}t^2 + \frac{4b^2\tau^2}{\tau^6}t^2 - \frac{4b^2\tau}{\tau^6}t^3 + \frac{b^2}{\tau^6}t^4 \\ &= \frac{4b^2}{\tau^2} - \frac{8b^2}{\tau^3}t + \frac{8b^2}{\tau^4}t^2 - \frac{4b^2}{\tau^5}t^3 + \frac{b^2}{\tau^6}t^4 \\ &= \frac{b^2}{\tau^6}(t^4 - 4\tau t^3 + 8\tau^2 t^2 - 8\tau^3 t + 4\tau^4)\end{aligned}$$

To find the maximum value of  $|\mathbf{v}|$  let's consider the time derivative of  $|\mathbf{v}|^2$

$$\begin{aligned}\frac{d|\mathbf{v}|^2}{dt} &= \frac{b^2}{\tau^6}(4t^3 - 12\tau t^2 + 16\tau^2 t - 8\tau^3) \\ &= \frac{4b^2}{\tau^6}(t^3 - 3\tau t^2 + 4\tau^2 t - 2\tau^3) \\ &= \frac{4b^2}{\tau^6}(t^3 - \tau t^2 - 2\tau t^2 + 2\tau^2 t + 2\tau^2 t - 2\tau^3) \\ &= \frac{4b^2}{\tau^6}(t(t^2 - 2\tau t + 2\tau^2) - \tau(t^2 - 2\tau t + 2\tau^2)) \\ &= \frac{4b^2}{\tau^6}(t - \tau)(t^2 - 2\tau t + 2\tau^2)\end{aligned}$$

Since  $t^2 - 2\tau t + 2\tau^2$  can be written as  $\tau^2 + (t - \tau)^2$  we see that this term no matter the value of  $t$  is always positive so

$$\frac{d|\mathbf{v}|^2}{dt} = \begin{cases} < 0 & \text{if } t < \tau \\ = 0 & \text{if } t = \tau \\ > 0 & \text{if } t > \tau \end{cases}$$

Hence  $|\mathbf{v}|$  achieves a minimum value when  $t = \tau$  so

$$\begin{aligned}|\mathbf{v}|^2 &= \frac{b^2}{\tau^6}\tau^4 \\ |\mathbf{v}| &= \frac{b}{\tau}\end{aligned}$$

Finally the acceleration is given by

$$\mathbf{a} = \left(-\frac{2b}{\tau^2} - \frac{2b}{\tau^3}t + \frac{b}{\tau^4}t^2\right)\hat{\mathbf{r}} + \left(\frac{4b}{\tau^2}\left(1 - \frac{t}{\tau}\right)\right)\hat{\boldsymbol{\theta}}$$

and when  $t = \tau$  we have that

$$\mathbf{a} = -\frac{3b}{\tau^2}\hat{\mathbf{r}}$$

□

*Solution. 2.10* The Lion's velocity can be calculated as

$$\mathbf{v} = \dot{r}\hat{\mathbf{r}} + r\dot{\theta}\hat{\boldsymbol{\theta}}$$

So calculating the speed of the Lion knowing that  $\dot{\theta}$  is the angular velocity and can be written as  $\dot{\theta} = \frac{u}{a}$  where  $u$  is the velocity of Daniel and  $a$  is the radius then we get that

$$\begin{aligned} U^2 &= \dot{r}^2 + (r\dot{\theta})^2 \\ &= \dot{r}^2 + \left(\frac{ru}{a}\right)^2 \end{aligned}$$

which can be re-written as

$$\begin{aligned} \dot{r}^2 &= U^2 - \left(\frac{ru}{a}\right)^2 \\ &= \frac{u^2}{a^2} \left( \frac{U^2 a^2}{u^2} - r^2 \right) \end{aligned}$$

This ODE can be solved as follows

$$\begin{aligned} \dot{r} &= \frac{u}{a} \sqrt{\frac{U^2 a^2}{u^2} - r^2} \\ \frac{dr}{dt} &= \frac{u}{a} \sqrt{\frac{U^2 a^2}{u^2} - r^2} \\ \int \frac{dr}{\sqrt{\frac{U^2 a^2}{u^2} - r^2}} &= \int \frac{u}{a} dt \\ \int \frac{dr}{\sqrt{\frac{U^2 a^2}{u^2} - r^2}} &= \frac{ut}{a} \\ \sin^{-1}\left(\frac{ur}{Ua}\right) + C &= \frac{ut}{a} \end{aligned}$$

the constant of integration  $C$  can be determined by knowing that  $r = 0$  when  $t = 0$  then  $C = 0$  and therefore

$$r = \frac{Ua}{u} \sin\left(\frac{ut}{a}\right)$$

For  $t = \pi a/2u$  we get that  $r = Ua/u$  and if  $U \geq u$  then  $\frac{U}{u} \geq 1$  so  $r = \frac{Ua}{u} \geq a$  which means that the Lion will caught Daniel in  $t = \pi a/2u$  if  $U \geq u$ .

In order to recognize the equation for  $r$  as a circle we multiply both sides by  $r$  so we get that

$$r^2 = \frac{Ua}{u} r \sin\left(\frac{ut}{a}\right)$$

now knowing that  $r^2 = x^2 + y^2$  and that  $r \sin(\frac{ut}{a}) = y$  then

$$\begin{aligned}x^2 + y^2 &= \frac{Ua}{u}y \\y^2 - \frac{Ua}{u}y &= -x^2 \\y^2 - \frac{2Ua}{2u}y + (\frac{Ua}{2u})^2 &= -x^2 + (\frac{Ua}{2u})^2 \\(y - \frac{Ua}{2u})^2 &= -x^2 + (\frac{Ua}{2u})^2 \\x^2 + (y - \frac{Ua}{2u})^2 &= (\frac{Ua}{2u})^2\end{aligned}$$

which is an equation of a circle with center at  $(0, \frac{Ua}{2u})$  and radius  $\frac{Ua}{2u}$ . When  $U = u$  we get that the Lion's path is described by

$$x^2 + (y - \frac{a}{2})^2 = (\frac{a}{2})^2$$

which is a circle with center at  $(0, \frac{a}{2})$  and radius  $\frac{a}{2}$ . □

*Solution. 2.11* We know that the velocity in a general form can be written as

$$\mathbf{v} = v\mathbf{t}$$

where  $\mathbf{t}$  is the unit tangential vector, and the acceleration can be written as

$$\mathbf{a} = \frac{dv}{dt}\mathbf{t} + \frac{v^2}{\rho}\mathbf{n}$$

where  $\mathbf{n}$  is the unit normal vector, since  $v$  is constant then  $\frac{dv}{dt} = 0$  so the acceleration can be written as

$$\mathbf{a} = \frac{v^2}{\rho}\mathbf{n}$$

which only has the normal component, therefore the acceleration is perpendicular to the velocity. □

*Solution. 2.14* Knowing that

$$r = b \cosh(\Omega t)$$

then

$$\begin{aligned}\dot{r} &= b\Omega \sinh(\Omega t) \\\ddot{r} &= b\Omega^2 \cosh(\Omega t)\end{aligned}$$

so from the polar velocity equation we have that

$$\begin{aligned}\mathbf{v} &= \dot{r}\hat{\mathbf{r}} + r\dot{\theta}\hat{\boldsymbol{\theta}} \\&= b\Omega \sinh(\Omega t)\hat{\mathbf{r}} + b\Omega \cosh(\Omega t)\hat{\boldsymbol{\theta}} \\&= b\Omega(\sinh(\Omega t)\hat{\mathbf{r}} + \cosh(\Omega t)\hat{\boldsymbol{\theta}})\end{aligned}$$

and the speed of the particle is then

$$\begin{aligned} |v|^2 &= b^2 \Omega^2 \sinh^2(\Omega t) + b^2 \Omega^2 \cosh^2(\Omega t) \\ &= b^2 \Omega^2 \cosh(2\Omega t) \end{aligned}$$

Finally, the acceleration is derived as follows

$$\begin{aligned} \mathbf{a} &= (\ddot{r} - r\dot{\theta}^2)\hat{\mathbf{r}} + (r\ddot{\theta} + 2\dot{r}\dot{\theta})\hat{\boldsymbol{\theta}} \\ &= [b\Omega^2 \cosh(\Omega t) - b\Omega^2 \cosh(\Omega t)]\hat{\mathbf{r}} + [0 + 2b\Omega^2 \sinh(\Omega t)]\hat{\boldsymbol{\theta}} \\ &= 2b\Omega^2 \sinh(\Omega t)\hat{\boldsymbol{\theta}} \end{aligned}$$

Therefore the acceleration direction is circumferential and is given by  $\hat{\boldsymbol{\theta}}$ .  $\square$

*Solution. 2.18* As the graph shows P is at  $(b \sin(\Omega t), 0)$  from there and by using Pythagoras theorem we can derive that Q is at  $(0, \sqrt{a^2 - b^2 \sin^2(\Omega t)})$  then C, the center of the link, has coordinates given  $(\frac{b}{2} \sin(\Omega t), \frac{\sqrt{a^2 - b^2 \sin^2(\Omega t)}}{2})$ . If we now move the origin  $(0, 0)$  to C then the coordinates of P are given by  $(\frac{b}{2} \sin(\Omega t), -\frac{\sqrt{a^2 - b^2 \sin^2(\Omega t)}}{2})$  so we have that the sine of the angle between the link and the negative Y-axis is given by

$$\sin(\theta) = \frac{\frac{b}{2} \sin(\Omega t)}{\frac{a}{2}} \quad \text{so} \quad \theta = \sin^{-1} \left( \frac{b \sin \Omega t}{a} \right)$$

now derivating this expression with respect to  $t$  we get the Angular velocity as

$$\dot{\theta} = \omega = \frac{b\Omega \cos(\Omega t)}{\sqrt{a^2 - b^2 \sin^2(\Omega t)}}$$

Finally, since we know the coordinates for C we can calculate the velocity as

$$\mathbf{v} = \frac{\Omega b}{2} \cos(\Omega t) \mathbf{i} - \frac{\Omega b^2 \sin(\Omega t) \cos(\Omega t)}{2\sqrt{a^2 - b^2 \sin^2(\Omega t)}} \mathbf{j}$$

so now we can calculate the speed for the link center by

$$\begin{aligned} |v|^2 &= \frac{\Omega^2 b^2}{4} \cos^2(\Omega t) + \frac{\Omega^2 b^4 \sin^2(\Omega t) \cos^2(\Omega t)}{4(a^2 - b^2 \sin^2(\Omega t))} \\ &= \frac{\Omega^2 b^2}{4} \cos^2(\Omega t) \left( 1 + \frac{b^2 \sin^2(\Omega t)}{a^2 - b^2 \sin^2(\Omega t)} \right) \\ &= \frac{\Omega^2 b^2}{4} \cos^2(\Omega t) \left( \frac{a^2}{a^2 - b^2 \sin^2(\Omega t)} \right) \\ &= \frac{\Omega^2 b^2 a^2 \cos^2(\Omega t)}{4(a^2 - b^2 \sin^2(\Omega t))} \end{aligned}$$

Finally

$$|v| = \frac{\Omega b a \cos(\Omega t)}{2\sqrt{a^2 - b^2 \sin^2(\Omega t)}}$$


$\square$

*Solution.* **2.22**


(i) Solving the equations given by computational means (Runge-kutta method)

$$\dot{X} = -\frac{v^D X}{\sqrt{X^2 + Y^2}} - v_x^H \quad \dot{Y} = -\frac{v^D Y}{\sqrt{X^2 + Y^2}} - v_y^H$$

when  $v^D = v^H$  we get the following plot




but if  $v^D > v^H$






- (ii) Now we solve the equations for the case where the Hare has a circular trajectory when  $v^D = v^H$  we get the following plot



but if  $v^D > v^H$



□