Hopemark Health

Two-Sided Flyer Content Brief

Purpose: Introduce Hopemark Health’s psychiatric services and referral partnerships on the front. Use the back to explain treatment-resistant depression, how common it is, and the treatment options Hopemark offers.

FRONT: Comprehensive Psychiatric Care

Content order: Logo and headline, introduction with a consultation photo, services and conditions, clinical team, clinic locations, and contact information.

Headline: Comprehensive Psychiatric Care

Subheading: Your Trusted Referral Partner

Introduction

Hopemark Health provides evidence-based psychiatric care for children, adolescents, and adults. Our multidisciplinary team supports accurate diagnosis, individualized treatment, and coordination with referring providers.

Clinical Services

- Psychiatric evaluations and medication management
- Psychotherapy
- Neuropsychological testing
- Child and adolescent psychiatry
- Sleep medicine
- Genetic testing - pharmacogenomics
- Substance use treatment and medically assisted recovery

Conditions We Treat

- Major depressive disorder
- Anxiety and panic disorders
- Post-traumatic stress disorder (PTSD)
- Bipolar disorder
- Obsessive-compulsive disorder (OCD)
- Substance use disorders

Our Clinical Team

- Psychiatrists
- Psychiatric mental health nurse practitioners
- Psychologists
- Physician assistants
- Licensed counselors and therapists

Hopemark Clinic Locations

- Chicago: Millennium Park and West Loop
- Hoffman Estates
- Naperville
- Oak Brook
- Orland Park

Contact Information

Referrals and appointments

Phone: 630.791.5923
Fax: 630.385.0290
Website: hopemarkhealth.com

Front imagery: Use the psychiatric consultation photo near the introduction. The supplied patient and clinic photographs are additional available image assets.

BACK: Treatment-Resistant Depression

Content order: TRD explanation and prevalence, how Hopemark can help, the three treatment options with their corresponding images, contact information, and the source footnote.

Headline: Treatment-Resistant Depression

Subheading: More options. Renewed hope.

What Is Treatment-Resistant Depression?

When depression has not improved enough after at least two antidepressants taken at an adequate dose for an adequate length of time, it may be treatment-resistant depression (TRD).

Symptoms such as persistent sadness, low energy, and loss of interest can continue to affect everyday life.

How Common Is It?

About 1 in 3*

A U.S. study estimated that approximately one-third of adults receiving medication for major depression had treatment-resistant depression.

How Hopemark Health Can Help

Our team reviews each patient’s symptoms and treatment history to build a personalized plan that may include novel therapies, medication management, and psychotherapy.

Deep TMS

Deep Transcranial Magnetic Stimulation

What it is and how it works: A noninvasive treatment that uses magnetic pulses to stimulate brain networks involved in mood and OCD. FDA-cleared for depression and as an additional treatment for OCD.

Potential benefits: May reduce persistent depression symptoms. An OCD-specific treatment protocol may reduce obsessive thoughts and compulsive behaviors.

Images: Deep TMS machine and the supplied before-and-after brain-activity visualization, placed with this treatment section.

Brain-image labels:

Before TMS

Potential response after TMS

Brain-image caption:

Educational visualization. Results vary.

BACK: Treatment Options and Closing Content

Spravato®

Esketamine Nasal Spray

What it is and how it works: An FDA-approved prescription nasal spray for adults with treatment-resistant depression. It acts on NMDA receptors in the brain’s glutamate system.

Potential benefits: May reduce depression symptoms, with improvement observed as early as approximately 24 hours in clinical trials. It can be used with or without an oral antidepressant for TRD.

Images: Official Spravato website logo alongside the nasal spray applicator, placed with this treatment section.

IV Ketamine

What it is and how it works: Low-dose ketamine delivered through an IV. It acts on NMDA receptors in the brain’s glutamate system and is used off-label for depression.

Potential benefits: May relieve depression symptoms within hours for some patients when standard treatments have not provided enough relief. Additional treatments may be needed to maintain benefits.

Image: An adult patient resting comfortably in a chair with an eye mask. Do not show an IV drip, bag, pole, tubing, or needle.

Closing Message

Discuss treatment options with our team.

Phone: 630.791.5923
Fax: 630.385.0290
Website: hopemarkhealth.com

Prevalence Footnote

In a 2021 U.S. study, an estimated 30.9% of adults with medication-treated major depression had treatment-resistant depression. Source: Zhdanava et al., Journal of Clinical Psychiatry. 2021;82(2):20m13699.

Placement note: List clinic locations on the front only. Include contact information on both sides.