

2042 MLM Mini Project (Spring 2020)

Group 1

May 13 2020

Team members and division of work

Frank Jiang, Lisa Song, Yuyue Hua, Seeun Jang, Tong Jin

```
set.seed(2042001)
```

Question 1

You will generate simulated data for a single school with 100 classrooms, each of which has 200 students.

- Outcome for student i in classroom j : Y_{ij} .
- There is a single predictor, $X_{ij} \sim U(0, 1)$ (uniform on $[0, 1]$)
- There is a classroom random effect, $\eta_j \sim N(0, \sigma_\eta^2)$, where $\sigma_\eta^2 = 2$.
- Subject level error, $\varepsilon_{ij} \sim N(0, \sigma_\varepsilon^2)$, where $\sigma_\varepsilon^2 = 2$.
- `set.seed(2042001)` once at the beginning of your code.
- Generate the random quantities in this order to ensure the same solution for everyone: X , η_j , ε_{ij}
- The outcome has the following form (DGP, given the modeling parameters above):

$$Y_{ij} = 0 + 1X_{ij} + \eta_j + \varepsilon_{ij}; \eta_j \sim N(0, \sigma_\eta^2), \varepsilon_{ij} \sim N(0, \sigma_\varepsilon^2), \text{indep.}$$

- Generate a single simulated dataset (you will need a “classid” variable to track classrooms); you can optionally assign a “studentid”) i. **Important:** construct classid such that classrooms appear consecutively within the dataframe. As per: `rep(1:J, each=n_j)`

```
# Compute variables related to the equation
N_j <- 100 # Total number of classrooms
n_i <- 200 # Number of students in each classroom
N_i <- N_j * 200 # Total number of students

X_ij <- runif( # Single predictor: Uniform on [0, 1]
  N_i,
  min = 0,
  max = 1
)
eta_j <- rnorm( # Classroom random effect: Normal on (0, 2)
```

```

N_j,
mean = 0,
sd = sqrt(2)
)
epsilon_ij <- rnorm( # Subject level error: Normal on (0, 2)
N_i,
mean = 0,
sd = sqrt(2)
)

# Create equation elements
eta_j <- rep( # Assign classroom random effect to students in each classroom
eta_j,
each = n_i
)
Y_ij <- 0 + 1 * X_ij + eta_j + epsilon_ij

classid <- rep(
1:N_j,
each = n_i
)
studentid <- seq(
1:N_i
)

# Create a dataframe to store all elements
classroom_sim <- data.frame(
outcome = Y_ij,
predictor = X_ij,
cls_raneff = eta_j,
subject_error = epsilon_ij,
classid = classid,
studentid = studentid,
row.names = studentid
)

```

Question 2

Fit the model corresponding to the DGP on your simulated data.

```

# Fit the model in Q1
fit_q2 <- lmer(outcome ~ predictor + (1 | classid), data = classroom_sim)
summary(fit_q2)

```

```

## Linear mixed model fit by REML. t-tests use Satterthwaite's method [
## lmerModLmerTest]
## Formula: outcome ~ predictor + (1 | classid)
## Data: classroom_sim
##
## REML criterion at convergence: 71227.3
##
## Scaled residuals:

```

```
##      Min      1Q  Median      3Q      Max
## -4.0143 -0.6761  0.0024  0.6711  3.7584
##
## Random effects:
##   Groups   Name      Variance Std.Dev.
##   classid (Intercept) 1.893    1.376
##   Residual              2.008    1.417
## Number of obs: 20000, groups:  classid, 100
##
## Fixed effects:
##              Estimate Std. Error      df t value Pr(>|t|)
## (Intercept) -7.493e-03  1.391e-01  1.022e+02 -0.054    0.957
## predictor    9.864e-01  3.496e-02  1.990e+04 28.216   <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##              (Intr)
## predictor -0.126
```

```
slope_X_q2 <- round(summary(fit_q2)$coefficients["predictor", "Estimate"], digits = 4)
```

a. Report coefficient estimate for slope on X.

Response: 0.9864.

b. Does a 95% confidence band for this coefficient estimate cover the “truth” that you used to generate the data?

```
CI_Q2_b <- confint(fit_q2, parm = "predictor", level = 0.95)
```

```
## Computing profile confidence intervals ...
```

```
lower <- round(CI_Q2_b[1], digits = 4)
upper <- round(CI_Q2_b[2], digits = 4)
```

****Response:**** Yes. The confidence band (95% of confidence) for this coefficient estimate is between 0.9

Question 3

3. Next, we simulate missing data in several ways. This is the first:

a. Make a copy of the data, then modify the copy following these instructions:

```
classroom_miss <- classroom_sim
```

b. Generate $Z_{ij} \sim \text{Bernoulli}(p)$, with $p = 0.5$.

c. Set Y to NA when $Z_{ij} == 1$. This should look a lot like “MCAR” missingness.

```
Z_ij <- rbinom(N_i, size = c(0, 1), prob = 0.5)
classroom_miss$missing <- Z_ij
classroom_miss$outcome <- ifelse(classroom_miss$missing == 1, yes = NA, no = classroom_miss$outcome)
```

- d. Refit the model on the new data and report the coefficient estimate for slope on X. Look at the other parameter estimates as well.

```
fit_q3 <- lmer(outcome ~ predictor + (1 | classid), data = classroom_miss)
summary(fit_q3)
```

```
## Linear mixed model fit by REML. t-tests use Satterthwaite's method [
## lmerModLmerTest]
## Formula: outcome ~ predictor + (1 | classid)
## Data: classroom_miss
##
## REML criterion at convergence: 53333.6
##
## Scaled residuals:
##      Min       1Q   Median       3Q      Max
## -4.0171 -0.6742  0.0023  0.6631  3.7668
##
## Random effects:
## Groups   Name                Variance Std.Dev.
## classid  (Intercept)  1.878      1.371
## Residual                    2.000      1.414
## Number of obs: 14963, groups: classid, 100
##
## Fixed effects:
##              Estimate Std. Error      df t value Pr(>|t|)
## (Intercept) 1.270e-02  1.390e-01 1.033e+02  0.091    0.927
## predictor   9.612e-01  4.029e-02 1.486e+04 23.856 <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Correlation of Fixed Effects:
##              (Intr)
## predictor -0.146
```

```
slope_X_q3 <- round(summary(fit_q3)$coefficients["predictor", "Estimate"], digits = 4)
```

```
# Calculate the confidence band of this model
```

```
CI_Q3_d <- confint(fit_q3, parm = "predictor", level = 0.95)
```

```
## Computing profile confidence intervals ...
```

```
lower <- round(CI_Q3_d[1], digits = 4)
```

```
upper <- round(CI_Q3_d[2], digits = 4)
```

- e. Do you see any real change in the β_X estimate?

Response: No. The β_X estimate of the model with missing data values is 0.9612, which is very close to the original estimate, 0.9864.

- i. Does a 95% confidence band for this coefficient estimate cover the “truth” that you used to generate the data?

Response: Yes. The confidence band (95% of confidence) for this coefficient estimate is between 0.8822 and 1.0402. This confirms that a 95% confidence band covers the “truth” of the slope, which is 1.

- f. What is the total sample size N used in the model fit?

Response: The total sample size used in this model fit is 14963.

Question 4:

Missing Data II: Make another copy of the original data, then modify the copy as follows: a. Generate $Z_{ij} \sim \text{Bernoulli}(X_{ij})$, with X_{ij} your predictor generated previously. b. Set Y to NA when $Z_{ij} == 1$. This should look a lot like “MAR” missingness.

```
# Insert code the generate your data
```

- c. Refit the model on the new data and report the coefficient estimate for slope on X. Look at the other parameter estimates as well. **comment**

```
# Insert code to fit model and compute confidence interval
```

Response:

- d. Do you see any real change in the β_X estimate?
- i. Does a 95% confidence band for this coefficient estimate cover the “truth” that you used to generate the data? **comment**

Response:

- e. What is the total sample size N used in the model fit? **comment**

Response:

Question 5:

Missing Data III: Make another copy of the original data, then modify the copy as follows:

```
# Insert code to make a copy of the original data
```

- a. First, define the expit function: `expit <- function(x) exp(x)/(1+exp(x))`

```
# Insert code to define expit function
```

- b. Generate $Z_{ij} \sim \text{Bernoulli}(\text{expit}(Y_{ij}))$, with Y_{ij} your *outcome* generated previously.
- c. Set Y to NA when $Z_{ij} == 1$. This should look like a violation of “MAR” missingness (missingness depends on outcome and cannot be *simply* predicted with the predictor set – Y should be correlated with X, though, so it might not be too bad a violation).

```
# Insert code the generate your data
```

- d. Refit the model on the new data and report the coefficient estimate for slope on X. Look at the other parameter estimates as well. **comment**

```
# Insert code to fit model and compute confidence interval
```

Response:

- e. Do you see any real change in the β_X estimate? **comment**
- i. Does a 95% confidence band for this coefficient estimate cover the “truth” that you used to generate the data? **comment**

Response:

- f. What is the total sample size N used in the model fit? **comment**

Response: