

Article

Cavity-Assisted Spin-Orbit Coupling of Ultracold atoms

Lin Dong ¹, Chuanzhou Zhu ¹ and Han Pu ^{1*}

¹ Department of Physics and Astronomy, Rice Quantum Institute, Rice University, Houston, Texas, 77251-1892, USA

* Author to whom correspondence should be addressed; hpu@rice.edu

Version March 5, 2015 submitted to Atoms. Typeset by L^AT_EX using class file mdpi.cls

Abstract: We investigate dynamical and static properties of ultracold atoms confined in an optical cavity, where two photon Raman process induces effective coupling between atom's internal degrees of freedom and center-of-mass motion. In the meantime, atomic dynamics exerts a back action to cavity photons. We adopt both mean field and master equation approach to tackle the problem and found surprising modifications to atomic dispersions and dynamical instabilities, arising from the intrinsic nonlinearity of the system. Correspondence between semi-classical and quantum limits is analyzed as well.

Keywords: cavity quantum electrodynamics; cold atoms; spin-orbit coupling

1. Introduction

When Jaynes and Cummings first studied the time evolution of a two-level atom in an electromagnetic field in a *fully* quantized way in 1960s [1], experimental realization of this ideal theoretical model was out of reach. It was made possible only with the advent of one-atom masers in late 1980s, by Rempe, Walther and Klein [2], to experimentally study the interaction of a single atom and a single resonant mode of electromagnetic field in a cavity. Jaynes-Cummings model (J-C Model) serves to understand the relationship between quantum theory of radiation and semi-classical theory of atom-light interaction. The field of cavity quantum electrodynamics (CQED) was further advanced by putting cold atoms into the high finesse optical cavities [3–5]. Unlike “hot” atoms, cold atoms’ center-of-mass motion (COM) can no longer be neglected in this “atom + cavity” system. One needs to seek a self-consistent solution for both light and atom by treating them on equal footing. Because intra-cavity photon and

atoms very frequently scatter off each other due to the geometric confinement, not only dipole force gets strongly enhanced but also atom's back-action *onto* light becomes significant. For Bose-Einstein condensate (BEC), atoms occupy the same motional quantum state, and because of long-range cavity photon mediation, the internal states (pseudo-spins) are infinitely coordinated. The most famous example is the Dicke model [6], which has been realized in CQED as well [7].

Another recent breakthrough in cold atoms stems from the realization of artificial (synthetic) gauge potentials for neutral atoms, first in bosonic systems [8,9] and later in fermionic counterparts [10,11]. Laser fields are properly aligned and designed in such a way that trapped atoms may mimic charged particles in a magnetic field with emergence of Lorentz-like force. The synthesis is achieved by inducing two-photon Raman transition between two hyperfine ground states. Using a group of degenerate (or quasi-degenerate) pseudo-spin eigenstates, non-abelian dynamics of cold atoms in light fields is generated, which effectively leads to the spin-orbit coupling (SOC) for cold atoms, simulating the electronic counterpart in condensed matter. Here, synthetic SOC refers to the coupling between atom's pseudo-spins (i.e. hyperfine states) and COM motion, rather than the generic interaction between electron's spin (or magnetic moment) and angular/linear momentum operator in quantum mechanics. SOC is essential in understanding numerous underlying novel quantum phenomena and particle physics [12], including *inter alia* topological insulators, Majorana and Weyl fermions, spin-Hall effects, etc [13–17].

In this work, we first briefly review our previous proposal [18], and theoretically explore the full quantum mechanical treatment beyond semi-classical mean field formalism, then investigate the correspondences in quantum and semi-classical regions. We consider a single atom (or an ensemble of \mathcal{N} non-interacting bosons) being confined by a single-mode unidirectional ring cavity, whose cavity mode together with an additional coherent laser beam form a pair of Raman beams that flips atomic transition between $|\uparrow\rangle$ and $|\downarrow\rangle$ while transferring recoil momentum of $\pm 2\hbar q_r \hat{z}$ from and/or to photon field. Hence, the so-realized effective coupling between atom's external and internal degrees of freedom is generated by the quantized light field, which is affected by atomic dynamics in return. We show that, at mean field level, the cavity-assisted SOC dramatically modifies the atomic dispersion relation, in particular, with emergence of loop structures under certain circumstances. We systematically characterize the atomic dispersion relation of atomic state and photon number, both as a function of atom's quasi-momentum. For given cavity parameters, we found with increasing Raman coupling strength Ω , dispersion curve changes from double minima to gapped double minima, looped structure, and gapless single minimum in sequence. Furthermore, we carry out the full quantum mechanical treatment by solving master equations of density operators, and find good agreement by comparing averaged photon number with mean field results in limiting parameter regions. The two distinctively different approaches give us an unified understanding of the atom-light effective non-linearity and induced dynamical instability in this system. **how do we cite papers by David Feder, Hui Zhai, Su Yi, etc in this paragraph?**

The article is organized as the following: After briefly reviewing key ideas of our previous work and semi-classical mean field approach in Sec. 2, we develop the full quantum mechanical formalism to the physical system of interest in Sec. 3 and discuss about the intimate correspondence between the two in Sec. 4, and finally conclude in Sec. 5.

2. Model Setup and Semi-classical Mean Field Formalism

We follow the effective model Hamiltonian proposed in [18],

$$\begin{aligned} \mathcal{H}_{\text{eff}} = & \sum_{\sigma} \int d\mathbf{r} \left[\psi_{\sigma}^{\dagger}(\mathbf{r}) \left(\frac{\mathbf{k}^2 + 2\alpha q_r k_z}{2m} + \alpha \delta \right) \psi_{\sigma}(\mathbf{r}) \right] + \frac{\Omega}{2} \int d\mathbf{r} \left[\psi_{\uparrow}^{\dagger}(\mathbf{r}) \psi_{\downarrow}(\mathbf{r}) c + h.c. \right] \\ & + i\varepsilon_p (c^{\dagger} - c) - \delta_c c^{\dagger} c - i\kappa c^{\dagger} c, \end{aligned} \quad (1)$$

where $\psi_{\sigma}(\mathbf{r})$ ($\sigma = \uparrow, \downarrow$) is the atomic operator after gauge transformation in rotating frame at pump frequency ω_p . $\alpha = \pm 1$ for $\sigma = \uparrow, \downarrow$, respectively. q_r denotes recoil momentum, δ represents the two-photon Raman detuning, ε_p refers to pumping rate, and δ_c is the cavity-pump detuning. Ω denotes the atom-photon coupling strength, however, the Raman coupling term $\frac{\Omega}{2} \int d\mathbf{r} [\psi_{\uparrow}^{\dagger}(\mathbf{r}) \psi_{\downarrow}(\mathbf{r}) c + h.c.]$ describes the cavity-assisted two-photon Raman transition processes, where cavity photon operators c and c^{\dagger} is explicitly taken into consideration. It is this coupling that renders the resulting SOC *dynamic*. Furthermore, in the semi-classical approach, we have treated the leakage of cavity photon phenomenologically by introducing a cavity decay rate κ .




Figure 1. Single particle eigen-energy spectrum “phase diagram”. For given parameter $\delta_c = \kappa$ and $\delta = 0$, the behavior of dispersion is divided into four categories which are represented by I to IV in (a). From (b) to (e), we fix $\varepsilon_p = \kappa$. In region I, the dispersion has double minima as shown in (b) with $\Omega = 0.03\kappa$; region II is enclosed by the red solid curve in (a) and we show the typical point in (c) ($\Omega = \kappa$) where only single minimum exists in the lower helicity band; region III is enclosed by the black dashed lines in (a) and it is a region where loop structure emerge, as in (d) with $\Omega = 5\kappa$; finally, in region IV we recover the double minimum dispersion although it's different from region I by closing the gap at $k_z = 0$, as in (e) with $\Omega = 8\kappa$. In the inset it shows a comparison between the region I boundary (red curve) and large pumping rate limit of classical result given by the function $4E_r \frac{\sqrt{\kappa^2 + \delta_c^2}}{\varepsilon_p}$.

From the Hamiltonian (1), one can easily obtain the EOM in Heisenberg picture. To make progress, we adopt the mean-field approximation by replacing the operators by their respective expectation values:

72 $c \rightarrow \langle c \rangle$, $\psi_\sigma(\mathbf{r}) \rightarrow \langle \psi_\sigma(\mathbf{r}) \rangle \equiv \varphi_\sigma(\mathbf{r})$. Assuming spatial homogeneity, we further take the plane-wave
 73 wave-function for the atomic modes $\varphi_\sigma(\mathbf{r}) = e^{i\mathbf{k}\cdot\mathbf{r}}\varphi_\sigma$ with the normalization condition $|\varphi_\uparrow|^2 + |\varphi_\downarrow|^2 = \mathcal{N}$.
 74 The steady-state solution for the photon field is obtained by taking the time derivative of the photon field
 75 to be zero, which is exact by itself without making further approximations. After some algebra, we write
 76 the coupled nonlinear time-dependent equations for the two spin components as,

$$i\dot{\varphi}_\uparrow = \left(\frac{\mathbf{k}^2}{2m} + q_r k_z + \delta \right) \varphi_\uparrow + \frac{\Omega \varepsilon_p - i\frac{\Omega}{2}\mathcal{N}\varphi_\downarrow^*\varphi_\uparrow}{2(\kappa - i\delta_c)} \varphi_\downarrow, \quad (2)$$

$$i\dot{\varphi}_\downarrow = \left(\frac{\mathbf{k}^2}{2m} - q_r k_z - \delta \right) \varphi_\downarrow + \frac{\Omega \varepsilon_p + i\frac{\Omega}{2}\mathcal{N}\varphi_\uparrow^*\varphi_\downarrow}{2(\kappa + i\delta_c)} \varphi_\uparrow. \quad (3)$$

For a given atomic quasi-momentum \mathbf{k} , we define eigenstate and eigenenergy as the solution of the time-independent version of Eqs. (2) and (3), by replacing $i(\partial/\partial t)$ with $\epsilon(\mathbf{k})$. We consider $\mathcal{N} = 1$ and always take $k_x = k_y = 0$ hereafter. After some lengthy nonetheless straightforward algebra, we find that $\epsilon(\mathbf{k})$ obeys a quartic equation

$$4\epsilon^4 + B\epsilon^3 + C\epsilon^2 + D\epsilon + E = 0, \quad (4)$$

77 where detailed derivations are better elaborated in the Supplementary Material of [18].

78 We can gain some insights about the general structure of the dispersion relation $\epsilon(\mathbf{k})$, e.g. the
 79 degeneracy condition and the appearance and disappearance of the loop. Simple analysis shows that
 80 there should be a total of four regimes. (i) As denoted by region I in Fig. 1(a) and (b), we show the
 81 dispersion curve exhibits a double minima structure. What is different from the degeneracy condition
 82 given by the classical laser induced SOC Hamiltonian, is that for arbitrarily small Ω , there *always* are
 83 double minima in the dispersion. (ii) In region II of Fig. 1(a) and (c), the dispersion curve has only single
 84 minimum. (iii) Region III corresponds to the region with loop structure and in Fig. 1(a) the two dashed
 85 black line are numerically obtained by examining the sign of second order derivative of the root with
 86 respect to k_z . When $\Omega_c^{(1)} \equiv 4\varepsilon_p \leq \Omega \leq 4\varepsilon_p \sqrt{1 + (\delta_c/\kappa)^2} \equiv \Omega_c^{(2)}$, at $k_z = 0$ there are total of four real
 87 roots allowed by the quartic equation — two degenerate roots at $\epsilon = 0$ and two additional roots with the
 88 same sign, see for instance Fig. 1(d). (iv) Finally, when $\Omega > \Omega_c^{(2)}$ in region IV, only the two degenerate
 89 roots at $\epsilon(k_z = 0) = 0$ exist and we show the double minimum dispersion curve and gapless point at
 90 $k_z = 0$.

91 We remark that the introduction of cavity photon feedback dramatically alters the dispersion relation,
 92 where one we could obtain from current proposals concerning synthetic SOC setups. As a matter
 93 of fact, due to cavity mediation, the emergent loop structure is a distinctive nonlinear feature of the
 94 system. Furthermore, upon studying the linearized perturbative expansion on top of fixed point solution
 95 to Eq. 2 and Eq. 3, we have found intriguing dynamical stability properties, where there *always* exist both
 96 dynamically stable and unstable branches regardless whether there is loop or not [18]. Nonetheless, if we
 97 consider the large limit of pumping rate ε_p , we will again be able to recover classical Raman SOC limit.
 98 In this scenario, cavity photon field becomes the coherent state. One can define the effective Raman
 99 coupling amplitude $|\tilde{\Omega}| = \frac{\Omega}{2} \frac{\varepsilon_p}{\sqrt{\kappa^2 + \delta_c^2}}$. Then one can recover the degeneracy condition for classical laser
 100 induced SOC Hamiltonian. When $|\tilde{\Omega}| < 2E_r$ or $\Omega < 4E_r \frac{\sqrt{\kappa^2 + \delta_c^2}}{\varepsilon_p}$ dispersion curve has degenerate double
 101 minimum; When $|\tilde{\Omega}| > 2E_r$ or $\Omega > 4E_r \frac{\sqrt{\kappa^2 + \delta_c^2}}{\varepsilon_p}$ we only have a single minimum in the dispersion curve.

As we show in the inset of Fig. 1, the cavity-assisted SOC determined boundary curve asymptotically approaches the black dashed line which is given by the classical correspondence of large pumping rate limit. Note, in large ε_p limit, dynamical instability decay rate γ is also strongly suppressed to a small value, such that the system fully recovers the synthetic SOC Hamiltonian.

As we will also emphasize in Sec. 4, the synthesis of cavity confinement and Raman coupling is not a trivial combination, but rather gives rise to nontrivial physics that deserves more scrutiny. In order to better address the intricate coupling between atom and photon, we dedicate the Sec. 3 and Sec. 4 to explore more on the quantum nature of this system.

3. Master Equation Approach: Full Quantum Mechanical Treatment

Semi-classical mean field approach gives an intuitive picture of understanding the atom-light interaction, as we have shown above. However, it ignores quantum fluctuations of both operators c and $\psi_\sigma(\mathbf{r})$. A more stringent approach is given by solving quantum master equation. The quantum master equations, in a nutshell, are differential equations for the density operators, including contributions from off-diagonal elements which represents quantum coherence characteristically. Master equation is generally considered to be more general than the Schrödinger equation, since it uses the density operator instead of a specific state vector and can therefore give statistical as well as quantum mechanical information.

Instead of treating the leakage of cavity photon phenomenologically in Eq. (1), we model the dissipation process by Liouvillian terms \mathcal{L} appearing in the Lindblad master equation for the atom-field density operator, i.e.,

$$\dot{\rho} = \frac{1}{i\hbar}[H_{\text{eff}}, \rho] + \mathcal{L}\rho. \quad (5)$$

where H_{eff} is the same as \mathcal{H}_{eff} in Eq. (1) by dropping the last term, i.e. $-i\kappa c^\dagger c$. Cavity loss is accommodated by the standard form of Lindblad superoperator [19,20],

$$\mathcal{L}\rho = \kappa(2c\rho c^\dagger - c^\dagger c\rho - \rho c^\dagger c). \quad (6)$$

Again, due to spatial homogeneity, we decouple momentum eigenstates by taking the plane-wave ansatz for the atomic modes as $\varphi_\sigma(\mathbf{r}) = e^{i\mathbf{k}\cdot\mathbf{r}}\varphi_\sigma$. Thereon, we are granted to work under the Hilbert subspace of a given momentum value \mathbf{k} . Here we explicitly write the commutator as,

$$\begin{aligned} [H_{\text{eff}}(\mathbf{k}), \rho] &= \left(\frac{\mathbf{k}^2}{2m} + \frac{q_r k_z}{m} + \delta \right) (\varphi_\uparrow^\dagger \psi_\uparrow \rho - \rho \varphi_\uparrow^\dagger \psi_\uparrow) + \left(\frac{\mathbf{k}^2}{2m} - \frac{q_r k_z}{m} - \delta \right) (\psi_\downarrow^\dagger \varphi_\downarrow \rho - \rho \psi_\downarrow^\dagger \varphi_\downarrow) \\ &+ \mathcal{N} \frac{\Omega}{2} (\varphi_\uparrow^\dagger \varphi_\downarrow c \rho + c^\dagger \varphi_\downarrow^\dagger \varphi_\uparrow \rho - \rho \varphi_\uparrow^\dagger \varphi_\downarrow c - \rho c^\dagger \varphi_\downarrow^\dagger \varphi_\uparrow) \\ &+ i\varepsilon_p (c^\dagger \rho - c \rho - \rho c^\dagger + \rho c) - \delta_c (c^\dagger c \rho - \rho c^\dagger c). \end{aligned} \quad (7)$$

To solve the operator equation, Eq. 5, we choose our basis state as direct product state $|n; \sigma\rangle$, where non-negative integer n denotes photon number and atomic pseudo-spin state is represented by $\sigma = \uparrow, \downarrow$. Our goal is to calculate the entire matrix elements of density operator under this set of basis states,

denoted by $\langle m; \sigma | \rho | n; \sigma' \rangle \equiv \rho_{mn}^{\sigma\sigma'}$. Without loss of generality, we have taken atom number $\mathcal{N} = 1$ to simplify discussions. We found the governing EOM can be written as,

$$\begin{aligned} \frac{d}{dt} \rho_{mn}^{\sigma\sigma'} = & -i \left(\frac{\mathbf{k}^2}{2m} + \frac{q_r k_z}{m} + \delta \right) (\delta_{\sigma\uparrow} - \delta_{\sigma'\uparrow}) \rho_{mn}^{\sigma\sigma'} - i \left(\frac{\mathbf{k}^2}{2m} - \frac{q_r k_z}{m} - \delta \right) (\delta_{\sigma\downarrow} - \delta_{\sigma'\downarrow}) \rho_{mn}^{\sigma\sigma'} \\ & + \frac{\Omega}{2i} (\delta_{\sigma\uparrow} \sqrt{m+1} \rho_{m+1n}^{\bar{\sigma}\sigma'} + \delta_{\sigma\downarrow} \sqrt{m} \rho_{m-1n}^{\bar{\sigma}\sigma'} - \delta_{\sigma'\uparrow} \sqrt{n+1} \rho_{mn+1}^{\sigma\bar{\sigma}'} - \delta_{\sigma'\downarrow} \sqrt{n} \rho_{mn-1}^{\sigma\bar{\sigma}'}) \\ & + \varepsilon_p \left(\sqrt{m} \rho_{m-1n}^{\sigma\sigma'} - \sqrt{m+1} \rho_{m+1n}^{\sigma\sigma'} + \sqrt{n} \rho_{mn-1}^{\sigma\sigma'} - \sqrt{n+1} \rho_{mn+1}^{\sigma\sigma'} \right) \\ & + i\delta_c (m-n) \rho_{mn}^{\sigma\sigma'} + \kappa \left(2\sqrt{m+1} \sqrt{n+1} \rho_{m+1n+1}^{\sigma\sigma'} - (m+n) \rho_{mn}^{\sigma\sigma'} \right), \end{aligned} \quad (8)$$

where $\bar{\sigma}$ represents the flip-spin value, i.e. $\bar{\uparrow} = \downarrow$ and $\bar{\downarrow} = \uparrow$.

With Eq. 8, we can study dynamical evolution of density operator ρ for a given initial state. For instance, we can initiate the system with a pure state $|0; \uparrow\rangle$, construct density operator $\rho = |0; \uparrow\rangle\langle 0; \uparrow|$, and let it evolve according to Eq. 8 until all components reach their respective steady state. Although at $t = 0$ we have $\text{Tr}[\rho^2] = 1$, at later times, we will always have $\text{Tr}[\rho^2] < 1$ for nonzero κ , because cavity decay term renders the system into mixed states.

Assuming the fate of time evolution gives the steady state solution, we can also solve the set of equations after equating the RHS of Eq. 8 to zero. Numerically, this is achieved by introducing a large truncation value N for maximum number of cavity photon under consideration, such that the problem reduces to simple linear algebra manipulations. In the following, we mainly focus on the steady-state solution of density operator and ignore the transient dynamics.

In order to further quantitatively characterize atom-photon interaction in this open system, we invoke the entanglement measure for mixed-state, the so-called negativity [21], defined as $\mathcal{N}(\rho) = \frac{\|\rho^{TA}\|_1 - 1}{2}$, where $\|\rho^{TA}\|_1$ denotes the trace norm of partial transpose of density operator with respect to atom party (the same is true for photon party). Density matrix ρ itself gives all positive definite eigenvalues and thus the trace norm $\|\rho\|_1 = \text{Tr}[\sqrt{\rho^\dagger \rho}] = \text{Tr}[\rho] = 1$. Although the partial transpose ρ^{TA} still satisfies $\text{Tr}[\rho^{TA}] = 1$, it does not necessarily guarantee positive definiteness in eigenvalues. The trace norm is written generally as $\|\rho^{TA}\|_1 = 1 + 2 \sum_i |\mu_i|$ where we have denoted negative eigenvalues as $\mu_i < 0$. Thus, by definition, the negativity $\mathcal{N}(\rho)$ is equal to $\sum_i |\mu_i|$, which measures by how much ρ^{TA} fails to be positive definite. An immediate consequence for any separable (unentangled) state ρ_s is that $\mathcal{N}(\rho_s) = 0$, while for unseparable mixed state, $\mathcal{N}(\rho)$ is believed to be a good entanglement measure.

4. Results and Discussions

With above preparations, we are now in a position to discuss about the results given by the two approaches, and address their relationship. As we have shown above and also in previous work [18], the cavity feedback dramatically modifies single particle dispersion relation. For instance, in intermediate region of atom-photon coupling strength of Ω , a loop structure emerge from the center tip of the eigenenergy spectrum. Additionally, in this effective nonlinear system, although atom-photon coupling is linear, dispersion spectrum possesses intriguing stability/instability properties. What we have shown in [18] also indicates that only part of the dispersion is stable for a given quasi-momentum state \mathbf{k} . The instability analysis prescribes a recipe to map out regions whether fluctuations around fixed point

157 solution would grow exponentially or not. Tout de suite, we apply the formalism developed in Sec. 2
 158 and Sec. 3 to calculate photon number expectation value inside the cavity.




Figure 2. Photon number comparison between semi-classical mean field result and full quantum mechanical master equation solution. From (a) to (c), $\Omega = 3\kappa, 5.6\kappa, 6\kappa$ and colorbar represents the renormalized decay rate γ/Ω of unstable states and red dashed lines are master equation solutions. Figures from (d) to (f) show the corresponding photon number fluctuation (blue solid curve) and negativity (green dashed line). Other parameters are $\varepsilon_p = \kappa$, $\delta_c = \kappa$, $\delta = 0$ and $\kappa = 1$ in the dimensionless unit. Although we have chosen $q_r = 0.22$ in our units (based on a realistic experimental parameter estimate) throughout all figures, we use arrows on vertical axis to denote results concerning $q_r = 0$ limit.

159 As we show in Fig. 2(a)-(c), we plot semi-classical mean field photon number $|\langle c \rangle|^2$ and quantum
 160 mechanical master equation result $\text{Tr}[\rho n]$, against atom's dimensionless quasi-momentum k_z/q_r . From
 161 semi-classical mean field result, with increasing magnitude of Ω , we sweep over regions with only two
 162 real roots, four real roots and only one double root (at $\epsilon = 0$) to the mean field solution of the quartic
 163 equation. We can further perform dynamical analysis [18] to pinpoint stable (black dots) and unstable
 164 branches of the semi-classical solution. For unstable states, we use colorbar to denote the renormalized
 165 decay rate γ/Ω where γ refers to the largest real eigenvalue of perturbed dynamical equation [18]. For
 166 comparison, with the same parameter set, we start from quantum master equation Eq. 8, and obtain steady
 167 state solution of density operator, and trace over the product of density operator and photon number, i.e.
 168 $\langle n \rangle = \text{Tr}[\rho n]$. Unlike semi-classical mean field theory, we can only have one unique solution of steady
 169 state density operator, thus only one branch of average photon number is found as a function of k_z , as
 170 represented by the red dashed line in Fig. 2(a)-(c).

171 We found, remarkably, that $\text{Tr}[\rho n]$ asymptotically approaches $|\langle c \rangle|^2$ value in dynamically stable
 172 branches at large $|k_z|$ limit. This agreement, first of all, further validates our previous dynamical
 173 stability analysis of semi-classical mean field result [18]. Due to momentum exchange of $\pm 2\hbar q_r \hat{z}$

in the Raman process, atomic states with small quasi-momentum $|k_z|$ are more coupled with photon field, in comparison to the states with large $|k_z|$. One would have naively thought cavity-assisted SOC should naturally provide more coupling between atom and photon field than the system with synthetic SOC, because atomic pseudo-spin state and COM are both coupled to cavity photon here. However, it is the SOC that renders this coupling *depends* on atomic quasi-momentum: (i) Momentum exchange plays a major role at small k_z where two SOC bands are avoided-crossing, and eigenstate is a strong hybridization between bare atomic pseudo-spin states. Thus atom and photon is strongly coupled in this region. (ii) While at large k_z limit, energy separation becomes so large that the two dressed states are more like two independent bare atomic pseudo-spin states $|\uparrow\rangle$ and $|\downarrow\rangle$. Atomic spin state, COM and photon field are all decoupled in this limit.

Second, the relationship between semi-classical mean field result and quantum master equation result can be understood from photon number fluctuation, which by definition is given by $\frac{\langle(\Delta n)^2\rangle}{\langle n\rangle} = \frac{\langle n^2\rangle - \langle n\rangle^2}{\langle n\rangle}$. In Fig. 2(d)-(f), on the left y -axis, we plot the renormalized fluctuation magnitude as a function of k_z , shown as blue solid curve. We found it degrades to one in large $|k_z|$ limit, and this is the limit where photon state is best modeled by the coherent state (Poissonian statistics) and atom's back-action *onto* photon becomes negligibly small. For small $|k_z|$ values, the majority part of fluctuation is larger than one, which implies a super-Poissonian photon number distribution, where cavity photon behaves more thermal like, which is far away from coherent state. If we have a positive definite probability distribution for photon number, by the application of Cauchy-Schwarz inequality, the fluctuation value would always have to be greater or equal to one. However, there are regions where renormalized fluctuation is smaller than one, e.g. in Fig. 2(d) at small negative k_z value, $\frac{\langle(\Delta n)^2\rangle}{\langle n\rangle} \sim 0.95$, which implies sub-Poissonian photon number distribution as a signature of system being genuinely non-classical.

Third, the degree of entanglement for a mixed-state can be quantitatively characterized by the negativity $\mathcal{N}(\rho)$ [21], which we briefly mentioned in Sec. 3. In Fig. 2(d)-(f), on the right y -axis, we plot the negativity against k_z/q_r for different Ω values using green dashed lines. We can see that when photon field approaches its coherent state at large $|k_z|$, negativity becomes very close to zero which means the “atom + photon” system becomes separable. When photon fluctuation deviates from one, negativity shows up a pronounced peak in a qualitatively similar manner.

Above observations conclude our discussion on the relationship between semi-classical theory and quantum mechanical master equation approach in the proposed cavity-assisted SOC system. Alternatively, we can also comment on the relationship between cold atom cavity QED physics and the current work. Conventionally, J-C Model is used to describe interaction between a two level atomic state and photon field by ignoring the rest degrees of freedom. In lieu of mimicking J-C Model, we set q_r to zero in the Hamiltonian and repeat the steady state solution to Eq. 5. Under this setting, atom's kinetic energy only contribute to a dynamical phase that does not affect expectation value of observable, e.g. photon number, fluctuation, etc. Then we choose an arbitrary value of k_z or ignore $k_z^2/2m$ term altogether, and compute $\text{Tr}[\rho n]$, $\frac{\langle(\Delta n)^2\rangle}{\langle n\rangle}$ and $\mathcal{N}(\rho)$ where we use arrows on the y -axis to denote their values in Fig. 2. As we can observe in Fig. 2 from (a) to (c), photon number in J-C Model limit is very different from the one by considering cavity-assisted SOC, *even* in large k_z asymptotic limit. We have $\text{Tr}[\rho n] \approx 0.49, 0.14, 0.11$ from (a) to (c), while we always have asymptotic value of $\frac{\varepsilon_p^2}{\kappa^2 + \delta_c^2}$ independent of Ω value in our system, which is 0.5 in Fig. 2. However, we are able to recover J-C model by

simultaneously setting both q_r and Ω to zero, where there is only generic atomic spin and photon coupling present. In Fig. 2 from (d) to (f), we have $\frac{\langle(\Delta n)^2\rangle}{\langle n\rangle} \approx 1.47, 1.91, 1.93$ and $\mathcal{N}(\rho) \approx 0.12, 0.06, 0.05$, respectively.

5. Conclusion and Outlook

In this work, we have studied spin-orbit coupled cold atoms inside a ring cavity system, employing both semi-classical mean field theory and full quantum mechanical master equation approach. By treating both light and atom on equal footing and seeking the self-consistent solution in both approaches, we have found cavity-assisted SOC dramatically modified atomic dispersion relation, intriguing dynamical instabilities, and atom's back-action onto light field also leads to non-trivial atom-photon coupling that are fundamentally different to the system with either synthetic SOC or J-C model type of interaction. We have also explored correspondence and discussed the relationship between the two approaches. We conclude that the synthesis of cavity QED and SOC is not a trivial combination and interesting new physics are emergent in this setting. The two distinctively different approaches give us an unified understanding of the atom-light effective non-linearity and induced dynamical instability in this system.

Although for simplicity reasons we have only considered one atom inside the cavity, the current mean field formalism and results can be easily generalized to \mathcal{N} identical non-interacting bosons, where we would expect similar physics involving atom and light couplings. However, due to anti-symmetrization constraint, for \mathcal{N} non-interacting fermions, it is less straightforward to bridge a connection. Furthermore, it would be interesting to consider atom-atom interactions for the many body case [cite Feder's paper here?](#) and study super-radiant Dicke phase transition [cite Hui Zhai's paper and Xiong-Jun Liu's super-radiant paper here?](#), using the master equation approach.

Acknowledgments

We acknowledge discussions with Zhengwei Zhou and XXX. H.P. is supported by the NSF and Welch Foundation (Grant No. C-1669 XXX);

Author Contributions

H.P. conceived the idea of the project, L.D. and C. Z. explored the theoretical and numerical aspects of the physics. All authors contributed to writing and revising the manuscript and participated in the discussions about this work.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. E.T. Jaynes, F.W. Cummings (1963). Proc. IEEE 51 (1): 89–109.
2. G. Rempe, H. Walther, and N. Klein (1987). Phys. Rev. Lett. 58 (4): 353–356.

3. Brennecke, F., Donner, T., Ritter, S., Bourdel, T., Kohl, M., and Esslinger, T., *Nature* **2007**, 450 268.
4. Colombe, Y., Steinmetz, T., Dubois, G., Linke, F., Hunger, D. and Reichel, J. *Nature* **2007**, 450 272.
5. Slama, S., Bux, S., Krenz, G., Zimmermann, C. and Courteille, Ph. W. *Phys. Rev. Lett.* **2007**, 98 053603.
6. Dicke, R. H. Coherence in spontaneous radiation processes. *Phys. Rev.* 93, 99–110 (1954).
7. Kristian Baumann, Christine Guerlin, Ferdinand Brennecke Tilman Esslinger. *Nature* 464, 1301–1306 (29 April 2010)
8. Y.-J. Lin, K. Jimenez-Garcia, and I. B. Spielman, *Nature* (London) **2011**, 471, 83.
9. Y.-J. Lin, R. L. Compton, K. Jimenez-Garcia, W. D. Phillips, J. V. Porto, and I. B. Spielman, *Nat. Phys.* **2011**, 7, 531.
10. P. Wang, Z.-Q. Yu, Z. Fu, J. Miao, L. Huang, S. Chai, H. Zhai, and J. Zhang, *Phys. Rev. Lett.* **2012**, 109, 095301.
11. L. W. Cheuk, A. T. Sommer, Z. Hadzibabic, T. Yefsah, W. S. Bakr, and M. W. Zwierlein, *Phys. Rev. Lett.* **2012**, 109, 095302.
12. V Galitski, IB Spielman, *Nature* **2013**, 494 7435.
13. Hasan, M. Z., Kane, C. L. *Rev. Mod. Phys.* **2010**, 82 3045–3067.
14. Sau, J. D., Lutchyn, R. M., Tewari, S. and Sarma, Das, S. *Phys. Rev. Lett.* **2010**, 104 040502.
15. Burkov, A. A. and Balents, L., *Phys. Rev. Lett.* **2011**, 107 127205.
16. Sinova, J., Cilcer, D., Niu, Q., Sinitsyn, N., Jungwirth, T., and MacDonald, A., *Phys. Rev. Lett.* **2004**, 92 126603.
17. Kato, Y. K., Myers, R. C., Gossard, A. C. and Awschalom, D. D., *Science* **2004**, 306 1910–1913.
18. Lin Dong, Lu Zhou, Biao Wu, B. Ramachandhran, and Han Pu, *Phys. Rev. A* **2014** 89, 011602(R).
19. Kossakowski, A. *Rep. Math. Phys.* **1972**, 3 (4): 247.
20. Lindblad, G. *Commun. Math. Phys.* **1976**, 48 (2): 119.
21. Vidal, G., Werner, R. F., *Phys. Rev. A* **2002**, 65 032314.

© March 5, 2015 by the authors; submitted to *Atoms* for possible open access publication under the terms and conditions of the Creative Commons Attribution license <http://creativecommons.org/licenses/by/4.0/>.