

Quantized Boundary Charges in 2D Spin-Orbit Coupled Crystals

Frank Schindler

University of Zurich
Kavli Institute for Theoretical Physics



University of
Zurich^{UZH}

Collaborators



Wladimir Benalcazar
The Pennsylvania State University



Marta Brzezinska
Wroclaw University of Science



Mikel Iraola
Donostia International Physics Center






Adrien Bouhon
Uppsala University

Stepan Tsirkin
University of Zurich


Maia Vergniory
Donostia International Physics Center

Titus Neupert
University of Zurich



Charge Fractionalization in 1D

Inversion symmetric
Su-Schrieffer-Heeger model






gapped bulk
quantized $1/2$ end charges




A	B	$Q_A + Q_B \in \mathbb{Z}, \quad Q_A = Q_B$ $Q_A = 0, 1/2 \pmod{1}$

0D midgap states are not protected by crystal symmetries,
but fractional boundary charges are!



Charge Fractionalization in 2D

Four-fold rotationally symmetric
obstructed atomic limit phase






$$Q_A + Q_B + Q_C + Q_D \in \mathbb{Z}, \quad Q_A = Q_B = Q_C = Q_D$$

$$Q_A = 0, 1/4, 2/4, 3/4 \pmod{1}$$

see also:

Phys. Rev. B 98, 081110(R) by **van Miert** and **Ortix**

arXiv:1809.02142 by **Benalcazar**, **Li** and **Hughes**








Spinful Time Reversal Symmetry

Introducing spin and time-reversal symmetry has two effects:

1) All charges are doubled: $Q_A \rightarrow 2Q_A$


2) We lose some topological invariants:



C_4 + TRS insulator in 2D




This phase was introduced in Phys. Rev. Lett. 119, 246402, by Z. Song, Z. Fang, C. Fang

$$H = (\mathbf{2DTI} \oplus \mathbf{2DTI}) \\ + C_4, T\text{-preserving coupling}$$






The only way to diagnose it so far has been via explicitly determining the maximally localized Wannier centers.

Our invariant: $(C_4)^4 = -1 \rightarrow C_4 = e^{\pm i\frac{\pi}{4}}, e^{\pm i\frac{3\pi}{4}}$
 $\rightarrow \mathbb{Z}_2$ **2DTI invariant in both C_4 subspaces**



Material Candidates

For the **first time**, we present material candidates for 2D charge fractionalization
Using **DFT**, we calculated the phase diagrams for **Sb, Bi, As** monolayers
as a function of tensile strain



Summary of Results

For 2D crystals with spatial symmetries and spinful time-reversal, we

1. Prove that fractionalized boundary charges are stable
2. Provide an exhaustive list of topological invariants
3. Map invariants to Wannier centres
4. Map Wannier centres to fractional boundary charges
5. Provide realistic material candidates

Thank you for your attention!

