

CANTIDAD DE MOVIMIENTO Y CHOQUES

$$\mathbf{p} = m\mathbf{v}, \quad \sum \mathbf{F} = \frac{d\mathbf{p}}{dt}$$
$$\mathbf{J} = \int \mathbf{F} dt = \Delta \mathbf{p} = \mathbf{F}_{\text{med}} \Delta t$$
$$\mathbf{r}_{\text{cm}} = \frac{\sum m_i \mathbf{r}_i}{M}, \quad \mathbf{P} = M \mathbf{v}_{\text{cm}}$$
$$\sum \mathbf{F}_{\text{ext}} = M \mathbf{a}_{\text{cm}}, \quad \sum \mathbf{F}_{\text{ext}} = 0 \Rightarrow \mathbf{P} = \text{cte}$$
$$m_1 \mathbf{v}_1 + m_2 \mathbf{v}_2 = m_1 \mathbf{v}'_1 + m_2 \mathbf{v}'_2$$
$$\mathbf{v}'_1 = \mathbf{v}'_2 = \frac{\mathbf{P}}{M}, \quad K_{\text{perd}} = \frac{1}{2} m_r v_{\text{rel}}^2$$
$$m_r = \frac{m_1 m_2}{m_1 + m_2}, \quad \mathbf{v}_{\text{rel}} = \mathbf{v}_A - \mathbf{v}_B$$
$$K = \frac{1}{2} M v_{\text{cm}}^2 + K_{\text{rel}}, \quad K = \frac{p^2}{2m}$$

COHETE

$$\mu = -\frac{dM}{dt}, \quad f = \mu |v_r| = I_{\text{sp}} g_0 \mu, \quad I_{\text{sp}} = \frac{|v_r|}{g_0}$$
$$M \frac{dV}{dt} = \mu |v_r| - Mg, \quad M(t^*) = \frac{\mu |v_r|}{g}$$
$$M(t) = M_0 - \mu t, \quad a(t) = \frac{\mu |v_r|}{M_0 - \mu t} - g$$
$$V(t) = V_0 + |v_r| \ln \frac{M_0}{M(t)} - gt$$
$$y(t) = y_0 + V_0 t - \frac{1}{2} g t^2 + |v_r| \left[t - \frac{M(t)}{\mu} \ln \frac{M_0}{M(t)} \right]$$
$$\Delta V = |v_r| \ln \frac{M_0}{M_f}, \quad \frac{M_0}{M_f} = \exp \left(\frac{\Delta V}{|v_r|} \right)$$
$$\Delta V_{\text{tot}} = \sum \Delta V_i, \quad \mathbf{F}_{\text{ext}} = \frac{d(M\mathbf{v})}{dt}$$

TRABAJO Y ENERGÍA

$$W = \int \mathbf{F} \cdot d\mathbf{l}, \quad W_{\text{tot}} = \Delta K, \quad P = \mathbf{F} \cdot \mathbf{v}$$
$$\mathbf{F} = -\nabla U, \quad E = K + U, \quad W_{\text{no cons}} = \Delta E$$

GRAVITACIÓN

$$\mathbf{F}_g = -\frac{GMm}{r^2} \hat{\mathbf{r}}, \quad U = -\frac{GMm}{r}$$
$$g_0 = \frac{GM}{R^2} = \frac{\mu}{R^2}, \quad g(h) = g_0 \left(\frac{R}{R+h} \right)^2$$
$$v_c = \sqrt{\frac{\mu}{r}}, \quad v_{\text{esc}} = \sqrt{2\mu/r} = \sqrt{2} v_c, \quad T = \frac{2\pi r}{v_c}$$
$$\Delta E_{R \rightarrow r} = \mu m \left(\frac{1}{R} - \frac{1}{r} \right)$$
$$\Delta E_{r_1 \rightarrow r_2} = \frac{\mu m}{2} \left(\frac{1}{r_1} - \frac{1}{r_2} \right)$$

MOMENTO ANGULAR

$$\mathbf{L}_O = \mathbf{r} \times m\mathbf{v}, \quad L = mrv \sin \varphi = mvd$$
$$\sum \tau_O = \frac{dL_O}{dt}, \quad \int \tau dt = \Delta L$$

$$h = rv_{\perp} = rv \cos \gamma = r^2 \dot{\theta} = \text{cte}, \quad \frac{dA}{dt} = \frac{h}{2}$$
$$r_P v_P = r_A v_A$$

DOS CUERPOS

$$\ddot{\mathbf{r}} = -\frac{\mu}{r^3} \mathbf{r}, \quad \mu = G(m_1 + m_2) \approx Gm_1$$
$$\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2, \quad \mathbf{r}_1 = \frac{m_2}{M} \mathbf{r}, \quad \mathbf{r}_2 = -\frac{m_1}{M} \mathbf{r}$$
$$E = \frac{1}{2} m \dot{r}^2 + \frac{L^2}{2mr^2} + U(r), \quad U_{\text{ef}} = U + \frac{L^2}{2mr^2}$$

ÓRBITAS

$$r = \frac{p}{1 + e \cos \nu}, \quad p = \frac{h^2}{\mu} = a(1 - e^2)$$
$$\cos \nu = \frac{1}{e} \left(\frac{p}{r} - 1 \right), \quad e \sin \nu = \frac{h v_r}{\mu}$$
$$\varepsilon = \frac{v^2}{2} - \frac{\mu}{r} = -\frac{\mu}{2a}, \quad v^2 = \mu \left(\frac{2}{r} - \frac{1}{a} \right)$$

$$e = \sqrt{1 + \frac{2\varepsilon h^2}{\mu^2}} = \frac{r_A - r_P}{r_A + r_P} = \frac{r_P v_P^2}{\mu} - 1$$
$$e^2 = \left(\frac{p}{r} - 1 \right)^2 + \left(\frac{h v_r}{\mu} \right)^2$$
$$e = \frac{r_2 - r_1}{r_1 \cos \nu_1 - r_2 \cos \nu_2}$$
$$h = rv \cos \gamma = r_P v_P = \sqrt{\mu p} = \sqrt{\mu a(1 - e^2)}$$
$$r_P = a(1 - e) = \frac{p}{1 + e}, \quad r_A = a(1 + e) = \frac{p}{1 - e}$$
$$a = \frac{r_P + r_A}{2}, \quad b = a\sqrt{1 - e^2} = \sqrt{r_P r_A}$$
$$\frac{1}{2} v_P^2 - \frac{\mu}{r_P} = \frac{1}{2} v_A^2 - \frac{\mu}{r_A}$$
$$v_P = \sqrt{\frac{2\mu r_A}{r_P(r_P + r_A)}}, \quad v_A = \frac{r_P}{r_A} v_P$$
$$v_P = \sqrt{\frac{\mu}{a} \frac{1 + e}{1 - e}}, \quad v_A = \sqrt{\frac{\mu}{a} \frac{1 - e}{1 + e}}$$
$$v_{\perp} = \frac{h}{r} = \frac{\mu}{h} (1 + e \cos \nu), \quad v_r = \frac{\mu}{h} e \sin \nu$$
$$\tan \gamma = \frac{v_r}{v_{\perp}} = \frac{e \sin \nu}{1 + e \cos \nu}$$

KEPLER

$$T = 2\pi \sqrt{\frac{a^3}{\mu}} = \frac{2\pi ab}{h}, \quad n = \frac{2\pi}{T} = \sqrt{\frac{\mu}{a^3}}$$
$$\tan \frac{E}{2} = \sqrt{\frac{1 - e}{1 + e}} \tan \frac{\nu}{2}, \quad \cos E = \frac{e + \cos \nu}{1 + e \cos \nu}$$
$$M_e = E - e \sin E = n(t - t_P)$$
$$r = a(1 - e \cos E)$$
$$T_{\text{sid}} = 24 \text{ h} \cdot \frac{360^\circ}{360,986^\circ} = 23,934 \text{ h}$$

MANIOBRAS

$$v_P = \sqrt{\frac{2\mu r_2}{r_1(r_1 + r_2)}}, \quad v_A = \frac{r_1}{r_2} v_P$$
$$\Delta v_1 = v_P - v_{c1}, \quad \Delta v_2 = v_{c2} - v_A, \quad v_{ci} = \sqrt{\frac{\mu}{r_i}}$$
$$a_t = \frac{r_1 + r_2}{2}, \quad t_v = \frac{T_t}{2} = \pi \sqrt{\frac{a_t^3}{\mu}}$$
$$\varphi = 180^\circ - n_2 t_v, \quad n_2 = 360^\circ / T_2$$
$$\frac{T'}{T} = 1 - \frac{\Delta \varphi}{360^\circ}, \quad a' = r \left(\frac{T'}{T} \right)^{2/3}$$
$$r_{P'} = 2a' - r$$

CONSTANTES

G	$6,674 \times 10^{-11} \text{ N}\cdot\text{m}^2/\text{kg}^2$
μ_T, R_T	$398\,600 \text{ km}^3/\text{s}^2, 6378 \text{ km}$
g_0	$9,80665 \text{ m/s}^2$
μ_{Luna}	$4903 \text{ km}^3/\text{s}^2, d_{\text{T-L}} \approx 384\,400 \text{ km}$
μ_{Sol}	$1,327 \times 10^{11} \text{ km}^3/\text{s}^2, 1 \text{ UA} = 149,6 \times 10^6 \text{ km}$

HIPÉRBOLA Y ESCAPE

$$e = 1: \quad \varepsilon = 0, \quad v = v_{\text{esc}} = \sqrt{\frac{2\mu}{r}}$$
$$\varepsilon = \frac{v_{\infty}^2}{2} = \frac{\mu}{2a}, \quad v^2 = v_{\text{esc}}^2 + v_{\infty}^2, \quad C_3 = v_{\infty}^2$$
$$a = \frac{p}{e^2 - 1} = \frac{\mu}{v_{\infty}^2}, \quad r_P = a(e - 1)$$
$$e = 1 + \frac{r_P v_{\infty}^2}{\mu}$$
$$\nu_{\infty} = \arccos \left(-\frac{1}{e} \right), \quad \delta = 2 \arcsin \left(\frac{1}{e} \right)$$
$$\Delta v_{\text{esc}} = \sqrt{\frac{2\mu}{r}} - v$$

ELEMENTOS ORBITALES

$$(\mathbf{r}, \mathbf{v}) \leftrightarrow (a, e, i, \Omega, \omega, \nu_0)$$

$$\mathbf{A} \times \mathbf{B} = \begin{vmatrix} \hat{I} & \hat{J} & \hat{K} \\ A_I & A_J & A_K \\ B_I & B_J & B_K \end{vmatrix}$$
$$= (A_J B_K - A_K B_J) \hat{I} - (A_I B_K - A_K B_I) \hat{J} + (A_I B_J - A_J B_I) \hat{K}$$
$$|\mathbf{A} \times \mathbf{B}| = AB \sin \alpha, \quad \mathbf{A} \times \mathbf{B} = -\mathbf{B} \times \mathbf{A}$$
$$\hat{I} \times \hat{J} = \hat{K}, \quad \hat{J} \times \hat{K} = \hat{I}, \quad \hat{K} \times \hat{I} = \hat{J}$$
$$\mathbf{A} \cdot \mathbf{B} = A_I B_I + A_J B_J + A_K B_K = AB \cos \alpha$$
$$\mathbf{r} \cdot \mathbf{v} = r v_r = r v \sin \gamma$$
$$\mathbf{h} = \mathbf{r} \times \mathbf{v}, \quad \mathbf{n} = \hat{K} \times \mathbf{h} = (-h_J, h_I, 0)$$
$$\mathbf{e} = \frac{1}{\mu} \left[\left(v^2 - \frac{\mu}{r} \right) \mathbf{r} - (\mathbf{r} \cdot \mathbf{v}) \mathbf{v} \right]$$
$$p = \frac{h^2}{\mu}, \quad e = |e|, \quad a = \frac{p}{1 - e^2} = -\frac{\mu}{2\varepsilon}$$
$$\cos i = \frac{h_K}{h}$$
$$\cos \Omega = \frac{n_I}{n} \quad (n_J < 0 \Rightarrow \Omega > 180^\circ)$$
$$\cos \omega = \frac{\mathbf{n} \cdot \mathbf{e}}{ne} \quad (e_K < 0 \Rightarrow \omega > 180^\circ)$$
$$\cos \nu_0 = \frac{\mathbf{e} \cdot \mathbf{r}}{er} \quad (\mathbf{r} \cdot \mathbf{v} < 0 \Rightarrow \nu_0 > 180^\circ)$$
$$i = 0: \quad \Pi = \Omega + \omega$$
$$\cos \Pi = \frac{e_I}{e} \quad (e_J < 0 \Rightarrow \Pi > 180^\circ)$$
$$e = 0: \quad u_0 = \omega + \nu_0$$
$$\cos u_0 = \frac{\mathbf{n} \cdot \mathbf{r}}{nr} \quad (r_K < 0 \Rightarrow u_0 > 180^\circ)$$
$$i = e = 0: \quad l_0 = \Omega + \omega + \nu_0$$
$$\cos l_0 = \frac{r_I}{r} \quad (r_J < 0 \Rightarrow l_0 > 180^\circ)$$
$$\mathbf{r} = r \cos \nu \hat{P} + r \sin \nu \hat{Q}$$
$$\mathbf{v} = \frac{\mu}{h} [-\sin \nu \hat{P} + (e + \cos \nu) \hat{Q}]$$

ROTACIÓN Y GIRÓSCOPO

$$v = \omega r, \quad a_t = \alpha r, \quad a_{\text{rad}} = \omega^2 r, \quad K = \frac{1}{2} I \omega^2$$
$$I = \sum m_i r_i^2, \quad I_P = I_{\text{cm}} + M d^2$$

varilla (cm)	$ML^2/12$	aro	MR^2
varilla (ext)	$ML^2/3$	disco	$MR^2/2$
esfera	$2MR^2/5$	esf. hueca	$2MR^2/3$

$$\tau = rF \sin \varphi, \quad \sum \tau_z = I \alpha_z, \quad L = I \omega$$
$$\sum \tau_{\text{ext}} = \frac{dL}{dt}$$
$$\sum \tau_{\text{ext}} = 0 \Rightarrow \sum I_i \omega_i = \text{cte}$$
$$K = \frac{1}{2} M v_{\text{cm}}^2 + \frac{1}{2} I_{\text{cm}} \omega^2, \quad v_{\text{cm}} = R \omega$$
$$\Delta L = \tau \Delta t, \quad \Omega = \frac{\tau}{L} = \frac{M g r}{I \omega}$$