

Binary Outcomes

Francisco Villamil

Applied Quantitative Methods II

IC3JM, Spring 2026

Today's goals

- Understand why OLS is problematic for binary outcomes

Today's goals

- Understand why OLS is problematic for binary outcomes
- Learn the linear probability model and its trade-offs

Today's goals

- Understand why OLS is problematic for binary outcomes
- Learn the linear probability model and its trade-offs
- Understand logistic regression and how to estimate it in R

Today's goals

- Understand why OLS is problematic for binary outcomes
- Learn the linear probability model and its trade-offs
- Understand logistic regression and how to estimate it in R
- Interpret logit results using predicted probabilities and marginal effects

Roadmap

The Problem with Binary Outcomes

Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):

Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):
 - Did someone vote?

Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):
 - Did someone vote?
 - Did a war break out?

Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):
 - Did someone vote?
 - Did a war break out?
 - Did a bill pass?

Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):
 - Did someone vote?
 - Did a war break out?
 - Did a bill pass?
 - Did a country democratize?

Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):
 - Did someone vote?
 - Did a war break out?
 - Did a bill pass?
 - Did a country democratize?

Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):
 - Did someone vote?
 - Did a war break out?
 - Did a bill pass?
 - Did a country democratize?
- Our outcome $Y \in \{0, 1\}$

Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):
 - Did someone vote?
 - Did a war break out?
 - Did a bill pass?
 - Did a country democratize?
- Our outcome $Y \in \{0, 1\}$
- We want to model: $P(Y = 1|X)$

What happens if we just use OLS?

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

- OLS gives us: $E[Y|X] = \beta_0 + \beta_1 X$

What happens if we just use OLS?

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

- OLS gives us: $E[Y|X] = \beta_0 + \beta_1 X$
- Since $Y \in \{0, 1\}$: $E[Y|X] = P(Y = 1|X)$

What happens if we just use OLS?

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

- OLS gives us: $E[Y|X] = \beta_0 + \beta_1 X$
- Since $Y \in \{0, 1\}$: $E[Y|X] = P(Y = 1|X)$

What happens if we just use OLS?

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

- OLS gives us: $E[Y|X] = \beta_0 + \beta_1 X$
- Since $Y \in \{0, 1\}$: $E[Y|X] = P(Y = 1|X)$
- So OLS is modeling a probability as a linear function of X

What happens if we just use OLS?

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

- OLS gives us: $E[Y|X] = \beta_0 + \beta_1 X$
- Since $Y \in \{0, 1\}$: $E[Y|X] = P(Y = 1|X)$
- So OLS is modeling a probability as a linear function of X
- This is the **linear probability model** (LPM)

The LPM: Simple and intuitive

- β_1 has a direct interpretation:

The LPM: Simple and intuitive

- β_1 has a direct interpretation:
 - A one-unit increase in X changes the probability of $Y = 1$ by β_1

The LPM: Simple and intuitive

- β_1 has a direct interpretation:
 - A one-unit increase in X changes the probability of $Y = 1$ by β_1

The LPM: Simple and intuitive

- β_1 has a direct interpretation:
 - A one-unit increase in X changes the probability of $Y = 1$ by β_1
- Easy to estimate: just `lm(y ~ x, data = df)`

The LPM: Simple and intuitive

- β_1 has a direct interpretation:
 - A one-unit increase in X changes the probability of $Y = 1$ by β_1
- Easy to estimate: just `lm(y ~ x, data = df)`
- Easy to interpret: same as OLS

The LPM: Simple and intuitive

- β_1 has a direct interpretation:
 - A one-unit increase in X changes the probability of $Y = 1$ by β_1
- Easy to estimate: just `lm(y ~ x, data = df)`
- Easy to interpret: same as OLS

The LPM: Simple and intuitive

- β_1 has a direct interpretation:
 - A one-unit increase in X changes the probability of $Y = 1$ by β_1
- Easy to estimate: just `lm(y ~ x, data = df)`
- Easy to interpret: same as OLS
- Many applied researchers use the LPM in practice

LPM limitations

- **Problem 1:** Predictions outside $[0, 1]$

LPM limitations

- **Problem 1:** Predictions outside $[0, 1]$
 - A linear function can produce $\hat{P} < 0$ or $\hat{P} > 1$

LPM limitations

- **Problem 1:** Predictions outside $[0, 1]$
 - A linear function can produce $\hat{P} < 0$ or $\hat{P} > 1$
 - Probabilities must be between 0 and 1!

LPM limitations

- **Problem 1:** Predictions outside $[0, 1]$
 - A linear function can produce $\hat{P} < 0$ or $\hat{P} > 1$
 - Probabilities must be between 0 and 1!

LPM limitations

- **Problem 1:** Predictions outside $[0, 1]$
 - A linear function can produce $\hat{P} < 0$ or $\hat{P} > 1$
 - Probabilities must be between 0 and 1!
- **Problem 2:** Heteroskedasticity by construction

LPM limitations

- **Problem 1:** Predictions outside $[0, 1]$
 - A linear function can produce $\hat{P} < 0$ or $\hat{P} > 1$
 - Probabilities must be between 0 and 1!
- **Problem 2:** Heteroskedasticity by construction
 - $\text{Var}(\varepsilon|X) = P(1 - P)$, which varies with X

LPM limitations

- **Problem 1:** Predictions outside $[0, 1]$
 - A linear function can produce $\hat{P} < 0$ or $\hat{P} > 1$
 - Probabilities must be between 0 and 1!
- **Problem 2:** Heteroskedasticity by construction
 - $\text{Var}(\varepsilon|X) = P(1 - P)$, which varies with X
 - Standard errors from OLS are wrong (use robust SEs)

LPM limitations

- **Problem 1:** Predictions outside $[0, 1]$
 - A linear function can produce $\hat{P} < 0$ or $\hat{P} > 1$
 - Probabilities must be between 0 and 1!
- **Problem 2:** Heteroskedasticity by construction
 - $\text{Var}(\varepsilon|X) = P(1 - P)$, which varies with X
 - Standard errors from OLS are wrong (use robust SEs)

LPM limitations

- **Problem 1:** Predictions outside $[0, 1]$
 - A linear function can produce $\hat{P} < 0$ or $\hat{P} > 1$
 - Probabilities must be between 0 and 1!
- **Problem 2:** Heteroskedasticity by construction
 - $\text{Var}(\varepsilon|X) = P(1 - P)$, which varies with X
 - Standard errors from OLS are wrong (use robust SEs)
- **Problem 3:** Non-linearity at the extremes

LPM limitations

- **Problem 1:** Predictions outside $[0, 1]$
 - A linear function can produce $\hat{P} < 0$ or $\hat{P} > 1$
 - Probabilities must be between 0 and 1!
- **Problem 2:** Heteroskedasticity by construction
 - $\text{Var}(\varepsilon|X) = P(1 - P)$, which varies with X
 - Standard errors from OLS are wrong (use robust SEs)
- **Problem 3:** Non-linearity at the extremes
 - True relationship between X and $P(Y = 1)$ is S-shaped

LPM limitations

- **Problem 1:** Predictions outside $[0, 1]$
 - A linear function can produce $\hat{P} < 0$ or $\hat{P} > 1$
 - Probabilities must be between 0 and 1!
- **Problem 2:** Heteroskedasticity by construction
 - $\text{Var}(\varepsilon|X) = P(1 - P)$, which varies with X
 - Standard errors from OLS are wrong (use robust SEs)
- **Problem 3:** Non-linearity at the extremes
 - True relationship between X and $P(Y = 1)$ is S-shaped
 - LPM forces it to be linear

When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)

When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
 - The linear approximation is reasonable here

When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
 - The linear approximation is reasonable here
- When you care about **average marginal effects**

When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
 - The linear approximation is reasonable here
- When you care about **average marginal effects**
 - LPM and logit often give similar AMEs

When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
 - The linear approximation is reasonable here
- When you care about **average marginal effects**
 - LPM and logit often give similar AMEs
- When simplicity of interpretation matters

When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
 - The linear approximation is reasonable here
- When you care about **average marginal effects**
 - LPM and logit often give similar AMEs
- When simplicity of interpretation matters

When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
 - The linear approximation is reasonable here
- When you care about **average marginal effects**
 - LPM and logit often give similar AMEs
- When simplicity of interpretation matters
- When is it **not** good enough?

When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
 - The linear approximation is reasonable here
- When you care about **average marginal effects**
 - LPM and logit often give similar AMEs
- When simplicity of interpretation matters
- When is it **not** good enough?
 - Many observations near 0 or 1

When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
 - The linear approximation is reasonable here
- When you care about **average marginal effects**
 - LPM and logit often give similar AMEs
- When simplicity of interpretation matters
- When is it **not** good enough?
 - Many observations near 0 or 1
 - You need predicted probabilities to be bounded

When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
 - The linear approximation is reasonable here
- When you care about **average marginal effects**
 - LPM and logit often give similar AMEs
- When simplicity of interpretation matters
- When is it **not** good enough?
 - Many observations near 0 or 1
 - You need predicted probabilities to be bounded
 - The relationship is clearly non-linear

Roadmap

Logistic Regression

The logistic function

$$P(Y = 1|X) = \frac{e^{X\beta}}{1 + e^{X\beta}} = \frac{1}{1 + e^{-X\beta}}$$

- This is an S-shaped (sigmoid) curve

The logistic function

$$P(Y = 1|X) = \frac{e^{X\beta}}{1 + e^{X\beta}} = \frac{1}{1 + e^{-X\beta}}$$

- This is an S-shaped (sigmoid) curve
- Output is always between 0 and 1

The logistic function

$$P(Y = 1|X) = \frac{e^{X\beta}}{1 + e^{X\beta}} = \frac{1}{1 + e^{-X\beta}}$$

- This is an S-shaped (sigmoid) curve
- Output is always between 0 and 1
- Steep in the middle, flat at the extremes

The logistic function

$$P(Y = 1|X) = \frac{e^{X\beta}}{1 + e^{X\beta}} = \frac{1}{1 + e^{-X\beta}}$$

- This is an S-shaped (sigmoid) curve
- Output is always between 0 and 1
- Steep in the middle, flat at the extremes
- A natural model for probabilities

The logit model

We can rearrange the logistic function:

$$\log\left(\frac{P}{1-P}\right) = \beta_0 + \beta_1 X$$

- The left side is the **log-odds** (or “logit”)

The logit model

We can rearrange the logistic function:

$$\log\left(\frac{P}{1-P}\right) = \beta_0 + \beta_1 X$$

- The left side is the **log-odds** (or “logit”)
- $\frac{P}{1-P}$ is the **odds** of the event

The logit model

We can rearrange the logistic function:

$$\log \left(\frac{P}{1-P} \right) = \beta_0 + \beta_1 X$$

- The left side is the **log-odds** (or “logit”)
- $\frac{P}{1-P}$ is the **odds** of the event
- The model is linear in the log-odds, not in the probability

Maximum likelihood estimation

- We can't use OLS for logistic regression

Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)

Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)

Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)
- The intuition:

Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)
- The intuition:
 - For each observation, the model predicts $P(Y_i = 1)$

Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)
- The intuition:
 - For each observation, the model predicts $P(Y_i = 1)$
 - MLE finds the coefficients that make the observed data most likely

Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)
- The intuition:
 - For each observation, the model predicts $P(Y_i = 1)$
 - MLE finds the coefficients that make the observed data most likely
 - The “likelihood” is the product of these predicted probabilities

Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)
- The intuition:
 - For each observation, the model predicts $P(Y_i = 1)$
 - MLE finds the coefficients that make the observed data most likely
 - The “likelihood” is the product of these predicted probabilities

Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)
- The intuition:
 - For each observation, the model predicts $P(Y_i = 1)$
 - MLE finds the coefficients that make the observed data most likely
 - The “likelihood” is the product of these predicted probabilities
- No need to derive this—R does it for us

Estimating logit in R

- The function: `glm(y ~ x, family = binomial, data = df)`

Estimating logit in R

- The function: `glm(y ~ x, family = binomial, data = df)`

Estimating logit in R

- The function: `glm(y ~ x, family = binomial, data = df)`
- `glm`: generalized linear model

Estimating logit in R

- The function: `glm(y ~ x, family = binomial, data = df)`
- `glm`: generalized linear model
- `family = binomial`: tells R to use logistic regression

Estimating logit in R

- The function: `glm(y ~ x, family = binomial, data = df)`
- `glm`: generalized linear model
- `family = binomial`: tells R to use logistic regression

Estimating logit in R

- The function: `glm(y ~ x, family = binomial, data = df)`
- `glm`: generalized linear model
- `family = binomial`: tells R to use logistic regression
- The syntax is identical to `lm()`, just change to `glm()`

Estimating logit in R

- The function: `glm(y ~ x, family = binomial, data = df)`
- `glm`: generalized linear model
- `family = binomial`: tells R to use logistic regression
- The syntax is identical to `lm()`, just change to `glm()`
- Works with `broom::tidy()`, `modelsummary()`, etc.

Interpreting logit output: Log-odds

- The direct output gives coefficients in **log-odds**

Interpreting logit output: Log-odds

- The direct output gives coefficients in **log-odds**

Interpreting logit output: Log-odds

- The direct output gives coefficients in **log-odds**
- $\beta_1 = 0.5$ means:

Interpreting logit output: Log-odds

- The direct output gives coefficients in **log-odds**
- $\beta_1 = 0.5$ means:
 - A one-unit increase in X increases the log-odds by 0.5

Interpreting logit output: Log-odds

- The direct output gives coefficients in **log-odds**
- $\beta_1 = 0.5$ means:
 - A one-unit increase in X increases the log-odds by 0.5

Interpreting logit output: Log-odds

- The direct output gives coefficients in **log-odds**
- $\beta_1 = 0.5$ means:
 - A one-unit increase in X increases the log-odds by 0.5
- This is hard to interpret!

Interpreting logit output: Log-odds

- The direct output gives coefficients in **log-odds**
- $\beta_1 = 0.5$ means:
 - A one-unit increase in X increases the log-odds by 0.5
- This is hard to interpret!
- Nobody thinks in log-odds

Interpreting logit output: Odds ratios

- Exponentiate the coefficient: $e^{\beta_1} = \text{odds ratio}$

Interpreting logit output: Odds ratios

- Exponentiate the coefficient: $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`

Interpreting logit output: Odds ratios

- Exponentiate the coefficient: $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`

Interpreting logit output: Odds ratios

- Exponentiate the coefficient: $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`
- $e^{0.5} \approx 1.65$ means:

Interpreting logit output: Odds ratios

- Exponentiate the coefficient: $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`
- $e^{0.5} \approx 1.65$ means:
 - A one-unit increase in X **multiplies** the odds by 1.65

Interpreting logit output: Odds ratios

- Exponentiate the coefficient: $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`
- $e^{0.5} \approx 1.65$ means:
 - A one-unit increase in X **multiplies** the odds by 1.65
 - Or: the odds increase by 65%

Interpreting logit output: Odds ratios

- Exponentiate the coefficient: $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`
- $e^{0.5} \approx 1.65$ means:
 - A one-unit increase in X **multiplies** the odds by 1.65
 - Or: the odds increase by 65%

Interpreting logit output: Odds ratios

- Exponentiate the coefficient: $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`
- $e^{0.5} \approx 1.65$ means:
 - A one-unit increase in X **multiplies** the odds by 1.65
 - Or: the odds increase by 65%
- Slightly more intuitive, but still not probabilities

Interpreting logit output: Odds ratios

- Exponentiate the coefficient: $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`
- $e^{0.5} \approx 1.65$ means:
 - A one-unit increase in X **multiplies** the odds by 1.65
 - Or: the odds increase by 65%
- Slightly more intuitive, but still not probabilities
- The change in probability depends on where you start

Why coefficients alone are not enough

- In OLS: β_1 = change in Y for one-unit change in X (always)

Why coefficients alone are not enough

- In OLS: β_1 = change in Y for one-unit change in X (always)

Why coefficients alone are not enough

- In OLS: β_1 = change in Y for one-unit change in X (always)
- In logit: the change in **probability** depends on:

Why coefficients alone are not enough

- In OLS: β_1 = change in Y for one-unit change in X (always)
- In logit: the change in **probability** depends on:
 - The current value of X

Why coefficients alone are not enough

- In OLS: β_1 = change in Y for one-unit change in X (always)
- In logit: the change in **probability** depends on:
 - The current value of X
 - The values of all other variables

Why coefficients alone are not enough

- In OLS: β_1 = change in Y for one-unit change in X (always)
- In logit: the change in **probability** depends on:
 - The current value of X
 - The values of all other variables

Why coefficients alone are not enough

- In OLS: β_1 = change in Y for one-unit change in X (always)
- In logit: the change in **probability** depends on:
 - The current value of X
 - The values of all other variables
- A coefficient of $\beta_1 = 0.5$ could mean:

Why coefficients alone are not enough

- In OLS: β_1 = change in Y for one-unit change in X (always)
- In logit: the change in **probability** depends on:
 - The current value of X
 - The values of all other variables
- A coefficient of $\beta_1 = 0.5$ could mean:
 - Going from $P = 0.01$ to $P = 0.016$ (tiny change)

Why coefficients alone are not enough

- In OLS: β_1 = change in Y for one-unit change in X (always)
- In logit: the change in **probability** depends on:
 - The current value of X
 - The values of all other variables
- A coefficient of $\beta_1 = 0.5$ could mean:
 - Going from $P = 0.01$ to $P = 0.016$ (tiny change)
 - Going from $P = 0.50$ to $P = 0.62$ (large change)

Why coefficients alone are not enough

- In OLS: β_1 = change in Y for one-unit change in X (always)
- In logit: the change in **probability** depends on:
 - The current value of X
 - The values of all other variables
- A coefficient of $\beta_1 = 0.5$ could mean:
 - Going from $P = 0.01$ to $P = 0.016$ (tiny change)
 - Going from $P = 0.50$ to $P = 0.62$ (large change)

Why coefficients alone are not enough

- In OLS: β_1 = change in Y for one-unit change in X (always)
- In logit: the change in **probability** depends on:
 - The current value of X
 - The values of all other variables
- A coefficient of $\beta_1 = 0.5$ could mean:
 - Going from $P = 0.01$ to $P = 0.016$ (tiny change)
 - Going from $P = 0.50$ to $P = 0.62$ (large change)
- We need better tools to interpret logit models

Roadmap

Interpreting Logit Results

Predicted probabilities

- The most intuitive way to interpret logit models

Predicted probabilities

- The most intuitive way to interpret logit models

Predicted probabilities

- The most intuitive way to interpret logit models
- “What is the predicted probability of $Y = 1$ for a person with these characteristics?”

Predicted probabilities

- The most intuitive way to interpret logit models
- “What is the predicted probability of $Y = 1$ for a person with these characteristics?”

Predicted probabilities

- The most intuitive way to interpret logit models
- “What is the predicted probability of $Y = 1$ for a person with these characteristics?”
- In R:

Predicted probabilities

- The most intuitive way to interpret logit models
- “What is the predicted probability of $Y = 1$ for a person with these characteristics?”
- In R:
 - `marginalEffects::predictions(model)`

Predicted probabilities

- The most intuitive way to interpret logit models
- “What is the predicted probability of $Y = 1$ for a person with these characteristics?”
- In R:
 - `marginalEffects::predictions(model)`
 - Returns predicted probabilities for each observation

Predicted probabilities

- The most intuitive way to interpret logit models
- “What is the predicted probability of $Y = 1$ for a person with these characteristics?”
- In R:
 - `marginalEffects::predictions(model)`
 - Returns predicted probabilities for each observation
 - Or at specific values: `predictions(model, newdata = ...)`

Average marginal effects (AME)

- The marginal effect varies across observations

Average marginal effects (AME)

- The marginal effect varies across observations
- The AME averages across all observations

Average marginal effects (AME)

- The marginal effect varies across observations
- The AME averages across all observations

Average marginal effects (AME)

- The marginal effect varies across observations
- The AME averages across all observations
- In R: `marginaleffects::avg_slopes(model)`

Average marginal effects (AME)

- The marginal effect varies across observations
- The AME averages across all observations
- In R: `marginaleffects::avg_slopes(model)`

Average marginal effects (AME)

- The marginal effect varies across observations
- The AME averages across all observations
- In R: `marginaleffects::avg_slopes(model)`
- Interpretation (like OLS):

Average marginal effects (AME)

- The marginal effect varies across observations
- The AME averages across all observations
- In R: `marginalEffects::avg_slopes(model)`
- Interpretation (like OLS):
 - “On average, a one-unit increase in X changes the probability of $Y = 1$ by ΔP ”

Average marginal effects (AME)

- The marginal effect varies across observations
- The AME averages across all observations
- In R: `marginalEffects::avg_slopes(model)`
- Interpretation (like OLS):
 - “On average, a one-unit increase in X changes the probability of $Y = 1$ by ΔP ”

Average marginal effects (AME)

- The marginal effect varies across observations
- The AME averages across all observations
- In R: `marginaleffects::avg_slopes(model)`
- Interpretation (like OLS):
 - “On average, a one-unit increase in X changes the probability of $Y = 1$ by ΔP ”
- This is often comparable to the LPM coefficient

Marginal effects at representative values

- Instead of averaging, evaluate at specific values

Marginal effects at representative values

- Instead of averaging, evaluate at specific values

Marginal effects at representative values

- Instead of averaging, evaluate at specific values
- Example: “What is the effect of education on voting for a 40-year-old woman?”

Marginal effects at representative values

- Instead of averaging, evaluate at specific values
- Example: “What is the effect of education on voting for a 40-year-old woman?”

Marginal effects at representative values

- Instead of averaging, evaluate at specific values
- Example: “What is the effect of education on voting for a 40-year-old woman?”
- In R:

Marginal effects at representative values

- Instead of averaging, evaluate at specific values
- Example: “What is the effect of education on voting for a 40-year-old woman?”
- In R:
 - `avg_slopes(model, newdata = datagrid(age = 40, gender = "F"))`

Marginal effects at representative values

- Instead of averaging, evaluate at specific values
- Example: “What is the effect of education on voting for a 40-year-old woman?”
- In R:
 - `avg_slopes(model, newdata = datagrid(age = 40, gender = "F"))`

Marginal effects at representative values

- Instead of averaging, evaluate at specific values
- Example: “What is the effect of education on voting for a 40-year-old woman?”
- In R:
 - `avg_slopes(model, newdata = datagrid(age = 40, gender = "F"))`
- Useful when the marginal effect varies a lot across the sample

Plotting predicted probabilities

- The best way to communicate logit results

Plotting predicted probabilities

- The best way to communicate logit results

Plotting predicted probabilities

- The best way to communicate logit results
- Show how $P(Y = 1)$ changes across values of X

Plotting predicted probabilities

- The best way to communicate logit results
- Show how $P(Y = 1)$ changes across values of X
- Include confidence bands

Plotting predicted probabilities

- The best way to communicate logit results
- Show how $P(Y = 1)$ changes across values of X
- Include confidence bands

Plotting predicted probabilities

- The best way to communicate logit results
- Show how $P(Y = 1)$ changes across values of X
- Include confidence bands
- In R:

Plotting predicted probabilities

- The best way to communicate logit results
- Show how $P(Y = 1)$ changes across values of X
- Include confidence bands
- In R:
 - `plot_predictions(model, condition = "x")`

Plotting predicted probabilities

- The best way to communicate logit results
- Show how $P(Y = 1)$ changes across values of X
- Include confidence bands
- In R:
 - `plot_predictions(model, condition = "x")`
 - Plots the S-curve with uncertainty

Plotting predicted probabilities

- The best way to communicate logit results
- Show how $P(Y = 1)$ changes across values of X
- Include confidence bands
- In R:
 - `plot_predictions(model, condition = "x")`
 - Plots the S-curve with uncertainty

Plotting predicted probabilities

- The best way to communicate logit results
- Show how $P(Y = 1)$ changes across values of X
- Include confidence bands
- In R:
 - `plot_predictions(model, condition = "x")`
 - Plots the S-curve with uncertainty
- Much more informative than a table of log-odds

Comparing LPM and logit

- In many cases, LPM and logit give similar results

Comparing LPM and logit

- In many cases, LPM and logit give similar results
 - Especially for average marginal effects

Comparing LPM and logit

- In many cases, LPM and logit give similar results
 - Especially for average marginal effects
 - Especially when probabilities are in the 0.2–0.8 range

Comparing LPM and logit

- In many cases, LPM and logit give similar results
 - Especially for average marginal effects
 - Especially when probabilities are in the 0.2–0.8 range

Comparing LPM and logit

- In many cases, LPM and logit give similar results
 - Especially for average marginal effects
 - Especially when probabilities are in the 0.2–0.8 range
- Where they differ:

Comparing LPM and logit

- In many cases, LPM and logit give similar results
 - Especially for average marginal effects
 - Especially when probabilities are in the 0.2–0.8 range
- Where they differ:
 - Predicted probabilities near 0 or 1

Comparing LPM and logit

- In many cases, LPM and logit give similar results
 - Especially for average marginal effects
 - Especially when probabilities are in the 0.2–0.8 range
- Where they differ:
 - Predicted probabilities near 0 or 1
 - LPM can go outside $[0, 1]$; logit cannot

Comparing LPM and logit

- In many cases, LPM and logit give similar results
 - Especially for average marginal effects
 - Especially when probabilities are in the 0.2–0.8 range
- Where they differ:
 - Predicted probabilities near 0 or 1
 - LPM can go outside $[0, 1]$; logit cannot
 - Marginal effects at extreme values

Comparing LPM and logit

- In many cases, LPM and logit give similar results
 - Especially for average marginal effects
 - Especially when probabilities are in the 0.2–0.8 range
- Where they differ:
 - Predicted probabilities near 0 or 1
 - LPM can go outside $[0, 1]$; logit cannot
 - Marginal effects at extreme values

Comparing LPM and logit

- In many cases, LPM and logit give similar results
 - Especially for average marginal effects
 - Especially when probabilities are in the 0.2–0.8 range
- Where they differ:
 - Predicted probabilities near 0 or 1
 - LPM can go outside $[0, 1]$; logit cannot
 - Marginal effects at extreme values
- A good practice: estimate both and compare

Model fit for logit

- No R^2 in the usual sense (MLE, not OLS)

Model fit for logit

- No R^2 in the usual sense (MLE, not OLS)

Model fit for logit

- No R^2 in the usual sense (MLE, not OLS)
- Alternative measures:

Model fit for logit

- No R^2 in the usual sense (MLE, not OLS)
- Alternative measures:
 - **Pseudo- R^2** : compares model to null model (McFadden)

Model fit for logit

- No R^2 in the usual sense (MLE, not OLS)
- Alternative measures:
 - **Pseudo- R^2** : compares model to null model (McFadden)
 - **AIC**: penalized likelihood (lower = better)

Model fit for logit

- No R^2 in the usual sense (MLE, not OLS)
- Alternative measures:
 - **Pseudo- R^2** : compares model to null model (McFadden)
 - **AIC**: penalized likelihood (lower = better)
 - **Classification**: what percent does the model correctly predict?

Model fit for logit

- No R^2 in the usual sense (MLE, not OLS)
- Alternative measures:
 - **Pseudo- R^2** : compares model to null model (McFadden)
 - **AIC**: penalized likelihood (lower = better)
 - **Classification**: what percent does the model correctly predict?
 - **ROC curve**: trade-off between true and false positives

Model fit for logit

- No R^2 in the usual sense (MLE, not OLS)
- Alternative measures:
 - **Pseudo- R^2** : compares model to null model (McFadden)
 - **AIC**: penalized likelihood (lower = better)
 - **Classification**: what percent does the model correctly predict?
 - **ROC curve**: trade-off between true and false positives

Model fit for logit

- No R^2 in the usual sense (MLE, not OLS)
- Alternative measures:
 - **Pseudo- R^2** : compares model to null model (McFadden)
 - **AIC**: penalized likelihood (lower = better)
 - **Classification**: what percent does the model correctly predict?
 - **ROC curve**: trade-off between true and false positives
- None is perfect; use them as rough guides

Model fit for logit

- No R^2 in the usual sense (MLE, not OLS)
- Alternative measures:
 - **Pseudo- R^2** : compares model to null model (McFadden)
 - **AIC**: penalized likelihood (lower = better)
 - **Classification**: what percent does the model correctly predict?
 - **ROC curve**: trade-off between true and false positives
- None is perfect; use them as rough guides
- Reported automatically by `modelsummary()` and `performance::r2()`

Roadmap

Practice

Worked example: LPM vs. logit

```
lpm <- lm(vote ~ age + income + educ, data = df)
logit <- glm(vote ~ age + income + educ,
             family = binomial, data = df)
```


Worked example: LPM vs. logit

```
lpm <- lm(vote ~ age + income + educ, data = df)
logit <- glm(vote ~ age + income + educ,
             family = binomial, data = df)
modelsummary(list("LPM" = lpm, "Logit" = logit))
```

Worked example: LPM vs. logit

```
lpm <- lm(vote ~ age + income + educ, data = df)
logit <- glm(vote ~ age + income + educ,
             family = binomial, data = df)
modelsummary(list("LPM" = lpm, "Logit" = logit))
avg_slopes(logit)                # compare to LPM coefficients
```

Worked example: LPM vs. logit

```
lpm <- lm(vote ~ age + income + educ, data = df)
logit <- glm(vote ~ age + income + educ,
             family = binomial, data = df)
modelsummary(list("LPM" = lpm, "Logit" = logit))
avg_slopes(logit)                # compare to LPM coefficients
plot_predictions(logit, condition = "income")
```

Decision tree: When to use which?

- **Use LPM when:**

Decision tree: When to use which?

- **Use LPM when:**

- You want simple, quick interpretation

Decision tree: When to use which?

- **Use LPM when:**

- You want simple, quick interpretation
- Probabilities are in the middle range

Decision tree: When to use which?

- **Use LPM when:**

- You want simple, quick interpretation
- Probabilities are in the middle range
- You mainly care about average effects

Decision tree: When to use which?

- **Use LPM when:**

- You want simple, quick interpretation
- Probabilities are in the middle range
- You mainly care about average effects

Decision tree: When to use which?

- **Use LPM when:**

- You want simple, quick interpretation
- Probabilities are in the middle range
- You mainly care about average effects

- **Use logit when:**

Decision tree: When to use which?

- **Use LPM when:**

- You want simple, quick interpretation
- Probabilities are in the middle range
- You mainly care about average effects

- **Use logit when:**

- You need bounded predicted probabilities

Decision tree: When to use which?

- **Use LPM when:**

- You want simple, quick interpretation
- Probabilities are in the middle range
- You mainly care about average effects

- **Use logit when:**

- You need bounded predicted probabilities
- Many observations have extreme probabilities (near 0 or 1)

Decision tree: When to use which?

- **Use LPM when:**

- You want simple, quick interpretation
- Probabilities are in the middle range
- You mainly care about average effects

- **Use logit when:**

- You need bounded predicted probabilities
- Many observations have extreme probabilities (near 0 or 1)
- You want to properly account for the binary nature of Y

Decision tree: When to use which?

- **Use LPM when:**

- You want simple, quick interpretation
- Probabilities are in the middle range
- You mainly care about average effects

- **Use logit when:**

- You need bounded predicted probabilities
- Many observations have extreme probabilities (near 0 or 1)
- You want to properly account for the binary nature of Y

Decision tree: When to use which?

- **Use LPM when:**

- You want simple, quick interpretation
- Probabilities are in the middle range
- You mainly care about average effects

- **Use logit when:**

- You need bounded predicted probabilities
- Many observations have extreme probabilities (near 0 or 1)
- You want to properly account for the binary nature of Y

- In practice: estimate both, report the one most appropriate

Summary: Key takeaways

- Binary outcomes require special treatment

Summary: Key takeaways

- Binary outcomes require special treatment
- LPM is simple but has known limitations

Summary: Key takeaways

- Binary outcomes require special treatment
- LPM is simple but has known limitations
- Logit bounds probabilities between 0 and 1

Summary: Key takeaways

- Binary outcomes require special treatment
- LPM is simple but has known limitations
- Logit bounds probabilities between 0 and 1
- Log-odds and odds ratios are not intuitive—use marginal effects

Summary: Key takeaways

- Binary outcomes require special treatment
- LPM is simple but has known limitations
- Logit bounds probabilities between 0 and 1
- Log-odds and odds ratios are not intuitive—use marginal effects
- Predicted probabilities and AMEs are the best way to interpret logit

Summary: Key takeaways

- Binary outcomes require special treatment
- LPM is simple but has known limitations
- Logit bounds probabilities between 0 and 1
- Log-odds and odds ratios are not intuitive—use marginal effects
- Predicted probabilities and AMEs are the best way to interpret logit
- Compare LPM and logit; they often agree on AMEs

For next week

- Read Urdinez & Cruz (2020), chapter 5 (§5.6)
- Read Gelman et al., chapters 11–12
- Complete Assignment 3

- Next session: Model interpretation and diagnostics
 - Beyond coefficient tables
 - Visualizing model results
 - Publication-quality tables
 - Key diagnostics

Questions?