

# Model Interpretation and Diagnostics

Francisco Villamil

Applied Quantitative Methods II  
MA in Social Sciences, Spring 2026

## Today's goals

- Move beyond coefficient tables to meaningful quantities

## Today's goals

- Move beyond coefficient tables to meaningful quantities
- Compute and plot predicted values, marginal effects, and first differences

## Today's goals

- Move beyond coefficient tables to meaningful quantities
- Compute and plot predicted values, marginal effects, and first differences
- Present results with publication-quality tables and coefficient plots


## Today's goals

- Move beyond coefficient tables to meaningful quantities
- Compute and plot predicted values, marginal effects, and first differences
- Present results with publication-quality tables and coefficient plots
- Understand simulation-based uncertainty

## Today's goals

- Move beyond coefficient tables to meaningful quantities
- Compute and plot predicted values, marginal effects, and first differences
- Present results with publication-quality tables and coefficient plots
- Understand simulation-based uncertainty
- Diagnose common regression problems: heteroskedasticity, non-linearity

# A motivating puzzle: Simpson's paradox




- Overall trend is **negative**, but within each group it's **positive**


Opt.50pt Ignoring  $Z$  gives the **wrong answer** — we need to account for it

## Controlling vs. interacting

**Control:**  $Y = \beta_1 X + \beta_2 Z$



**Interaction:**  $+ \beta_3 X \cdot Z$



- **Controlling:** adjusts the level (intercept) — the effect of  $X$  stays the same
- **Interacting:** allows the **slope** of  $X$  to differ across groups



# Roadmap

Beyond Coefficient Tables

Predicted Values and Marginal Effects

Presenting Results

Diagnostics

Wrap-up

## The interpretation problem

- We know how to estimate models and read output

## The interpretation problem

- We know how to estimate models and read output
- But what do the numbers actually **mean**?

## The interpretation problem

- We know how to estimate models and read output
- But what do the numbers actually **mean**?
- Raw coefficients are often not directly informative:

## The interpretation problem

- We know how to estimate models and read output
- But what do the numbers actually **mean**?
- Raw coefficients are often not directly informative:
  - Logit: coefficients are in log-odds (last week)

## The interpretation problem

- We know how to estimate models and read output
- But what do the numbers actually **mean**?
- Raw coefficients are often not directly informative:
  - Logit: coefficients are in log-odds (last week)
  - Interactions: the coefficient on  $X_1$  depends on  $X_2$

## The interpretation problem


- We know how to estimate models and read output
- But what do the numbers actually **mean**?
- Raw coefficients are often not directly informative:
  - Logit: coefficients are in log-odds (last week)
  - Interactions: the coefficient on  $X_1$  depends on  $X_2$
  - Log transformations: coefficients are elasticities or semi-elasticities

## The interpretation problem

- We know how to estimate models and read output
- But what do the numbers actually **mean**?
- Raw coefficients are often not directly informative:
  - Logit: coefficients are in log-odds (last week)
  - Interactions: the coefficient on  $X_1$  depends on  $X_2$
  - Log transformations: coefficients are elasticities or semi-elasticities
  - Different scales: is  $\hat{\beta} = 0.003$  big or small?



## Coefficients vs. quantities of interest



You estimate a model with GDP (in dollars), population (in millions), and an interaction between them.

What does the coefficient on GDP tell you?

# Roadmap

Beyond Coefficient Tables

Predicted Values and Marginal Effects

Presenting Results

Diagnostics

Wrap-up

Predicted values:  $E[Y \mid X]$

- The most intuitive quantity: “What does the model predict for this scenario?”

Predicted values:  $E[Y \mid X]$

- The most intuitive quantity: “What does the model predict for this scenario?”
- Set predictors to specific values, compute  $\hat{Y}$

Predicted values:  $E[Y \mid X]$

- The most intuitive quantity: “What does the model predict for this scenario?”
- Set predictors to specific values, compute  $\hat{Y}$
- Example: “What is predicted income for a 35-year-old with a college degree?”

Predicted values:  $E[Y \mid X]$

- The most intuitive quantity: “What does the model predict for this scenario?”
- Set predictors to specific values, compute  $\hat{Y}$
- Example: “What is predicted income for a 35-year-old with a college degree?”
- In R:

Predicted values:  $E[Y \mid X]$

- The most intuitive quantity: “What does the model predict for this scenario?”
- Set predictors to specific values, compute  $\hat{Y}$
- Example: “What is predicted income for a 35-year-old with a college degree?”
- In R:
  - `predictions(model)`



Predicted values:  $E[Y \mid X]$

- The most intuitive quantity: “What does the model predict for this scenario?”
- Set predictors to specific values, compute  $\hat{Y}$
- Example: “What is predicted income for a 35-year-old with a college degree?”
- In R:
  - `predictions(model)`
  - `predictions(model,  
                  newdata = datagrid(age = 35, college = 1))`

## Plotting predictions

- The gold standard for communicating model results

## Plotting predictions

- The gold standard for communicating model results
- `plot_predictions(model, condition = "age")`

## Plotting predictions

- The gold standard for communicating model results
- `plot_predictions(model, condition = "age")`
  - Shows  $\hat{Y}$  across values of age

## Plotting predictions

- The gold standard for communicating model results
- `plot_predictions(model, condition = "age")`
  - Shows  $\hat{Y}$  across values of age
  - Includes 95% confidence band (for the mean, not individual predictions)

## Plotting predictions

- The gold standard for communicating model results
- `plot_predictions(model, condition = "age")`
  - Shows  $\hat{Y}$  across values of age
  - Includes 95% confidence band (for the mean, not individual predictions)
  - Holds other variables at their means (or modes)

## Plotting predictions

- The gold standard for communicating model results
- `plot_predictions(model, condition = "age")`
  - Shows  $\hat{Y}$  across values of age
  - Includes 95% confidence band (for the mean, not individual predictions)
  - Holds other variables at their means (or modes)
- With two conditions:

## Plotting predictions

- The gold standard for communicating model results
- `plot_predictions(model, condition = "age")`
  - Shows  $\hat{Y}$  across values of age
  - Includes 95% confidence band (for the mean, not individual predictions)
  - Holds other variables at their means (or modes)
- With two conditions:
  - `plot_predictions(model, condition = c("age", "female"))`



## Plotting predictions

- The gold standard for communicating model results
- `plot_predictions(model, condition = "age")`
  - Shows  $\hat{Y}$  across values of age
  - Includes 95% confidence band (for the mean, not individual predictions)
  - Holds other variables at their means (or modes)
- With two conditions:
  - `plot_predictions(model, condition = c("age", "female"))`
  - Separate lines/panels for each group

## Marginal effects

$$\text{Marginal effect of } X_1 = \frac{\partial E[Y | X]}{\partial X_1}$$

- “How much does  $Y$  change for a small change in  $X_1$ ?”

## Marginal effects

$$\text{Marginal effect of } X_1 = \frac{\partial E[Y | X]}{\partial X_1}$$

- “How much does  $Y$  change for a small change in  $X_1$ ?”
- For OLS without interactions: marginal effect =  $\hat{\beta}_1$  (constant)

## Marginal effects

$$\text{Marginal effect of } X_1 = \frac{\partial E[Y | X]}{\partial X_1}$$

- “How much does  $Y$  change for a small change in  $X_1$ ?”
- For OLS without interactions: marginal effect =  $\hat{\beta}_1$  (constant)
- For logit, interactions, polynomials: marginal effect **varies**

## Marginal effects

$$\text{Marginal effect of } X_1 = \frac{\partial E[Y | X]}{\partial X_1}$$

- “How much does  $Y$  change for a small change in  $X_1$ ?”
- For OLS without interactions: marginal effect =  $\hat{\beta}_1$  (constant)
- For logit, interactions, polynomials: marginal effect **varies**
- **Average marginal effect (AME)**: average across all observations


## Marginal effects

$$\text{Marginal effect of } X_1 = \frac{\partial E[Y | X]}{\partial X_1}$$

- “How much does  $Y$  change for a small change in  $X_1$ ?”
- For OLS without interactions: marginal effect =  $\hat{\beta}_1$  (constant)
- For logit, interactions, polynomials: marginal effect **varies**
- **Average marginal effect (AME)**: average across all observations  
→ `avg_slopes(model)`

## Marginal effects with interactions

$$Y = \beta_0 + \beta_1 \text{Education} + \beta_2 \text{Female} + \beta_3 (\text{Education} \times \text{Female}) + \varepsilon$$



## Conditional marginal effects in R

- Marginal effect of  $X_1$  at specific values of  $X_2$ :



## Conditional marginal effects in R

- Marginal effect of  $X_1$  at specific values of  $X_2$ :
  - `slopes(model, variables = "education",  
          newdata = datagrid(female = c(0, 1)))`

## Conditional marginal effects in R

- Marginal effect of  $X_1$  at specific values of  $X_2$ :
  - `slopes(model, variables = "education",  
          newdata = datagrid(female = c(0, 1)))`
- Plot how the marginal effect varies:

## Conditional marginal effects in R

- Marginal effect of  $X_1$  at specific values of  $X_2$ :
  - `slopes(model, variables = "education",  
          newdata = datagrid(female = c(0, 1)))`
- Plot how the marginal effect varies:
  - `plot_slopes(model, variables = "education",  
              condition = "female")`

## Conditional marginal effects in R

- Marginal effect of  $X_1$  at specific values of  $X_2$ :
  - `slopes(model, variables = "education",  
          newdata = datagrid(female = c(0, 1)))`
- Plot how the marginal effect varies:
  - `plot_slopes(model, variables = "education",  
              condition = "female")`
- This replaces the old manual approach of computing  $\beta_1 + \beta_3 \cdot X_2$  by hand

## First differences (comparisons)

- “How different is  $\hat{Y}$  between two scenarios?”

## First differences (comparisons)

- “How different is  $\hat{Y}$  between two scenarios?”
- Example: predicted income for college vs. no college

## First differences (comparisons)

- “How different is  $\hat{Y}$  between two scenarios?”
- Example: predicted income for college vs. no college  
→  $\Delta = E[Y \mid \text{college} = 1] - E[Y \mid \text{college} = 0]$

## First differences (comparisons)

- “How different is  $\hat{Y}$  between two scenarios?”
- Example: predicted income for college vs. no college  
→  $\Delta = E[Y \mid \text{college} = 1] - E[Y \mid \text{college} = 0]$
- In R:



## First differences (comparisons)

- “How different is  $\hat{Y}$  between two scenarios?”
- Example: predicted income for college vs. no college
  - $\Delta = E[Y \mid \text{college} = 1] - E[Y \mid \text{college} = 0]$
- In R:
  - `comparisons(model,  
                  variables = list(college = c(0, 1)))`

## First differences (comparisons)

- “How different is  $\hat{Y}$  between two scenarios?”
- Example: predicted income for college vs. no college
  - $\Delta = E[Y \mid \text{college} = 1] - E[Y \mid \text{college} = 0]$
- In R:
  - `comparisons(model,`  
                  `variables = list(college = c(0, 1)))`
  - `avg_comparisons(model, variables = "college")`

## First differences (comparisons)

- “How different is  $\hat{Y}$  between two scenarios?”
- Example: predicted income for college vs. no college
  - $\Delta = E[Y \mid \text{college} = 1] - E[Y \mid \text{college} = 0]$
- In R:
  - `comparisons(model,`  
                  `variables = list(college = c(0, 1)))`
  - `avg_comparisons(model, variables = "college")`
- Especially useful for:

## First differences (comparisons)

- “How different is  $\hat{Y}$  between two scenarios?”
- Example: predicted income for college vs. no college
  - $\Delta = E[Y \mid \text{college} = 1] - E[Y \mid \text{college} = 0]$
- In R:
  - `comparisons(model,`  
                  `variables = list(college = c(0, 1)))`
  - `avg_comparisons(model, variables = "college")`
- Especially useful for:
  - Binary/categorical predictors

## First differences (comparisons)

- “How different is  $\hat{Y}$  between two scenarios?”
- Example: predicted income for college vs. no college
  - $\Delta = E[Y \mid \text{college} = 1] - E[Y \mid \text{college} = 0]$
- In R:
  - `comparisons(model,`  
                  `variables = list(college = c(0, 1)))`
  - `avg_comparisons(model, variables = "college")`
- Especially useful for:
  - Binary/categorical predictors
  - Comparing meaningful scenarios (e.g., min vs. max)

## Choosing the right quantity

Quantity	Question	R function
Predicted value	What does the model predict here?	<code>predictions()</code>
Marginal effect	How much does $Y$ change per unit of $X$ ?	<code>slopes()</code>
Average ME	What is the average effect across the sample?	<code>avg_slopes()</code>
First difference	How different are two scenarios?	<code>comparisons()</code>

- All from the `marginalEffects` package (Arel-Bundock et al.)

## Choosing the right quantity

Quantity	Question	R function
Predicted value	What does the model predict here?	<code>predictions()</code>
Marginal effect	How much does $Y$ change per unit of $X$ ?	<code>slopes()</code>
Average ME	What is the average effect across the sample?	<code>avg_slopes()</code>
First difference	How different are two scenarios?	<code>comparisons()</code>

- All from the `marginalEffects` package (Arel-Bundock et al.)
- All compute standard errors and CIs automatically

## Choosing the right quantity

Quantity	Question	R function
Predicted value	What does the model predict here?	<code>predictions()</code>
Marginal effect	How much does $Y$ change per unit of $X$ ?	<code>slopes()</code>
Average ME	What is the average effect across the sample?	<code>avg_slopes()</code>
First difference	How different are two scenarios?	<code>comparisons()</code>

- All from the `marginalEffects` package (Arel-Bundock et al.)
- All compute standard errors and CIs automatically
- All work with `lm()`, `glm()`, and many other model types



# Roadmap

Beyond Coefficient Tables

Predicted Values and Marginal Effects

Presenting Results

Diagnostics

Wrap-up

Publication-quality tables with

# `modelsummary()`

- We already know `modelsummary()` from Session 2

Publication-quality tables with

# `modelsummary()`

- We already know `modelsummary()` from Session 2
- Key options:

## Publication-quality tables with `modelsummary()`

- We already know `modelsummary()` from Session 2
- Key options:
  - `stars = TRUE` for significance stars

## Publication-quality tables with `modelsummary()`

- We already know `modelsummary()` from Session 2
- Key options:
  - `stars = TRUE` for significance stars
  - `vcov = "robust"` for robust SEs

## Publication-quality tables with `modelsummary()`

- We already know `modelsummary()` from Session 2
- Key options:
  - `stars = TRUE` for significance stars
  - `vcov = "robust"` for robust SEs
  - `coef_rename` to clean up variable names

## Publication-quality tables with `modelsummary()`

- We already know `modelsummary()` from Session 2
- Key options:
  - `stars = TRUE` for significance stars
  - `vcov = "robust"` for robust SEs
  - `coef_rename` to clean up variable names
  - `gof_map` to select goodness-of-fit statistics

## Publication-quality tables with `modelsummary()`

- We already know `modelsummary()` from Session 2
- Key options:
  - `stars = TRUE` for significance stars
  - `vcov = "robust"` for robust SEs
  - `coef_rename` to clean up variable names
  - `gof_map` to select goodness-of-fit statistics
- Multiple models side by side:



## Publication-quality tables with `modelsummary()`

- We already know `modelsummary()` from Session 2
- Key options:
  - `stars = TRUE` for significance stars
  - `vcov = "robust"` for robust SEs
  - `coef_rename` to clean up variable names
  - `gof_map` to select goodness-of-fit statistics
- Multiple models side by side:
  - `modelsummary(list("OLS" = m1, "Logit" = m2))`

## Publication-quality tables with `modelsummary()`

- We already know `modelsummary()` from Session 2
- Key options:
  - `stars = TRUE` for significance stars
  - `vcov = "robust"` for robust SEs
  - `coef_rename` to clean up variable names
  - `gof_map` to select goodness-of-fit statistics
- Multiple models side by side:
  - `modelsummary(list("OLS" = m1, "Logit" = m2))`
- Output to LaTeX, Word, HTML, or the console

# Coefficient plots with `modelplot()`

- Visual alternative to coefficient tables

# Coefficient plots with `modelplot()`

- Visual alternative to coefficient tables
- `modelplot(model)`

# Coefficient plots with `modelplot()`

- Visual alternative to coefficient tables
- `modelplot(model)`
  - Point = estimate, line = 95% CI

# Coefficient plots with `modelplot()`

- Visual alternative to coefficient tables
- `modelplot(model)`
  - Point = estimate, line = 95% CI
  - Vertical line at zero = “no effect”

# Coefficient plots with `modelplot()`

- Visual alternative to coefficient tables
- `modelplot(model)`
  - Point = estimate, line = 95% CI
  - Vertical line at zero = “no effect”
- When are coefficient plots better than tables?

# Coefficient plots with `modelplot()`

- Visual alternative to coefficient tables
- `modelplot(model)`
  - Point = estimate, line = 95% CI
  - Vertical line at zero = “no effect”
- When are coefficient plots better than tables?
  - Many predictors



# Coefficient plots with `modelplot()`

- Visual alternative to coefficient tables
- `modelplot(model)`
  - Point = estimate, line = 95% CI
  - Vertical line at zero = “no effect”
- When are coefficient plots better than tables?
  - Many predictors
  - Comparing across model specifications


# Coefficient plots with `modelplot()`

- Visual alternative to coefficient tables
- `modelplot(model)`
  - Point = estimate, line = 95% CI
  - Vertical line at zero = “no effect”
- When are coefficient plots better than tables?
  - Many predictors
  - Comparing across model specifications
  - Audience that finds tables intimidating

# Coefficient plots with `modelplot()`

- Visual alternative to coefficient tables
- `modelplot(model)`
  - Point = estimate, line = 95% CI
  - Vertical line at zero = “no effect”
- When are coefficient plots better than tables?
  - Many predictors
  - Comparing across model specifications
  - Audience that finds tables intimidating
- `modelplot(list("Model 1" = m1, "Model 2" = m2))`

## Prediction plots: the gold standard



- Shows the **full relationship**, not just one number
- Any audience can read it

## Tables vs. plots: when to use which

	<b>Table</b>	<b>Plot</b>
Exact numbers needed	✓	
Many models side by side	✓	
Conveying one key relationship		✓
Non-specialist audience		✓
Interactions / non-linearities		✓
Appendix / robustness checks	✓	

- Best practice: prediction plots in the main text, tables in the appendix

## Tables vs. plots: when to use which

	<b>Table</b>	<b>Plot</b>
Exact numbers needed	✓	
Many models side by side	✓	
Conveying one key relationship		✓
Non-specialist audience		✓
Interactions / non-linearities		✓
Appendix / robustness checks	✓	

- Best practice: prediction plots in the main text, tables in the appendix
- Both are easy to produce with `modelsummary` and `marginaleffects`

## Simulation-based uncertainty

- All quantities of interest have **uncertainty**

## Simulation-based uncertainty

- All quantities of interest have **uncertainty**
- Standard approach: delta method (a linear approximation for SEs of nonlinear functions)



## Simulation-based uncertainty

- All quantities of interest have **uncertainty**
- Standard approach: delta method (a linear approximation for SEs of nonlinear functions)
- Alternative: **simulation**

## Simulation-based uncertainty

- All quantities of interest have **uncertainty**
- Standard approach: delta method (a linear approximation for SEs of nonlinear functions)
- Alternative: **simulation**
  - Draw  $\tilde{\beta}$  from  $N(\hat{\beta}, \hat{V})$  many times

## Simulation-based uncertainty

- All quantities of interest have **uncertainty**
- Standard approach: delta method (a linear approximation for SEs of nonlinear functions)
- Alternative: **simulation**
  - Draw  $\tilde{\beta}$  from  $N(\hat{\beta}, \hat{V})$  many times
  - Compute the quantity of interest for each draw

## Simulation-based uncertainty

- All quantities of interest have **uncertainty**
- Standard approach: delta method (a linear approximation for SEs of nonlinear functions)
- Alternative: **simulation**
  - Draw  $\tilde{\beta}$  from  $N(\hat{\beta}, \hat{V})$  many times
  - Compute the quantity of interest for each draw
  - Use the distribution of results as the uncertainty estimate

## Simulation-based uncertainty

- All quantities of interest have **uncertainty**
- Standard approach: delta method (a linear approximation for SEs of nonlinear functions)
- Alternative: **simulation**
  - Draw  $\tilde{\beta}$  from  $N(\hat{\beta}, \hat{V})$  many times
  - Compute the quantity of interest for each draw
  - Use the distribution of results as the uncertainty estimate
- Why simulate?

## Simulation-based uncertainty

- All quantities of interest have **uncertainty**
- Standard approach: delta method (a linear approximation for SEs of nonlinear functions)
- Alternative: **simulation**
  - Draw  $\tilde{\beta}$  from  $N(\hat{\beta}, \hat{V})$  many times
  - Compute the quantity of interest for each draw
  - Use the distribution of results as the uncertainty estimate
- Why simulate?
  - Works for any quantity, no matter how complex

## Simulation-based uncertainty


- All quantities of interest have **uncertainty**
- Standard approach: delta method (a linear approximation for SEs of nonlinear functions)
- Alternative: **simulation**
  - Draw  $\tilde{\beta}$  from  $N(\hat{\beta}, \hat{V})$  many times
  - Compute the quantity of interest for each draw
  - Use the distribution of results as the uncertainty estimate
- Why simulate?
  - Works for any quantity, no matter how complex
  - Easy to combine multiple quantities

## Simulation-based uncertainty

- All quantities of interest have **uncertainty**
- Standard approach: delta method (a linear approximation for SEs of nonlinear functions)
- Alternative: **simulation**
  - Draw  $\tilde{\beta}$  from  $N(\hat{\beta}, \hat{V})$  many times
  - Compute the quantity of interest for each draw
  - Use the distribution of results as the uncertainty estimate
- Why simulate?
  - Works for any quantity, no matter how complex
  - Easy to combine multiple quantities
  - Confidence intervals are automatic



## Simulation: the logic



## Worked example: predicted values with uncertainty

```
m = lm(income ~ age + education + female, data = df)

# Using marginaleffects (delta method)
predictions(m,
  newdata = datagrid(age = 40, education = 16, female = 1))

# Using simulation
library(MASS)
sim_b = mvrnorm(1000, coef(m), vcov(m))
x = c(1, 40, 16, 1) # intercept, age, educ, female
sim_pred = sim_b %*% x
quantile(sim_pred, c(0.025, 0.5, 0.975))
```

# Roadmap

Beyond Coefficient Tables

Predicted Values and Marginal Effects

Presenting Results

Diagnostics

Wrap-up

## What can go wrong?

- OLS assumes:

## What can go wrong?

- OLS assumes:
  - Linear relationship between  $X$  and  $E[Y]$

## What can go wrong?

- OLS assumes:
  - Linear relationship between  $X$  and  $E[Y]$
  - Constant error variance (**homoskedasticity**)

## What can go wrong?

- OLS assumes:
  - Linear relationship between  $X$  and  $E[Y]$
  - Constant error variance (**homoskedasticity**)
  - No extreme outliers driving results

## What can go wrong?

- OLS assumes:
  - Linear relationship between  $X$  and  $E[Y]$
  - Constant error variance (**homoskedasticity**)
  - No extreme outliers driving results
- When these fail:



## What can go wrong?

- OLS assumes:
  - Linear relationship between  $X$  and  $E[Y]$
  - Constant error variance (**homoskedasticity**)
  - No extreme outliers driving results
- When these fail:
  - Coefficients may be biased (non-linearity)

## What can go wrong?

- OLS assumes:
  - Linear relationship between  $X$  and  $E[Y]$
  - Constant error variance (**homoskedasticity**)
  - No extreme outliers driving results
- When these fail:
  - Coefficients may be biased (non-linearity)
  - Standard errors may be wrong (heteroskedasticity)



## What can go wrong?

- OLS assumes:
  - Linear relationship between  $X$  and  $E[Y]$
  - Constant error variance (**homoskedasticity**)
  - No extreme outliers driving results
- When these fail:
  - Coefficients may be biased (non-linearity)
  - Standard errors may be wrong (heteroskedasticity)
  - Results may be driven by a few observations (outliers)

## What can go wrong?

- OLS assumes:
  - Linear relationship between  $X$  and  $E[Y]$
  - Constant error variance (**homoskedasticity**)
  - No extreme outliers driving results
- When these fail:
  - Coefficients may be biased (non-linearity)
  - Standard errors may be wrong (heteroskedasticity)
  - Results may be driven by a few observations (outliers)
- How do we check? **Residual diagnostics**

## Residuals vs. fitted values



- In R: `plot(model, which = 1)`
- Look for patterns: funnel shape, curves, clusters

You plot residuals vs. fitted values and see a clear funnel shape.

What does this tell you about your model?

What would you do about it?

## Heteroskedasticity

- **Heteroskedasticity:**  $\text{Var}(\varepsilon \mid X)$  is not constant

## Heteroskedasticity

- **Heteroskedasticity:**  $\text{Var}(\varepsilon \mid X)$  is not constant
- Consequences:



## Heteroskedasticity

- **Heteroskedasticity:**  $\text{Var}(\varepsilon \mid X)$  is not constant
- Consequences:
  - Coefficients are still unbiased

## Heteroskedasticity

- **Heteroskedasticity:**  $\text{Var}(\varepsilon \mid X)$  is not constant
- Consequences:
  - Coefficients are still unbiased
  - But standard errors are **wrong**

## Heteroskedasticity

- **Heteroskedasticity:**  $\text{Var}(\varepsilon \mid X)$  is not constant
- Consequences:
  - Coefficients are still unbiased
  - But standard errors are **wrong**
  - So confidence intervals and p-values are unreliable

## Heteroskedasticity

- **Heteroskedasticity:**  $\text{Var}(\varepsilon \mid X)$  is not constant
- Consequences:
  - Coefficients are still unbiased
  - But standard errors are **wrong**
  - So confidence intervals and p-values are unreliable
- Very common in practice:

## Heteroskedasticity

- **Heteroskedasticity:**  $\text{Var}(\varepsilon \mid X)$  is not constant
- Consequences:
  - Coefficients are still unbiased
  - But standard errors are **wrong**
  - So confidence intervals and p-values are unreliable
- Very common in practice:
  - Income models (richer countries, more variance)

## Heteroskedasticity

- **Heteroskedasticity:**  $\text{Var}(\varepsilon \mid X)$  is not constant
- Consequences:
  - Coefficients are still unbiased
  - But standard errors are **wrong**
  - So confidence intervals and p-values are unreliable
- Very common in practice:
  - Income models (richer countries, more variance)
  - Any outcome with natural bounds

# Heteroskedasticity

- **Heteroskedasticity:**  $\text{Var}(\varepsilon \mid X)$  is not constant
- Consequences:
  - Coefficients are still unbiased
  - But standard errors are **wrong**
  - So confidence intervals and p-values are unreliable
- Very common in practice:
  - Income models (richer countries, more variance)
  - Any outcome with natural bounds
  - Binary outcomes (always heteroskedastic, as in LPM)

## Robust standard errors

- The fix: **heteroskedasticity-consistent (HC) standard errors**



## Robust standard errors

- The fix: **heteroskedasticity-consistent (HC) standard errors**
- Also called “robust” or “White” standard errors

## Robust standard errors

- The fix: **heteroskedasticity-consistent (HC) standard errors**
- Also called “robust” or “White” standard errors
- In R (multiple options):

## Robust standard errors

- The fix: **heteroskedasticity-consistent (HC) standard errors**
- Also called “robust” or “White” standard errors
- In R (multiple options):
  - `modelsummary(model, vcov = "robust")`

## Robust standard errors

- The fix: **heteroskedasticity-consistent (HC) standard errors**
- Also called “robust” or “White” standard errors
- In R (multiple options):
  - `modelsummary(model, vcov = "robust")`
  - `modelsummary(model, vcov = "HC3")`

## Robust standard errors

- The fix: **heteroskedasticity-consistent (HC) standard errors**
- Also called “robust” or “White” standard errors
- In R (multiple options):
  - `modelsummary(model, vcov = "robust")`
  - `modelsummary(model, vcov = "HC3")`
  - `estimatr::lm_robust(y ~ x, data = df)`

## Robust standard errors

- The fix: **heteroskedasticity-consistent (HC) standard errors**
- Also called “robust” or “White” standard errors
- In R (multiple options):
  - `modelsummary(model, vcov = "robust")`
  - `modelsummary(model, vcov = "HC3")`
  - `estimatr::lm_robust(y ~ x, data = df)`
- Best practice: **always use robust SEs**

## Robust standard errors

- The fix: **heteroskedasticity-consistent (HC) standard errors**
- Also called “robust” or “White” standard errors
- In R (multiple options):
  - `modelsummary(model, vcov = "robust")`
  - `modelsummary(model, vcov = "HC3")`
  - `estimatr::lm_robust(y ~ x, data = df)`
- Best practice: **always use robust SEs**
  - If homoskedastic: robust SEs  $\approx$  classical SEs

## Robust standard errors

- The fix: **heteroskedasticity-consistent (HC) standard errors**
- Also called “robust” or “White” standard errors
- In R (multiple options):
  - `modelsummary(model, vcov = "robust")`
  - `modelsummary(model, vcov = "HC3")`
  - `estimatr::lm_robust(y ~ x, data = df)`
- Best practice: **always use robust SEs**
  - If homoskedastic: robust SEs  $\approx$  classical SEs
  - If heteroskedastic: robust SEs are correct, classical are not



## Log transformations: when and why

- Another common fix for heteroskedasticity and non-linearity:  
**transform the variables**

## Log transformations: when and why

- Another common fix for heteroskedasticity and non-linearity:  
**transform the variables**
- Many variables are right-skewed:

## Log transformations: when and why

- Another common fix for heteroskedasticity and non-linearity:  
**transform the variables**
- Many variables are right-skewed:
  - Income, GDP, population, firm size, casualties

## Log transformations: when and why

- Another common fix for heteroskedasticity and non-linearity:  
**transform the variables**
- Many variables are right-skewed:
  - Income, GDP, population, firm size, casualties
- Taking  $\log(X)$  compresses the right tail

## Log transformations: when and why

- Another common fix for heteroskedasticity and non-linearity:  
**transform the variables**
- Many variables are right-skewed:
  - Income, GDP, population, firm size, casualties
- Taking  $\log(X)$  compresses the right tail
- Makes the distribution more symmetric

## Log transformations: when and why

- Another common fix for heteroskedasticity and non-linearity:  
**transform the variables**
- Many variables are right-skewed:
  - Income, GDP, population, firm size, casualties
- Taking  $\log(X)$  compresses the right tail
- Makes the distribution more symmetric
- Can reduce heteroskedasticity

## Log transformations: when and why

- Another common fix for heteroskedasticity and non-linearity:  
**transform the variables**
- Many variables are right-skewed:
  - Income, GDP, population, firm size, casualties
- Taking  $\log(X)$  compresses the right tail
- Makes the distribution more symmetric
- Can reduce heteroskedasticity
- But interpretation changes!

## Interpreting logs

Model	Equation	Interpretation of $\beta_1$
Level-level	$Y = \beta_0 + \beta_1 X$	$\Delta X = 1 \Rightarrow \Delta Y = \beta_1$
Log-level	$\log(Y) = \beta_0 + \beta_1 X$	$\Delta X = 1 \Rightarrow \% \Delta Y \approx 100 \cdot \beta_1$
Level-log	$Y = \beta_0 + \beta_1 \log(X)$	$\% \Delta X = 1 \Rightarrow \Delta Y \approx \beta_1 / 100$
Log-log	$\log(Y) = \beta_0 + \beta_1 \log(X)$	$\% \Delta X = 1 \Rightarrow \% \Delta Y \approx \beta_1$

- Log-log:  $\beta_1$  is the **elasticity**



## Interpreting logs

Model	Equation	Interpretation of $\beta_1$
Level-level	$Y = \beta_0 + \beta_1 X$	$\Delta X = 1 \Rightarrow \Delta Y = \beta_1$
Log-level	$\log(Y) = \beta_0 + \beta_1 X$	$\Delta X = 1 \Rightarrow \% \Delta Y \approx 100 \cdot \beta_1$
Level-log	$Y = \beta_0 + \beta_1 \log(X)$	$\% \Delta X = 1 \Rightarrow \Delta Y \approx \beta_1 / 100$
Log-log	$\log(Y) = \beta_0 + \beta_1 \log(X)$	$\% \Delta X = 1 \Rightarrow \% \Delta Y \approx \beta_1$

- Log-log:  $\beta_1$  is the **elasticity**
- Log-level:  $\beta_1$  is a semi-elasticity

## Interpreting logs

Model	Equation	Interpretation of $\beta_1$
Level-level	$Y = \beta_0 + \beta_1 X$	$\Delta X = 1 \Rightarrow \Delta Y = \beta_1$
Log-level	$\log(Y) = \beta_0 + \beta_1 X$	$\Delta X = 1 \Rightarrow \% \Delta Y \approx 100 \cdot \beta_1$
Level-log	$Y = \beta_0 + \beta_1 \log(X)$	$\% \Delta X = 1 \Rightarrow \Delta Y \approx \beta_1 / 100$
Log-log	$\log(Y) = \beta_0 + \beta_1 \log(X)$	$\% \Delta X = 1 \Rightarrow \% \Delta Y \approx \beta_1$

- Log-log:  $\beta_1$  is the **elasticity**
- Log-level:  $\beta_1$  is a semi-elasticity
- Most common in practice: log-level and log-log

## Interpreting logs: a worked example

$$\log(\text{Income}) = 8.2 + 0.05 \cdot \text{Education}$$

- This is a **log-level** model

## Interpreting logs: a worked example

$$\log(\text{Income}) = 8.2 + 0.05 \cdot \text{Education}$$

- This is a **log-level** model
- One additional year of education  $\Rightarrow \approx 5\%$  higher income

## Interpreting logs: a worked example

$$\log(\text{Income}) = 8.2 + 0.05 \cdot \text{Education}$$

- This is a **log-level** model
- One additional year of education  $\Rightarrow \approx 5\%$  higher income
- What about 10 extra years?

## Interpreting logs: a worked example

$$\log(\text{Income}) = 8.2 + 0.05 \cdot \text{Education}$$

- This is a **log-level** model
- One additional year of education  $\Rightarrow \approx 5\%$  higher income
- What about 10 extra years?
  - $\rightarrow$  Approximation:  $10 \times 5\% = 50\%$

## Interpreting logs: a worked example

$$\log(\text{Income}) = 8.2 + 0.05 \cdot \text{Education}$$

- This is a **log-level** model
- One additional year of education  $\Rightarrow \approx 5\%$  higher income
- What about 10 extra years?
  - $\rightarrow$  Approximation:  $10 \times 5\% = 50\%$
  - $\rightarrow$  Exact:  $e^{0.05 \times 10} - 1 = e^{0.5} - 1 \approx 65\%$

## Interpreting logs: a worked example

$$\log(\text{Income}) = 8.2 + 0.05 \cdot \text{Education}$$

- This is a **log-level** model
- One additional year of education  $\Rightarrow \approx 5\%$  higher income
- What about 10 extra years?
  - $\rightarrow$  Approximation:  $10 \times 5\% = 50\%$
  - $\rightarrow$  Exact:  $e^{0.05 \times 10} - 1 = e^{0.5} - 1 \approx 65\%$
- The approximation breaks down for large  $\beta_1 \cdot \Delta X$




## Interpreting logs: a worked example

$$\log(\text{Income}) = 8.2 + 0.05 \cdot \text{Education}$$


- This is a **log-level** model
- One additional year of education  $\Rightarrow \approx 5\%$  higher income
- What about 10 extra years?
  - $\rightarrow$  Approximation:  $10 \times 5\% = 50\%$
  - $\rightarrow$  Exact:  $e^{0.05 \times 10} - 1 = e^{0.5} - 1 \approx 65\%$
- The approximation breaks down for large  $\beta_1 \cdot \Delta X$
- Rule of thumb: use approximation when  $\beta_1 \cdot \Delta X < 0.1$

## Residual plots: checking linearity



- A U-shape in residuals  $\Rightarrow$  missing non-linearity

## Residual plots: checking linearity



- A U-shape in residuals  $\Rightarrow$  missing non-linearity
- Fixes: add  $X^2$ , use  $\log(X)$ , or use a more flexible model

## Influential observations

- Some observations can disproportionately influence results

## Influential observations

- Some observations can disproportionately influence results
- **Cook's distance**: measures how much the model changes if you remove observation  $i$

## Influential observations

- Some observations can disproportionately influence results
- **Cook's distance**: measures how much the model changes if you remove observation  $i$ 
  - In R: `plot(model, which = 4)`

## Influential observations

- Some observations can disproportionately influence results
- **Cook's distance**: measures how much the model changes if you remove observation  $i$ 
  - In R: `plot(model, which = 4)`
  - Observations with Cook's  $D > 4/n$ : investigate further

## Influential observations

- Some observations can disproportionately influence results
- **Cook's distance**: measures how much the model changes if you remove observation  $i$ 
  - In R: `plot(model, which = 4)`
  - Observations with Cook's  $D > 4/n$ : investigate further
- What to do with influential observations?



## Influential observations

- Some observations can disproportionately influence results
- **Cook's distance**: measures how much the model changes if you remove observation  $i$ 
  - In R: `plot(model, which = 4)`
  - Observations with Cook's  $D > 4/n$ : investigate further
- What to do with influential observations?
  - Do NOT automatically delete them

## Influential observations

- Some observations can disproportionately influence results
- **Cook's distance**: measures how much the model changes if you remove observation  $i$ 
  - In R: `plot(model, which = 4)`
  - Observations with Cook's  $D > 4/n$ : investigate further
- What to do with influential observations?
  - Do NOT automatically delete them
  - Check if they are data errors

## Influential observations

- Some observations can disproportionately influence results
- **Cook's distance**: measures how much the model changes if you remove observation  $i$ 
  - In R: `plot(model, which = 4)`
  - Observations with Cook's  $D > 4/n$ : investigate further
- What to do with influential observations?
  - Do NOT automatically delete them
  - Check if they are data errors
  - Re-run the model without them as a robustness check

## Influential observations

- Some observations can disproportionately influence results
- **Cook's distance**: measures how much the model changes if you remove observation  $i$ 
  - In R: `plot(model, which = 4)`
  - Observations with Cook's  $D > 4/n$ : investigate further
- What to do with influential observations?
  - Do NOT automatically delete them
  - Check if they are data errors
  - Re-run the model without them as a robustness check
  - Report both results

# Roadmap

Beyond Coefficient Tables

Predicted Values and Marginal Effects

Presenting Results

Diagnostics

Wrap-up

## Summary: key takeaways

- Raw coefficients are rarely enough — compute **quantities of interest**

## Summary: key takeaways

- Raw coefficients are rarely enough — compute **quantities of interest**
- Predicted values, marginal effects, and first differences (all via `marginal effects`)

## Summary: key takeaways

- Raw coefficients are rarely enough — compute **quantities of interest**
- Predicted values, marginal effects, and first differences (all via `marginalEffects`)
- Present with plots (main text) and tables (appendix)



## Summary: key takeaways

- Raw coefficients are rarely enough — compute **quantities of interest**
- Predicted values, marginal effects, and first differences (all via `marginaleffects`)
- Present with plots (main text) and tables (appendix)
- Always use robust standard errors and check residual diagnostics

## Summary: key takeaways

- Raw coefficients are rarely enough — compute **quantities of interest**
- Predicted values, marginal effects, and first differences (all via `marginalEffects`)
- Present with plots (main text) and tables (appendix)
- Always use robust standard errors and check residual diagnostics
- Log-transform skewed variables (but interpret carefully)

## For next week

- Complete Assignment 4
- Next session: Best Practices in Computing
  - Reproducible workflows
  - Project organization
  - Writing clean R code

Questions?