

# Binary Outcomes

Francisco Villamil

Applied Quantitative Methods II  
MA in Social Sciences, Spring 2026

# Today's goals

- Understand why OLS is problematic for binary outcomes

# Today's goals

- Understand why OLS is problematic for binary outcomes
- Learn the linear probability model and its trade-offs

# Today's goals

- Understand why OLS is problematic for binary outcomes
- Learn the linear probability model and its trade-offs
- Understand logistic regression and how to estimate it in R

# Today's goals

- Understand why OLS is problematic for binary outcomes
- Learn the linear probability model and its trade-offs
- Understand logistic regression and how to estimate it in R
- Interpret logit results using marginal effects and predicted probabilities

# Today's goals

- Understand why OLS is problematic for binary outcomes
- Learn the linear probability model and its trade-offs
- Understand logistic regression and how to estimate it in R
- Interpret logit results using marginal effects and predicted probabilities
- Compare LPM and logit in practice

# Roadmap

# Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):



# Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):
  - Did someone vote?

# Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):
  - Did someone vote?
  - Did a war break out?

# Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):
  - Did someone vote?
  - Did a war break out?
  - Did a bill pass?

# Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):
  - Did someone vote?
  - Did a war break out?
  - Did a bill pass?
  - Did a country democratize?

# Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):
  - Did someone vote?
  - Did a war break out?
  - Did a bill pass?
  - Did a country democratize?
- Our outcome  $Y \in \{0, 1\}$

# Binary outcomes are everywhere

- Many outcomes in social science are binary (yes/no):
  - Did someone vote?
  - Did a war break out?
  - Did a bill pass?
  - Did a country democratize?
- Our outcome  $Y \in \{0, 1\}$
- We want to model:  $\Pr(Y = 1 \mid X)$

# What happens if we just use OLS?

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

- OLS gives us:  $E[Y | X] = \beta_0 + \beta_1 X$

# What happens if we just use OLS?

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

- OLS gives us:  $E[Y | X] = \beta_0 + \beta_1 X$
- Since  $Y \in \{0, 1\}$ :  $E[Y | X] = \Pr(Y = 1 | X)$



# What happens if we just use OLS?

$$Y = \beta_0 + \beta_1 X + \varepsilon$$


- OLS gives us:  $E[Y | X] = \beta_0 + \beta_1 X$
- Since  $Y \in \{0, 1\}$ :  $E[Y | X] = \Pr(Y = 1 | X)$
- So OLS is modeling a probability as a **linear function** of  $X$

# What happens if we just use OLS?

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

- OLS gives us:  $E[Y | X] = \beta_0 + \beta_1 X$
- Since  $Y \in \{0, 1\}$ :  $E[Y | X] = \Pr(Y = 1 | X)$
- So OLS is modeling a probability as a **linear function** of  $X$
- This is the **linear probability model** (LPM)

# The LPM in pictures



# The LPM: Simple and intuitive

- $\beta_1$  has a direct interpretation:

# The LPM: Simple and intuitive

- $\beta_1$  has a direct interpretation:
  - A one-unit increase in  $X$  changes the probability of  $Y = 1$  by  $\beta_1$

# The LPM: Simple and intuitive

- $\beta_1$  has a direct interpretation:
  - A one-unit increase in  $X$  changes the probability of  $Y = 1$  by  $\beta_1$
- Easy to estimate: just `lm(y ~ x, data = df)`

# The LPM: Simple and intuitive

- $\beta_1$  has a direct interpretation:
  - A one-unit increase in  $X$  changes the probability of  $Y = 1$  by  $\beta_1$
- Easy to estimate: just `lm(y ~ x, data = df)`
- Easy to interpret: same as OLS

# The LPM: Simple and intuitive

- $\beta_1$  has a direct interpretation:
  - A one-unit increase in  $X$  changes the probability of  $Y = 1$  by  $\beta_1$
- Easy to estimate: just `lm(y ~ x, data = df)`
- Easy to interpret: same as OLS
- Many applied researchers use the LPM in practice



# The LPM: Simple and intuitive

- $\beta_1$  has a direct interpretation:
  - A one-unit increase in  $X$  changes the probability of  $Y = 1$  by  $\beta_1$
- Easy to estimate: just `lm(y ~ x, data = df)`
- Easy to interpret: same as OLS
- Many applied researchers use the LPM in practice
- Especially common in economics and political science

# LPM limitations

- **Problem 1:** Predictions outside  $[0, 1]$

# LPM limitations

- **Problem 1:** Predictions outside  $[0, 1]$ 
  - A linear function can produce  $\hat{P} < 0$  or  $\hat{P} > 1$

# LPM limitations

- **Problem 1:** Predictions outside  $[0, 1]$ 
  - A linear function can produce  $\hat{P} < 0$  or  $\hat{P} > 1$
  - Probabilities must be between 0 and 1

# LPM limitations

- **Problem 1:** Predictions outside  $[0, 1]$ 
  - A linear function can produce  $\hat{P} < 0$  or  $\hat{P} > 1$
  - Probabilities must be between 0 and 1
- **Problem 2:** Heteroskedasticity by construction

# LPM limitations

- **Problem 1:** Predictions outside  $[0, 1]$ 
  - A linear function can produce  $\hat{P} < 0$  or  $\hat{P} > 1$
  - Probabilities must be between 0 and 1
- **Problem 2:** Heteroskedasticity by construction
  - $\text{Var}(\varepsilon | X) = P(1 - P)$ , which varies with  $X$

# LPM limitations

- **Problem 1:** Predictions outside  $[0, 1]$ 
  - A linear function can produce  $\hat{P} < 0$  or  $\hat{P} > 1$
  - Probabilities must be between 0 and 1
- **Problem 2:** Heteroskedasticity by construction
  - $\text{Var}(\varepsilon | X) = P(1 - P)$ , which varies with  $X$
  - Always use robust SEs with the LPM

# LPM limitations

- **Problem 1:** Predictions outside  $[0, 1]$ 
  - A linear function can produce  $\hat{P} < 0$  or  $\hat{P} > 1$
  - Probabilities must be between 0 and 1
- **Problem 2:** Heteroskedasticity by construction
  - $\text{Var}(\varepsilon | X) = P(1 - P)$ , which varies with  $X$
  - Always use robust SEs with the LPM
- **Problem 3:** Non-linearity at the extremes



# LPM limitations

- **Problem 1:** Predictions outside  $[0, 1]$ 
  - A linear function can produce  $\hat{P} < 0$  or  $\hat{P} > 1$
  - Probabilities must be between 0 and 1
- **Problem 2:** Heteroskedasticity by construction
  - $\text{Var}(\varepsilon | X) = P(1 - P)$ , which varies with  $X$
  - Always use robust SEs with the LPM
- **Problem 3:** Non-linearity at the extremes
  - True relationship between  $X$  and  $\Pr(Y = 1)$  is S-shaped

# LPM limitations

- **Problem 1:** Predictions outside  $[0, 1]$ 
  - A linear function can produce  $\hat{P} < 0$  or  $\hat{P} > 1$
  - Probabilities must be between 0 and 1
- **Problem 2:** Heteroskedasticity by construction
  - $\text{Var}(\varepsilon | X) = P(1 - P)$ , which varies with  $X$
  - Always use robust SEs with the LPM
- **Problem 3:** Non-linearity at the extremes
  - True relationship between  $X$  and  $\Pr(Y = 1)$  is S-shaped
  - LPM forces it to be linear

## When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)

# When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
  - The linear approximation is reasonable here

# When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
  - The linear approximation is reasonable here
- When you care about **average marginal effects**

# When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
  - The linear approximation is reasonable here
- When you care about **average marginal effects**
  - LPM and logit often give similar AMEs

# When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
  - The linear approximation is reasonable here
- When you care about **average marginal effects**
  - LPM and logit often give similar AMEs
- When simplicity of interpretation matters

# When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
  - The linear approximation is reasonable here
- When you care about **average marginal effects**
  - LPM and logit often give similar AMEs
- When simplicity of interpretation matters
- When is it **not** good enough?



# When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
  - The linear approximation is reasonable here
- When you care about **average marginal effects**
  - LPM and logit often give similar AMEs
- When simplicity of interpretation matters
- When is it **not** good enough?
  - Rare events (many observations near 0)

# When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
  - The linear approximation is reasonable here
- When you care about **average marginal effects**
  - LPM and logit often give similar AMEs
- When simplicity of interpretation matters
- When is it **not** good enough?
  - Rare events (many observations near 0)
  - You need predicted probabilities to be bounded

# When is the LPM “good enough”?

- When probabilities are in the middle range (0.2–0.8)
  - The linear approximation is reasonable here
- When you care about **average marginal effects**
  - LPM and logit often give similar AMEs
- When simplicity of interpretation matters
- When is it **not** good enough?
  - Rare events (many observations near 0)
  - You need predicted probabilities to be bounded
  - The relationship is clearly non-linear

You estimate an LPM predicting civil war onset. You find that 8% of predicted probabilities are negative.

Is this a problem? What would you do?

# Roadmap

# The logistic function

$$\Pr(Y = 1 \mid X) = \frac{e^{\beta_0 + \beta_1 X}}{1 + e^{\beta_0 + \beta_1 X}} = \frac{1}{1 + e^{-(\beta_0 + \beta_1 X)}}$$

- This is an S-shaped (sigmoid) curve

# The logistic function

$$\Pr(Y = 1 \mid X) = \frac{e^{\beta_0 + \beta_1 X}}{1 + e^{\beta_0 + \beta_1 X}} = \frac{1}{1 + e^{-(\beta_0 + \beta_1 X)}}$$

- This is an S-shaped (sigmoid) curve
- Output is always between 0 and 1

# The logistic function

$$\Pr(Y = 1 \mid X) = \frac{e^{\beta_0 + \beta_1 X}}{1 + e^{\beta_0 + \beta_1 X}} = \frac{1}{1 + e^{-(\beta_0 + \beta_1 X)}}$$

- This is an S-shaped (sigmoid) curve
- Output is always between 0 and 1
- Steep in the middle, flat at the extremes




# The logistic function

$$\Pr(Y = 1 \mid X) = \frac{e^{\beta_0 + \beta_1 X}}{1 + e^{\beta_0 + \beta_1 X}} = \frac{1}{1 + e^{-(\beta_0 + \beta_1 X)}}$$

- This is an S-shaped (sigmoid) curve
- Output is always between 0 and 1
- Steep in the middle, flat at the extremes
- A natural model for probabilities

# The sigmoid curve



# The logit transformation

We can rearrange the logistic function:

$$\log\left(\frac{P}{1-P}\right) = \beta_0 + \beta_1 X$$

- The left side is the **log-odds** (or “logit”)

# The logit transformation

We can rearrange the logistic function:

$$\log\left(\frac{P}{1-P}\right) = \beta_0 + \beta_1 X$$

- The left side is the **log-odds** (or “logit”)
- $\frac{P}{1-P}$  is the **odds** of the event

# The logit transformation

We can rearrange the logistic function:

$$\log\left(\frac{P}{1-P}\right) = \beta_0 + \beta_1 X$$

- The left side is the **log-odds** (or “logit”)
- $\frac{P}{1-P}$  is the **odds** of the event
- The model is linear in the log-odds, not in the probability

# The logit transformation

We can rearrange the logistic function:

$$\log\left(\frac{P}{1-P}\right) = \beta_0 + \beta_1 X$$

- The left side is the **log-odds** (or “logit”)
- $\frac{P}{1-P}$  is the **odds** of the event
- The model is linear in the log-odds, not in the probability
- This is why we call it “logistic regression” or “logit”

# Maximum likelihood estimation

- We can't use OLS for logistic regression

# Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)



# Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)
- The intuition:

# Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)
- The intuition:
  - For each observation, the model predicts  $\Pr(Y_i = 1)$

# Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)
- The intuition:
  - For each observation, the model predicts  $\Pr(Y_i = 1)$
  - MLE finds the coefficients that make the observed data most likely

# Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)
- The intuition:
  - For each observation, the model predicts  $\Pr(Y_i = 1)$
  - MLE finds the coefficients that make the observed data most likely
- No need to derive this — R does it for us

# Maximum likelihood estimation

- We can't use OLS for logistic regression
- Instead, we use **maximum likelihood estimation** (MLE)
- The intuition:
  - For each observation, the model predicts  $\Pr(Y_i = 1)$
  - MLE finds the coefficients that make the observed data most likely
- No need to derive this — R does it for us
- The math is different, but the workflow is the same

# Estimating logit in R

- The function: `glm(y ~ x, family = binomial, data = df)`

# Estimating logit in R

- The function: `glm(y ~ x, family = binomial, data = df)`
- `glm`: generalized linear model

# Estimating logit in R

- The function: `glm(y ~ x, family = binomial, data = df)`
- `glm`: generalized linear model
- `family = binomial`: tells R to use logistic regression



# Estimating logit in R

- The function: `glm(y ~ x, family = binomial, data = df)`
- `glm`: generalized linear model
- `family = binomial`: tells R to use logistic regression
- The syntax is identical to `lm()`, just swap to `glm()`

# Estimating logit in R

- The function: `glm(y ~ x, family = binomial, data = df)`
- `glm`: generalized linear model
- `family = binomial`: tells R to use logistic regression
- The syntax is identical to `lm()`, just swap to `glm()`
- Works with `broom::tidy()`, `modelsummary()`, etc.

## Interpreting logit output: Log-odds

- The direct output gives coefficients in **log-odds**

## Interpreting logit output: Log-odds

- The direct output gives coefficients in **log-odds**
- $\beta_1 = 0.5$  means:

# Interpreting logit output: Log-odds

- The direct output gives coefficients in **log-odds**
- $\beta_1 = 0.5$  means:
  - A one-unit increase in  $X$  increases the log-odds by 0.5

# Interpreting logit output: Log-odds

- The direct output gives coefficients in **log-odds**
- $\beta_1 = 0.5$  means:
  - A one-unit increase in  $X$  increases the log-odds by 0.5
- This is hard to interpret!

# Interpreting logit output: Log-odds

- The direct output gives coefficients in **log-odds**
- $\beta_1 = 0.5$  means:
  - A one-unit increase in  $X$  increases the log-odds by 0.5
- This is hard to interpret!
- Nobody thinks in log-odds

## Interpreting logit output: Odds ratios

- Exponentiate the coefficient:  $e^{\beta_1} = \text{odds ratio}$



## Interpreting logit output: Odds ratios

- Exponentiate the coefficient:  $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`

## Interpreting logit output: Odds ratios

- Exponentiate the coefficient:  $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`
- $e^{0.5} \approx 1.65$  means:

# Interpreting logit output: Odds ratios

- Exponentiate the coefficient:  $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`
- $e^{0.5} \approx 1.65$  means:
  - A one-unit increase in  $X$  **multiplies** the odds by 1.65

## Interpreting logit output: Odds ratios

- Exponentiate the coefficient:  $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`
- $e^{0.5} \approx 1.65$  means:
  - A one-unit increase in  $X$  **multiplies** the odds by 1.65
  - Or: the odds increase by 65%

# Interpreting logit output: Odds ratios

- Exponentiate the coefficient:  $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`
- $e^{0.5} \approx 1.65$  means:
  - A one-unit increase in  $X$  **multiplies** the odds by 1.65
  - Or: the odds increase by 65%
- Slightly more intuitive, but still not probabilities

# Interpreting logit output: Odds ratios

- Exponentiate the coefficient:  $e^{\beta_1} = \text{odds ratio}$
- In R: `exp(coef(model))`
- $e^{0.5} \approx 1.65$  means:
  - A one-unit increase in  $X$  **multiplies** the odds by 1.65
  - Or: the odds increase by 65%
- Slightly more intuitive, but still not probabilities
- The change in probability depends on where you start

## Worked example: From log-odds to probability

$$\text{Logit model: } \log\left(\frac{P}{1-P}\right) = -2 + 0.5 \cdot \text{Education}$$

- For someone with Education = 4:

## Worked example: From log-odds to probability

$$\text{Logit model: } \log\left(\frac{P}{1-P}\right) = -2 + 0.5 \cdot \text{Education}$$

- For someone with Education = 4:  
→ Log-odds =  $-2 + 0.5 \times 4 = 0$



## Worked example: From log-odds to probability

$$\text{Logit model: } \log\left(\frac{P}{1-P}\right) = -2 + 0.5 \cdot \text{Education}$$

- For someone with Education = 4:
  - Log-odds =  $-2 + 0.5 \times 4 = 0$
  - Odds =  $e^0 = 1$  (50-50 chance)

## Worked example: From log-odds to probability

$$\text{Logit model: } \log\left(\frac{P}{1-P}\right) = -2 + 0.5 \cdot \text{Education}$$

- For someone with Education = 4:
  - Log-odds =  $-2 + 0.5 \times 4 = 0$
  - Odds =  $e^0 = 1$  (50-50 chance)
  - Probability =  $\frac{1}{1+e^0} = 0.50$

## Worked example: From log-odds to probability

$$\text{Logit model: } \log\left(\frac{P}{1-P}\right) = -2 + 0.5 \cdot \text{Education}$$

- For someone with Education = 4:
  - Log-odds =  $-2 + 0.5 \times 4 = 0$
  - Odds =  $e^0 = 1$  (50-50 chance)
  - Probability =  $\frac{1}{1+e^0} = 0.50$
- For someone with Education = 8:

## Worked example: From log-odds to probability

$$\text{Logit model: } \log\left(\frac{P}{1-P}\right) = -2 + 0.5 \cdot \text{Education}$$

- For someone with Education = 4:
  - Log-odds =  $-2 + 0.5 \times 4 = 0$
  - Odds =  $e^0 = 1$  (50-50 chance)
  - Probability =  $\frac{1}{1+e^0} = 0.50$
- For someone with Education = 8:
  - Log-odds =  $-2 + 0.5 \times 8 = 2$

## Worked example: From log-odds to probability

$$\text{Logit model: } \log\left(\frac{P}{1-P}\right) = -2 + 0.5 \cdot \text{Education}$$

- For someone with Education = 4:
  - Log-odds =  $-2 + 0.5 \times 4 = 0$
  - Odds =  $e^0 = 1$  (50-50 chance)
  - Probability =  $\frac{1}{1+e^0} = 0.50$
- For someone with Education = 8:
  - Log-odds =  $-2 + 0.5 \times 8 = 2$
  - Odds =  $e^2 \approx 7.4$

## Worked example: From log-odds to probability

$$\text{Logit model: } \log\left(\frac{P}{1-P}\right) = -2 + 0.5 \cdot \text{Education}$$

- For someone with Education = 4:
  - Log-odds =  $-2 + 0.5 \times 4 = 0$
  - Odds =  $e^0 = 1$  (50-50 chance)
  - Probability =  $\frac{1}{1+e^0} = 0.50$
- For someone with Education = 8:
  - Log-odds =  $-2 + 0.5 \times 8 = 2$
  - Odds =  $e^2 \approx 7.4$
  - Probability =  $\frac{1}{1+e^{-2}} \approx 0.88$

## Worked example: From log-odds to probability

$$\text{Logit model: } \log\left(\frac{P}{1-P}\right) = -2 + 0.5 \cdot \text{Education}$$

- For someone with Education = 4:
  - Log-odds =  $-2 + 0.5 \times 4 = 0$
  - Odds =  $e^0 = 1$  (50-50 chance)
  - Probability =  $\frac{1}{1+e^0} = 0.50$
- For someone with Education = 8:
  - Log-odds =  $-2 + 0.5 \times 8 = 2$
  - Odds =  $e^2 \approx 7.4$
  - Probability =  $\frac{1}{1+e^{-2}} \approx 0.88$
- Going from 4 to 8 years:  $P$  goes from 0.50 to 0.88

## Worked example: From log-odds to probability

$$\text{Logit model: } \log\left(\frac{P}{1-P}\right) = -2 + 0.5 \cdot \text{Education}$$

- For someone with Education = 4:
  - Log-odds =  $-2 + 0.5 \times 4 = 0$
  - Odds =  $e^0 = 1$  (50-50 chance)
  - Probability =  $\frac{1}{1+e^0} = 0.50$
- For someone with Education = 8:
  - Log-odds =  $-2 + 0.5 \times 8 = 2$
  - Odds =  $e^2 \approx 7.4$
  - Probability =  $\frac{1}{1+e^{-2}} \approx 0.88$
- Going from 4 to 8 years:  $P$  goes from 0.50 to 0.88
- The same 4-unit change gives  $\Delta P = 0.38$



## Why coefficients alone are not enough

- In OLS:  $\beta_1$  = change in  $Y$  for one-unit change in  $X$  (always)

## Why coefficients alone are not enough

- In OLS:  $\beta_1$  = change in  $Y$  for one-unit change in  $X$  (always)
- In logit: the change in **probability** depends on:

# Why coefficients alone are not enough

- In OLS:  $\beta_1$  = change in  $Y$  for one-unit change in  $X$  (always)
- In logit: the change in **probability** depends on:
  - The current value of  $X$

# Why coefficients alone are not enough

- In OLS:  $\beta_1$  = change in  $Y$  for one-unit change in  $X$  (always)
- In logit: the change in **probability** depends on:
  - The current value of  $X$
  - The values of all other variables

# Why coefficients alone are not enough

- In OLS:  $\beta_1$  = change in  $Y$  for one-unit change in  $X$  (always)
- In logit: the change in **probability** depends on:
  - The current value of  $X$
  - The values of all other variables
- A coefficient of  $\beta_1 = 0.5$  could mean:

# Why coefficients alone are not enough

- In OLS:  $\beta_1$  = change in  $Y$  for one-unit change in  $X$  (always)
- In logit: the change in **probability** depends on:
  - The current value of  $X$
  - The values of all other variables
- A coefficient of  $\beta_1 = 0.5$  could mean:
  - Going from  $P = 0.01$  to  $P = 0.016$  (tiny change)

# Why coefficients alone are not enough

- In OLS:  $\beta_1$  = change in  $Y$  for one-unit change in  $X$  (always)
- In logit: the change in **probability** depends on:
  - The current value of  $X$
  - The values of all other variables
- A coefficient of  $\beta_1 = 0.5$  could mean:
  - Going from  $P = 0.01$  to  $P = 0.016$  (tiny change)
  - Going from  $P = 0.50$  to  $P = 0.62$  (large change)

# Why coefficients alone are not enough

- In OLS:  $\beta_1$  = change in  $Y$  for one-unit change in  $X$  (always)
- In logit: the change in **probability** depends on:
  - The current value of  $X$
  - The values of all other variables
- A coefficient of  $\beta_1 = 0.5$  could mean:
  - Going from  $P = 0.01$  to  $P = 0.016$  (tiny change)
  - Going from  $P = 0.50$  to  $P = 0.62$  (large change)
- We need better tools to interpret logit models



# Roadmap

A logit model estimates  $\hat{\beta}_1 = 0.8$  for education.

Your colleague says: “Education increases the probability of voting by 0.8.”

What’s wrong with this statement?

# Predicted probabilities

- The most intuitive way to interpret logit models

# Predicted probabilities

- The most intuitive way to interpret logit models
- “What is the predicted probability of  $Y = 1$  for a person with these characteristics?”

# Predicted probabilities

- The most intuitive way to interpret logit models
- “What is the predicted probability of  $Y = 1$  for a person with these characteristics?”
- In R:

# Predicted probabilities

- The most intuitive way to interpret logit models
- “What is the predicted probability of  $Y = 1$  for a person with these characteristics?”
- In R:
  - `marginalEffects::predictions(model)`

# Predicted probabilities

- The most intuitive way to interpret logit models
- “What is the predicted probability of  $Y = 1$  for a person with these characteristics?”
- In R:
  - `marginalEffects::predictions(model)`
  - Returns predicted probabilities for each observation

# Predicted probabilities

- The most intuitive way to interpret logit models
- “What is the predicted probability of  $Y = 1$  for a person with these characteristics?”
- In R:
  - `marginalEffects::predictions(model)`
  - Returns predicted probabilities for each observation
  - Or at specific values: `predictions(model, newdata = datagrid(...))`



## Average marginal effects (AME)

- The marginal effect of  $X$  on  $\Pr(Y = 1)$  varies across observations

## Average marginal effects (AME)

- The marginal effect of  $X$  on  $\Pr(Y = 1)$  varies across observations
- The **AME** averages across all observations in the sample

## Average marginal effects (AME)

- The marginal effect of  $X$  on  $\Pr(Y = 1)$  varies across observations
- The **AME** averages across all observations in the sample
- In R: `marginaleffects::avg_slopes(model)`

## Average marginal effects (AME)

- The marginal effect of  $X$  on  $\Pr(Y = 1)$  varies across observations
- The **AME** averages across all observations in the sample
- In R: `marginaleffects::avg_slopes(model)`
- Interpretation (just like OLS):


# Average marginal effects (AME)

- The marginal effect of  $X$  on  $\Pr(Y = 1)$  varies across observations
- The **AME** averages across all observations in the sample
- In R: `marginalEffects::avg_slopes(model)`
- Interpretation (just like OLS):
  - “On average, a one-unit increase in  $X$  changes  $\Pr(Y = 1)$  by  $\Delta P$ ”

# Average marginal effects (AME)

- The marginal effect of  $X$  on  $\Pr(Y = 1)$  varies across observations
- The **AME** averages across all observations in the sample
- In R: `marginaleffects::avg_slopes(model)`
- Interpretation (just like OLS):
  - “On average, a one-unit increase in  $X$  changes  $\Pr(Y = 1)$  by  $\Delta P$ ”
- AMEs from logit are often similar to LPM coefficients

# Why marginal effects vary



## Marginal effects at representative values

- Instead of averaging, evaluate at specific values



# Marginal effects at representative values

- Instead of averaging, evaluate at specific values
- Example: “What is the effect of education on voting for a 40-year-old woman?”

# Marginal effects at representative values

- Instead of averaging, evaluate at specific values
- Example: “What is the effect of education on voting for a 40-year-old woman?”
- In R:

# Marginal effects at representative values

- Instead of averaging, evaluate at specific values
- Example: “What is the effect of education on voting for a 40-year-old woman?”
- In R:
  - `avg_slopes(model, newdata = datagrid(age = 40, female = 1))`

# Marginal effects at representative values

- Instead of averaging, evaluate at specific values
- Example: “What is the effect of education on voting for a 40-year-old woman?”
- In R:
  - `avg_slopes(model, newdata = datagrid(age = 40, female = 1))`
- Useful when the effect varies substantially across groups

# Marginal effects at representative values

- Instead of averaging, evaluate at specific values
- Example: “What is the effect of education on voting for a 40-year-old woman?”
- In R:
  - `avg_slopes(model, newdata = datagrid(age = 40, female = 1))`
- Useful when the effect varies substantially across groups
- E.g., the effect may be larger for young people than for old people

# Plotting predicted probabilities

- The best way to communicate logit results to any audience

## Plotting predicted probabilities

- The best way to communicate logit results to any audience
- Show how  $\Pr(Y = 1)$  changes across values of  $X$

# Plotting predicted probabilities

- The best way to communicate logit results to any audience
- Show how  $\Pr(Y = 1)$  changes across values of  $X$
- Include confidence bands



# Plotting predicted probabilities

- The best way to communicate logit results to any audience
- Show how  $\Pr(Y = 1)$  changes across values of  $X$
- Include confidence bands
- In R:

# Plotting predicted probabilities

- The best way to communicate logit results to any audience
- Show how  $\Pr(Y = 1)$  changes across values of  $X$
- Include confidence bands
- In R:
  - `plot_predictions(model, condition = "x")`

# Plotting predicted probabilities

- The best way to communicate logit results to any audience
- Show how  $\Pr(Y = 1)$  changes across values of  $X$
- Include confidence bands
- In R:
  - `plot_predictions(model, condition = "x")`
  - Plots the S-curve with 95% confidence interval

# Plotting predicted probabilities

- The best way to communicate logit results to any audience
- Show how  $\Pr(Y = 1)$  changes across values of  $X$
- Include confidence bands
- In R:
  - `plot_predictions(model, condition = "x")`
  - Plots the S-curve with 95% confidence interval
- Much more informative than a table of log-odds

# Comparing LPM and logit

- In many cases, LPM and logit give similar results

# Comparing LPM and logit

- In many cases, LPM and logit give similar results
  - Especially for average marginal effects

# Comparing LPM and logit

- In many cases, LPM and logit give similar results
  - Especially for average marginal effects
  - Especially when probabilities are in the 0.2–0.8 range

# Comparing LPM and logit

- In many cases, LPM and logit give similar results
  - Especially for average marginal effects
  - Especially when probabilities are in the 0.2–0.8 range
- Where they differ:



# Comparing LPM and logit

- In many cases, LPM and logit give similar results
  - Especially for average marginal effects
  - Especially when probabilities are in the 0.2–0.8 range
- Where they differ:
  - Predicted probabilities near 0 or 1

# Comparing LPM and logit

- In many cases, LPM and logit give similar results
  - Especially for average marginal effects
  - Especially when probabilities are in the 0.2–0.8 range
- Where they differ:
  - Predicted probabilities near 0 or 1
  - LPM can go outside  $[0, 1]$ ; logit cannot

# Comparing LPM and logit

- In many cases, LPM and logit give similar results
  - Especially for average marginal effects
  - Especially when probabilities are in the 0.2–0.8 range
- Where they differ:
  - Predicted probabilities near 0 or 1
  - LPM can go outside  $[0, 1]$ ; logit cannot
  - Marginal effects at extreme values

# Comparing LPM and logit

- In many cases, LPM and logit give similar results
  - Especially for average marginal effects
  - Especially when probabilities are in the 0.2–0.8 range
- Where they differ:
  - Predicted probabilities near 0 or 1
  - LPM can go outside  $[0, 1]$ ; logit cannot
  - Marginal effects at extreme values
- A good practice: estimate both and compare

## Model fit for logit

- No  $R^2$  in the usual sense (MLE, not OLS)

# Model fit for logit

- No  $R^2$  in the usual sense (MLE, not OLS)
- Alternative measures:

# Model fit for logit

- No  $R^2$  in the usual sense (MLE, not OLS)
- Alternative measures:
  - **Pseudo- $R^2$** : compares model to null model (McFadden's)

# Model fit for logit

- No  $R^2$  in the usual sense (MLE, not OLS)
- Alternative measures:
  - **Pseudo- $R^2$** : compares model to null model (McFadden's)
  - **AIC / BIC**: penalized likelihood (lower = better)



# Model fit for logit

- No  $R^2$  in the usual sense (MLE, not OLS)
- Alternative measures:
  - **Pseudo- $R^2$** : compares model to null model (McFadden's)
  - **AIC / BIC**: penalized likelihood (lower = better)
  - **Classification accuracy**: percent correctly predicted

# Model fit for logit

- No  $R^2$  in the usual sense (MLE, not OLS)
- Alternative measures:
  - **Pseudo- $R^2$** : compares model to null model (McFadden's)
  - **AIC / BIC**: penalized likelihood (lower = better)
  - **Classification accuracy**: percent correctly predicted
  - **ROC / AUC**: trade-off between true and false positives

# Model fit for logit

- No  $R^2$  in the usual sense (MLE, not OLS)
- Alternative measures:
  - **Pseudo- $R^2$** : compares model to null model (McFadden's)
  - **AIC / BIC**: penalized likelihood (lower = better)
  - **Classification accuracy**: percent correctly predicted
  - **ROC / AUC**: trade-off between true and false positives
- None is perfect; use as rough guides

# Model fit for logit

- No  $R^2$  in the usual sense (MLE, not OLS)
- Alternative measures:
  - **Pseudo- $R^2$** : compares model to null model (McFadden's)
  - **AIC / BIC**: penalized likelihood (lower = better)
  - **Classification accuracy**: percent correctly predicted
  - **ROC / AUC**: trade-off between true and false positives
- None is perfect; use as rough guides
- Reported automatically by `modelsummary()` and `performance::r2()`

# A note on probit

- Logit uses the logistic CDF as the link function

## A note on probit

- Logit uses the logistic CDF as the link function
- **Probit** uses the normal (Gaussian) CDF instead

## A note on probit

- Logit uses the logistic CDF as the link function
- **Probit** uses the normal (Gaussian) CDF instead
- In R: `glm(y ~ x, family = binomial(link = "probit"), data = df)`

## A note on probit

- Logit uses the logistic CDF as the link function
- **Probit** uses the normal (Gaussian) CDF instead
- In R: `glm(y ~ x, family = binomial(link = "probit"), data = df)`
- In practice, logit and probit give very similar results



# A note on probit

- Logit uses the logistic CDF as the link function
- **Probit** uses the normal (Gaussian) CDF instead
- In R: `glm(y ~ x, family = binomial(link = "probit"), data = df)`
- In practice, logit and probit give very similar results
  - Probit coefficients  $\approx$  logit coefficients  $\times 0.625$

# A note on probit

- Logit uses the logistic CDF as the link function
- **Probit** uses the normal (Gaussian) CDF instead
- In R: `glm(y ~ x, family = binomial(link = "probit"), data = df)`
- In practice, logit and probit give very similar results
  - Probit coefficients  $\approx$  logit coefficients  $\times 0.625$
  - Predicted probabilities are nearly identical

# A note on probit

- Logit uses the logistic CDF as the link function
- **Probit** uses the normal (Gaussian) CDF instead
- In R: `glm(y ~ x, family = binomial(link = "probit"), data = df)`
- In practice, logit and probit give very similar results
  - Probit coefficients  $\approx$  logit coefficients  $\times 0.625$
  - Predicted probabilities are nearly identical
- Logit is more common in political science; probit in economics

# A note on probit

- Logit uses the logistic CDF as the link function
- **Probit** uses the normal (Gaussian) CDF instead
- In R: `glm(y ~ x, family = binomial(link = "probit"), data = df)`
- In practice, logit and probit give very similar results
  - Probit coefficients  $\approx$  logit coefficients  $\times 0.625$
  - Predicted probabilities are nearly identical
- Logit is more common in political science; probit in economics
- Use whichever is conventional in your field

# Roadmap

# Complete workflow in R

```
lpm <- lm(vote ~ age + income + educ, data = df)
logit <- glm(vote ~ age + income + educ,
             family = binomial, data = df)
```

## Complete workflow in R

```
lpm <- lm(vote ~ age + income + educ, data = df)
logit <- glm(vote ~ age + income + educ,
             family = binomial, data = df)
modelsummary(list("LPM" = lpm, "Logit" = logit),
             vcov = list("robust", NULL))
```

## Complete workflow in R

```
lpm <- lm(vote ~ age + income + educ, data = df)
logit <- glm(vote ~ age + income + educ,
             family = binomial, data = df)
modelsummary(list("LPM" = lpm, "Logit" = logit),
             vcov = list("robust", NULL))
avg_slopes(logit)                                # compare AMEs to LPM
```



## Complete workflow in R

```
lpm <- lm(vote ~ age + income + educ, data = df)
logit <- glm(vote ~ age + income + educ,
             family = binomial, data = df)
modelsummary(list("LPM" = lpm, "Logit" = logit),
             vcov = list("robust", NULL))
avg_slopes(logit)                                # compare AMEs to LPM
plot_predictions(logit, condition = "income")
```

## Decision tree: When to use which?

- **Use LPM when:**

# Decision tree: When to use which?

- **Use LPM when:**
  - You want simple, direct interpretation

# Decision tree: When to use which?

- **Use LPM when:**
  - You want simple, direct interpretation
  - Probabilities are in the middle range

# Decision tree: When to use which?

- **Use LPM when:**

- You want simple, direct interpretation
- Probabilities are in the middle range
- You mainly care about average effects

# Decision tree: When to use which?

- **Use LPM when:**

- You want simple, direct interpretation
- Probabilities are in the middle range
- You mainly care about average effects

- **Use logit when:**

# Decision tree: When to use which?

- **Use LPM when:**

- You want simple, direct interpretation
- Probabilities are in the middle range
- You mainly care about average effects

- **Use logit when:**

- You need bounded predicted probabilities

# Decision tree: When to use which?

- **Use LPM when:**

- You want simple, direct interpretation
- Probabilities are in the middle range
- You mainly care about average effects

- **Use logit when:**

- You need bounded predicted probabilities
- Many observations have extreme probabilities (near 0 or 1)



# Decision tree: When to use which?

- **Use LPM when:**

- You want simple, direct interpretation
- Probabilities are in the middle range
- You mainly care about average effects

- **Use logit when:**

- You need bounded predicted probabilities
- Many observations have extreme probabilities (near 0 or 1)
- You want to properly account for the binary nature of  $Y$

# Decision tree: When to use which?

- **Use LPM when:**

- You want simple, direct interpretation
- Probabilities are in the middle range
- You mainly care about average effects

- **Use logit when:**

- You need bounded predicted probabilities
- Many observations have extreme probabilities (near 0 or 1)
- You want to properly account for the binary nature of  $Y$

- In practice: estimate both, report the one most appropriate

# Decision tree: When to use which?

- **Use LPM when:**
  - You want simple, direct interpretation
  - Probabilities are in the middle range
  - You mainly care about average effects
- **Use logit when:**
  - You need bounded predicted probabilities
  - Many observations have extreme probabilities (near 0 or 1)
  - You want to properly account for the binary nature of  $Y$
- In practice: estimate both, report the one most appropriate
- Always report marginal effects, not just log-odds

## Summary: Key takeaways

- Binary outcomes require special treatment

## Summary: Key takeaways

- Binary outcomes require special treatment
- The LPM is simple but has known limitations

## Summary: Key takeaways

- Binary outcomes require special treatment
- The LPM is simple but has known limitations
- Logistic regression bounds probabilities between 0 and 1

## Summary: Key takeaways

- Binary outcomes require special treatment
- The LPM is simple but has known limitations
- Logistic regression bounds probabilities between 0 and 1
- Log-odds are not intuitive — use marginal effects and predicted probabilities

## Summary: Key takeaways

- Binary outcomes require special treatment
- The LPM is simple but has known limitations
- Logistic regression bounds probabilities between 0 and 1
- Log-odds are not intuitive — use marginal effects and predicted probabilities
- AMEs from logit are often similar to LPM coefficients



## Summary: Key takeaways

- Binary outcomes require special treatment
- The LPM is simple but has known limitations
- Logistic regression bounds probabilities between 0 and 1
- Log-odds are not intuitive — use marginal effects and predicted probabilities
- AMEs from logit are often similar to LPM coefficients
- Always estimate both and compare; always plot predicted probabilities

# For next week

- Read Gelman et al., chapters 11–12
- Read Arel-Bundock, Greifer, and Heiss (2025), chapters 1–4
- Complete Assignment 3
  
- Next session: Model interpretation and diagnostics
  - Beyond coefficient tables
  - Visualizing model results
  - Residual diagnostics
  - Influence and outliers

Questions?